

FOREST WILDLIFE POPULATIONS

Forest Wildlife Populations and Research Group
1201 East Highway 2
Grand Rapids, MN 55744
(218) 327-4432



2014 MINNESOTA SPRING GROUSE SURVEYS

Charlotte Roy, Forest Wildlife Populations and Research Group

SUMMARY OF FINDINGS

Each spring, the Minnesota DNR coordinates statewide ruffed grouse (*Bonasa umbellus*) and sharp-tailed grouse (*Tympanuchus phasianellus*) surveys with the help of wildlife managers, cooperating agencies, and organizations (e.g., tribal agencies, U.S. Forest Service, college wildlife clubs). In 2014, ruffed grouse surveys were conducted between 21 April and 28 May, which was later than usual, but it allowed the peak of drumming activity to be captured during the late spring. Mean ruffed grouse drums per stop (dps) were 1.1 (95% confidence interval = 0.9–1.3) and increased 34% from the previous year. This increase occurred in the northern portion of the state; increases were not observed in southern regions. This may indicate the beginning of an upswing in the grouse cycle, which has been in the declining phase since 2009.

Sharp-tailed grouse surveys were conducted between 28 March and 28 May 2014, with 1,771 birds observed at 181 leks. The mean numbers of sharp-tailed grouse/lek were 5.4 (4.5–6.4) in the East Central (EC) survey region, 10.9 (9.8–12.1) in the Northwest (NW) region, and 9.8 (8.8–10.9) statewide. Comparisons between leks observed in consecutive years (2013 and 2014) were higher statewide ($t = 2.2$, $P = 0.04$) but increases were not significant in regional comparisons ($P > 0.05$).

INTRODUCTION

The ruffed grouse (*Bonasa umbellus*) is the most popular game bird in Minnesota, with an annual harvest averaging >500,000 birds (~150,000 -1.4 million birds). Ruffed grouse hunter numbers have been as high as 92,000 during the last decade, although hunter numbers did not peak with the recent peak in grouse numbers, as they have traditionally. Sharp-tailed grouse (*Tympanuchus phasianellus*) are also popular among hunters, with an annual harvest of 6,000–22,000 birds since the early-1990s and 5,000–10,000 hunters in Minnesota.

The Minnesota DNR coordinates grouse surveys each year to monitor changes in grouse populations through time. These surveys provide a reasonable index to population trends, when the primary source of variation in counts among years is change in densities. However, weather, habitat conditions, observer ability, and grouse behavior, also vary over time and can influence survey counts. Thus, making inferences from survey data over short time periods (e.g., a few years) can be tenuous. Nevertheless, over longer time periods and when large changes in index values occur, these surveys can provide a reasonable index to long-term grouse population trends. Spring surveys, in combination with hunter harvest statistics, provide evidence that the ruffed grouse population cycles at approximately 10-year intervals.

The first surveys of ruffed grouse in Minnesota occurred in the mid-1930s, and the first spring survey routes were established along roadsides in 1949. By the mid-1950s, ~50 routes were established with ~70 more routes added during the late-1970s and early-1980s. Since that time, spring drumming counts have been conducted annually to survey ruffed grouse in the forested regions of the state where ruffed grouse habitat occurs. Drumming is a low sound produced by males as they beat their wings rapidly and in increasing frequency to signal the location of their territory. These drumming displays also attract females that are ready to begin

nesting, so the frequency of drumming increases in the spring during the breeding season. The sound produced when male grouse drum is easy to hear and thus drumming counts are a convenient way to survey ruffed grouse populations in the spring.

Sharp-tailed grouse were first surveyed in Minnesota between the early-1940s and 1960. The current survey is based on counts at dancing grounds during the spring and was first conducted in 1976. Male sharp-tailed grouse display, or dance, together in open areas to attract females in the spring. This display consists of the males stomping their feet with out-stretched wings. Females visit the dancing grounds to select males for breeding. These dancing grounds, or leks, are reasonably stable in location from year to year, allowing surveyors to visit and count individuals each spring. Surveys are conducted in openland portions of the state where sharp-tailed grouse persist, although they were formerly much more widely distributed in Minnesota at the early part of the 20th century.

METHODS

Ruffed Grouse

Surveys for ruffed grouse were conducted along 121 established routes throughout the state. Each route consisted of 10 listening stops at approximately 1.6-km (1-mile) intervals. The placement of routes on the landscape was determined from historical survey routes, which were originally placed near ruffed grouse habitat in low traffic areas. Annual sampling of these historical routes provides information about temporal changes along the routes, but may not be representative of the counties or regions where the routes occurred.

Survey observers were solicited from among state, federal, tribal, private, and student biologists. Each observer was provided a set of instructions and route location information. No formal survey training was conducted but all observers had a professional background in wildlife science, and most had previously participated in the survey. Participants were asked to conduct surveys at sunrise during peak drumming activity (in April or May) on days that had little wind and no precipitation. Each observer drove the survey route once and listened for drumming at each stop for 4 minutes. Observers recorded the number of drums heard at each stop (not necessarily the number of individual grouse), along with information about phenology and weather at the time of the survey.

The number of drums heard per stop (dps) was used as the survey index value. I determined the mean dps for each route, for each of 4 survey regions (Figure 1), and for the entire state. For each survey region, I calculated the mean of route-level means for all routes partially or entirely within the region. Routes that traversed regional boundaries were included in the means for both regions. Because the number of routes within regions was not related to any proportional characteristic, I used the weighted mean of index values for the 4 Ecological Classification Sections (ECS) in the Northeast region and the 7 ECS sections in the state. The geographic area of the section was used as the weight for each section mean (i.e., Lake Agassiz, Aspen Parklands = 11,761 km², Northern Minnesota and Ontario Peatlands = 21,468 km², Northern Superior Uplands = 24,160 km², Northern Minnesota Drift and Lake Plains = 33,955 km², Western Superior Uplands = 14,158 km², Minnesota and Northeast Iowa Morainal (MIM) = 20,886 km², and Paleozoic Plateau (PP) = 5,212 km²). The area used to weight drum index means for the MIM and PP sections was reduced to reflect the portion of these areas within ruffed grouse range (~50%) using subsection boundaries. A 95% confidence interval (CI) was calculated to convey

the uncertainty of each mean index value using 10,000 bootstrap samples of route-level means for survey regions and the whole state. Confidence interval boundaries were defined as the 2.5th and 97.5th percentiles of bootstrap frequency distributions.

Sharp-tailed Grouse

Wildlife Managers and volunteers surveyed known sharp-tailed grouse lek locations in their work areas in the Northwest (NW) and East Central (EC) portions of the state (Figure 2). The NW region consisted of Lake Agassiz & Aspen Parklands, Northern Minnesota & Ontario Peatlands, and Red River Valley ECS sections. The EC region consisted of selected subsections of the Northern Minnesota Drift & Lake Plains, Western Superior Uplands, and Southern Superior Uplands sections. Some leks may have been missed, but most managers believed that they included most of the leks in their work area. Given the uncertainty in the proportion of leks missed, especially those occurring outside traditional areas, the survey may not necessarily reflect sharp-tailed grouse numbers in larger areas such as counties or regions.

Each cooperator was provided with instructions and asked to conduct surveys on ≥ 1 day in an attempt to obtain a maximum count of male sharp-tailed grouse attendance at each lek. Observers were asked to conduct surveys within 2.5 hours of sunrise under clear skies and during low winds (< 16 km/hr, or 10 mph) when lek attendance and ability to detect leks were expected to be greatest. Data recorded during each lek visit included the number of males, females, and birds of unknown sex.

The number of sharp-tailed grouse per dancing ground was used as the index value and was averaged for the NW region, the EC region, and statewide, using known males and birds of unknown sex. Observations of just 1 grouse were not included in the index. Data from former survey years were available for comparison, however, survey effort and success varied among years rendering comparisons of the full survey among years invalid. Therefore, to make valid comparisons between 2 consecutive years, only counts of birds from dancing grounds that were surveyed during both years were considered. Paired t-tests were used to test the significance of comparisons among years. Confidence intervals (95%) were calculated using 10,000 bootstrap samples of lek counts for each region and statewide.

RESULTS & DISCUSSION

Ruffed Grouse

Observers from 11 cooperating organizations surveyed routes between 21 April and 28 May 2014. Most routes (75%) were surveyed between 4 May and 16 May, with the median date (7 May) earlier than last year (May 10) but comparatively late (April 23 and 25 in 2010 and 2012, and May 1 and 3 in 2009 and 2011, respectively). Excellent (41%), Good (50%), and Fair (9%) survey conditions were reported for 116 routes reporting conditions, which is notable as the only time that more people reported good than excellent conditions in the last decade. However, the guidance provided was to survey during the peak of drumming activity in each area, if conditions would allow.

Statewide counts of ruffed grouse drums averaged 1.1 dps (95% confidence interval = 1.0–1.3 dps) during 2014 (Figure 3). Drum counts were 1.3 (1.1–1.5) dps in the Northeast ($n = 98$

routes), 1.2 (0.7–2.1) dps in the Northwest ($n = 8$), 0.8 (0.4–1.2) dps in the Central Hardwoods ($n = 13$), and 0.3 (0.1–0.5) dps in the Southeast ($n = 7$) regions (Figure 4a-d).

Statewide drum counts increased 34% this year. Increases were driven by changes in the northern portion of the state, in the prime ruffed grouse range. This increase is consistent with changes expected with the 10-year cycle, with the most recent peak in drum counts during 2009. The cycle is less pronounced in the more southern regions of the state, near the edge of their range.

Sharp-tailed Grouse

A total of 1,771 male sharp-tailed grouse and grouse of unknown sex was counted at 181 leks (Table 1) during 28 March - 28 May 2014. More leks (30%) were observed in 2014 than during 2013, in part due to the filling of several DNR Wildlife staff vacancies in northwestern Minnesota which permitted greater effort this year. Leks with ≥ 2 grouse were observed an average of 1.8 times.

The statewide index value of 9.8 (8.8–10.9) was centrally located among values observed since 1980 (Figure 5). In the EC survey region, 201 grouse were counted on 37 leks, and 1,570 grouse were counted on 144 leks in the NW region. The index value (i.e., grouse/lek) was higher statewide and in both regions compared to 2014, but confidence intervals overlapped those from the last few years (Table 1). Counts at leks observed during both years increased statewide from 2013 ($t = 2.2$, $P = 0.04$), but changes by region were not significant ($P > 0.05$) in either region (Table 2). These changes may indicate the beginning of an upswing in the cycle concordant with that of ruffed grouse. Sharp-tailed grouse population index values peaked with those for ruffed grouse in 2009, and appear to have troughed with them in 2013, although sharp-tailed grouse peaks can follow those of ruffed grouse by as much as 2 years.

ACKNOWLEDGEMENTS

The ruffed grouse survey was accomplished this year through the combined efforts of staff and volunteers at Chippewa and Superior National Forests (USDA Forest Service); Fond du Lac, Red Lake, and White Earth Reservations; 1854 Treaty Authority; Agassiz and Tamarac National Wildlife Refuges (U.S. Fish & Wildlife Service); Vermilion Community College; Cass County Land Department; and DNR staff at Aitkin, Baudette, Bemidji, Brainerd, Cambridge, Carlos Avery Wildlife Management Area (WMA), Cloquet, Crookston, Detroit Lakes, Fergus Falls, Grand Rapids, International Falls, Karlstad, Little Falls, Mille Lacs WMA, Park Rapids, Red Lake WMA, Rochester, Roseau River WMA, Sauk Rapids, Thief Lake WMA, Thief River Falls, Tower, Two Harbors, Whitewater WMA, and Winona work areas. I would like to thank DNR staff and volunteers at Aitkin, Baudette, Bemidji, Cambridge, Cloquet, Karlstad, International Falls, Tower, Thief River Falls, and Thief Lake work areas, staff and volunteers at Red Lake and Roseau River WMAs, and partners at Agassiz National Wildlife Refuge for participating in sharp-tailed grouse surveys. Laura Gilbert helped enter ruffed grouse data. Gary Drotts, John Erb, and Rick Horton organized an effort to enter the ruffed grouse survey data for 1982–2004, and Doug Mailhot and another volunteer helped enter the data. I would also like to thank Mike Larson for his assistance in the transition coordinating the surveys and for making helpful comments on this report. This work was funded in part through the Federal Aid in Wildlife Restoration Act.



Table 1. Sharp-tailed grouse / lek (≥ 2 males) at all leks observed during spring surveys each year in Minnesota.

Year	Statewide			Northwest ^a			East Central ^a		
	Mean	95% CI ^b	<i>n</i> ^c	Mean	95% CI ^b	<i>n</i> ^c	Mean	95% CI ^b	<i>n</i> ^c
2004	11.2	10.1–12.3	183	12.7	11.3–14.2	116	8.5	7.2–9.9	67
2005	11.3	10.2–12.5	161	13.1	11.5–14.7	95	8.8	7.3–10.2	66
2006	9.2	8.3–10.1	161	9.8	8.7–11.1	97	8.2	6.9–9.7	64
2007	11.6	10.5–12.8	188	12.7	11.3–14.1	128	9.4	8.0–11.0	60
2008	12.4	11.2–13.7	192	13.6	12.0–15.3	122	10.4	8.7–12.3	70
2009	13.6	12.2–15.1	199	15.2	13.4–17.0	137	10.0	8.5–11.7	62
2010	10.7	9.8–11.7	202	11.7	10.5–12.9	132	8.9	7.5–10.5	70
2011	10.2	9.5–11.1	216	11.2	10.2–12.2	156	7.8	6.7–8.9	60
2012	9.2	8.2–10.3	153	10.7	9.3–12.3	100	6.3	5.4–7.3	53
2013	9.2	8.2–10.2	139	10.5	9.3–11.7	107	4.8	3.8–5.9	32
2014	9.8	8.8–10.9	181	10.9	9.8–12.1	144	5.4	4.5–6.4	37

^a Survey regions; see Figure 1.

^b 95% CI = 95% confidence interval

^c *n* = number of leks in the sample.

Table 2. Difference in the number of sharp-tailed grouse / lek observed during spring surveys of the same lek in consecutive years in Minnesota.

Comparison ^b	Statewide			Northwest ^a			East Central ^a		
	Mean	95% CI ^c	<i>n</i> ^d	Mean	95% CI ^c	<i>n</i> ^d	Mean	95% CI ^c	<i>n</i> ^d
2004 - 2005	-1.3	-2.2– -0.3	186	-2.1	-3.5– -0.8	112	0.0	-1.0– 1.1	74
2005 - 2006	-2.5	-3.7– -1.3	126	-3.6	-5.3– -1.9	70	-1.1	-2.6– -0.6	56
2006 - 2007	2.6	1.5– 3.8	152	3.3	1.7– 5.1	99	1.2	0.1– 2.3	53
2007 - 2008	0.4	-0.8– 1.5	166	0.0	-1.6– 1.6	115	1.2	0.1– 2.5	51
2008 - 2009	0.9	-0.4– 2.3	181	1.8	-0.1– 3.8	120	-0.8	-2.1– -0.6	61
2009 - 2010	-0.6	-1.8– -0.6	179	-0.8	-2.6– 1.0	118	-0.1	-1.2– 1.0	61
2010 - 2011	-1.7	-2.7– -0.8	183	-1.8	-3.1– -0.5	124	-1.5	-2.8– -0.3	59
2011 - 2012	-2.0	-2.9– -1.1	170	-1.7	-2.9– -0.4	112	-2.4	-3.3– -1.6	58
2012 - 2013	-0.8	-2.0– -0.4	140	0.4	-1.3– 2.3	88	-2.9	-4.2– -1.8	52
2013 - 2014	1.4	0.1– 2.7	121	1.6	-0.3– 3.5	79	1.1	-0.1– 2.3	42

^a Survey regions; see Figure 1.

^b Consecutive years for which comparable leks were compared.

^c 95% CI = 95% confidence interval

^d *n* = number of leks in the sample. Here, a lek can have a 0 count in 1 of the 2 years and still be considered.




Figure 1. Survey regions for **ruffed grouse** in Minnesota. Northwest (NW), Northeast (NE), Central Hardwoods (CH), and Southeast (SE) survey regions are depicted relative to county boundaries (dashed lines) and influenced by the Ecological Classification System.




Figure 2. Survey regions for **sharp-tailed grouse** in Minnesota. Northwest (NW) and East Central (EC) survey regions are depicted relative to county boundaries (dashed lines) and influenced by Ecological Classification System Subsections boundaries.








Figure 3. Statewide ruffed grouse population index values in Minnesota. Bootstrap (95%) confidence intervals (CI) are provided after 1981, but different analytical methods were used prior to this and thus CI are not available for earlier years. The difference between 1981 and 1982 is biological and not an artifact of the change in analysis methods.

a.



b.

c.



d.

Figure 4a, b, c, d. Ruffed grouse population index values in the **Northeast** (a), **Northwest** (b), **Central Hardwoods** (c), and **Southeast** (d) survey regions of Minnesota. The mean for 1984-2004 is indicated by the dashed line. Bootstrap (95%) confidence intervals are provided for each mean. In the bottom panel, the CI for 1986 extends beyond area depicted in the figure.




Figure 5. **Sharp-tailed grouse** counted in Spring lek surveys statewide during 1980–2014. Bootstrap (95%) confidence intervals are provided for recent years. Annual means are not connected by lines because the same leks were not surveyed every year.



2014 MINNESOTA PRAIRIE-CHICKEN SURVEY

Charlotte Roy, Forest Wildlife Populations and Research Group

SUMMARY OF FINDINGS

Greater prairie-chickens (*Tympanuchus cupido pinnatus*) were surveyed in 16 of 17 survey blocks during the spring of 2014. Observers located 138 booming grounds and counted 1,245 male prairie-chickens and 101 birds of unknown sex. Estimated densities of 0.10 (0.07-0.13) booming grounds/km² and 9.8 (8.4-11.2) males/booming ground within the survey blocks were similar to densities during recent years and during the 10 years preceding modern hunting seasons (i.e., 1993–2002).

INTRODUCTION

Historically, greater prairie-chicken (*Tympanuchus cupido pinnatus*) range in Minnesota was restricted to the southeastern portion of the state. However, dramatic changes in their range occurred in the 19th century as settlers expanded and modified the landscape with farming and forest removal, providing abundant food sources and access to new areas. However, as grass continued to be lost from the landscape, prairie-chicken populations began to decline, their range contracted, and hunting closed after 1942. In an attempt to bolster populations and expand prairie-chicken range, the Minnesota Department of Natural Resources (DNR) conducted a series of translocations in the Upper Minnesota River Valley during 1998-2006. Today, the beach ridges of glacial Lake Agassiz hold most of Minnesota's prairie-chickens, but their populations do extend southward (Figure 1). Hunting was re-opened using a limited-entry season in 2003, and approximately 120 prairie-chickens are now harvested annually.

With the opening of the new hunting season, the DNR had a greater interest in the monitoring of prairie-chicken populations, which the Minnesota Prairie-Chicken Society (MPCS) had been coordinating since 1974. The DNR, in collaboration with MPCS members, began coordinating prairie-chicken surveys and adopted a standardized survey design in 2004. These surveys are conducted at small open areas called leks, or booming grounds, where male prairie-chickens display for females in the spring and make a low-frequency booming vocalization that can be heard for miles.

Prairie-chickens continue to be surveyed to monitor changes in population densities over time. However, density estimates can be costly and difficult to obtain, so instead we count individuals and make the assumption that changes in density are the primary source of variation in counts among years. If true, counts should provide a reasonable index to long-term trends in prairie-chicken populations. However, counts are also influenced by weather, habitat conditions, observer ability, and bird behavior among other factors, which make it difficult to make inferences over short periods of time (e.g., a few annual surveys) or from small changes in index values. Nevertheless, over long time periods and when changes in index values are large, inferences from prairie-chicken surveys are more likely to be valid.

METHODS

Cooperating biologists and volunteers surveyed booming grounds in 16 of 17 designated survey blocks in western Minnesota (Figure 2) during late-March through mid-May. Each survey block was nonrandomly selected so that surveys would be conducted in areas where habitat was expected to be good (i.e., grassland was relatively abundant) and booming grounds were known to occur. Each surveyor attempted to find and observe each booming ground repeatedly in his/her assigned block, which comprised 4 sections of the Public Land Survey (approximately 4,144 ha). We obtained multiple counts at each booming ground in the morning because male attendance at leks varies throughout the season and throughout the day.

During each survey, observers obtained visual counts of males, females, and birds of unknown sex from a distance with binoculars. Sex was determined through behavior; males display conspicuously, and females do not. If no birds were displaying during the survey period, then sex was recorded as unknown. When a reliable count could not be obtained visually because vegetation or topography prevented it, birds were flushed for counts and sex was recorded as unknown. Most birds for which sex was unknown were likely male because female attendance at leks is sporadic, and they are less conspicuous during lek attendance than displaying males.

In the analysis, I used counts of males and unknowns at each booming ground but not females. Booming grounds were defined as having ≥ 2 males, so observations of single males were not counted as leks. Data were summarized by hunting permit area and spring survey block. The survey block data were separated into a core group and a periphery group for analysis. The core group had a threshold density of approximately 1.0 male/km² during 2010, and was located proximally to other such blocks (Figure 2). I compared densities of leks and prairie-chickens to estimated densities from previous years.

I also encouraged surveyors to submit observations of booming grounds outside the survey blocks because these observations may provide additional information that is helpful to prairie-chicken management. These data were included in estimates of minimum abundance of prairie-chickens. However, these data were not used in the analysis of lek and prairie-chicken densities because effort and methods may have differed from those used in the survey blocks.

RESULTS & DISCUSSION

Observers from DNR Division of Fish and Wildlife, the U.S. Fish & Wildlife Service, and The Nature Conservancy, as well as many unaffiliated volunteers counted prairie-chickens between 24 March and 23 May 2014. Observers located 138 booming grounds and observed 1,245 male prairie-chickens and 101 birds of unknown sex within and outside survey blocks during 2014 (Table 1). These counts represent a minimum number of prairie-chickens in Minnesota during 2014, but because survey effort outside of survey blocks is not standardized among years, these counts should not be compared among years or permit areas.

Table 1. Minimum abundance of prairie-chickens within and outside hunting permit areas in Minnesota during spring 2014. Lek and bird counts are not comparable among permit areas or years.

Permit Area	Area (km ²)	Leks	Males	Unk ^a
803A	1,411	18	163	0
804A	435	NA	NA	NA
805A	267	14	168	0
806A	747	10	60	0
807A	440	25	151	26
808A	417	19	248	0
809A	744	13	152	0
810A	505	7	83	0
811A	706	8	37	27
812A	914	10	30	28
813A	925	3	56	0
PA subtotal	7,511	127	1,148	81
Outside PAs ^b	NA ^c	11	97	20
Grand total	NA ^c	138	1,245	101

^a Unk = prairie-chickens for which sex was unknown, but which were probably males.

^b Counts done outside permit areas (PA).

^c NA = not applicable because the area outside permit areas was not defined.

Within the standardized survey blocks, 669 males and birds of unknown sex were counted on 68 booming grounds during 2014 (Table 2). Each lek was observed an average of 1.8 times (median = 1), with 53% of booming grounds observed just once. Densities of prairie-chickens in the 10 core survey blocks were 0.11 (0.07–0.14) booming grounds/km² and 10.9 (9.1–12.7) males/booming ground (Table 2, Figure 2). In 6 of the 7 peripheral survey blocks, densities were 0.08 (0.03–0.14) booming grounds/km² and 7.8 (5.9–9.6) males/booming ground.

The density of 0.10 (0.07–0.13) booming grounds/km² in all survey blocks during 2014 was similar to densities during recent years (Table 2, Figure 3) and the average of 0.08 (0.06–0.09) booming grounds/km² during the 10 years preceding recent hunting seasons (i.e., 1993–2002). Similarly, the density of 9.8 (8.4–11.2) males/booming ground in all survey blocks during 2014 was comparable to densities during recent years and similar to the average of 11.5 (10.1–12.9) males/booming ground observed during 1993–2002 (Table 2, Figure 3). These counts should not be regarded as estimates of abundance because detection probabilities of leks and birds have not been estimated. However, if we assume that detection probabilities are similar among years, then this index can be used to monitor changes in abundance among years.

Table 2. Prairie-chicken counts within survey blocks in Minnesota.

Range ^b	Survey Block	Area (km ²)	2014		Change from 2013 ^a	
			Booming grounds	Males ^c	Booming grounds	Males ^c
Core	Polk 1	41.2	6	49	-1	-13
	Polk 2	42.0	6	97	-8	-51
	Norman 1	42.0	1	13	-1	-3
	Norman 2	42.2	2	33	-5	-37
	Norman 3	41.0	9	49	4	-9
	Clay 1	46.0	6	73	0	-24
	Clay 2	41.0	2	43	0	-6
	Clay 3	42.0	5	51	-1	-35
	Clay 4	39.0	3	27	1	0
	Wilkin 1	40.0	4	47	-1	-20
	Core subtotal	415.0	44	482	-12	-198
Periphery	Mahnomen	41.7	3	37	1	21
	Becker 1	41.4	10	58	NA ^d	NA ^d
	Becker 2	41.7	4	33	2	-1
	Wilkin 2	41.7	2	20	0	5
	Wilkin 3	42.0	3	25	-1	-4
	Otter Tail 1	41.0	2	14	-1	-6
	Otter Tail 2	40.7	NA	NA	NA	NA
		Periphery subtotal	290.6	24 ^e	187 ^e	11 ^e
	Grand total	705.5	68 ^e	669 ^e	-1 ^e	-125 ^e

^a The 2013 count was subtracted from the 2014 count, so positive values indicate increases.

^b Survey blocks were categorized as within the core or periphery of the Minnesota prairie-chicken range based upon bird densities and geographic location.

^c Includes birds recorded as being of unknown sex but excludes lone males.

^d Surveys were not conducted in this block during 2013.

^e These totals only reflect blocks for which count data were available.

ACKNOWLEDGMENTS

I would like to thank cooperators who conducted and helped coordinate the prairie-chicken survey. Cooperators within the DNR included Ross Hier, Emily Hutchins, Brian Torgusson, and Michael Oehler; cooperators with The Nature Conservancy included Brian Winter, Travis Issendorf, and volunteers Pat Beauzay, Rick Julian, Dennis Thielen, Matt Mecklenburg, Candis Sommerfeld, Bob O'Connor, and Tony Nelson; cooperators with the US Fish and Wildlife Service included Doug Wells, Shawn Papon, Chad Raitz, Maria Fosado, Larry Hanson, Stacy Salvevold, Jessica Dowler, Jacob Kaplan, and Trina Brennan; and numerous additional volunteers participated including Steve Bommersbach, Dan Svedarsky, Tom Kucera, and Doug Hedtke. This survey was funded in part by the Wildlife Restoration (Pittman-Robertson) Program W-69-S-13 Project #16. Mike Larson provided assistance and comments which improved this report.






Figure 1. Primary greater prairie-chicken range in Minnesota (shaded area) relative to county boundaries. The range boundary was based on Ecological Classification System Land Type Associations and excludes some areas known to be occupied by prairie-chickens.




Figure 2. Prairie-chicken lek survey blocks (41 km², labeled squares) and hunting permit areas (thick grey lines) in western Minnesota. Survey blocks were either in the core (black) or periphery (white) of the range with a threshold of 1.0 male/km² in 2010, and were named after their respective counties (thin black lines). Permit areas were revised in 2013 to eliminate 801A and 802A, modify 803A, and add 812A and 813A. See previous reports for former permit area boundaries.




Figure 3. Mean prairie-chicken males/booming ground (circles connected by solid line) and booming grounds/km² (triangles connected by dashed line) in survey blocks in Minnesota with 95% confidence intervals. Counts for 6 of the survey blocks in 2011, including 4 blocks in the core, were not available for this report.



2014 AERIAL MOOSE SURVEY

Glenn D. DelGiudice, Forest Wildlife Populations and Research Group

INTRODUCTION

Each year, we conduct an aerial survey in northeastern Minnesota in an effort to monitor moose (*Alces alces*) numbers and fluctuations in the overall status of Minnesota's largest deer species. The primary objectives of this annual survey are to estimate moose numbers, calf:cow and bull:cow ratios. We use these data to determine and examine the population's trend and composition, to contribute to our understanding of moose ecology, and to set the harvest quota for the subsequent hunting season when applicable.

METHODS

We estimated moose numbers, age and sex ratios by flying transects within a stratified random sample of survey plots (Figure 1). All survey plots are reviewed and re-stratified as low, medium, or high density about every 5 years based on past survey observations of moose, locations of harvested moose in past years, and extensive field experience of moose managers and researchers. For the most recent re-stratification (November 2013), survey plots were classified as low, medium, or high based on whether < 2 , 3-7, or ≥ 8 moose, respectively, would be expected to occur in a specific plot. Stratification is most important to optimizing precision of our survey estimates. In 2012, we added a 4th stratum to represent a series of 9 plots which have undergone disturbance by wild fire, prescribed burning, and timber harvest. Each year since, these same 9 plots are surveyed in an effort to evaluate the effect of disturbance on moose density over time.

As in previous years, all survey plots were rectangular (5 x 2.67 mi.) and oriented east to west with 8 transects. Minnesota Department of Natural Resources (MNDNR) Enforcement pilots flew the Bell Jet Ranger (OH-58) helicopters used to conduct the survey. We sexed moose using the presence of antlers or the presence of a vulval patch (Mitchell 1970), nose coloration, bell size and shape, and identified calves on the basis of size and behavior. We used the program DNRSurvey on Toughbook[®] tablet style computers to record survey data. DNRSurvey allowed us to display transect lines superimposed on a background of aerial photography, observe the aircraft's flight path over this background in real time, and record data using a tablet pen with a menu-driven data entry form. Two of the primary strengths of this survey are the consistency and standardization of the methods since 2005 and the long-term consistency of the survey team personnel.

We accounted for visibility bias by using a sightability model (Giudice et al. 2012). We developed this model between 2004 and 2007 using moose that were radiocollared as part of research on the dynamics of the northeastern moose population (Lenarz et al. 2009). Logistic regression indicated that the covariate "visual obstruction" (VO) was the most important covariate in determining whether radiocollared moose were observed. We defined VO as the proportion of vegetation within a circle (30-ft. radius or roughly 4 moose lengths) that would prevent you from seeing a moose when circling that spot from an oblique angle. If we observed more than 1 moose at a location, VO was based on the first moose sighted. We used uncorrected

estimates (no visibility bias correction) of bulls, cows, and calves, adjusted for sampling, to calculate the bull:cow and calf:cow ratios (i.e., using the combined ratio estimator; Cochran 1977:165).




Figure 1. Moose survey area and sample plots flown in the 2014 aerial moose survey. The study area for ongoing MNDNR moose research also is shown.

RESULTS AND DISCUSSION

The survey was conducted from 7 to 18 January 2014. It consisted of 9 actual survey days and included 52 survey plots. Snow depths were consistently greater than the minimum 8” desired for the survey (92% of plots had >16”, 8% had 8-16”), with 96% of the plot surveys conducted under good survey conditions and 2% under fair conditions. Overall, survey conditions were notably better than during the past several years. During the survey flights, 419 moose were observed on 41 (79%) of the 52 plots flown (694 mi²), including 176 bulls, 174 cows, 65 calves, and 4 unclassified moose. This apparent occupancy of plots compares to a 10-year average of 82%. An average of 10.2 moose were observed per “occupied” plot (range = 1-51 moose) compared to a 10-year average of 12.2 moose. Estimates of the calf:cow and bull:cow ratios were 0.44 and 1.24, respectively, both among the highest ratios since 2005 (Table 1).

After adjusting for sampling and sightability, we estimated the population in northeastern Minnesota at 4,350 (3,220–6,210) moose (Table 1). Based on the log rate of change (0.456;

90% CI: 0.086, 0.827), the 2014 population estimate was significantly higher (58%; 90% CI: 9-128%) than the 2013 estimate of 2760 moose, but similar to the 2012 and 2011 estimates of 4,230 and 4,900 moose, respectively. As can be noted from the 90% confidence limits associated with the population point estimates (3,220-6,210; Table 1, Figure 2), statistical uncertainty inherent in aerial wildlife surveys, even of large, relatively conspicuous animals such as moose during the winter, can be quite large due to the varied (1) occurrence of dense vegetation, (2) habitat use by moose, (3) behavioral responses to aircraft, (4) effects of annual environmental conditions (e.g., snow depth) on their movements, and (5) the interaction of these factors. Past aerial survey and research results have indicated that the trend of the population of northeastern Minnesota has been declining since 2006 (Lenarz et al. 2010, DelGiudice 2013). Despite this year's higher point estimate, the downward trend persists ($r^2 = 0.82$, $P = 0.0003$, Figure 2). Lenarz et al. (2010) used simulation modeling to integrate survival and reproductive rates measured between 2002 and 2008 and found that the population was decreasing approximately 15% per year over the long-term. The 2013 estimate (2,760 moose) indicated a 35% decrease from 2012 and a 52% decrease in the population since 2010, not inconsistent with the declining trend, but exceeding the projected rate of change (Table 1). It is likely that the population was underestimated in 2013 and that with almost optimum snow and survey conditions this year, more moose were observed and the estimate is more reflective of actual moose numbers, although the variability associated with the estimate is large due to atypically high numbers of moose being observed in low and medium density plots.

Table 1. Estimated moose numbers, 90% confidence intervals, and calf:cow ratios, percent calves, percent cows with twins, and bull:cow ratios estimated from aerial surveys in northeastern Minnesota, 2005-2014.

Survey	Estimate	90% Confidence Interval	Calf: Cow	% Calves	% Cows w/ twins	Bull: Cow
2005	8,160	5,960 – 11,170	0.52	19	9	1.04
2006	8,840	6,670 – 11,710	0.34	13	5	1.09
2007	6,860	5,230 – 9,000	0.29	13	3	0.89
2008	7,890	5,970 – 10,420	0.36	17	2	0.77
2009	7,840	6,190 – 9,910	0.32	14	2	0.94
2010	5,700	4,480 – 7,250	0.28	13	3	0.83
2011	4,900	3,810 – 6,290	0.24	13	1	0.64
2012	4,230	3,190 – 5,600	0.36	15	6	1.08
2013	2,760	2,120 – 3,580	0.33	13	3	1.23
2014	4,350	3,220 – 6,210	0.44	15	3	1.24

Estimated calf recruitment from this year's survey remained *relatively* high and similar to last year's estimate (Table 1). The calf:cow ratio in mid-January 2014 was 0.44, up slightly from last year's survey (0.33), and calves represented 15% of the total moose observed, also slightly elevated from last year's estimate (Table 1). Like last year, only 3% of the cow moose were accompanied by twins (Table 1). Based on survey results calf survival through to mid-January 2014 appears relatively high. However, an ongoing study of GPS-collared moose calves

indicates that calf survival is low (Severud and DelGiudice, unpublished data), and *annual* recruitment of the calves is not actually determined until the next spring calving season when winter survey-observed calves become yearlings. At this point, little is known about the survival rates of moose calves during the period between the annual winter survey and subsequent spring calving. It also is important to note that adult moose survival has the most significant impact on annual changes in the moose population (Lenarz et al. 2010), and elevated annual mortality of adult moose has continued during the past year (~21%, Carstensen et al., unpublished data).

The estimated bull:cow ratio (Table 1; Figure 4) was similar to last year's estimate and is the highest since 2005. Further, the past two year's estimated bull:cow ratios indicate that adult bulls may outnumber adult females, although there is a great deal of variability associated with these annual ratio estimates. Consequently, there is no clear upward or downward long-term trend (2005-2014) in bull:cow ratios.




Figure 2. Point estimates, 90% confidence intervals, and trend line of estimated moose numbers in northeastern Minnesota, 2005-2014. (Note: The 2005 survey was the first to be flown with helicopters, and to include a sightability model and a uniform grid of east-west oriented rectangular 5 x 2.67 mi plots).




Figure 3. Estimated calf:cow ratios (solid diamonds, dashed trend line) and percent calves (solid squares, solid trend line) from aerial moose surveys in northeastern Minnesota, 2005-2014.




Figure 4. Estimated bull:cow ratios, 90% confidence intervals, and trend line from aerial moose surveys in northeastern Minnesota, 2005-2014.

ACKNOWLEDGMENTS

This survey is an excellent partnership between the Division of Enforcement, the Division of Fish and Wildlife, the Fond du Lac Band of Lake Superior Chippewa and the 1854 Treaty Authority. In particular, I would like to thank Thomas Buker, Chief Pilot, for coordinating all of the aircraft and pilots; Tom Rusch for coordinating flights and survey crews; and Mike Schrage (Fond du Lac Band of Lake Superior Chippewa) and Andy Edwards (1854 Treaty Authority) for securing supplemental survey funding from their respective groups. Enforcement pilots, Brad Maas and John Heineman, skillfully piloted the aircraft during the surveys, and Tom Rusch, Andy Edwards, Mike Schrage, and Nancy Hansen flew as observers; their efforts are gratefully appreciated. I also want to thank John Giudice who continues to provide critical statistical consultation and analyses. Thanks to Barry Sampson for creating the process to generate the GIS survey maps and GPS coordinates for the transect lines and for his work on re-stratification of the survey plots, as well as to Bob Wright, Brian Haroldson, and Chris Pouliot for the creation of the program DNRSurvey. Bob also modifies the software as needed and each year provides refresher training for survey observers using DNRSurvey.

LITERATURE CITED

- Cochran, W. G. 1977. Sampling techniques. Third edition. Wiley and Sons, New York, USA.
- DelGiudice, G. D. 2013. 2013 Aerial moose survey. Minnesota Department of Natural Resources, Section of Wildlife, unpublished report. St. Paul, Minnesota. 6pp.
- Fieberg, J. 2012. Estimating population abundance using sightability models: R sightability model package. *Journal of Statistical Software* 51:1-20.
- Gasaway, W. C., and S. D. DuBois. 1987. Estimating moose population parameters. *Swedish Wildlife Research (Supplement)* 1:603–617.
- Giudice, J. H., J. R. Fieberg, M. S. Lenarz. 2012. Spending degrees of freedom in a poor economy: a case study of building a sightability model for moose in northeastern Minnesota. *Journal of Wildlife Management* 76:75-87.
- Lenarz, M. S., M. E. Nelson, M. W. Schrage, and A. J. Edwards. 2009. Temperature mediated moose survival in northeastern Minnesota. *Journal of Wildlife Management* 73:503-510.
- Lenarz, M. S., J. Fieberg, M. W. Schrage, and A. J. Edwards. 2010. Living on the edge: viability of moose in northeastern Minnesota. *Journal of Wildlife Management*. 74:1013-1023.
- Mitchell, H.B. 1970. Rapid aerial sexing of antlerless moose in British Columbia. *Journal Wildlife Management*. 34: 645-646.
- R Development Core Team. 2011, R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Version 2.13.1, ISBN 3-900051-07-0 <http://www.r-project.org/>.



REGISTERED FURBEARER POPULATION MODELING, 2014 Report



Drawing by Gilbert Proulx

John Erb, Forest Wildlife Populations and Research Group

INTRODUCTION

For populations of secretive carnivores, obtaining field-based estimates of population size remains a challenging task (Hochachka et al. 2000; Wilson and Delehay 2001; Conn et al. 2004). This is particularly true when one is interested in annual estimates, multiple species, or large areas. Nevertheless, population estimates are desirable to assist in making management or harvest decisions. Population modeling is a valuable tool for synthesizing our knowledge of population demography, predicting outcomes of management decisions, and approximating population size.

In the late 1970s, Minnesota developed population models for 4 species of carnivores (fisher, marten, bobcat, and otter) to help ‘estimate’ population size and track population changes. All are deterministic accounting models that do not currently incorporate density-dependence. However, annual adjustments to demographic inputs are often made for bobcats, fishers, and martens in response to the known or assumed influence of factors such as prey fluctuations and winter conditions, or competitor/predator density. Modeling projections are interpreted in conjunction with harvest data and results from any annual field-based track surveys.

METHODS

Primary model inputs include the estimated 1977 ‘starting’ population size, estimates of age-specific survival and reproduction, and sex- and age-specific harvest data. Reproductive inputs are based largely on carcass data collected in the early 1980s, and for bobcats, additional data collected in 1992 and from 2003-present. Initial survival inputs were based on a review of published estimates in the literature, but are periodically adjusted as noted above. In some cases, parameter adjustments for previous years are delayed until additional data on prey abundance trends is available. Hence, population estimates reported in previous reports may not always match those reported in current reports. Obtaining updated Minnesota-specific survival and reproductive estimates remains a goal of ongoing research.

Harvest data is obtained through mandatory furbearer registration. A detailed summary of 2013 harvest information is available in a separate report. Bobcat, marten, and fisher age data is obtained via x-ray examination of pulp cavity width or microscopic counts of cementum annuli from teeth of harvested animals. Although the population models only utilize data for the 3 age-classes (juvenile, yearling, adult), cementum annuli counts have periodically been collected for all non-juveniles either to examine age-specific reproductive output (bobcats) or to obtain periodic information on year-class distribution for selected species. In years where age data is not obtained for a given species, harvest age proportions are approximated using averages computed from the most recent period when data was collected.

For comparison to model projections, field-based track survey indices are presented in this report as running 3-year (t-1, t, t+1) averages of the observed track index, with the most recent year’s average computed as $(2/3 \times \text{current index} + 1/3 \times \text{previous index})$. More detailed descriptions of scent post and winter track survey methods and results are available in separate reports.

RESULTS AND DISCUSSION

Bobcat. The 2013 registered DNR trapping and hunting harvest declined 45% to 1,038 (Table 1). Total modeled harvest, which includes reported tribal take, was 1,138. At this time, age and reproductive analyses from harvested bobcats have not been completed. Data from previous years is presented in Table 1 and Figure 1.

Based on projections from the population model, 20% of the fall 2013 population was harvested. As a result of the high harvests in 2011 and 2012, plus an assumed reduction in survival from two severe winters and reduced ungulate prey, population modeling projects a 7% decline in the bobcat population (Figure 3), with an estimated 2014 spring population size of ~ 4,000 (Figure 3). Both track indices remain near the upper end of their previously recorded range (Figure 3).

Fisher. Over the past 7 years, the fisher harvest season has become progressively more conservative, with the 2013 season lasting only 6 days and a per trapper limit of 2 (identical to the 2012 season). Fisher harvest this year under the DNR framework decreased ~ 11% to 1,146 (Table 2). Modeled harvest, which includes reported tribal take, was 1,295.

After a 15-year lapse, fisher carcass collections were resumed in 2010 to collect current information on harvest age distribution. A total of 1,040 carcasses were collected in 2013 (Table 2). Juveniles accounted for 51% of the total harvest, similar to the previous 3-year average (a period of estimated population decline), but well below the average (64%) from 1977-1994 (generally a period of population growth). Similarly, the juvenile:adult female ratio was 3.4, slightly below the recent average (3.8), and well below the 1977-1994 average of 6.4 (Table 2). Average age of harvested males and females was 1.5 and 1.9, respectively, with the harvest being comprised of few fishers over the age of 1.5 (Figures 4 and 5).

Based on model projections, 17% of the fall fisher population was harvested during the 2013 season. In spite of the conservative seasons in recent years, this year's harvest may still have exceeded the current sustainable level, and the 3-year-averaged winter track index for fisher once again declined to its lowest level (Figure 6). It remains possible that last winter's track index could be biased low due to the deep snow and cold weather that may have reduced fisher activity. Furthermore, the population model inputs (and winter track index) are assumed to best reflect the fisher population only in the historically 'core areas' of northern Minnesota. Along the southern and western periphery of fisher range, harvest data and other anecdotal information clearly indicate a population increase over the past 5-10 years, though these areas represent a comparatively small portion of overall fisher range. Acknowledging these caveats, modeling projects a 1.5% decline in the fisher population with an estimated 2014 spring population size of ~ 6,000 fishers (Figure 6).

Marten. As with fishers, the marten harvest season has become progressively more conservative in recent years, with the 2013 season lasting 6 days and a per trapper limit of 2 martens. The 2012 season was also 6 days, but with a limit of 5 martens per trapper. Harvest this year under the DNR framework was 1,014, down 31% from last year and the lowest since 1991 (Table 3). Modeled harvest, which includes reported tribal take, was 1,323.

Juveniles accounted for 43% of the total harvest with a juvenile:adult female ratio of 3.5 (Table 3, Figure 7). Both numbers are similar to their 2002-12 averages (3.8; 43%) when modeling projects the population to have been in decline, and well below levels estimated from the 1986 – 2001 period (10; 61%) when the population is projected to have increased. Average age of both harvested males and females was 2.1 (Figures 8 and 9).

Based on projections from the marten population model, 15% of the fall 2013 population was harvested. This represents the lowest estimated harvest rate since 2001 (Table 3). Although modeling projects that conservative seasons have slowed or periodically stopped the population decline, collective data has yet to suggest a multi-year increase (Figure 10). In spite of the reduced 2013 marten harvest, the estimated harvest rate was still close to the projected sustainable level. In addition, downward adjustments have been made to juvenile survival inputs the past 2 years because of an apparent low in small mammal abundance. Modeling projects a 1% decline in the population from last year, with an estimated 2014 spring population size of ~ 7,500 martens.

Otter. From 1977 - 2007, otter harvest was only allowed in the northern part of the state. From 2007-2009, otter harvest was allowed in 2 separate zones with differing limits (4 otter in the north zone, 2

in the southeast zone). Beginning in 2010, otter harvest was allowed statewide with a consistent limit of 4 otter per trapper. Statewide otter harvest in 2013 under the DNR framework decreased 11% to 2,824 (Table 4). Modeled statewide otter harvest, which includes tribal take, was 2,993 (Table 4).

An estimated 21% of the fall 2013 otter population was harvested. Carcass collections ended in 1986 so no age or reproductive data are available, and no harvest-independent otter survey is currently established. Because demographic parameters in the otter model are typically held constant, annual differences in population trajectory are largely a function of varying harvest levels. Harvest levels exceeding ~3,000 for consecutive years typically predict population declines. After the population declined and then rebounded from 2002-12 as a result of a cycle in fur prices and harvests, modeling indicates the population has once again declined the past 2 years as a result of higher harvests and fur prices. Nevertheless, the population remains near the high end of levels estimated over the past 35 years (Figure 11). The 2014 spring population is estimated to be ~ 11,300, a 2.6% decline from last year.

LITERATURE CITED

CAUGHLEY, G. 1974. Interpretation of age ratios. *Journal of Wildlife Management* 38: 557-562.

CONN, P. B., L. L. BAILEY, and J. R. SAUER. 2004. Indexes as surrogates to abundance for low-abundance species. Pages 59-76 *in* W. L. Thompson, editor. *Sampling rare or elusive species: Concepts, designs, and techniques for estimating population parameters*. Island Press, Washington, D.C., USA.

HARRIS, N. C., M. J. KAUFFMAN, and L. S. MILLS. 2008. Inferences about ungulate population dynamics derived from age ratios. *Journal of Wildlife Management* 72: 1143-1151.

HOCHACHKA, W. M., K. MARTIN, F. DOYLE, and C. J. KREBS. 2000. Monitoring vertebrate populations using observational data. *Canadian Journal of Zoology* 78:521-529.

WILSON, G. J., and R. J. DELEHAY. 2001. A review of methods to estimate the abundance of terrestrial carnivores using field signs and observation. *Wildlife Research* 28:151-164.

Table 1. Bobcat harvest data, 1984 to 2013.

Year	DNR Harvest	Modeled Harvest ¹	% Autumn Pop. Taken ²	Carcasses Examined	% juveniles	% yearlings	% adults	Juv: Ad. Female ratio	% male juveniles	% male yearlings	% male adults	Overall % males	Mean Pelt Price ³
1984	280	288	15	288	37	13	50	1.4	52	66	44	51	\$76
1985	119	121	6	99	33	19	48	1.2	41	41	43	42	\$70
1986	160	160	8	132	26	17	57	0.9	53	32	51	51	\$120
1987	214	229	12	163	33	16	51	1.3	44	52	48	48	\$101
1988	140	143	7	114	40	18	42	1.7	58	62	46	54	\$68
1989	129	129	6	119	39	17	44	2.0	49	53	56	53	\$48
1990	84	87	4	62	20	34	46	0.8	58	80	44	59	\$43
1991	106	110	5	93	35	33	32	3.5	59	55	70	61	\$37
1992	167	167	7	151	28	22	50	1.2	55	45	53	53	\$28
1993	201	210	8	161	32	20	48	1.4	51	45	52	50	\$43
1994	238	270	11	187	26	16	58	0.8	64	43	45	50	\$36
1995	134	152	6	96	31	15	54	2.7	57	71	79	71	\$32
1996	223	250	10	164	35	20	45	1.8	51	30	49	46	\$33
1997	364	401	16	270	35	16	49	1.4	60	37	43	48	\$30
1998	103	107	4	77	29	26	45	1.6	59	60	60	60	\$28
1999	206	228	8	163	18	24	58	0.8	55	59	62	60	\$24
2000	231	250	8	183	31	26	43	1.4	54	59	50	53	\$33
2001	259	278	8	213	30	21	49	1.3	46	45	47	46	\$46
2002	544	621	15	475	27	25	48	1.1	68	51	48	54	\$72
2003	483	518	13	425	25	13	62	0.9	62	48	54	55	\$96
2004	631	709	14	524	28	34	38	1.7	52	40	55	49	\$99
2005	590	638	13	485	25	13	62	0.8	51	48	47	48	\$96
2006	890	983	18	813	26	17	57	1.1	60	51	58	57	\$101
2007	702	758	14	633	34	14	52	1.2	55	60	47	52	\$93
2008	853	928	15	714	26	25	49	1.1	55	52	50	52	\$75
2009	884	942	15	844	24	22	54	0.9	57	46	51	51	\$43
2010	1012	1042	15	955	38	16	46	1.4	62	55	42	52	\$71
2011	1711	1898	26	1626	23	21	55	0.8	61	73	47	56	\$98
2012	1875	2026	29	1744	25	19	56	1.0	63	53	54	56	\$144
2013	1038	1138	20	634				data not yet available				56	\$89

¹Includes DNR and Tribal harvests

²Estimated from population model; includes estimated non-reported harvest of 10%.

³Average pelt price based on a survey of in-state fur buyers only.

Bobcat Harvest Age-Classes




Figure 1. Age-class distribution of bobcats harvested in Minnesota 1977 - 2012.




Figure 2. Pregnancy rates for yearling and adult bobcats in Minnesota, 2003-2012.




Figure 3. Bobcat populations, harvests, and survey indices, 1977-2014. Harvests include an estimate of non-reported take.

Table 2. Fisher harvest data, 1984 to 2013.

Year	DNR harvest	Modeled Harvest ¹	% Autumn Pop. Harvested ²	Carcasses examined	% juveniles	% yearlings	% adults	Juv: Ad. Female ratio	% male juveniles	% male yearlings	% male adults	% males overall	Pelt price Males ³	Pelt price Females ³
1984	1285	1332	18	1270	63	20	17	6.6	52	45	45	49	\$70	\$122
1985	678	735	10	712	63	20	18	5.5	46	40	34	43	\$74	\$130
1986	1068	1186	15	1186	59	24	18	5.2	48	50	37	46	\$84	\$162
1987	1642	1749	22	1534	63	15	22	4.7	46	40	37	43	\$84	\$170
1988	1025	1050	14	805	70	15	15	6.7	48	45	33	45	\$54	\$100
1989	1243	1243	16	1024	64	19	17	5.8	47	47	36	45	\$26	\$53
1990	746	756	9	592	65	14	21	4.4	44	55	30	43	\$35	\$46
1991	528	528	6	410	66	21	13	7.5	50	52	35	48	\$21	\$48
1992	778	782	8	629	58	21	21	4.8	42	55	45	46	\$16	\$29
1993	1159	1192	10	937	59	22	19	6.0	47	37	42	44	\$14	\$28
1994	1771	1932	15	1360	56	18	26	4.0	47	54	44	48	\$19	\$30
1995	942	1060	8	-	-	-	-	-	-	-	-	45	\$16	\$25
1996	1773	2000	14	-	-	-	-	-	-	-	-	45	\$25	\$34
1997	2761	2974	20	-	-	-	-	-	-	-	-	45	\$31	\$34
1998	2695	2987	20	-	-	-	-	-	-	-	-	45	\$19	\$22
1999	1725	1880	13	-	-	-	-	-	-	-	-	45	\$19	\$20
2000	1674	1900	13	-	-	-	-	-	-	-	-	45	\$20	\$19
2001	2145	2362	15	-	-	-	-	-	-	-	-	54	\$23	\$23
2002	2660	3028	20	-	-	-	-	-	-	-	-	54	\$27	\$25
2003	2521	2728	19	-	-	-	-	-	-	-	-	55	\$27	\$26
2004	2552	2753	20	-	-	-	-	-	-	-	-	52	\$30	\$27
2005	2388	2454	19	-	-	-	-	-	-	-	-	52	\$36	\$31
2006	3250	3500	29	-	-	-	-	-	-	-	-	51	\$76	\$68
2007	1682	1811	18	-	-	-	-	-	-	-	-	52	\$63	\$48
2008	1712	1828	19	-	-	-	-	-	-	-	-	52	\$22	\$37
2009	1259	1323	15	-	-	-	-	-	-	-	-	53	\$35	\$34
2010	903	951	11	759	52	25	23	4.5	55	54	50	54	\$38	\$37
2011	1473	1651	19	1314	47	28	25	3.2	59	53	42	53	\$48	\$40
2012	1293	1450	18	1108	51	24	25	3.7	59	53	45	54	\$62	\$63
2013	1146	1295	17	1040	51	24	25	3.4	55	56	42	52	\$74	\$68

¹ Includes DNR and Tribal harvests

² Estimated from population model, includes estimated non-reported harvest of 20% 1977-1992, and 10% from 1993-present.

³ Average pelt price based on a survey of in-state fur buyers only.




Figure 4. Age structure of female fishers in the 2013 harvest.




Figure 5. Age structure of male fishers in the 2013 harvest.




Figure 6. Fisher populations, harvests, and survey indices, 1977-2014. Harvests include an estimate of non-reported take.

Table 3. Marten harvest data, 1985 to 2013.

Year	DNR harvest	Modeled Harvest ¹	% Autumn Pop. Harvested ²	Carcasses Examined ³	% juveniles	% yearlings	% adults	Juv: Ad. Female ratio	% male juveniles	% male yearlings	% male adults	% males overall	Pelt price Males ⁴	Pelt price Females ⁴
1985	430	430	6	507	73	18	9	49.2	69	68	82	70	\$30	\$28
1986	798	798	9	884	64	21	15	23.1	65	71	81	69	\$36	\$27
1987	1363	1363	15	1754	66	18	16	16.7	65	67	75	67	\$43	\$39
1988	2072	2072	18	1977	66	11	23	8.8	58	50	66	59	\$50	\$43
1989	2119	2119	19	1014	68	12	20	9.9	57	63	65	59	\$48	\$47
1990	1349	1447	13	1375	48	18	34	3.6	59	54	61	59	\$44	\$41
1991	686	1000	8	716	74	9	17	13.5	69	71	72	70	\$40	\$27
1992	1602	1802	14	1661	65	18	17	14.8	63	70	75	66	\$28	\$25
1993	1438	1828	14	1396	57	20	23	7.6	61	71	67	64	\$36	\$30
1994	1527	1846	13	1452	58	15	27	6.5	62	76	67	66	\$34	\$28
1995	1500	1774	13	1393	60	18	22	8.2	63	68	66	65	\$28	\$21
1996	1625	2000	14	1372	48	22	30	4.9	62	69	67	65	\$34	\$29
1997	2261	2762	20	2238	61	13	26	6.2	60	60	63	61	\$28	\$22
1998	2299	2795	21	1577	57	18	25	6.5	62	66	65	63	\$20	\$16
1999	2423	3000	21	2013	67	12	21	9.9	65	66	67	66	\$25	\$21
2000	1629	2050	15	1598	56	25	19	8.8	62	69	66	64	\$28	\$21
2001	1940	2250	15	1895	62	15	23	10.7	65	73	74	69	\$24	\$23
2002	2839	3192	20	2451	38	30	32	3.3	59	65	62	62	\$28	\$27
2003	3214	3548	23	2391	49	16	35	4.2	59	66	68	64	\$30	\$27
2004	3241	3592	24	2776	26	28	46	1.4	54	67	59	60	\$31	\$27
2005	2653	2873	22	1992	53	16	31	5.1	64	63	65	65	\$37	\$32
2006	3788	4120	32	1914	64	17	20	9.5	67	68	67	67	\$74	\$66
2007	2221	2481	21	1355	30	29	41	1.6	60	68	54	60	\$59	\$50
2008	1823	1953	18	1095	40	21	39	2.4	62	64	57	60	\$31	\$28
2009	2073	2250	20	1252	56	15	29	5.1	67	49	63	63	\$27	\$30
2010	1842	1977	18	1202	47	25	28	4.4	71	56	62	65	\$40	\$37
2011	2525	2744	25	1615	39	25	36	2.7	64	64	60	62	\$42	\$39
2012	1472	1610	17	1260	34	30	36	2.6	67	57	64	63	\$57	\$54
2013	1014	1323	15	942	43	20	37	3.5	59	62	68	63	\$74	\$71

¹ Includes DNR and Tribal harvests

² Estimated from population model; includes estimated non-reported harvest of 40% in 1985-1987 and 1991, 20% in 1988-1990 and 1992-1998, and 10% from 1999-present.

³ Starting in 2005, the number of carcasses examined represents a random sample of ~ 70% of the carcasses collected in each year.

⁴ Average pelt price based on a survey of in-state fur buyers only

Marten Harvest Age-Classes




Figure 7. Age-class distribution of martens harvested in Minnesota, 1985 - 2013.




Figure 8. Age structure of female martens in the 2013 harvest.




Figure 8. Age structure of male martens in the 2013 harvest.




Figure 10. American marten populations, harvests, and survey indices, 1979-2014. Harvests include an estimate of non-reported take.

Table 4. Otter harvest data¹, 1984 to 2013. Carcasses were only collected from 1980-86.

Year	DNR harvest	Modeled Harvest ¹	% Autumn Pop. Harvested ²	Carcasses examined	% juveniles	% yearlings	% adults	Juv:ad. females	% male juveniles	% male yearlings	% male adults	% males overall	Pelt price Otter ³	Pelt price Beaver ³
1984	529	561	7	549	48	23	29	3.2	47	50	49	49	\$22	\$12
1985	559	572	6	572	43	23	34	2.2	53	50	43	51	\$21	\$15
1986	777	777	8	745	45	23	32	2.7	45	48	46	47	\$24	\$20
1987	1386	1484	15	-	-	-	-	-	-	-	-	52	\$23	\$17
1988	922	922	9	-	-	-	-	-	-	-	-	52	\$22	\$14
1989	1294	1294	12	-	-	-	-	-	-	-	-	52	\$22	\$12
1990	888	903	8	-	-	-	-	-	-	-	-	52	\$24	\$9
1991	855	925	8	-	-	-	-	-	-	-	-	51	\$25	\$9
1992	1368	1365	10	-	-	-	-	-	-	-	-	52	\$30	\$7
1993	1459	1368	10	-	-	-	-	-	-	-	-	52	\$43	\$10
1994	2445	2708	18	-	-	-	-	-	-	-	-	52	\$48	\$14
1995	1435	1646	12	-	-	-	-	-	-	-	-	52	\$39	\$12
1996	2219	2500	17	-	-	-	-	-	-	-	-	52	\$39	\$19
1997	2145	2313	16	-	-	-	-	-	-	-	-	52	\$40	\$17
1998	1946	2139	15	-	-	-	-	-	-	-	-	52	\$34	\$13
1999	1635	1717	12	-	-	-	-	-	-	-	-	52	\$41	\$11
2000	1578	1750	12	-	-	-	-	-	-	-	-	52	\$51	\$14
2001	2301	2531	17	-	-	-	-	-	-	-	-	57	\$46	\$13
2002	2145	2390	15	-	-	-	-	-	-	-	-	59	\$61	\$10
2003	2766	2966	19	-	-	-	-	-	-	-	-	57	\$85	\$12
2004	3450	3700	24	-	-	-	-	-	-	-	-	56	\$87	\$14
2005	2846	3018	22	-	-	-	-	-	-	-	-	58	\$89	\$15
2006	2720	2873	21	-	-	-	-	-	-	-	-	56	\$43	\$17
2007	1861	1911	15	-	-	-	-	-	-	-	-	55	\$29	\$16
2008	1938	1983	15	-	-	-	-	-	-	-	-	59	\$24	\$12
2009	1544	1578	12	-	-	-	-	-	-	-	-	59	\$36	\$13
2010	1814	1830	13	-	-	-	-	-	-	-	-	57	\$35	\$13
2011	2294	2490	17	-	-	-	-	-	-	-	-	58	\$51	\$17
2012	3171	3377	22	-	-	-	-	-	-	-	-	60	\$72	\$16
2013	2824	2993	21	-	-	-	-	-	-	-	-	48	\$61	\$17

¹ Includes DNR and Tribal harvests

² Estimated from population model. Incl. estimated non-reported harvest of 30% to 1991, 22% from 1992-2001, and 15% from 2002-present.

³ Weighted average of spring (beaver only) and fall prices based on a survey of in-state fur buyers.




Figure 11. Otter populations and harvests, 1977-2014. Harvests include an estimate of non-reported take.

