

Review Draft (August 1, 2025)

**Minnesota's Wildlife Action Plan 2025-2035:
Conserving Habitats and Biodiversity**

Acknowledgements

We would like to thank more than 250 people who contributed to the development of this State Wildlife Action Plan (SWAP) throughout our revision process over the past two years. Everyone's varied perspectives and expertise has made this a better plan and will carry on into the next ten years of conservation action for Minnesota's biodiversity and vulnerable wildlife.

Funding

The SWAP revision has been funded through U.S. Fish and Wildlife Service State Wildlife Grants as well as matching funds from private donations to the Nongame Wildlife Fund and Reinvest in Minnesota funds. We also received funding from the Environmental and Natural Resources Trust Fund as administered through the Legislative-Citizen Committee on Minnesota Resources.

NONGAME WILDLIFE PROGRAM

500 Lafayette Road

St. Paul, MN 55155-4040

888-646-6367 or 651-296-6157

DNR.gov

The Minnesota Department of Natural Resources (DNR) prohibits discrimination in its programs and services based on race, color, creed, religion, national origin, sex, marital or familial status, disability, public assistance status, age, sexual orientation, and local human rights commission activity. Individuals with a disability who need a reasonable accommodation to access or participate in DNR programs and services, including those who would like to request this document in an alternative format, should contact the DNR ADA Title II Coordinator at info.dnr@state.mn.us or 651-296-6157. We welcome calls from Telecommunications Relay Service (TRS) users. For assistance in other languages, please call 651-296-6157 or 888-MINNDNR (646-6367). Discrimination inquiries should be sent to Minnesota DNR, 500 Lafayette Road, St. Paul, MN 55155-4049; or Office of Diversity, Inclusion and Civil Rights, U.S. Department of the Interior, 1849 C. Street NW, Washington, DC 20240.

This document contains maps that are available in alternative formats to individuals with disabilities by calling (651) 259-5149 or emailing the SWAP Coordinator at alison.cariveau@state.mn.us.

Table of Contents

How to Use Minnesota’s Wildlife Action Plan 2025-2035	1
Roadmap to the Eight Elements	2
Executive Summary	3
Chapter 1. Foundation	7
Chapter 2. Species in Greatest Conservation Need	18
Amphibians	31
Birds	53
Fish	81
Mammals	99
Reptiles	123
Aquatic Invertebrates	146
Terrestrial Invertebrates	169
Plants and Lichens	192
Chapter 3. Habitats	206
Prairie and Other Grasslands	223
Savanna	260
Upland Conifer Forest and Woodlands (Fire-dependent)	273
Upland Deciduous Forests and Woodlands (Fire-dependent)	297
Mesic Hardwood Forests	320
Deciduous Wet Forest	346
Coniferous Forest Wetland	360
Riparian and Floodplain Forest	383
Non-forested Wetlands	403
Rivers and Streams	426
Lakes	450

Vernal Pools.....	475
Cliff, Talus, and Rock Outcrop	492
Caves	505
Urban and Other Developed Lands.....	520
Chapter 4. Public Engagement	549
Information Sharing	551
Environmental Education	557
Wildlife Viewing	560
Responsible Recreation	565
Habitat Stewardship	568
Participatory Science	573
Financial Support.....	577
Chapter 5. Monitoring and Research	585
Chapter 6. Implementation	613
Conservation Action Network	613
Conservation Opportunity Areas.....	626
Climate Adaptation.....	631
APPENDIX A – SGCN Designation Flowchart	660
APPENDIX B – SGCN and SNI Lists	663
Amphibians	664
Birds.....	664
Fishes.....	667
Mammals.....	668
Reptiles.....	670
Bees (Class Insecta; Order Hymenoptera).....	670
Beetles (Class Insecta; Order Coleoptera).....	678

Butterflies, Moths (Class Insecta; Order Lepidoptera)	679
Flies (Class Insecta; Order Diptera)	691
Grasshoppers, Crickets (Class Insecta; Order Orthoptera)	691
Millipedes (Class Diplopoda)	691
Snails (Class Gastropoda)	692
Spiders (Class Arachnida; Order Araneae)	693
Springtails (Class Collembola)	693
True Bugs, Cicadas, Leafhoppers (Class Insecta; Order Hemiptera)	693
Caddisflies (Class Insecta; Order Trichoptera)	694
Dragonflies and Damselflies (Class Insecta; Order Odonata)	697
Mayflies (Class Insecta; Order Ephemeroptera)	698
Mussels (Class Bivalvia)	701
Shrimp (Class Branchiopoda)	702
Stoneflies (Class Insecta; Order Plecoptera)	703
Vascular Plants	705
Lichens	723
APPENDIX C – Animal SGCN Habitat Associations	733
Amphibians	733
Birds	734
Fish	739
Mammals	740
Reptiles	742
Beetles (Class Insecta; Order Coleoptera)	748
Butterflies, Moths (Class Insecta; Order Lepidoptera)	749
Snails (Class Gastropoda)	753
Spiders (Class Arachnida; Order Araneae)	754

True Bugs, Cicadas, Leafhoppers (Class Insecta; Order Hemiptera).....	754
Caddisflies (Class Insecta; Order Trichoptera).....	754
Dragonflies and Damselflies (Class Insecta; Order Odonata).....	756
Mayflies (Class Insecta; Order Ephemeroptera).....	757
Mussels (Class Bivalvia)	758
Stoneflies (Class Insecta; Order Plecoptera)	759
APPENDIX D - Plant SGCN Associations to Primary Habitat Types	760
Prairie and Other Grasslands.....	761
Savanna	764
Upland Conifer Forest and Woodlands	764
Upland Deciduous Forest and Woodlands	765
Mesic Hardwood Forest	766
Deciduous Wet Forest	768
Coniferous Forest Wetland.....	768
Riparian and Floodplain Forest	769
Non-forested Wetlands	770
Rivers and Streams.....	771
Lakes.....	772
Cliff, Talus, and Rock Outcrop	773

How to Use Minnesota's Wildlife Action Plan 2025-2035

Minnesota's Wildlife Action Plan 2025-2035 has been designed in a modularized way with the information seeker – the viewer – in mind. The Plan is optimized for online access, based on feedback we received in a State Wildlife Action Plan (SWAP) awareness survey (2023), in which user groups and partners expressed their suggestions to make the SWAP more accessible and searchable. The Plan will eventually be presented online in an interactive website that enables visitors to find, read, and download the information they seek. Presenting information in this way makes the large volume of information contained in the Plan more accessible and able to be applied into various conservation contexts – for researchers, teachers, land managers, interpreters, private landowners, and the public. Access to Plan information will be guided online along several primary pathways:

By Topic or Chapter. If you are interested in learning about the Species in Greatest Conservation Need, refer to Chapter 2. If you are interested in Public Engagement, select Chapter 4, and so on.

By Taxonomic Group. If you are interested in mammals, mussels, or plants, you can directly select to read about a group of species. These reports will bring together information about stressors and actions for these species.

By Habitat. A landowner or land manager might like to learn about the forest or stream on their land. Information will be provided for 15 habitats, including a general description, primary stressors in those environments, associated native plant communities and SGCN, and conservation actions in that environment to benefit SGCN.

By Geography. If possible, we envision the capability for people interested in a specific region within the state, such as a county or ecoregional section, to select those geographical areas on an interactive map to gain information about the habitats and associated species in that area.

On a Map. A spatial data viewer will enable viewers to interact with several key spatial resources. The Conservation Action Network highlights the most important areas for biodiversity and connectivity. Conservation Opportunity Areas highlight places prioritized for on-the-ground partnerships. The viewer will also include layers depicting the habitats in the state, climate resources, and native plant communities.

This Draft Plan is compiled into one document for review and will be available later for those who wish to have a complete reference. The next section, *Roadmap to the Eight Elements*, explains where in the Plan each of the congressionally required elements of State Wildlife Action Plans can be found.

Roadmap to the Eight Elements

State Wildlife Action Plans are required for states to secure funding under the State and Tribal Wildlife Grants (STWG) Program, established in 2001. The plans have eight required elements, depicted in the table below along with where the material can be found within this Plan.

Element and Description	Location in this SWAP
Element 1. Identify Species of Greatest Conservation Need (SGCN) including species that have low populations or are in decline.	Chapter 2. SGCN
Element 2. Identify the habitats essential to the conservation of those SGCN.	Chapter 3. Habitat
Element 3. Identify threats to SGCN and associated habitats.	Chapters 2 and 3
Element 4. Identify the conservation actions needed to address those threats.	Chapters 2 and 3
Element 5. Describe a plan for monitoring SGCN and their habitats, and the effectiveness of conservation actions in order to adaptively manage them	Chapter 5. Monitoring
Element 6. Review and revise the SWAP at least every 10 years.	Chapter 1. Foundation
Element 7. Develop, review, and implement SWAPs in conjunction with conservation partners including Tribes and federal, state, and local agencies that manage lands.	Chapter 1. Foundation
Element 8. Seek broad public participation in the development and implementation of the SWAP.	Chapters 1 and 4. Public Engagement

Executive Summary

Minnesota's Wildlife Action Plan 2025-2035: Conserving Habitats and Biodiversity, is Minnesota's third State Wildlife Action Plan (Plan), updating information and strategies initiated in the 2005 and 2015 plans. **The overarching goal of this Plan is to guide, facilitate, and inspire conservation actions that sustain the long-term health and viability of Minnesota's biodiversity, with a focus on species that are rare, declining, or vulnerable (Species in Greatest Conservation Need) and their habitats.**

Vision: working together to conserve Minnesota's rare, declining, and vulnerable species and their habitats

State Wildlife Action Plans provide strategic guidance for conserving Species in Greatest Conservation Need (SGCN) and the habitats they depend on within each state. State Wildlife Action Plans are non-regulatory guides

for all Minnesotans that outline proactive and voluntary approaches to conserving wildlife, designed for implementation by broad arrays of partners within states. These plans reflect a shared commitment to maintaining a healthy environment that includes all animals and plants. By guiding conservation efforts before species become critically imperiled, SWAPs help prevent the need for listing animals or plants as endangered and reduce the need for costly and restrictive measures to recover species from the brink of extinction.

State Wildlife Action Plans also enable states to access State Wildlife Grant funds, the primary federal funding targeted toward the conservation of non-hunted or fished wildlife species. These funds in Minnesota cover 75% of the vertebrate animals including amphibians, birds, fish, mammals, and reptiles. When including invertebrates such as insects, spiders, and mussels, these funds cover approximately 96% of Minnesota's animal species.

Three Primary Plan Goals

This Plan includes three primary goals, each detailed within a chapter. For each goal, a set of strategies is described, with each strategy supported by a set of potential conservation actions. These primary goals are:

- 1) **Species:** Conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions.
- 2) **Habitat:** Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN.
- 3) **Public Engagement:** Inspire, motivate, and empower Minnesotans to engage in conservation actions that support SGCN and their habitats.

Species in Greatest Conservation Need

Species in Greatest Conservation Need are the animal and plant species that are vulnerable or at risk of decline or extirpation and species of stewardship or tribal significance in Minnesota.

To identify these species, we completed a large-

scale systematic conservation status assessment for the more than 7,000 species in the state for which there was sufficient information. We evaluated rarity, trends, and threats, following an internationally adopted process, to generate S-rank scores that classify species from secure to highly imperiled. These standardized evaluations then formed the foundation for revising our state list of SGCN, now in this update, totaling 1,155

SGCN are animal and plant species that are vulnerable or at risk of decline or extirpation and species of stewardship or tribal significance in Minnesota.

species. For the first time, we have included plants and lichens, which account for 447 and 151 of the SGCN species, respectively. Another 844 plant and animal species were identified as Species in Need of Information (SNI), species believed to be rare in Minnesota, but for which we do not have enough information to categorize them as SGCN.

Throughout the Plan we discuss stressors, factors that pose challenges to vulnerable plant and wildlife species, and actions to address them to direct conservation attention to where it is needed most. We profile 24 stressors following an international threats classification system, specifically tailored to species groups and habitats in the sub-chapters described below. The stressor of climate change is discussed in further detail with strategies and resources for climate adaptation within Chapter 6. Implementation. The species goal is implemented with three strategies:

1. Survey, monitoring, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.
2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnerships with agencies, non-governmental organizations, and private landowners, and
3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management practices.

Detailed sets of conservation actions are described under these strategies within sub-chapters for eight groups of species: amphibians, reptiles, birds, fishes, mammals, aquatic invertebrates, terrestrial invertebrates, and plants and lichens.

Habitats

Habitat loss and degradation are primary factors for species declines. Directing conservation at the habitat, community, and ecosystem levels is fundamental to conserving SGCN. The habitat chapter outlines five strategies to guide habitat conservation for Minnesota's biodiversity:

1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.
2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.
3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.
4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to stressors, natural disturbances, conservation actions, and climatic conditions.
5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

A set of 15 primary habitat types are described in sub-chapters that provide conservation context, an assessment of the stressors for these habitats, and actions nested under strategies to address those stressors.

Public Engagement

The Plan's third goal of public engagement emphasizes the role that all Minnesotans can play in stewardship. This chapter articulates seven strategies for inspiring, motivating, and empowering Minnesotans to practice conservation for SGCN and their habitats. The importance and rationale for each strategy is discussed, accompanied by a set of recommended actions to implement it. The strategies are:

1. Develop and distribute informational resources and media on SGCN, their habitats, and threats.
2. Support environmental education on SGCN and their habitats on a statewide level.
3. Provide and promote wildlife viewing opportunities and programs that inspire curiosity, caring, and connection with Minnesota's SGCN.
4. Foster responsible recreation practices that reduce recreational impacts to SGCN and their habitats while allowing for safe and satisfying access to the outdoors.
5. Engage and empower communities to improve wildlife habitats where they live, work, and recreate.
6. Offer participatory science programming for Minnesotans to take part in conservation volunteerism that addresses SGCN data needs.
7. Cultivate financial support for the conservation of SGCN through the development and promotion of public funding opportunities and public-private partnerships.

Following these three main goals in the Plan, two additional chapters provide more tools and approaches in support of the overall goals: Chapter 5. Monitoring and Research and Chapter 6. Implementation.

Monitoring and Research

Though discussed throughout the Plan, Chapter 5. Monitoring and Research emphasizes the critical role of surveys, monitoring, and research in generating foundational ecological data. These activities support surveillance and conservation effectiveness monitoring and help assess the impacts of stressors on Minnesota's plant and animal species, habitats, and ecological communities. The chapter defines these tools and explains their application within the framework of the State Wildlife Action Plan. It features a range of projects and programs led by the DNR and its partners, spanning diverse species, habitats, and conservation strategies. The chapter concludes with a practical guide on study design and key statistical principles essential for collecting rigorous, defensible, and repeatable scientific data. These principles are especially valuable in the context of adaptive management, where objectives and scientific understanding evolve over time through iterative learning and refinement.

Implementation

The Implementation Chapter contains information for conservation practitioners about the Conservation Action Network, Conservation Opportunity Areas, and the topic of Climate Adaptation.

The Conservation Action Network (CAN) is a mapped resource representing the best core areas in Minnesota for habitat quality, biodiversity (high numbers of SGCN), and locations of SGCN (see Figure 1.1). It also represents buffers and corridors to further protect and connect them and addresses the growing impacts of climate change by identifying areas that allow for habitat resilience and the movement of species across the landscape. The purpose of the CAN is to guide and prioritize conservation actions given limited resources. The CAN is based on the Wildlife Action Network developed in the 2015 Plan, renamed to CAN to include plant SGCN.

Conservation Opportunity Areas (COAs) are places where partnerships for on-the-ground conservation actions are concentrated to benefit SGCN and their habitats (see Figure 1.2). The locations are based on the core areas of high biodiversity, highest quality habitats, and corridors identified by the CAN. However, many COAs expand beyond those areas to incorporate locations where established partnerships are actively collaborating to deliver habitat management projects to benefit SGCN. Formerly named Conservation Focus Areas under the 2015 Plan, the new name of Conservation Opportunity Areas reflects the expanded boundaries that capitalize on partnership opportunities and increase the breadth of ecological benefits at the landscape scale.

In the Climate Adaptation section we share resources to understand trends in Minnesota's climate as well as for managers to work to facilitation adaptation for the ecosystems they manage.

To summarize, this Plan provides information, strategies, and example actions to benefit Minnesota's biodiversity, its Species in Greatest Conservation Need, and the habitats that support them. This Plan is for all Minnesotans. Achieving our vision of working together to conserve Minnesota's rare, declining, and vulnerable species and their habitats depends on strong, collaborative partnerships statewide, including private landowners; the 11 federally recognized Tribes within Minnesota's borders; local, state, and federal land managers; academic institutions; and non-governmental conservation organizations. We appreciate your interest and participation in the conservation of these valuable and vulnerable animals, plants, and habitats.

Chapter 1. Foundation

The Significance of State Wildlife Action Plans

State Wildlife Action Plans (SWAP) provide strategic guidance for conserving rare, declining, and vulnerable animal and plant species and the habitats they depend on within each state. These plans also serve a critical function by unlocking access to U.S. Fish and Wildlife Service State Wildlife Grant funds (SWG), the primary source of federal funding dedicated to the conservation of non-hunted or fished wildlife species. In Minnesota, game and sportfish receive targeted federal support through separate programs, but species with take restrictions account for only about 25% of the state's vertebrates¹. This means SWG funds are essential for the conservation of the remaining 75% of Minnesota's vertebrate species, including amphibians, birds, fish, mammals, and reptiles. When invertebrates such as insects, spiders, and mussels are included (3,789 species assessed for this Plan), the coverage of SWG funding extends to approximately 96% of Minnesota's animal species. These figures highlight the vital role that SWAPs and SWG play in supporting biodiversity conservation across Minnesota.

Across the nation, SWG funds have distributed over 1.4 billion dollars to 56 states, territories, commonwealths and the District of Columbia since 2000. Minnesota has received nearly \$26 million from 2002 through 2024. These critical funds have supported wildlife conservation projects dedicated to conserving rare and declining species across the state. Examples include surveys of rare salamanders and fish, restorations of native prairies and oak savannas, and propagating rare mussels for reintroduction.

State Wildlife Action Plans are non-regulatory guides that outline proactive and voluntary approaches to conserving wildlife, designed for implementation by broad arrays of partners within states. These plans reflect a shared commitment to maintaining a healthy environment that includes all animals and plants. By guiding conservation efforts before species become critically imperiled, SWAPs help prevent the need for listing animals or plants as endangered and reduce reliance on costly, reactive, and restrictive measures to recover species from the brink of extinction.

State Wildlife Action Plans focus on Species in Greatest Conservation Need (SGCN); their habitats; the threats they face; and the conservation actions needed to support their survival, including monitoring and research. These elements are core requirements of each Plan (see Roadmap) and among them, habitat conservation warrants particular emphasis. As

Species in Greatest Conservation Need (SGCN) are animal and plant species that are vulnerable or at risk of decline or extirpation and species of stewardship or tribal significance in Minnesota.

¹ An additional 18% of Minnesota's vertebrate species are unregulated game or fish species. There is also some overlap between game species and SGCN.

our knowledge of species ecology and conservation challenges deepens and as the SGCN list continues to grow, it becomes increasingly clear that species-by-species conservation efforts are often impractical. In many cases the most effective conservation strategies will occur at the ecosystem level, where actions can benefit multiple species simultaneously.

The conservation of Minnesota's SGCN depends on strong, collaborative partnerships across the state, including private landowners; the 11 federally recognized Tribes within Minnesota's borders; local, state, and federal land managers; academic institutions; and non-governmental conservation organizations. This Plan serves as a resource to inform and support the collective efforts of this diverse community of people committed to safeguarding Minnesota's SGCN and their habitats.

Goals of Minnesota's Wildlife Action Plan 2025-2035

Vision: Working together to conserve Minnesota's rare, declining, and vulnerable species and their habitats.

Overarching Goal of the SWAP: To guide, facilitate, and inspire conservation actions that sustain the long-term health and viability of Minnesota's biodiversity, with a focus on species that are rare, declining, or vulnerable (Species in Greatest Conservation Need, SGCN) and their habitats.

Three Primary Goals

1. **Species:** Conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions. (see Chapter 2. Species in Greatest Conservation Need)
2. **Habitat:** Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN. (see Chapter 3. Habitats)
3. **Public Engagement:** Inspire, motivate, and empower Minnesotans to engage in conservation actions that support SGCN and their habitats. (see Chapter 4. Public Engagement)

Building on the Strengths of the 2005 and 2015 Plans

Minnesota published two previous ten-year State Wildlife Action Plans; this document represents the third revision and is consistent with the earlier versions. Each of the earlier plans provided valuable guidance that continues to inform conservation efforts today. This updated Plan builds upon those foundations by incorporating current approaches, new information, and refined strategies. The key strengths of the previous plans are briefly summarized below.

2005-2015 Plan: Tomorrow's Habitat for the Wild and Rare

SGCN Identification and Distribution

Minnesota's initial State Wildlife Action Plan, [*Tomorrow's Habitat for the Wild and Rare*](#) (DNR, 2005) established criteria to objectively identify Species in Greatest Conservation Need. A total of 292 SGCN were identified, including 189 vertebrates and 103 invertebrates.

Habitat Focus

The 2005 Plan identified that the primary threat to SGCN was the significant loss and degradation of habitat. It emphasized that effective conservation depends on the ability to identify and protect the habitats these species require. The Plan identified 16 key habitats and associated SGCN with key habitats.

Ecoregional Profiles

To provide greater specificity, the Plan also included summaries for each of Minnesota's 25 ecoregional subsections, as defined by the state's Ecological Classification System. Each subsection profile featured maps illustrating the distribution of key habitats and SGCN richness by township, summaries of the key habitats used by SGCN and which habitats are most important to the greatest number of SGCN. In addition, ten-year goals, management challenges, strategies, and conservation actions were outlined for each subsection.

Minnesota's Wildlife Action Plan 2015-2025

Species in Greatest Conservation Need Update

For Minnesota's second Plan, [*Minnesota's Wildlife Action Plan 2015-2025*](#) (DNR, 2015), technical advisory teams conducted a comprehensive review and update of the 2005 SGCN list, adding 111 species and removing 59 species, resulting in a revised total of 346 SGCN. As part of the review, stressors affecting each species were assessed to determine which SGCN would benefit from a habitat-based conservation approach and which required more targeted actions.

Wildlife Action Network

For species that would most benefit from a habitat-based approach, the 2015 Plan introduced the Wildlife Action Network (WAN), a map to illustrate the distribution of high-quality habitats for both terrestrial and aquatic SGCN. To prioritize the implementation of on-the-ground conservation efforts, sites within the WAN were assigned priority rankings based on five metrics, including SGCN population viability and species richness.

Conservation Focus Areas

Acknowledging the limited fiscal and human resources available to implement conservation objectives, a prioritization system was developed to guide habitat conservation efforts more effectively. The WAN rankings helped identify 37 Conservation Focus Areas (CFAs)—specific locations where resources such as partnerships and funding were available—to support habitat restoration and management efforts across organizational and jurisdictional boundaries. CFAs serve as a valuable framework for prioritizing on-the-ground actions that directly benefit SGCN and their habitats. The CFA approach has proven to be an effective strategy for fostering partnerships and leveraging shared resources to address the habitat needs of SGCN across Minnesota.

Strategies, Stressors, and Actions

Strategies for implementation of the Plan are described within each of the chapters that address the three primary goals (Chapter 2. Species, Chapter 3. Habitats, and Chapter 4. Public Engagement).

Stressors, or threats, are factors that create conservation challenges for habitats or species. These are first explained in Chapter 2. Species, then specifically elaborated within each of 15 habitat sub-chapters and 8 species group sub-chapters. They are depicted with a set of icons that are used to connect to Conservation Actions.

Conservation Actions are identified to address stressors, fill information gaps, or describe engagement opportunities. These are organized by Strategies and tagged with icons to relate them to Stressors or to Chapter 4. Engagement or Chapter 5. Monitoring and Research.

Monitoring and Research

Chapter 5. Monitoring and Research emphasizes the critical role of surveys, monitoring, and research in generating foundational ecological data. These activities support surveillance and conservation effectiveness monitoring and help assess the impacts of stressors on Minnesota's plant and animal species, habitats, and ecological communities. The chapter defines these tools and explains their application within the framework of the State Wildlife Action Plan. It features a range of projects and programs led by the DNR and its partners, spanning diverse species, habitats, and conservation strategies. The chapter concludes with a practical guide on study design and key statistical principles essential for collecting rigorous, defensible, and repeatable scientific data. These principles are especially valuable in the context of adaptive management, where objectives and scientific understanding evolve over time through iterative learning and refinement.

Implementation

Chapter 6. Implementation explains in detail three elements that assist in our delivery of the Plan in Minnesota: the Conservation Action Network, Conservation Opportunity Areas, and Climate Adaptation.

Conservation Action Network (CAN)

The 2025 Conservation Action Network (CAN) is built from a set of geographic information system (GIS) layers representing the best core areas in Minnesota for habitat quality, biodiversity, and locations of SGCN. It is a mapped resource representing important areas to the SWAP, including core areas of high biodiversity (high numbers of SGCN), highest quality habitats, and buffers and corridors to further protect and connect them (see Figure 1.1). The purpose of the CAN is to guide and prioritize conservation actions given limited resources. It also addresses the growing impacts of climate change by identifying areas that allow for essential ecosystem processes habitat resilience, and the movement of species across the landscape.

Then, a connectivity analysis was conducted to identify the best paths of connection between core areas represented in the CAN. Finally, the network was scored using a subset of layers to prioritize where multiple layer features overlap resulting in the CAN map (Figure 1.1).




Figure 1.1 The Conservation Action Network built from a series of data layers including richness of SGCN and areas of outstanding biological significance and places with strong landscape connectivity. A higher CAN score represents areas where more of the GIS layers overlap and indicates highest priority for conservation actions.

Conservation Opportunity Areas (COAs)

Conservation Opportunity Areas (COAs) are based on the core areas of high biodiversity (high numbers of SGCN), highest quality habitats, and buffers and corridors identified by the Conservation Action Network (CAN; see Figure 1.2). However, many of the COAs expand beyond those areas identified in the CAN to incorporate areas where established partnerships are actively collaborating to deliver habitat management projects that have the potential to benefit SGCN. These partnership efforts present unique opportunities to capitalize on established working relationships and shared conservation goals.

COAs are not meant to represent all of the important areas in the CAN for SGCN. The goal is to identify the highest priority areas for conservation work to occur within the larger partnership framework that exists within a COA. This allows our COA Coordinators to focus their efforts and capitalize on existing partnerships that are already addressing similar habitat concerns.

Potential projects within the COAs will be prioritized based on their location related to the priority areas designated by the CAN. Focusing habitat management, restoration, and protection within the CAN will enhance core areas of habitat and connections between habitats. This focus will also allow land managers/owners to incorporate existing partnerships while focusing their efforts on priority areas. This will increase efficiency by facilitating the coordination of management, prioritizing outreach efforts, and targeting technical assistance.

Conservation Opportunity Areas may contain multiple partnership groups that are actively collaborating on project implementation. Each of these groups may have a slightly different approach to their efforts and/or a slightly different set of habitat types in which they work. Rather than having a set of standards that each COA must meet to be considered active, the 2025 Minnesota Wildlife Action Plan will consider a COA active if there is at least one partnership group present, regularly meeting with the intent of implementing habitat improvement projects.

It is hoped that the COA approach will increase the number of active partnership groups that the COA Coordinators can engage with as well as the coordination of those partnership groups across a broader geographic area.




Figure 1.2. Map of Conservation Opportunity Areas (COAs) for 2025-2035, developed with input from internal and external partners to make revisions to the 2015-2025 Conservation Focus Areas. The number of COA's grew from 37 to 39. Two additional COAs proposed in north central Minnesota require further consultation with Tribes for delineation.

Climate Adaptation

The Climate Adaptation section presents scientific data, concepts, and strategies to support effective conservation in the face of a changing climate. While climate change was prominently featured in the 2015 Plan, current trends in temperature and precipitation continue to intensify, as clearly illustrated through updated charts and statistics. The impacts of rising temperatures and shifts in rainfall patterns, including increased flooding, shorter winters, and altered growing seasons, are being felt across all ecosystems. The climate-related stressors are addressed throughout the habitat sub-chapters, as they influence nearly every aspect of ecological function. Climate stress also interacts heavily with many other stressors, such as the spread of invasive species, fire regime alterations, and the emergence of diseases and pathogens. In this section of Chapter 6. Implementation, we provide background information, actionable strategies, and illustrate case studies to help conservation practitioners better understand and incorporate climate adaptation into their work.

Public and Partner Input to the Revision

A key emphasis of this revision of the State Wildlife Action Plan (the Plan) has been broad engagement across a diverse array of people to help shape its content and enhance its relevance for on-the-ground conservation. The network of individuals and organizations who contributed to the development of the Plan will also play a vital role in sharing its vision and connecting others to implement its conservation actions.

We began the revision of the Plan in 2022 with a survey of individuals who use the SWAP. We asked what they valued most about previous Plans and what they hoped to see most in the new revision. Of the 136 responses received, 89 came from DNR staff and 47 from a broad range of conservation partners, including conservation organizations, federally recognized Tribes, federal agencies, local governments, and academic institutions. The feedback showed overwhelming support for the SGCN list, which continues to serve as a cornerstone for conservation planning and implementation statewide. Respondents also emphasized the lasting value of the habitat-based information introduced in the 2005 Plan and interest in refreshing that information. From the 2015 Plan, they highlighted the following elements as especially helpful: the Wildlife Action Network, the identification of Conservation Focus Areas, and the Plan's clear orientation toward action. A new request for this revision was the development of an interactive, online version of the Plan to allow users to easily access and tailor information to their specific needs.

In 2023, we invited a broad coalition of partners including Tribal representatives, conservation organizations, government agencies, academics, educators, and staff from all DNR divisions to participate in the Plan revision process. This collaboration took shape through a series of seven in-person workshops held within each region of the state. Prior to the workshops, a questionnaire was distributed to gather specific input from potential participants. Through these workshops, a total of 161 participants, representing more than 25 organizations, contributed their perspective and expertise to help shape foundational content on the most critical stressors and actions to be highlighted in this updated Plan. During the workshops, participants shared insights about the stressors affecting vulnerable species and ecosystems, identified conservation actions to address those threats, and discussed barriers to effective implementation. They also worked together to propose solutions for overcoming those challenges, helping ensure that this Plan reflects both on-the-ground realities and shared conservation goals.

Following the workshops, we assembled volunteer teams to assist in developing of content for the revised Plan. More than 210 individuals participated across 11 teams, contributing their expertise to identify Primary Stressors and Priority Conservation Actions for key habitat types and taxonomic groups of plants and animals. Contributions were made during online working meetings. In addition, many of these volunteers also assisted in drafting chapters, collaborating through shared online documents to ensure the Plan reflects a broad base of knowledge and practical experience.

We also engaged more than 50 technical species experts to support the conservation status assessment (S-rank) process for all regularly occurring animal and plant species in Minnesota with sufficient data. Relevant datasets were compiled, species were evaluated, and S-rank scores were assigned to inform the identification of SGCN (see Chapter 2. Species in Greatest Conservation Need) and Species in Need of Information (SNI; see *New for 2025* section below). The draft lists of SGCN and SNI were subsequently reviewed by species experts, volunteer team members, and DNR staff, all of whom were invited to provide feedback and comments.

To gather additional input for the development of Chapter 5. Public Engagement, including strategies to meaningfully engage Minnesotans in the stewardship of the state’s rich natural resources (see *New for 2025* section below), two final workshops were held in December 2024 and January 2025. These events brought together a total of 90 participants from 29 organizations, including strong representation of environmental educators and conservation outreach professionals, to identify barriers to reaching Minnesota’s increasingly diverse population and to explore strategies for more effective engagement with local partners and communities. Follow-up meetings were held with a smaller working group, during which participants collaborated to craft the public engagement strategies and actions presented in this Plan.

To ensure input from Tribal Nations, we maintained regular communication with representatives from Tribal natural resource departments, meeting with them in some cases, and inviting their review of various sections of the Plan, including the draft list of draft SGCN.

The final stage of the public review involved posting the full draft Plan online during July and August 2025, during which comments were collected through an online submission form.

New for 2025

For the 2025-2035 Plan we have added several main new elements.

Conservation Status Assessments

We employed a systematic, nationally standardized approach to assess the conservation needs of over 7,000 species of animals and plants in Minnesota, often referred to as S-ranks (see Chapter 2. SGCN). This method offers a repeatable approach for future reassessments and provides consistent, comparable information to support conservation at regional, national, and international conservation scales.

Plants and Lichens

For the first time, the list of SGCN has been expanded to include plants and lichens (see Chapter 2. SGCN). This reflects a growing recognition of the need for closer integration between plant and animal conservation efforts, as encouraged by national initiatives such as the Restoring America’s Wildlife Act, which also

recognized the importance of plants. While plants and fungi are not currently eligible for SWG funding, their inclusion in this Plan lays the groundwork for a more comprehensive biodiversity conservation strategy. Plants are foundational to ecosystems and form the structure of habitats, serve as essential resources for specialist animal species, and provide critical ecosystem services that support life across the landscape.

Species in Need of Information

Through the systematic review of all Minnesota plant and animal species with available data, it became evident that an additional classification of species was needed. Species in Need of Information (SNI) are those for which insufficient data currently exist to determine whether they warrant designation as SGCN (see Chapter 2. SGCN). This new category highlights the need for targeted survey and monitoring efforts to fill critical information gaps.

Species of Tribal Significance

We introduced new criteria for identifying SGCN to include species of cultural significance to Tribes in Minnesota, recognizing the deep and enduring connections that Tribal Nations have with the plants and animals with whom we share the land. This addition allows Tribes to nominate species they consider vulnerable, even if those species do not meet the criteria for SGCN designation. As a result, wild rice (*Zizania palustris*) and snowshoe hare (*Lepus americanus*) were added to the list. Wild rice—*manoomin* in Ojibwe and *psin* in Dakota—is a culturally and spiritually important food traditionally harvested by Indigenous peoples, and it is vulnerable to many disturbances in the lakes where it grows.

Public Engagement

In this Plan revision, the topic of public engagement was significantly expanded into a full chapter including seven strategies and many recommended actions. Chapter 5. Public Engagement highlights the vital role people play in conservation. It reflects the DNR's Nongame Wildlife Program commitment to better understanding and connecting with Minnesotans and our partners in conservation science, outreach, and education. This chapter emphasizes the importance of sharing information broadly and expanding the community of individuals and organizations actively working to conserve SGCN and their habitats.

Conservation Action Network

Although not new to this Plan, the Conservation Action Network (CAN), formerly known as the Wildlife Action Network (or WAN), has been renamed to better reflect our expanded commitment to biodiversity conservation, including plant species as SGCN. The content of the CAN also has been updated with current data and new connectivity analyses. For complete information about the CAN, see Chapter 6. Implementation.

Conservation Opportunity Areas (COAs)

As with the Conservation Action Network, this component is not new in the 2025 Plan but has been renamed from Conservation Focus Areas to Conservation Opportunity Areas (COAs; see Figure 1.2). Updated and expanded boundaries to COAs were informed by extensive input gathered during in-person regional workshops, ten targeted mapping meetings with local partner groups, DNR staff, and external regional partners across the state. These discussions brought up the need to expand existing boundaries to capitalize

on current partnership opportunities and to increase ecological connectivity. The new name, Conservation Opportunity Areas, better reflects these expanded boundaries and the broader, landscape-scale conservation opportunities they represent. It also aligns more closely with where partners are located, thereby facilitating collaboration and providing connectivity among core areas of the CAN. For more information refer to Chapter 6. Implementation.

Chapter 2. Species in Greatest Conservation Need

Overview

This Chapter gives an overview of Species in Greatest Conservation Need (SGCN), animal and plant species that are vulnerable or at risk of decline or extirpation, or species of stewardship or Tribal significance in Minnesota. We also introduce Species in Need of Information (SNI), species believed to be rare in

Minnesota, but for which we do not have enough information to categorize them as SGCN. Also new this year is the inclusion of plants and lichen to the SGCN and SNI list.

Species in Greatest Conservation Need (SGCN) are animal and plant species that are vulnerable or at risk of decline or extirpation and species of stewardship or tribal significance in Minnesota.

SGCN include species that are very imperiled, such as those that are facing extirpation or even extinction. All species listed as endangered, threatened, or special concern by the state or federally are designated as SGCN. Per Minnesota Statute 84.0895, a species is designated as endangered if it is threatened with extinction throughout all or a significant portion of its range, threatened if the species is likely to become endangered within the foreseeable future throughout all or a significant portion of its range, and species of special concern if it is extremely uncommon in the state or has unique habitat requirements. Legal protections exist for SGCN that are designated as endangered or threatened species, with the goal of preventing the extinction of species. Many of these species also have species-specific plans directing conservation efforts for their recovery. The DNR endangered species program is responsible for administering a permitting program to ensure that activities that may cause take of state endangered or threatened species are not detrimental to the species. The DNR cooperates with USFWS on the protection and conservation of federally listed species.

In addition to these imperiled species, SGCN include species that are more common. An important function of the SWAP is to identify early conservation actions for at-risk species that can reduce their need for listing as endangered or threatened species and improve the probability of their recovery to sustainable levels. Designating more abundant species as SGCN brings attention to declines or high levels of threats so that conservation measures may be enacted before they become so rare that recovery efforts are more costly and less likely to succeed.

Topics in this chapter include the Goal, three Strategies, and how these relate to conservation actions. We also touch on the evolution of SGCN across the three Plan revisions and the current definition of SGCN. We explain the conservation status assessment (S-rank process) underlying this Plan's list of SGCN and explain our approach to discussing stressors (threats) relating to species and their habitats. We provide a summary table of the 2025 list of SGCN and SNI sorted by taxonomic groups. This is then followed by a set of 8 sub-chapters for amphibians, birds, fishes, mammals, reptiles, aquatic invertebrates, terrestrial invertebrates, and plants (with lichens).

SWAP Species Goal: Conserve rare, declining, and vulnerable wildlife and plant Species in Greatest Conservation Need (SGCN) through targeted actions.

Species Strategies

1. **Survey, monitoring, and research** to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.
2. **Collaborate to deliver conservation actions** that support resilient populations of SGCN and their habitats in partnerships with agencies, non-governmental organizations, and private landowners.
3. **Develop and disseminate informational material** to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management practices.



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.

The Plan is an informational resource that provides scientific findings that can be applied to conservation actions. Surveys, monitoring, and research are essential for knowing which animals and plants are vulnerable. Information collected may focus on rarity, answering the question of how extensively the species is found within the state. Monitoring is employed to determine if there are changes in the species' abundance or distribution, often requiring longer term efforts over broader areas, to determine if a trend can be detected. Research into the threats that cause SGCN populations to decline as well as the effectiveness of conservation practices intended to help those species are also critical. Well-designed surveys, monitoring, and research projects help us to learn about species' specific habitat requirements, and conditions of that habitat where the species flourish. For more information on this strategy, please also see *Chapter 5. Monitoring and Research*.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnerships with agencies, Tribes, non-governmental organizations, and private landowners.

Collaborating to implement conservation actions is an essential strategy for SGCN conservation. The number of SGCN is so great, and the level of need so high, that any one entity will make little impact on their own. As documented in the stressor sections, the extent to which our environments are impacted by human activities, both past and present, requires special attention to protecting, restoring, enhancing, and maintaining the habitats we have left, and optimizing their conditions to support these species in need.

While the DNR is the lead facilitator of efforts guided by this Wildlife Action Plan, implementation of actions is undertaken every year by numerous entities throughout the state. The role of the DNR Nongame Wildlife

Program is to facilitate partnerships with a broad array of entities so that we are working collaboratively. Collective actions when coordinated across entities such as municipalities, soil and watershed conservation districts, private landowners, Tribes, other state agencies, national agencies, educational institutions, and non-profit organizations will have the most impact. Most of work we do is in partnerships.

Conservation Opportunity Areas (COAs) are places where on-the-ground conservation actions are concentrated to benefit SGCN and their habitats. COAs build on the Conservation Action Network, which depicts areas of high biodiversity and high habitat quality, linking them with buffers and corridors. The COAs are delineated to highlight areas specifically for partnerships to focus efforts together on habitat conservation projects designed to benefit SGCN. For more information, please see the Conservation Opportunity Areas section in Chapter 6. Implementation.

While actions that are specifically related to developing partnerships will be listed under Strategy 2, collaboration as an approach can apply to all actions in the Plan, as all conservation actions would benefit from broader implementation by partners.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management practices.

This Plan is meant to help put science into action. We identify threats and outline conservation actions to take, and we share case studies of successful conservation programs and links to existing resources. However, there is still a great demand for information to guide conservation efforts by conservation practitioners throughout the state. In some cases, research or investigations are needed to develop the management guidelines, avoidance measures, or habitat best management practices (see *Chapter 5. Monitoring and Research*). In other cases, there may be good content coming out of the research community or becoming known through applied conservation practice that requires translation into easily applied materials and resources. There is a huge opportunity to investigate, collate, and write up recovery and management guidance for more SGCN.

There are also opportunities to build on resources for species accounts, such as DNR's [Rare Species Guide](#) that covers state and federally listed species. Similar information, recovery or action plans, could be developed for additional SGCN, such as the Wood Turtle Plan, which is a great resource but species-specific plans like this only exist for few species in our state.

This strategy also relates to communications that help more people know about these informational resources. Workshops, field trips, webpages, webinars, and one-pagers can be used to spread the word when new research, threats assessments or best management practices become available. This strategy is analogous to Strategy 5 in the *Chapter 3. Habitats* and Strategy 1 detailed in *Chapter 4. Public Engagement*.

Conservation Actions

Strategies are implemented by conservation actions, which are described in the SGCN sub-chapters. Conservation actions are also described in the habitat sub-chapters, so referring to these sections in tandem is recommended; for example, looking at the lakes and river chapters when reading about fish will provide more comprehensive information. Conservation actions are related to stressors through a set of icons, as depicted in the stressor section below.

Building on the two Prior Plans

The definition of Species in Greatest Conservation Need has remained consistent across Minnesota's SWAPs, with minor additions and updates in selection processes over time.

2005-2015 Tomorrow's Habitat for the Wild and Rare

Minnesota's original State Wildlife Action Plan, *Tomorrow's Habitat for the Wild and Rare*, defined Species in Greatest Conservation Need as "animals whose populations are rare, declining, or vulnerable to decline and are below levels desirable to ensure their long-term health and stability." The criteria for SGCN included species whose populations were rare, declining, or vulnerable to decline, species at risk because of dependence on rare, declining, or vulnerable habitats, species subject to other threats that make them vulnerable, species with certain characteristics that make them vulnerable, or species whose Minnesota populations are stable but are declining in a substantial part of their range outside of Minnesota. In addition, federally or state-listed species, species with global conservation status of G1-G3, species of concern by regional lists such as the USFWS Region 3 Species of Concern, were also included. Vagrant and extirpated species not expected to return within 10 years were excluded. The selection process identified 292 SGCN, including 189 vertebrates and 103 invertebrates (DNR 2005). These were then related to 16 key habitats as well as to 24 ecoregional subsections.

Minnesota's Wildlife Action Plan 2015-2025

In *Minnesota's Wildlife Action Plan 2015-2025*, technical advisory teams updated the 2005 list, adding 111 species and removing 59, for a new total of 346 SGCN (DNR 2015). This included 185 vertebrate and 161 invertebrate species. Unique life history traits were identified along with the primary threats, or stressors, to the long-term sustainability of each species (Appendix C of 2015-2025 Plan). In addition, the distribution and abundance of "viable/persistent" SGCN populations were mapped for 156 SGCN with sufficient data and used to develop the Wildlife Action Network.

Species in Greatest Conservation Need (SGCN) Definition

The definition of Species in Greatest Conservation Need (SGCN) is consistent with the prior versions of Minnesota's Wildlife Action Plan, this year with an expansion to include plants, lichen and Tribally nominated species of cultural significance. ***SGCN are animal and plant species¹ that are vulnerable or at risk of decline or extirpation, or species of stewardship or Tribal significance in Minnesota.*** This set of species provides focus to the entirety of the State Wildlife Action Plan. SGCN are native species whose populations are rare, declining, or vulnerable to decline and are below levels desirable to ensure their long-term health and stability in our state. Also included are species for which Minnesota has stewardship responsibility - those for which populations in Minnesota represent a significant portion of their North American population, or species for which Minnesota populations are stable while populations outside of Minnesota have declined or are declining in a substantial part of their range.

In this Plan revision, we also added the eligibility of species of Tribal significance to be SGCN. Including tribal significance as a factor for designating SGCN status enables Tribes to nominate species they believe to be vulnerable even if those species would not otherwise be selected via the other SGCN selection factors. In 2025, this resulted in the addition of the snowshoe hare (*Lepus americanus*) and wild rice (*Zizania palustris*). Wild rice, *manoomin* (in Ojibwe) or *psin* (in Dakota), is an important food that is traditionally harvested and vulnerable to many disturbances in the lakes in which it is found. It should be noted that other tribally important species such as moose were already included in the SGCN list.

SGCN include:

- Species on Minnesota's list of endangered, threatened, or special concern species and species on the federal Endangered Species list as Endangered, Threatened, candidate, or recently de-listed species found in Minnesota.
- Species with that are Rare, Declining or Vulnerable to Threats, with species conservation status of critically imperiled (S1), imperiled (S2), or vulnerable (S3) with additional factors for concern
- Stewardship Species - Species for which populations in Minnesota represent a significant portion of their North American population or whose Minnesota populations are stable, while populations outside of Minnesota have declined or are declining in a substantial part of their range.
- Tribally Significant Species - Species of cultural significance to Tribes in Minnesota which are believed to be vulnerable to decline due to other factors in the SGCN process.

Species in Need of Information (SNI) Definition

New for the 2025 Plan, we have added another category known as Species in Need of Information (SNI). ***SNI are species believed to be rare in Minnesota, but for which we do not have enough information to categorize them as SGCN.*** Species were designated as SNI if they are thought to be a potential future SGCN, and there is a desire to conduct surveys or monitoring for them within the next ten years. A designation of SNI enables State Wildlife Grant funds to be allocated toward survey or monitoring for those species.

SGCN and SNI List Development

Updating the SGCN list for the 2025-2035 SWAP included a few steps. First, we implemented a standardized, repeatable, comprehensive conservation status assessment for all regularly occurring species in our state for which we had sufficient data (see Conservation Status Assessment section below). Second, we established criteria for the SGCN list based on the S-ranks as well as additional factors, including plants and lichens in the SGCN list process for the first time. Third, we defined a list of Species in Need of Information (SNI).

Conservation Status Assessments

As a foundation to updating the SGCN list for this Plan, we conducted conservation status assessments for all native, regularly occurring (non-vagrant) species of animals and plants in Minnesota for which there was sufficient information, more than 7,000 species. We followed the [NatureServe Conservation Status Assessment](#) process, which is a rigorous and repeatable method adopted widely by states and provinces throughout North America (Faber-Langendoen et al. 2012, Master et al. 2012). The assessments generate conservation status ranks from 1 (critically imperiled) to 5 (secure) (see inset for definitions of the ranks). Assessments are done at global (G-ranks), national (N-ranks), and sub-national levels (S-ranks). We calculated S-ranks for the species in Minnesota.

The conservation status assessment process incorporated eight status factors in three categories, highlighting three main ways in which species may become vulnerable:

- **Rarity** assesses the distribution of the species in the state, tabulating the number of known occurrences. It is based on range extent, area of occupancy (the subset of that range containing the habitat that is occupied), population size (when known), and number of occurrences. Range extent and area of occupancy represent the idea that when occurrences are spread widely they are less vulnerable to threats such as from development, fire, or climate. Occurrences are detections or records grouped together so that near detections comprise an occurrence, analogous to a population. Although infrequent, the number of occurrences with good viability was incorporated when known.
- **Trends** in population size, abundance, occurrences, or area of occupancy are considered, both long-term and short-term. Long-term is considered from the time of European colonization or around 200 years, and short-term is considered 10 years or 3 generations, whichever is longer (up to a maximum of 100 years). Trends are important for assessing species status, but they are known only for few species. When state level trends are available those are used. If those are not available, regional trends are applied; followed by continental trends only if more local trends are unavailable. There is always uncertainty in whether trends at a continental or regional level are equivalent to trends within Minnesota. When trend in population or abundance is not available, we examined changes in area of occupancy as a surrogate.
- **Threats** were assessed for direct and indirect impacts to species. We considered 10 threats from the IUCN threats classification system (level-one threats), such as development or climate change, which we describe in detail in the Stressors section below (of IUCN's 11 level-one threats, we omitted 'geological events'). Each threat was assessed by taxonomic expert panels for scope and severity. Scope describes how extensive the overlap of the threat with the species' occurrences, such as what proportion of the population is subjected to a particular threat. For this we found relevant spatial data layers, when available, and provided information to reviewers about the proportion of range that

intercepted those mapped threat layers. Severity refers to the degree to which the threat damages the species (such as what proportion of the exposed population will be killed).

Conservation Status Ranks

Conservation Status Ranks indicate the risk of extinction for species ([NatureServe Conservation Status Assessment](#)). S-ranks consider species status within the state of Minnesota. Each species received one of the following ranks:

- SX – Presumed Extirpated - Species or ecosystem is believed to be extirpated from the jurisdiction (i.e., nation, or state/province). Not located despite intensive searches of historical sites and other appropriate habitat, and virtually no likelihood that it will be rediscovered.
- SH – Possibly Extirpated - Known from only historical records but still some hope of rediscovery. There is evidence that the species or ecosystem may no longer be present in the jurisdiction, but not enough to state this with certainty. Examples of such evidence include (1) that a species has not been documented in approximately 20-40 years despite some searching and/or some evidence of significant habitat loss or degradation; (2) that a species or ecosystem has been searched for unsuccessfully, but not thoroughly enough to presume that it is no longer present in the jurisdiction.
- S1 – Critically Imperiled - At very high risk of extirpation in the jurisdiction due to very restricted range, very few populations or occurrences, very steep declines, severe threats, or other factors
- S2 – Imperiled - — At high risk of extirpation in the jurisdiction due to restricted range, few populations or occurrences, steep declines, severe threats, or other factors.
- S3 – Vulnerable - — At moderate risk of extirpation in the jurisdiction due to a fairly restricted range, relatively few populations or occurrences, recent and widespread declines, threats, or other factors.
- S4 – Apparently Secure - At a fairly low risk of extirpation in the jurisdiction due to an extensive range and/or many populations or occurrences, but with possible cause for some concern as a result of local recent declines, threats, or other factors.
- S5 – Secure - At very low or no risk of extirpation in the jurisdiction due to a very extensive range, abundant populations or occurrences, with little to no concern from declines or threats.
- SU – Unrankable (for species that we have compiled data and considered, but there is inadequate information available to rank)
- SNA – Not Applicable (for species that are not an appropriate target of conservation efforts, such as non-native species or vagrants)
- SNR – Species Not Ranked (for species we know have occurred in MN but we have not considered in this effort due to lack of information, e.g., all of our species of ants)
- S#M- Migratory species that regularly occur in Minnesota, but do not breed here are ranked, but will be given a modifier (e.g., S2M).

Prior to calculating S-ranks, we requested and incorporated species observation data from partners, including academic institutions, other governmental entities, non-profit organizations, tribes, and community science driven databases. These datasets were compiled for use in assessing the conservation status factors. Using the NatureServe S-rank Calculator, we calculated preliminary S-ranks based on rarity, trends, and threats. These preliminary ranks were reviewed by taxonomic experts to assess any data deficiencies, discrepancies, or gaps, and add additional information particularly regarding threats as based on expert knowledge. S-ranks were then

revised and reviewed once again after those updates were completed. Once the S-rank was established species were then assessed for SGCN and SNI designation. Information compiled for status assessments and SGCN identification will also be used in the state's endangered and threatened species list revision process.

SGCN and SNI Selection

We developed a flowchart (see Appendix A) to determine whether species meet criteria for inclusion on our SGCN list, summarized as:

- SGCN include species ranked in Minnesota as Critically Imperiled (S1) or Imperiled (S2) in our conservation status assessments, species ranked globally as G1, G2, or G3 (Vulnerable), those federally listed as Endangered or Threatened under the Endangered Species Act (or recently de-listed or candidate species on the five-year work plan for listing), and those state listed in Minnesota as Threatened, Endangered, or Special Concern in 2013 and thought to remain on the future state list.
- In addition, other species (particularly those that ranked as Vulnerable, S3) were considered for inclusion on the SGCN list, particularly if they met some of these additional factors:
- On the 2015 SGCN list or meet the 2015 criteria including for stewardship species
- On the 2022 [Midwest Regional SGCN \(RSGCN\) list](#), which are species classified as SGCN by at least one Midwest Association of Fish and Wildlife Agencies (MAFWA) state and meet regional stewardship criteria (% distribution in MAFWA states, core population in MAFWA states, etc.) and conservation concern criteria (federal listing, state listing, imperiled state and global NatureServe conservation status ranks, etc.)
- On the Superior or Chippewa National Forest [Regional Forester Sensitive Species \(RFSS\) list](#).
- Tribally significant species (based on expert knowledge and recommendations by Tribes in Minnesota)

Draft lists of SGCN and SNI were sent out widely for review, within DNR and to experts who were invited to submit S-rank comments and participants on the plant or animal SGCN volunteer team assembled from regional workshops. If the state's list of threatened and endangered species changes during the term of the 2025 SGCN List, we will update SGCN materials to reflect those updates accordingly.

2025 SGCN List Summary

The 2025-2035 SGCN list contains 1,155 species and 844 SNI species. Plants comprise 447 of the SGCN, followed by lichens (151 species), butterflies and moths (122), and bees (117). Plants and lichen together comprise 52% of the SGCN list. Vertebrate species (amphibians, birds, fishes, mammals, and reptiles) total 167 or 15% of the list. Refer to Table 2.1 for a summary of numbers of SGCN and SNI species by taxonomic groups. See Appendix B for the list of SGCN and SNI animal, plant and lichen species.

Table 2.1. Numbers of species of SGCN and SNI, the number of species assessed and the number of species that changed status from 2015, grouped first by vertebrate species, terrestrial invertebrates, aquatic invertebrates, plants, and lichen.

Taxa	# SGCN	# SNI	# Neither	Total Assessed	# SGCN removed	# SGCN added	# SNI added	% SGCN	% SNI
Amphibians	7	1	14	22	1	0	1	32%	5%
Birds	79	1	366	446	21	8	1	18%	0%
Fishes	36	5	124	165	8	2	5	22%	3%
Mammals	32	3	51	86	0	5	3	37%	3%
Reptiles	13	1	17	31	2	0	1	42%	3%
Bees (Class Insecta; Order Hymenoptera)	117	125	273	515	0	112	125	23%	24%
Beetles (Class Insecta; Order Coleoptera)	12	10	68	90	1	4	10	13%	11%
Butterflies, Moths (Class Insecta; Order Lepidoptera)	122	257	1708	2087	6	95	257	6%	12%
Flies (Class Insecta; Order Diptera)	0	5	62	67	0	0	5	0%	7%
Grasshoppers, Crickets (Class Insecta; Order Orthoptera)	0	6	5	11	0	0	6	0%	55%
Millipedes (Class Diplopoda)	0	1	0	1	0	0	1	0%	100%
Snails (Class Gastropoda)	9	23	120	152	0	4	23	6%	15%
Spiders (Class Arachnida; Order Araneae)	7	0	521	528	6	3	0	1%	0%
Springtails (Class Collembola)	0	3	0	3	0	0	3	0%	100%
True Bugs, Cicadas, Leafhoppers (Class Insecta; Order Hemiptera)	2	8	2	12	2	1	8	17%	67%
Caddisflies (Class Insecta; Order Trichoptera)	50	51	213	314	3	29	51	16%	16%
Dragonflies, Damselflies (Class Insecta; Order Odonata)	25	8	112	145	20	5	8	17%	6%
Mayflies (Class Insecta; Order Ephemeroptera)	8	83	72	163	0	8	83	5%	51%
Mussels (Class Bivalvia)	33	1	17	51	0	3	1	65%	2%
Shrimp (Class Branchiopoda)	0	1	8	9	0	0	1	0%	11%
Stoneflies (Class Insecta; Order Plecoptera)	5	34	37	76	0	5	34	7%	45%
Plants	447	87	1933	2467	0	447	87	18%	4%
Lichens	151	130	757	1038	0	151	130	15%	13%
TOTAL	1155	844	6480	8479	70	882	844		

Species-Habitat Associations

To associate SGCN with habitat types, species experts classified the SGCN by habitat types they use based on professional knowledge. Vertebrate species and some invertebrate species were classified as using habitats as primary or secondary habitat. Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently. Plants were only associated with a primary single habitat type. Animals with more general habitat requirements were associated with multiple habitat types, while specialists were associated with one or few. A detailed table associating each animal SGCN with the 15 habitats identified in this Plan can be found in Appendix C. Appendix D lists the plant SGCN associated with each of the primary habitats.

Stressors and Actions

Throughout Minnesota, over time, habitats have been lost and degraded due to pressures that result from human inhabitation, subsistence, livelihoods, and recreation. **Thus, habitat loss or alteration is the primary threat for most, if not all, SGCN.** However, an approach has been taken at the international level to classify threats, or ‘stressors’ as we call them in this Plan, to shine light on more of the proximate activities that continue to degrade our habitats or impact our SGCN directly. See also *Chapter 3. Habitats*.

We identify stressors as a way to direct conservation attention to where it is needed most. Conservation actions are explicitly linked to address the stressors, presented as a set of conservation opportunities. Stressors are factors that pose challenges to vulnerable plant and wildlife species, either directly or indirectly. Some stressors can be considered direct threats to SGCN (e.g., the stressor of ‘wind and solar energy’ and collisions with wind turbines). Others may operate indirectly through habitat conditions (e.g., the stressor of ‘invasive species’ causes a loss of streamside sandy nesting habitat due to cover by reed canary grass). Many stressors operate both directly and indirectly, such as roads, trails, and railroads causing both vehicular strikes (direct) and habitat fragmentation (indirect).

It should be noted that sometimes factors or activities labeled as ‘stressors’ can also be applied to advance conservation goals. Broad language such as ‘livestock management’ and ‘fire management’ is an attempt to remind readers of this – that these practices may sometimes be stressors but in other cases are applied as conservation tools. It is a matter of ‘how’ and ‘where’ these practices take place that determines if they are advancing ecosystem health and resilience as conservation actions (such as prescribed fire) or if they put stress on the ecological community, habitat, or species (such as a catastrophic wildfire following a long period of fire suppression). Throughout the Plan you will find language clarifying when a stressor may be a challenge and when it can be applied as part of a conservation solution.

We base the stressors listed in this Plan on an internationally employed threats lexicon developed by the International Union for the Conservation of Nature (IUCN) Species Survival Commission and the Conservation Measures Partnership (Salafsky et al., 2024). First developed in 2008 (Salafsky et al., 2008) and updated in 2024, version 4.0 of this hierarchical system describes eleven Level 1 (higher level) threats and then within those, 41 Level 2 (more specific) threats. We have selected a subset of 24 of these to discuss in relation to the species and habitats in Minnesota. We have shortened and re-named some for ease of understanding (see

Table 2.2). We developed a set of icons that we use throughout the Plan to depict stressors visually and to connect them to related Actions.

Table 2.2. Icons and the names of stressors on species and habitats referenced throughout the SWAP, along with threat level information from the IUCN classification (Salafsky et al., 2024).

ICON	STRESSOR IN OUR PLAN	LEVEL1_2	LEVEL 1 THREAT (IUCN CLASSIFICATION)
	Development	1	Residential, Commercial, & Recreation Areas
	Crop Production	2_1	Agriculture & Aquaculture
	Tree Plantations	2_2	Agriculture & Aquaculture
	Livestock Management	2_3	Agriculture & Aquaculture
	Mining and Quarrying	3_2	Energy Production & Mining
	Wind and Solar Energy Infrastructure	3_3	Energy Production & Mining
	Roads, Trails, and Railroads	4_1	Transportation, Service, & Security Corridors
	Utility Corridors	4_2	Transportation, Service, & Security Corridors
	Shipping Lanes	4_3	Transportation, Service, & Security Corridors
	Hunting and Collecting Animals	5_1	Biological Resource Use & Control
	Gathering Plants	5_2	Biological Resource Use & Control
	Timber harvest	5_3	Biological Resource Use & Control
	Fishing	5_4	Biological Resource Use & Control
	Recreation	6_1	Human intrusions & disturbance
	Fire Management	7_1	Natural System Management & Modifications
	Dams and Water Management	7_2	Natural System Management & Modifications
	Invasive Species (Problematic Non-native Species)	8_1	Invasive/Other Problematic Species, Genes & Pathogens
	Problematic Native Species	8_2	Invasive/Other Problematic Species, Genes & Pathogens
	Diseases and Pathogens	8_4	Invasive/Other Problematic Species, Genes & Pathogens
	Water-borne Pollution	9_1	Pollution

	Air-borne Pollution	9_3	Pollution
	Light and Noise Pollution	9_4	Pollution
	Changes in Temperature related to Climate	11_2	Climate Change
	Changes in Precipitation and Hydrology related to Climate	11_3	Climate Change

Also note that within each habitat or species sub-chapter, we only include primary stressors pertinent to that habitat or group of species rather than all of them, omitting the others for focus.

SGCN Sub-chapters

Information on Species in Greatest Conservation Need and Stressors is presented in 8 subchapters, one for each of the following groups of species:

1. Amphibians
2. Reptiles
3. Birds
4. Fishes
5. Mammals
6. Terrestrial Invertebrates
7. Aquatic Invertebrates
8. Plants and Lichens

Each of these taxonomic subchapters includes the following information:

Overview

Each subchapter begins with several paragraphs that summarize relevant information about each taxonomic group of species, including unique aspects of the species, the depth of information available, and/or the ecological importance of the species.

SGCN and SNI Summary Information

A brief summary of the number of SGCN and SNI identified is provided and the reader is directed to Appendix B with extensive detailed information.

Species-Habitat Associations

The association of each SGCN with the 15 habitats has been delineated and described in the 2025-2035 SWAP and are briefly summarized with details for animals provided in Appendix C and details for plants in Appendix D.

Stressors

As noted above, the primary stressors that may negatively affect SGCN in each group are identified and briefly described. Each stressor is depicted by an 'icon' to help the reader connect content from stressor to actions.

Priority Conservation Strategies

In each sub-chapter we reference the three Strategies for implementation, using these to group Conservation Actions.

Potential Conservation Actions

Conservation Actions, organized under Strategies, address primary stressors and are tagged by stressor icons to relate them. Frequently a conservation action can address more than one Stressor. In those cases, the Conservation Action is identified by multiple Stressor Icons.

Case Studies, Key Partnerships and Resources.

In these sections we highlight case studies, key partnerships, and provide links to key resources that may be related to the species group in that sub-chapter. Case studies feature examples of conservation projects that effectively address one or more of the stressors, including a brief summary of the project as well as links to read additional information. Key partnerships are described throughout the narrative; these are not an exhaustive list but rather key partnerships or programs that may assist in the implementation of conservation actions.

References

Master, L. L., D. Faber-Langendoen, R. Bittman, G. A. Hammerson, B. Heidel, L. Ramsay, K. Snow, A. Teucher, and A. Tomaino. 2012. NatureServe Conservation Status Assessments: Factors for Evaluating Species and Ecosystem Risk. NatureServe, Arlington, VA.

Faber-Langendoen, D., J. Nichols, L. Master, K. Snow, A. Tomaino, R. Bittman, G. Hammerson, B. Heidel, L. Ramsay, A. Teucher, and B. Young. 2012. NatureServe Conservation Status Assessments: Methodology for Assigning Ranks. NatureServe, Arlington, VA.

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, e14434.

<https://doi.org/10.1111/cobi.14434>

Salafsky, N., Salzer, D., Stattersfield, A. J., Hilton-Taylor, C., Neugarten, R., Butchart, S. H. M., Collen, B., Cox, N., Master, L. L., O'Connor, S., & Wilkie, D. (2008). A standard lexicon for biodiversity conservation: Unified classifications of threats and actions. *Conservation Biology*, 22, 897–911.

Amphibians

Overview

Minnesota is home to 22 species of amphibians – 14 frogs and toads, and 8 salamanders. In 1944 Walter Breckenridge published *Reptiles and Amphibians of Minnesota*, marking the state’s beginning of modern herpetology. At the time only 13 species of frogs and toads and 5 species of salamanders had been documented in the state. Breckenridge’s publication was considered Minnesota’s standard herpetological reference until *Amphibians and Reptiles Native to Minnesota* was published by Oldfield and Moriarty in 1994, followed by an update in 2014 (Moriarty and Hall). In the late 1970s, the Minnesota DNR’s Nongame Wildlife Program was established and has coordinated breeding frog call surveys and funded studies on Northern leopard frogs and Blanchard’s cricket frogs. The Minnesota Biological Survey, initiated in 1987, surveyed counties across the state targeting rare wildlife and collecting records on common species. Their work has helped establish Minnesota’s list of amphibian SGCNs and has added thousands of records of amphibians to the DNR’s Natural Heritage database.

Amphibians are the most threatened vertebrate group worldwide due to the compounding effects of habitat conversion, contaminants, invasive species, disease, and climate change, with 40% of species at risk of extinction (Campbell et al., 2024; Luedtke et al., 2023). This is especially concerning given the wide range of benefits amphibians provide. In their ecosystems, amphibians function as both predator and prey, assist with energy and nutrient flow between terrestrial and aquatic systems, and enhance soil aeration and productivity (IUCN SSC Amphibian Specialist Group, 2024). Amphibians also decrease the transmission of dangerous pathogens to humans through their predation on mosquitoes and flies and are frequently used to further the field of human medicine (IUCN SSC Amphibian Specialist Group, 2024). They are often sensitive indicators of environmental contaminants and serve as indicators of environmental health. Lastly, encountering amphibians in the wild at a young age can foster a positive attitude toward conservation. Many people have fond memories of catching frogs in a local pond or finding a salamander under a log on a hike through the forest. These moments can be the first spark that instills an interest in conservation action.

Despite all of this, lack of funding for amphibian conservation is a persistent issue (Bishop et al., 2012), likely because people often prefer charismatic large animals and some may fear “slimy” creatures like frogs and salamanders. Many amphibian species in Minnesota live cryptic lives and are not frequently encountered, so people might not even know they have some living in their own backyard. Our frogs, toads, and salamanders use a variety of terrestrial and aquatic habitats and exhibit a multitude of life history strategies. Therefore, we need to take a wide range of conservation actions to address the stressors detailed below, including monitoring and public engagement efforts, such as the Minnesota Frog and Toad Calling Survey (see inset) that actively involves citizen scientists in the goal of learning more about our amphibian fauna. The [Minnesota Herpetological Society](#) is another important venue for public education and promoting the study and conservation of both reptiles and amphibians. On a larger scale, [Midwest Partners in Amphibian and Reptile Conservation](#) (PARC) is an important network dedicated to the conservation of reptiles and amphibians that provides an avenue for collaborative conservation initiatives, outreach and education resources, and more.

Case Study: Engaging the Public in Frog and Toad Monitoring

A volunteer-based community science program, the [MN Frog and Toad Calling Survey](#) is designed to monitor the distribution and abundance of all 14 of the state's frogs and toads across the state. Over 100 routes along roads have been established, each with approximately 10 stops where volunteers stop to listen and record the species that are calling. Routes are sampled three times during the breeding season to cover the breeding seasons of all species. The original survey ran from 1994 through 2017, collecting thousands of records. The survey was re-launched in 2023 with updated tools and protocols and continues to develop and grow. The survey not only provides critically important information on a group of species that is often overlooked, but it has become a vital part of many volunteers' spring season. As one volunteer, Nick Krueger, reflected: "The frogs and toads add to the diversity of nature here, but they're also part of the beauty for us, the beauty of the soundscape. It's good to take time to be amazed at the world."

One of our rarest frogs, the Blanchard's cricket frog, has a unique history in Minnesota. Thought to have been lost from Minnesota by the late 1980s, the species was rediscovered in the late 1990s. Upon rediscovery, the Blanchard's cricket frog remained at very low population levels until new populations began to turn up in the 2010s and 2020s (Moriarty and Hall 2014, Smith 2018). As a state-listed, protected species, it is critical for regulators and scientists to understand occurrence and occupancy patterns for this tiny frog. Given the challenges with fitting surveys for Blanchard's cricket frogs into busy workloads, DNR staff searched for alternative methods to track what we hope is this species' recovery. From this need was born a partnership between DNR Nongame Wildlife Program staff and a community science, non-profit organization called [HerpMapper](#). HerpMapper is designed to gather and share information about amphibian and reptile observations across the planet. The [Blanchard's Cricket Frog Survey](#) asks Minnesotans to help find and document Blanchard's cricket frogs on the HerpMapper website or smartphone app. This effort combines community science with cutting edge species study and recovery efforts to help this enigmatic rare species click-and clack their way across Minnesota again.

SGCN and SNI Summary Information

Seven amphibian species have been designated Species in Greatest Conservation Need (SGCN) - two frogs, one toad, and four salamanders. Four of these are state-listed as Species of Special Concern, and one, Blanchard's cricket frog (*Acris blanchardi*), is State Endangered. Most of the amphibian SGCN species are also Midwest Regional SGCN and have documented declines and significant stressors impacting populations across their range in Minnesota. One species that was listed as a SGCN in the 2015 SWAP, the Eastern red-backed salamander (*Plethodon cinereus*) was removed given that populations currently appear to be stable in Minnesota's northern forests. One species, the Western tiger salamander (*Ambystoma mavortium*), has been designated a Species in Need of Information (SNI). The Western tiger salamander is visually similar to the Eastern tiger salamander (*Ambystoma tigrinum*), and their ranges overlap in Minnesota. More research, likely using molecular techniques, is needed to determine the extent of Western tiger salamander populations across the state. A complete list of both SGCN and SNI species can be found in Appendix B.

Habitat Associations

The amphibian SGCN utilize a variety of habitats, but all are associated with some type of aquatic habitat due to their life histories. Table 2.3 provides a summary of their association with the habitats discussed in the 2025-2035 SWAP. Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently. Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Detailed tables associating each SGCN with the primary 15 habitats in the 2025-2035 SWAP can be found in Appendix C.

Table 2.3. Numbers of species of amphibian SGCN associated with each of the habitat types (only showing habitats with amphibian SGCN associations).

Habitats	Primary Habitat	Secondary Habitat	Total
Prairie and Other Grasslands	1	0	1
Upland Conifer Forest and Woodland	0	3	3
Upland Deciduous Forest and Woodland	3	0	3
Mesic Hardwood Forest	2	0	2
Deciduous Wet Forest	0	3	3
Riparian and Floodplain Forest	2	0	2
Non-forested Wetlands	4	1	5
Rivers and Streams	3	0	3
Lakes	2	1	3
Vernal Pools	3	0	3
Urban and Other Developed Lands	0	1	1

Minnesota's seven amphibian SGCN were associated with an average of 4 primary habitats each. Several species use upland habitats in addition to wet habitats, and overall amphibians are associated with 11 of the primary 15 habitats. The habitat associated with the greatest number of species was non-forested wetlands supporting five of the seven SGCN.

The [Great Plains toad \(*Anaxyrus cognatus*\)](#) is the only amphibian SGCN that is primarily associated with open grassland habitats. It may be found in remnant prairies, wet meadows, and even agricultural fields, where it breeds in shallow wetlands and flooded fields that form after heavy rain in the early summer. The toads spend much of their time in underground burrows throughout the rest of the year.

Blanchard's cricket frogs are strongly associated with permanent water bodies like lakes, ponds, streams, and rivers, utilizing both the aquatic habitat and the adjacent shoreline. Pickerel frogs (*Lithobates palustris*) are found in similar habitats, though prefer cool streams and often forage in wet meadows and grasslands in the summer.

Three of the salamander SGCN – [spotted \(*Ambystoma maculatum*\)](#), [four-toed \(*Hemidactylium scutatum*\)](#), and eastern newt (*Notophthalmus viridescens*) – are strongly associated with mature forest habitats. They primarily occupy deciduous forests but will also occur in mixed deciduous-conifer forest types. A variety of

forested wetlands including vernal pools are critical for these species, which use them in the spring to breed. Eggs are laid in or near the water, and larvae remain in the pools until metamorphosis occurs. Four-toed salamanders typically lay their eggs in moss hummocks adjacent to open water. Eastern newts use a wide range of aquatic habitats, including lakes, ponds, and streams. Their unique life history includes the development of egg to larvae in aquatic habitat followed by a lengthy terrestrial juvenile stage that lasts a few years. Upon reaching maturity, adults undergo additional transformations as they revert to being primarily aquatic.

The [mudpuppy \(*Necturus maculosus*\)](#) is Minnesota's only fully aquatic salamander and is restricted to medium or large rivers and lakes. They take cover under submerged rocks or logs, which are important components of their habitat. Females attach their eggs to the underside of such structures and guard them until they hatch.

Primary Stressors for Amphibians

Stressors are factors that pose direct or indirect challenges to vulnerable plant and wildlife species. Habitat loss and degradation are generally considered the primary threat for most SGCN, and some stressors reduce the quality of those habitats. Other stressors may operate more directly on the SGCN, such as disease and pathogens. In some cases, stressors may operate both directly and indirectly. For example, roads, trails, and railroads can directly harm SGCN through vehicular strikes, or operate indirectly through habitat fragmentation and reduced landscape connectivity. The stressor list below is adapted from an internationally recognized threats classification developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For more information, see the “Stressors” section of Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some factors listed as “stressors” can also be used to advance conservation goals. Broad terms such as “fire management” reflect the dual nature of these factors as they may function as stressors in some contexts (e.g. catastrophic wildfire following a prolonged period of fire suppression) while serving as valuable conservation tools in others (e.g., appropriately planned and applied prescribed fire).

Information about a subset of primary stressors specifically impacting this species group is included below, followed by a set of conservation actions addressing those stressors.



Development

Development for residential communities, recreation, or commercial and industrial areas eliminates, degrades and fragments amphibian habitat. This leads to a variety of negative impacts that affect survival and reproduction of amphibians, such as desiccation, reduced reproductive output, exposure to higher temperatures, reduced dispersal ability, increased exposure to contaminants, and reduced foraging success (Eakin et al., 2019; IUCN SSC Amphibian Specialist Group, 2024). It also makes traversing the landscape more difficult and dangerous (see also Roads, Trails, and Railroads).



Crop Production

Conversion of amphibian habitat to agriculture eliminates, degrades and fragments amphibian habitat. This can lead to a variety of negative impacts on amphibians, including desiccation, reduced reproductive output, exposure to higher temperatures, reduced dispersal ability, and reduced foraging success (IUCN SSC Amphibian Specialist Group, 2024). Drainage also plays a key role in row crop agriculture but has adverse effects on aquatic and terrestrial habitats that amphibians depend upon (Blann et al., 2009; Schottler et al., 2013; Schulz et al., 2015). The loss of wetlands due to agricultural practices can eliminate important wetland habitats, resulting in wetlands being separated beyond the species’ normal dispersal distance. The presence of grassland and wet meadow habitats between wetlands provides cover and foraging habitat for dispersing frogs, improving the potential for successful migration.



Livestock Management

Grazing by livestock can help to maintain open canopy habitats for amphibian species dependent on those. Nevertheless, the overall effects of livestock operations depend on management details such as stocking rates and whether livestock have direct access to water bodies. Amphibians are generally

sensitive to water quality parameters such as levels of dissolved oxygen and nitrate concentration. Amphibians may be impacted by livestock operations where waterways are not buffered, impacting shoreline habitat as well as water quality from manure, fertilizer, pesticide, and sediment runoff (Howell et al., 2019).



Mining and Quarrying

Habitat changes caused by mining, such as changes in water quality, hydrology, or the physical structure of wetlands, can affect amphibian species composition and abundance. Further, mining can contribute to the introduction of pollutants that may cause amphibian malformations (Sasaki et al., 2015, Frelich, 2019; Azizishirazi et al., 2021).



Wind and Solar Energy Infrastructure

There is insufficient research on how wind and solar facilities affect amphibians and the need to identify effective measures to reduce impacts. Wind and solar farms may impact amphibians through physical habitat alteration and noise and light pollution (de Oliveira et al., 2025; Chock et al., 2020).



Roads, Trails, and Railroads

The proliferation of roads fragments and destroys amphibian habitat. This reduces the dispersal capabilities of individuals and may result in genetic isolation of populations on either side of the road. Many amphibian species attempt to cross roads, especially during breeding seasons, and experience high levels of road mortality. Roads also increase human access to natural habitats and increase the spread of invasive species that may degrade the habitat or outcompete native amphibian species (Andrews et al., 2008, Schmidt & Zumbach, 2008). Research by Fahrig et al. (1995) indicated that traffic mortality had a significant negative impact on local anuran populations.



Utility Corridors

Amphibians are underrepresented in studies addressing direct and indirect impacts and response thresholds to pipelines and powerlines (Richardson et al., 2017). There are existing and proposed transport pipelines in Minnesota with potential to impact amphibians via habitat loss, degradation and fragmentation (Brown et al., 2023). Amphibians may also be impacted in the event of pipeline spills (see Water-borne Pollution).



Hunting and Collecting Animals

In Minnesota, frogs, toads, and salamanders are classified under “protected wild animals.” A person may not take, buy, sell, transport, or possess a protected wild animal unless allowed by the game and fish laws (M.S. 97A). There are stricter rules for state endangered or threatened species (M.S. 84.0895). Minnesota fishing regulations allow people with a fishing license to take, use, buy, and sell an unlimited number of frogs up to 6” long for bait, with the exception of the state-endangered Blanchard’s cricket frog. A special license is needed to take frogs for any purpose other than bait. Current fishing regulations do not allow taking of any native salamander species as bait.

Even though these rules are in place, illegal collection still occurs. People might not be aware of the rules or purposefully break them. Anecdotal reports of tadpoles mixed with minnows at bait shops indicate that there

are cases where unintentional movement of frogs occurs. It is very difficult to monitor illegal take and appropriately enforce the rules.



Timber Harvest

Timber harvest can have negative impacts on forest-dwelling amphibians, especially salamanders. Logging alters microhabitats, increases soil compaction and desiccation, and reduces habitat complexity. Clearcuts are often the most damaging; some studies suggest it may take decades for populations to recover after a clearcut (Pough et al., 1987, deMaynadier and Hunter, 1995).

Forest-dwelling salamanders like spotted salamanders, four-toed salamanders, and eastern newts prefer high levels of canopy cover, up to 90% (Faccio, 2003). One study found that spotted salamanders were absent where forest cover was reduced to below 30%, and eastern newts were absent where forest cover was reduced to below 50% (Gibbs, 1998).

Salamanders require both forested upland and wetland habitat to complete their life cycles. Many small wetlands (i.e. vernal pools) are undocumented and unprotected, and identification is time-consuming as they cannot be easily identified from aerial imagery. Therefore, it is difficult to prevent destruction of these critical habitats during timber harvest operations. (See Case Study at the end of this subchapter on monitoring the impacts of forest management on the four-toed salamander.)



Fishing

Mudpuppies are occasionally caught by anglers, and may be killed accidentally, or purposefully by individuals who mistakenly believe mudpuppies are poisonous or venomous. Reports submitted to the DNR indicate that most mudpuppies are released unharmed.



Recreation

Motorized vehicles can destroy sensitive habitat or run over small, slow-moving amphibians. Bonfires, party sites, camping or fishing spots, and picnic areas can pollute habitats with garbage and attract predators such as raccoons, skunks, and foxes. Habitat can be degraded when recreational activity includes rolling logs or lifting rocks to locate secretive animals. These provide important shelter for amphibians and should carefully be replaced to their original position.



Fire Management

Fire suppression increases the growth and spread of woody vegetation in open upland habitats used by some amphibian species like the Great Plains toad. It can also lead to larger wildfires due to a buildup of vegetation on the landscape. Amphibians are slow-moving and may find it difficult to escape wildfires if they are on the surface; however, since many amphibian species utilize underground burrows, most individuals might be able to avoid harm. More research is needed on the response of amphibians to fire on the landscape (Hossack and Pilliod, 2011).



Dams and Water Management

Water management is comprised of many diverse actions such as water level manipulation, dams, drainage, dredging, water deposition and transfer, and more. These can greatly alter hydrology, stream

geomorphology, and water quality which in turn can affect amphibians. Dredging and even beneficial practices like stream restoration often involve construction, heavy equipment use, and excavation which can result in direct injury or mortality of amphibians. Increased drainage can cause downstream flooding and high flows, erosion, and runoff of chemicals and nutrients (Schottler et al., 2013). Dams reduce connectivity to key habitats and separate populations so they can no longer breed with each other. Water management can convert or degrade wetland and lake habitats. Ephemeral prairie wetlands have been drained for row crop production, severely reducing native breeding habitats for the Great Plains toad. Removal and placement of dredged material should avoid potential impacts to amphibian SGCN, such as the Blanchard's Cricket Frog. Placing dredge spoil materials into breeding sites and/or altering the hydrology associated with overwintering sites could cause major impacts to local populations.



Invasive Species (Problematic Non-native Species)

In Minnesota, the American bullfrog is considered to be non-native and invasive outside of the extreme southeastern corner of the state. Populations occur in many other counties, likely due to intentional releases. This could eventually displace other native amphibian species in aquatic habitats throughout Minnesota due to predation and competition for resources, if left unchecked (Bucciarelli et al., 2014; Moriarty and Hall 2014). Research has found that invasive plants such as buckthorn reduce hatching success and larval survival of some amphibians and may contribute to amphibian declines (Sacerdote & King, 2014).



Problematic Native Species

Fish releases into wetlands and shallow lakes—particularly previously fishless water bodies-- is known to reduce abundance, density, and/or species richness of amphibians (e.g. release of unused bait fish; intentional fish-rearing and stocking.) Fish may also enter these systems through other means such as flooding and ditching. These effects can be from fish that are either native or non-native and are not limited to predatory fish (Hecnar & M'Closkey, 1997; Herwig et al., 2013; Holbrook & Dorn, 2015).



Diseases and Pathogens

Infectious diseases are one of the top causes of amphibian decline globally (IUCN SSC Amphibian Specialist Group 2024). Ranavirus is transmitted through contaminated water, physical contact, and ingestion of infected tissues and may present as abnormal behavior, lethargy, swelling, fluid accumulation, and skin hemorrhage or ulcers (Cornell University, 2025a). Chytridiomycosis is caused by the waterborne fungus *Batrachochytrium dendrobatidis* and infects the keratin layer of the skin, causing weight loss, lethargy, skin shedding, red skin, and abnormal behavior (Cornell University, 2025b). Another more recently discovered fungus, *Batrachochytrium salamandrivorans*, has decimated salamander populations in Europe. It also infects the skin, causing reddening and ulcers, as well as lethargy and weight loss (Cornell University, 2025c). This fungus has not yet been found in North America, but if it does spread here, it could have major impacts on our salamander populations. Of note, one of Minnesota's SGCN, the eastern newt (*Notophthalmus viridescens*), has been found to be highly susceptible to the fungus in research trials (Cornell University 2025c). Major contributors to the spread of pathogens like these include the pet trade, biological supplies industry, and people using contaminated equipment or footwear. Challenges to addressing this threat include lack of funding and the high workload and short response window needed to research amphibian diseases.



Water-borne Pollution

Amphibians are very susceptible to water-borne pollutants, including salt from roads and water softeners, endocrine disrupters, hormones from birth control, herbicides, fertilizers, and more.

These can impact the growth, development, and survival of eggs and larvae through direct and sublethal effects that influence survival (Ocampo et al., 2022; Thompson, 2022; Benli, 2024; Tornabene et al., 2024). Frog malformations have been a topic of concern in Minnesota after high rates of malformed frogs were found in a variety of ponds and wetlands in the 1990s. The cause of malformations is likely related to combinations of chemical, biological and physical factors (U.S. Geological Survey, 2001). Impacts associated with acid rain include direct toxicity and developmental abnormalities, as well as indirect impacts on food availability and aquatic vegetation (Croteau et al., 2008).



Light and Noise Pollution

Reproductive behaviors of amphibian species can be altered by artificial lights at night. Studies have found that it can inhibit amphibian movement to and from breeding areas and decrease mating call activity (Longcore & Rich, 2004.) A quick increase in the amount of light can cause a reduction in vision that can last for minutes or hours (Longcore & Rich, 2004).

Noise pollution is also problematic for frogs and toads, which rely on acoustic communication. Noises from many sources (urbanized areas, traffic, energy production) influence calling activity, alter mate selection, and cause physiological changes such as increased stress, suppressed immune function, and coloration change (Zaffaroni-Caorsi et al., 2022).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

Amphibians are micro-habitat specialists and thus are very sensitive to changes in temperature. Current studies generally suggest that extreme heat stress can suppress amphibian immune responses. Elevated temperatures can also alter microbial communities in amphibian skin and gut, which may reduce resistance to

pathogens. Studies also suggest that increasing temperatures may cause reductions in body size, which may then lower reproductive success (Sheridan & Bickford, 2011; Hernández-Pacheco et al., 2020).



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota’s average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al. 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

More specifically, amphibians are very sensitive to changes in precipitation and moisture. Droughts can suppress amphibian immune responses (Rollins-Smith & Le Sage, 2023). Blanchard’s cricket frogs are negatively impacted by river flooding in the winter because they cannot survive inundation in anoxic water for more than 24 hours nor in initially oxygenated water past 10 days (Irwin et al., 1999). If precipitation increases due to climate change and results in the inundation of overwintering sites, Blanchard’s cricket frog mortality will also increase. In the long-term, climate change could result in drastic changes to the landscape. Extensive periods of drought and/or elevated temperatures could eliminate shallow wetlands critical as amphibian breeding sites and recruitment of young (McMenamin et al., 2008).

Priority Conservation Strategies

To implement the SGCN Goal of this Plan, to conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions, three strategies were identified:



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnerships with agencies, Tribes, non-governmental organizations, and private landowners.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management practices.

Examples of conservation actions are grouped below under these three strategies and tagged with icons for the stressor(s) that they address. Some actions combine multiple strategies, in which case we present it under the one it fits best. Some actions such as those focused on monitoring might not always be tied to a specific stressor; these are labeled with 'n/a' in the stressor column.

Potential Conservation Actions for Amphibians




Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.

Stressor	Action
	Develop citizen science programs to identify stretches of road that are mortality hotspots for amphibians, especially during spring and fall breeding seasons.
	Conduct wetland surveys focusing on identifying and mapping ephemeral wetlands like vernal pools (Hofmeister et al., 2022). Create a citizen science program or app for reporting locations of ephemeral wetlands.
	Monitor population trends of rare forest species. Monitor impact of forest management activities on rare forest species.
	Support research on effects of recreation on SGCN amphibians, including thresholds for response and avoidance of significant adverse effects. Prioritize specific recreation types and habitats based

	on assessment of need in environmental review. Identify sensitive areas and avoid promoting recreational use and development in those areas. Enforce closures in sensitive areas.
	Monitor the status of bullfrog populations to help establish where they may be a problem for native species in central and northern Minnesota.
	Prioritize amphibian disease monitoring to understand impacts on populations and to facilitate early detection of <i>Batrachochytrium salamandrivorans</i> (Bsal) if it arrives in North America. Increase capacity of amphibian disease surveillance and response through collaboration between DNR and external partners, such as the USGS National Wildlife Health Center and its Wildlife Health Information Sharing Partnership (WHISPers) and the North American Bsal Task Force .
	Investigate mudpuppy mortality events. Conduct surveys to better understand mudpuppy habitat use. Work with state and federal partners to develop and test hypotheses related to stressors. Engage citizens by soliciting reports of amphibian mortality events.
n/a	Engage the public in community science to help survey and monitor amphibians. Example opportunities include: Minnesota Frog & Toad Calling Survey and Minnesota Amphibian & Reptile Survey .
n/a	Conduct rigorous effectiveness monitoring to assess responses of SGCN to specific management activities in order to develop improved evidence-based management recommendations and avoidance measures for SGCN (Binley et al., 2025) as part of adaptive management.
n/a	Increase collaboration with universities to further research and monitoring of selected amphibian SGCN.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.

Stressor	Action
	Work with MnDOT, cities, and counties to develop under-road wildlife crossing tunnels and surmountable curbs in areas close to amphibian breeding sites such as ponds and vernal pools, especially those identified as road mortality hotspots. Consider additional options such as road closure or speed reduction during amphibian breeding seasons and signage informing drivers of amphibian crossings (Gunson et al., 2016; Marcelino et al., 2024).
	Maintain vegetated buffers comprised of locally native vegetation adjacent to waterways. Avoid or reduce mowing, burning, grazing or chemical application on these buffers. Support and promote state and federal programs incentivizing buffers and other best practices for lakes, streams, wetlands and ponds (Semlitsch & Bodie, 2003).
	Promote protection and restoration of native shoreline habitats (Restore Your Shore ; Rivers and Streams).
	Increase representation of all stakeholder groups in forest management decisions. Incorporate ecological values into sustainable timber harvest model and evaluate proposed harvest rates for impacts on wildlife habitat.
	Work with road authorities to implement culvert designs that improve stream connectivity and aquatic organism passage (Hernick et al., 2019).
	Restore wetlands to benefit a variety of amphibians, including seasonal and temporary wetlands that do not sustain fish.
	Support intra and inter-agency drainage teams focused on evidence-based best practices such as: - Control drainage to strategically manage water levels and mimic natural drainage and adapt to climate change to improve water quality, reduce flooding and support ecological functions - Restore and protect wetlands for water filtration, flood control and wildlife habitat - Reduce runoff - Incorporate native perennial plantings and cover crops - Establish shallow drainage systems to help maintain wetlands - Establish two-stage ditches designed to slow and filter runoff, reduce erosion and improve water quality - Restore river and stream connectivity via properly designed passage structures and removing dams

Stressor	Action
	To mitigate impacts of dams and changes in water management regimes, focus on dam removal and stream restorations that avoid or minimize unintended negative impacts – for example, employ special precautions and conduct at a time of year that is less detrimental to stream-dwelling amphibians. Stream restoration should include naturally associated off-channel habitats (e.g. oxbows) and stream bank specifications conducive to rare amphibians, where indicated. Because SGCN amphibians can co-occur with SGCN from other wildlife groups, it is important to consider their needs as well, for a more integrated approach.
	Incentivize renewable energy that “piggybacks” on existing infrastructure versus developing new facilities on currently undeveloped sites (e.g. example, atop buildings, parking lots and ramps, etc.). Explore examples from other countries.
	Limit creation of new utility line corridors in important wetland and shallow lake areas. Develop best management practices for creation and maintenance of utility lines. Reduce chemical vegetative control methods.
	Maintaining and improving landscape connectivity is a key strategy to support amphibians in the face of climate change. For example, maintaining and restoring a diverse set of interconnected, healthy aquatic and terrestrial habitats and ensuring there are safe movement corridors that may also allow them to track shifting conditions (Uden et al., 2014; Souza et al., 2023). SGCN such as Blanchard’s cricket frog (Badje et al., 2021) and Great Plains toad (Bragg, 1940) are among the many amphibian species that would benefit from improved dispersal options to access overwintering, migration, and breeding habitats. For a given amphibian SGCN, ensure that uplands adjacent to their aquatic habitats are equally suitable and permeable (Youngquist & Boone, 2014).



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Stressor	Action
	Increase enforcement of nongame illegal take issues; educate conservation officers on the issues and increase monitoring of important areas for poaching.
	Increase public awareness of state rules and regulations around take, possession, purchasing, and selling of native amphibian species and the reasons for these laws. Promote alternative ways to appreciate Minnesota's amphibians in their native habitats.
	Educate land managers, including private landowners, about habitat needs of rare forest species.
	Increase public education and outreach about how releases of captive animals (including wild-caught individuals) can spread disease.
	Increased public education and outreach about best management practices for artificial lights at night, published by Dark Sky International . Advocate for increased regulation of light and noise pollution, especially near important amphibian breeding sites.

Case Study: Monitoring the Response of Four-toed Salamanders to Forest Management

Understanding how a species, and their habitat, responds to forest management is a vital piece to conserving rare forest-dependent wildlife. Initiated in 2022, the DNR's Nongame Wildlife Program began a study to explore how four-toed salamander populations respond to forest management activities in proximity to their preferred nesting habitat: vernal pools. This study utilizes a Before-After-Control-Impact (BACI) study design in an effort to tease apart natural annual variation in salamander abundance from the impact that forest management may have on these populations. Within four-toed salamander range in Minnesota, 45 vernal pools were identified for long-term monitoring with approximately half of these vernal pools scheduled for forest management activities within 350 meters of the pool boundary (i.e., treatment vernal pools). The other half of the survey pools are identified as controls and are not planned for forest management. Survey pools were selected based on historical and recent observations of four-toed salamanders or pools that consisted of suitable nesting habitat. Survey work involves conducting nesting surveys that consist of 30 minutes of "active" search time or until all suitable nesting habitat within a vernal pool has been searched, whichever comes first. "Active" search time is defined as any time spent searching suitable nesting habitat (moss hummocks, woody debris, sedges clumps, etc.) within the highwater mark of a vernal pool for female four-toed salamanders and/or eggs. Relative abundance is determined by counting the occurrence of nesting activity (i.e., presence of salamander guarding eggs, gravid female salamanders, or unattended egg masses) within each wetland. All wetlands are surveyed at least once during the nesting season (May 1 – June 15).

Started in 2024, vegetation surveys and trail cameras deployment occurred within several survey pools (both treatments and controls) to better understand vegetation composition and hydroperiod of vernal pools, respectively. Vegetation surveys were conducted in August after the drying period within these vernal pools to allow for ample growing season and to minimize impacts to salamander larval development. These surveys employed a relevé sampling structure and documented species abundance, diversity, and other vernal pool characteristics. The goals of these surveys are to document vegetation composition/diversity within vernal pools and to develop a classification structure for this understudied wetland type. Additionally, these data establish a baseline of vegetation composition and provide the opportunity to examine changes in vegetation after forest management has occurred. The objectives of the trail camera deployment are to record spatial and temporal variation in hydroperiod across most of the study vernal pools, and to determine whether hydroperiod significantly changed after forest management operations. These cameras are programmed to record timelapse photos of the wetlands and identify how many days standing water was present/visible. The deepest location of the wetlands was located during nesting surveys and a measuring post and camera was installed to monitor this location.

As of 2024, four-toed salamanders were detected in approximately 90% of the study vernal pools. Over 500 nests have been detected with over 290 female four-toed salamanders observed during nesting surveys. The mean egg mass size is 39.5 eggs (StDev: 53.2). The majority of vernal pool depths ranged from 24 – 35 cm at the time of nesting surveys. With May 1 as the start day for hydroperiod monitoring, most vernal pools held water for 74 to 152 day (July 14 and September 30, respectively) with a mean hydroperiod of 103 days (August 12). To date, few vernal pools have experienced the planned forest management activities; thus, not enough data exists to make any inference on the impacts of forest management on four-toed salamander abundance. As this study continues for the next several years, additional data will yield valuable information on how best to protect these sensitive vernal pools and the species that rely on them.

References for Amphibians

- Andrews, K. M., Gibbons, J. W., Jochimsen, D. M., & Mitchell, J. (2008). Ecological effects of roads on amphibians and reptiles: a literature review. *Herpetological Conservation*, 3, 121–143.
- Azizishirazi, A., Klemish, J. L., & Pyle, G. G. (2021). Sensitivity of amphibians to copper. *Environmental Toxicology and Chemistry*, 40(7), 1808–1819. <https://doi.org/10.1002/etc.5049>
- Badje, A., Brandt, T., Bergeson, T., Paloski, R. & Kapfer, J. (2021). Spring movement ecology of Blanchard's cricket frog (*Acris blanchardi*) in southwestern Wisconsin, USA. *Herpetological Conservation and Biology*, 16(2), 271-280.
- Benli, P. P. (2024). Responses of Fishes, Amphibians, and Reptiles to Neonicotinoids. In Singh, R., Singh, V.K., Kumar, A., Tripathi, S., Bhadouria, R. (eds) *Neonicotinoids in the Environment*. Sustainable Landscape Planning and Natural Resources Management. Springer, Cham. https://doi.org/10.1007/978-3-031-45343-4_6
- Binley, A. D., Haddaway, L., Buxton, R., Lalla, K. M., Lesbarreres, D., Smith, P. A., & Bennett, J. R. (2025). Endangered species lack research on the outcomes of conservation action. *Conservation Science and Practice*. <https://doi.org/10.1111/csp2.13304>
- Bishop, P. J., Angulo, A., Lewis, J. P., Moore, R. D., Rabb, G. B., & Garcia Moreno, J. (2012). The amphibian extinction crisis - what will it take to put the action into the Amphibian Conservation Action Plan? *S.A.P.I.E.N.S.*, 5(2), 97-111.
- Blann, K. L., Anderson, J. L., Sands, G. R., & Vondracek, B. (2009). Effects of Agricultural drainage on Aquatic ecosystems: a review. *Critical Reviews in Environmental Science and Technology*, 39(11), 909–1001. <https://doi.org/10.1080/10643380801977966>
- Bragg A.N. (1940). Observations on the ecology and natural history of Anura. I. habits, habitat and breeding of *Bufo cognatus* say. *American Naturalist* 74(753), 424–438.
- Breckenridge, W. (1944). *Reptiles and Amphibians of Minnesota*. University of Minnesota Press.
- Brown, D. J., Knopka, S. C., Grushecky, S. T., Owen, S. F., & Edwards, J. W. (2023). Influence of natural gas pipeline Right-of-Ways on eastern red-backed salamander occurrence in the Northern Appalachians. *Journal of Fish and Wildlife Management*, 14(2), 303–314. <https://doi.org/10.3996/jfwm-22-032>
- Bucciarelli, G. M., Blaustein, A. R., Garcia, T. S., & Kats, L. B. (2014). Invasion complexities: The diverse impacts of nonnative species on amphibians. *Copeia*, 2014(4), 611–632. <https://doi.org/10.1643/ot-14-014>
- Campbell, K. S., Baltensperger, A. P., & Kerby, J. L. (2024). Random Frogs: using future climate and land-use scenarios to predict amphibian distribution change in the Upper Missouri River Basin. *Landscape Ecology*, 39(3). <https://doi.org/10.1007/s10980-024-01841-z>

- Chock, R. Y., Clucas, B., Peterson, E. K., Blackwell, B. F., Blumstein, D. T., et al. (2020). Evaluating potential effects of solar power facilities on wildlife from an animal behavior perspective. *Conservation Science and Practice*, 3(2). <https://doi.org/10.1111/csp2.319>
- Cornell University Wildlife Health Lab. (2025a). "Ranavirus." <https://cwhl.vet.cornell.edu/disease/ranavirus>
- Cornell University Wildlife Health Lab. (2025b). "Chytridiomycosis." <https://cwhl.vet.cornell.edu/disease/chytridiomycosis>
- Cornell University Wildlife Health Lab. (2025c). "Salamander Chytrid." <https://cwhl.vet.cornell.edu/disease/salamander-chytrid>
- Croteau, M.C., Hogan, N., Gibson, J.C., Lean, D., & Trudeau, V. (2008). Toxicological threats to amphibians and reptiles in urban environments. In J.C. Mitchell & R.E. Jung Brown (Eds.), *Herpetological Conservation* (vol. 3, pp. 197-209).
- De Oliveira, R. F., De Araujo Lira, A. F., Zaffaroni-Caorsi, V., Feitosa, M. L. B., & De Moura, G. J. B. (2025). Wind farm noise negatively impacts the calling behavior of three frogs in Caatinga dry forests. *PLoS ONE*, 20(3), e0318517. <https://doi.org/10.1371/journal.pone.0318517>
- deMaynadier, P. G., & Hunter, M. L., Jr. (1995). The relationship between forest management and amphibian ecology: a review of the North American literature. *Environmental Reviews*, 3(3–4), 230–261. <https://doi.org/10.1139/a95-012>
- Eakin, C.J., Calhoun, A.J. & Hunter, M.L. (2019). Effects of suburbanizing landscapes on reproductive effort of vernal pool-breeding amphibians. *Herpetological Conservation and Biology*, 14(2), 515-532. http://www.vernalpools.me/wp-content/uploads/2020/02/Eakin_et al_20192.pdf
- Faccio, S. D. (2003). Postbreeding emigration and habitat use by Jefferson and spotted salamanders in Vermont. *Journal of Herpetology*, 37(3), 479–489. <https://doi.org/10.1670/155-02a>
- Fahrig, L., Pedlar, J. H., Pope, S. E., & et al. (1995). Effect of road traffic on amphibian density. *Biological Conservation*, 177–182. <https://escholarship.org/content/qt22t1h3q1/qt22t1h3q1.pdf>
- Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Frelich, L. E. (2019). Terrestrial ecosystem Impacts of sulfide mining: Scope of Issues for the Boundary Waters Canoe Area Wilderness, Minnesota, USA. *Forests*, 10(9), 747. <https://doi.org/10.3390/f10090747>
- Gibbs, J.P. (1998). Distribution of woodland amphibians along a forest fragmentation gradient. *Landscape Ecology* 13, 263-268.

- Gunson, K., Seburn, D., Kintsch, J. & Crowley, J. (2016). Best management practices for mitigating the effects of roads on amphibian and reptile species at risk in Ontario. Ontario Ministry of Natural Resources and Forestry, Queen Printer for Ontario, Ontario, pp. 7-8. https://files.ontario.ca/bmp_herp_2016_final_final_resized.pdf
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Hecnar, S. J., & M'Closkey, R. T. (1997). The effects of predatory fish on amphibian species richness and distribution. *Biological Conservation*, 79(2–3), 123–131. [https://doi.org/10.1016/s0006-3207\(96\)00113-9](https://doi.org/10.1016/s0006-3207(96)00113-9)
- Hernández-Pacheco, R., Plard, F., Grayson, K. L., & Steiner, U. K. (2020). Demographic consequences of changing body size in a terrestrial salamander. *Ecology and Evolution*, 11(1), 174–185. <https://doi.org/10.1002/ece3.6988>
- Hernick, M., Lenhart, C., Kozarek, J. and Nieber, J. (2019). Minnesota guide for stream connectivity and aquatic organism passage through culverts. Minnesota Department of Transportation. <https://conservancy.umn.edu/bitstreams/be794e36-0663-490f-9432-be1dea584904/download>
- Herwig, B. R., Schroeder, L. W., Zimmer, K. D., Hanson, M. A., Staples, D. F., Wright, R. G., & Younk, J. A. (2013). Fish influences on amphibian presence and abundance in prairie and parkland landscapes of Minnesota, USA. *Journal of Herpetology*, 47(3), 489–497. <https://doi.org/10.1670/11-283>
- Hofmeister, K. L., Eggert, S. L., Palik, B. J., Morley, D., Creighton, E., Rye, M., & Kolka, R. K. (2022). The identification, mapping, and management of seasonal ponds in forests of the Great Lakes region. *Wetlands*, 42(1). <https://doi.org/10.1007/s13157-021-01526-2>
- Holbrook, J. D., & Dorn, N. J. (2015). Fish reduce anuran abundance and decrease herpetofaunal species richness in wetlands. *Freshwater Biology*, 61(1), 100–109. <https://doi.org/10.1111/fwb.12683>
- Hossack, B. R., & Pilliod, D. S. (2011). Amphibian Responses to Wildfire in the Western United States: Emerging Patterns from Short-Term Studies. *Fire Ecology*, 7(2), 129–144. <https://doi.org/10.4996/fireecology.0702129>
- Howell, H.J., Mothes, C.C., Clements, S.L., Catania, S.V., Rothermel, B.B. & Searcy, C.A. (2019). Amphibian responses to livestock use of wetlands: new empirical data and a global review. *Ecological Applications*, 29(8), p.e01976. https://caitlinmothes.com/files/Howell_et_al-2019-Ecological_Applications.pdf
- Irwin, J. T., Costanzo, J. P., & Lee, R. E., Jr. (1999). Terrestrial hibernation in the northern cricket frog, *Acris crepitans*. *Canadian Journal of Zoology*, 77(8), 1240–1246. <https://doi.org/10.1139/z99-087>
- IUCN SSC Amphibian Specialist Group. (2024). Amphibian conservation action plan: A status review and roadmap for global amphibian conservation. Wren, S., Borzée, A., Marcec-Greaves, R. & Angulo, A. (Eds.). IUCN SSC Occasional Paper, No 57. Gland, Switzerland: IUCN.

- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Longcore, T. & Rich, C. (2004). Ecological light pollution. *Frontiers in Ecology and the Environment*, 2(4), 191-198.
- Luedtke, J. A., Chanson, J., Neam, K., Hobin, L., Maciel, A. O., et al. (2023). Ongoing declines for the world's amphibians in the face of emerging threats. *Nature*, 622(7982), 308–314. <https://doi.org/10.1038/s41586-023-06578-4>
- Marcelino, M. R., Parren, S. G., & Mosher, B. A. (2024). Assessing the Efficacy of Wildlife Underpasses in Mitigating Amphibian Road Mortality: A Case Study from the Northeastern United States. *Research Square (Research Square)*. <https://doi.org/10.21203/rs.3.rs-5551430/v1>
- McMenamin, S. K., Hadly, E. A., & Wright, C. K. (2008). Climatic change and wetland desiccation cause amphibian decline in Yellowstone National Park. *Proceedings of the National Academy of Sciences*, 105(44), 16988–16993. <https://doi.org/10.1073/pnas.0809090105>
- Moriarty, J.J. & Hall, C.D. (2014). *Amphibians and Reptiles in Minnesota*. University of Minnesota Press.
- Ocampo, M., Chuirazzi, C., & Takahashi, M. K. (2022). The effects of road salt (NaCl), predation, and competition on the growth and community interactions of spotted salamanders (*Ambystoma maculatum*) and wood frogs (*Lithobates sylvaticus*). *Environmental Pollution*, 315, 120349. <https://doi.org/10.1016/j.envpol.2022.120349>
- Oldfield, B. & Moriarty, J.J. (1994). *Amphibians and Reptiles Native to Minnesota*. University of Minnesota Press.
- Pough, F. H., Smith, E. M., Rhodes, D. H., & Collazo, A. (1987). The abundance of salamanders in forest stands with different histories of disturbance. *Forest Ecology and Management*, 20(1–2), 1–9. [https://doi.org/10.1016/0378-1127\(87\)90146-0](https://doi.org/10.1016/0378-1127(87)90146-0)
- Richardson, M.L., Wilson, B.A., Aiuto, D.A.S., Crosby, J.E., Alonso, A., Dallmeier, F. & Golinski, K.G. (2017). A review of the impact of pipelines and power lines on biodiversity and strategies for mitigation. *Biodiversity and Conservation*, 26, 1801–1815. <https://doi.org/10.1007/s10531-017-1341-9>
- Rollins-Smith, L. A., & Sage, E. H. L. (2023). Heat stress and amphibian immunity in a time of climate change. *Philosophical Transactions of the Royal Society B Biological Sciences*, 378(1882). <https://doi.org/10.1098/rstb.2022.0132>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>

- Sacerdote, A.B. & King, R.B. (2014). Direct Effects of an Invasive European Buckthorn Metabolite on Embryo Survival and Development in *Xenopus laevis* and *Pseudacris triseriata*. *Journal of Herpetology*, 48(1), 51-58. <https://doi.org/10.1670/12-066>
- Sasaki, K., Lesbarrères, D., Watson, G., & Litzgus, J. (2015). Mining-caused changes to habitat structure affect amphibian and reptile population ecology more than metal pollution. *Ecological Applications*, 25(8), 2240–2254. <https://doi.org/10.1890/14-1418.1>
- Salafsky, N., Relton, C., Young, B.E., Lamarre, P., Bohm, M., et al. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*. 39(3). <https://doi.org/10.1111/cobi.14434>
- Schmidt, B. R. & Zumbach, S. (2008). Amphibian road mortality and how to prevent it: a review. In J. C. Mitchell, R. E. Jung Brown, & B. Bartholomew (Eds.), *Urban Herpetology* (pp. 157–167). St. Louis, Missouri: Society for the Study of Amphibians and Reptiles. <https://doi.org/10.5167/uzh-10142>
- Schottler, S. P., Ulrich, J., Belmont, P., Moore, R., Lauer, J. W., Engstrom, D. R., & Almendinger, J. E. (2013). Twentieth century agricultural drainage creates more erosive rivers. *Hydrological Processes*, 28(4), 1951–1961. <https://doi.org/10.1002/hyp.9738>
- Schulz, R., Bundschuh, M., Gergs, R., Brühl, C. A., Diehl, D., et al. (2015). Review on environmental alterations propagating from aquatic to terrestrial ecosystems. *The Science of the Total Environment*, 538, 246–261. <https://doi.org/10.1016/j.scitotenv.2015.08.038>
- Semlitsch, R. D., & Bodie, J. R. (2003). Biological Criteria for Buffer Zones around Wetlands and Riparian Habitats for Amphibians and Reptiles. *Conservation Biology*, 17(5), 1219–1228. <https://doi.org/10.1046/j.1523-1739.2003.02177.x>
- Sheridan, J. A., & Bickford, D. (2011). Shrinking body size as an ecological response to climate change. *Nature Climate Change*, 1(8), 401–406. <https://doi.org/10.1038/nclimate1259>
- Souza, K. S., Fortunato, D. S., Jardim, L., Terribile, L. C., Lima-Ribeiro, M. S., et al. (2023). Evolutionary rescue and geographic range shifts under climate change for global amphibians. *Frontiers in Ecology and Evolution*, 11. <https://doi.org/10.3389/fevo.2023.1038018>
- Thompson, C.M. (2022). Carryover Effects of Hydroperiod Length, Neonicotinoid Pesticide Exposure, and Predation Risk in a Pond-Breeding Amphibian. Ohio University. https://etd.ohiolink.edu/acprod/odb_etd/ws/send_file/send?accession=ohiou1660314558287638&disposition=inline
- Tornabene, B.J., Smalling, K.L. & Hossack, B.R. (2024). Effects of Harmful Algal Blooms on Amphibians and Reptiles are Under-Reported and Under-Represented. *Environmental Toxicology and Chemistry*, 43(9), 1936-1949. <https://academic.oup.com/etc/article-pdf/43/9/1936/60762915/etc5941.pdf>
- U.S. Geological Survey. (2001). Malformed frogs in Minnesota: an update. <https://pubs.usgs.gov/fs/fs-043-01/pdf/fs-043-01.pdf>

Uden, D. R., Hellman, M. L., Angeler, D. G., & Allen, C. R. (2014). The role of reserves and anthropogenic habitats for functional connectivity and resilience of ephemeral wetlands. *Ecological Applications*, 24(7), 1569–1582. <https://doi.org/10.1890/13-1755.1>

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Youngquist, M. B., & Boone, M. D. (2014). Movement of amphibians through agricultural landscapes: The role of habitat on edge permeability. *Biological Conservation*, 175, 148–155. <https://doi.org/10.1016/j.biocon.2014.04.028>

Zaffaroni-Caorsi, V., Both, C., Márquez, R., Llusia, D., Narins, P., Debon, M., & Borges-Martins, M. (2022). Effects of anthropogenic noise on anuran amphibians. *Bioacoustics*, 32(1), 90–120. <https://doi.org/10.1080/09524622.2022.2070543>

Birds

Overview

Birds are perhaps the most popular and intensively studied animals in Minnesota and throughout the world. Most species are easily visible, display beautiful plumages, and communicate with unique calls and songs. From Native Americans who first occupied North America to the European settlers who arrived, people have observed, listened, consumed, collected, studied, and written about birds for hundreds of years. Indeed, Minnesota is fortunate to have a spectacular legacy of information regarding our avifauna, beginning with the oral legends of Native Americans (e.g. Adikwan, 2019) to articles, pamphlets, peer-reviewed papers, and numerous books describing their life history, distribution, and abundance.

Minnesota is fortunate to have the recent [Minnesota Breeding Bird Atlas](#), a project led by Audubon Minnesota, which documents the breeding status and locations of all birds nesting in Minnesota at a local scale and provides in-depth species accounts (Pfanmuller et al., 2017). Earlier foundational works include the two-volume tome by T.S. Roberts in 1932, followed with books by Green and Janssen (1975), Janssen (1987; 2019), and Pfanmuller, Niemi, and Green (2024).

Multiple other accounts, projects, journal articles, and publications by resource agencies and the [Minnesota Ornithologists' Union](#) supplement these sources and provide a wealth of information on the 452 species that have been reported in the state (MOU, 2024). These include the [Blueprint for Minnesota Bird Conservation](#), the [North American Bird Conservation Initiative](#), the [Minnesota National Forest Breeding Bird Monitoring Program](#), and the [North American Breeding Bird Survey \(BBS\)](#). The latter has monitored the population trends of North American birds since 1967 providing an abundance of valuable data on the status of all birds for more than 50 years. This information is now augmented by [eBird](#) data collected by recreational birders and compiled by the Cornell Laboratory of Ornithology, which has recently produced information for state-level [eBird Status and Trends](#). Some birds, such as colonial nesting waterbirds, secretive marshbirds, raptors, and nocturnal birds will likely still require additional efforts to track their numbers. Several of Minnesota's SGCN fall in these categories, such as the Franklin's gull, black-crowned night-heron, and Eastern whip-poor-will. It is important to find cost-effective and efficient means to monitor these species. It is important that monitoring efforts of all these types are supported throughout the future.

Given the depth of attention they have received over the years, our knowledge of the threats and challenges Minnesota birds face are well-documented. The threats are numerous but three stand out as primary concerns: 1) the significant population declines of many species that occur in a variety of habitats, from the Eastern screech owl (*Megascops asio*) and Eastern whip-poor-will (*Antrostomus vociferus*) to the Connecticut warbler (*Oporornis agilis*) and [purple martin](#) (*Progne subis*); 2) the especially steep decline of grassland species such as [marbled godwit](#) (*Limosa fedoa*), bobolink (*Dolichonyx oryzivorus*), and grasshopper sparrow (*Ammodramus savannarum*); and 3) the predicted stress of a changing climate and even potential extirpation of many of our northern boreal forest species as temperatures continue to rise ([National Audubon Society 2019](#); see case study later in this chapter). Despite these pervasive threats, there are remarkable success stories. Due largely to targeted conservation efforts some species, such as the [Trumpeter Swan](#) (*Cygnus buccinator*) (see inset), [Peregrine Falcon](#) (*Falco peregrinus*), and [Bald Eagle](#) (*Haliaeetus leucocephalus*), have recovered from the brink of state extirpation. It is not uncommon, for example, to sight a Bald Eagle not only in

the dense forests of northern Minnesota, but also in the center of the Twin Cities or in the agricultural landscapes across southern and western Minnesota.

SGCN and SNI Summary Information

The 2025-2035 list of Species in Greatest Conservation Need identified 79 Minnesota bird species as SGCN and one species, the Eastern Screech Owl, as a SNI (Species in Need of Information). A total of 21 species were removed from the SGCN list in the 2015-2025 SWAP primarily because their numbers have increased and/or they are now known to be more common and widespread than biologists thought in 2015 (see for example Case Study box below). Eight species were added: Canvasback (*Aythya calisineria*), Bufflehead (*Bucephala albeola*), Buff-breasted Sandpiper (*Calidris subruficollis*), Solitary Sandpiper (*Tringa solitaria*), Great Gray Owl (*Strix nebulosa*), American Barn Owl (*Tyto furcata*), American Three-toed Woodpecker (*Picoides dorsalis*), and Swainson's Thrush (*Catharus ustulatus*)

Among the 79 Minnesota birds that are designated SGCN, the [piping plover \(*Charadrius melodus*\)](#) is federally endangered, the red knot (*Calidris canutus*) is federally threatened, and the golden-winged warbler (*Vermivora chrysoptera*) is under review for federal listing. Nine species are state listed as Endangered, 2 Threatened, and 19 are listed as Special Concern. A complete list of both SGCN and SNI species can be found in Appendix B.

Case Study: The Successful Recovery of the Trumpeter Swan in Minnesota

When Roberts compiled his treatise on Minnesota birds in 1932, he wrote, "The days of the Trumpeter Swan as a bird of Minnesota have long since passed." Unfortunately, this large white bird was an easy target for the state's early European settlers and market hunters. The young birds were tender and tasty while the older birds were either shot for sport or for their skin and feathers. By 1932 only 69 birds were believed to remain, surviving in the remote mountain valleys of Montana, Idaho, and Wyoming. These birds, along with those from a breeding population found in Alaska in the 1950s, became breeding stock for introductions farther east, in areas formerly occupied by the species, including the LaCreek National Wildlife Refuge in western South Dakota.

Interest in reintroducing Trumpeter Swans to Minnesota began in 1966 by the Hennepin County Park Reserve District (now Three Rivers Park District). Their initial efforts to establish a breeding population by releasing adult birds secured from Montana failed. The tide turned when the district initiated its own captive breeding program, caring for the young cygnets and not releasing them until they were two years old. The first release of young raised by this program occurred in 1978, and the first successful nesting occurred in 1979. By 1996, more than 245 young birds had been raised and were successfully nesting within the park district and beyond. The park district's program continued for another thirteen years, until it was retired in 2009. In 1982, the DNR's Nongame Wildlife Program outlined an initiative to build on the work of the park district by reintroducing birds in northwestern, north-central, and south-central Minnesota, working collaboratively with state and federal agencies, provinces, universities, tribes, and the Trumpeter Swan Society. Together, these two initiatives by the Park District and DNR were so successful that by 2019, the Minnesota Waterfowl Breeding Population Survey generated a statewide population estimate of 23,200 swans. It is now common to find breeding pairs of Trumpeter Swans in wetlands across the state. *Excerpted from Pfannmuller et al., 2024.*

Habitat Associations

SGCN birds rely upon many different habitats across Minnesota. Table 2.4 provides a summary of their associations with the 15 habitats discussed in the 2025-2035 SWAP. Three habitats that provide primary habitat for the largest number of SGCN are prairie and other grasslands (22 SGCN), followed by non-forested wetlands (18 SGCN), and lakes (16 SGCN). Collectively, forested habitats (upland, lowland, and riparian) provide primary habitat for 27 bird SGCN. Primary habitats are those where the species is found most consistently; the species is most reliant on the habitat and, should the habitat be degraded or lost, it would have the greatest effect on the species. Secondary habitats are those where a species can be found some of the time, but less frequently than a primary habitat. Some species, especially those with more general habitat requirements, may be assigned to multiple primary or secondary habitats. A complete list of all primary and secondary habitats for each SGCN can be found in Appendix C.

Table 2.4. Habitat Associations for Bird SGCN.

Habitats	Primary Habitat	Secondary Habitat	Total
Prairie and Other Grasslands	22	7	29
Savanna	3	4	7
Upland Conifer Forest and Woodland	7	9	16
Upland Deciduous Forest and Woodland	3	9	12
Mesic Hardwood Forest	5	2	7
Deciduous Wet Forest	0	4	4
Coniferous Forest Wetland	9	3	12
Riparian and Floodplain Forest	3	3	6
Non-forested Wetlands	18	12	30
Rivers and Streams	1	0	1
Lakes	16	6	22
Vernal Pools	0	1	1
Cliff, Talus, and Rock Outcrops	4	0	4
Urban and Other Developed Lands	3	6	9

Primary Stressors for Birds

Stressors are factors that pose direct or indirect challenges to vulnerable plant and wildlife species. Habitat loss and degradation are generally considered the primary threat for most SGCN, and some stressors reduce the quality of those habitats. Other stressors may operate more directly on the SGCN, such as disease and pathogens. In some cases, stressors may operate both directly and indirectly. For example, roads, trails, and railroads can directly harm SGCN through vehicular strikes, or operate indirectly through habitat fragmentation and reduced landscape connectivity. The stressor list below is adapted from an internationally recognized threats classification developed by the International Union for the Conservation of Nature (Salafsky et al.,

2024). For more information, see the “Stressors” section of Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts (e.g. catastrophic wildfire following a prolonged period of fire suppression) while serving as valuable conservation tools in others (e.g., appropriately planned and applied prescribed fire).

Information about a subset of primary stressors specifically impacting this species group is included below, followed by a set of conservation actions addressing those stressors.



Development

Collisions with towers and buildings, especially windows, are a well-documented threat to bird populations (e.g. Loss et al., 2015; Loss et al., 2014). Birds often mistake reflective glass that mirrors vegetation or sky as open habitat, leading them to fly directly into it, often resulting in injury or death. Long-distance migratory species are particularly vulnerable to these collisions (Arnold & Zink, 2011; Machtans et al., 2013). Among the 10 migrants that were disproportionately vulnerable to building collisions was the Minnesota SGCN, the blue-throated warbler (*Setophaga caerulescens*; Runwal, 2020). Communication towers also pose a significant hazard, especially during migration on foggy nights or when cloud ceilings are low (Loss et al., 2015).

Birds that nest or feed along lake and river shorelines are especially vulnerable to the increasing rate of shoreline development and habitat degradation. These areas are considered prime real estate for summer homes and recreational use, leading to widespread alteration. A recent report estimates that Minnesota has lost 40-50% of its natural shoreline and continues to lose an additional 1 to 2% per decade (Minnesota Natural Shoreline Partnership, 2023). This ongoing loss impacts species like the common loon (*Gavia immer*) that requires undisturbed shorelines for nesting. Additionally, during migration, many migrants depend on shoreline habitats rich in insects to refuel, making these areas essential stopover habitat.

The loss and degradation of forests, especially across the intensively developed Eastern Broadleaf Forest Province, is a notable factor in the decline of forest-dependent bird species such as the [cerulean warbler](#) (*Setophaga cerulea*), [Acadian flycatcher](#) (*Empidonax vireescens*), and eastern whip-poor-will (Pfannmuller et al., 2024).

In addition, unsupervised domestic and feral cats can pose a serious threat to local bird populations. Annual bird mortality caused by cats is estimated at 1.4 to 4 billion birds in the U.S. alone, though these figures include significant uncertainties (Loss et al., 2013a). Feral cats in particular may be a greater threat than free-roaming domestic cats (Loss et al. 2015; 2022).



Crop Production

Grassland birds, including sedge wrens (*Cistothorus stellaris*), bobolinks, grasshopper sparrows, and [short-eared owls](#) (*Asio flammeus*) are affected by loss of habitat caused by conversion to rowcrop agriculture on the landscape. Since 1970, birds dependent on grassland habitats have declined an

average of 43% as the Great Plains continues to lose approximately 1 to 2 million acres of grassland per year ([State of the Birds Report](#)).

In addition to habitat loss, extensive tiling of agricultural lands increases runoff to wetlands embedded in the agricultural landscape. Spring and summer storms increase runoff, which can suddenly flood the nests of waterfowl, shorebirds, and rails. Over time, the resulting sediment load can accumulate and gradually fill in wetlands (see also Dams and Water Management stressor below).



Tree Plantations

Tree plantations typically support lower plant species diversity and reduced structural complexity, resulting in lower-quality habitat for forest-dependent birds. The lack of a well-developed understory, for example, can significantly reduce the suite of forest bird species, often reducing it to 6 to 15 species that primarily occupy the conifer canopy. In contrast, plantations that include deciduous shrubs and saplings may support an average of 20 to 35 bird species (Green, 1995).



Livestock Management

Grazing by large ungulates, especially bison, was historically an important natural disturbance that helped maintain grassland habitats and the bird species that depend on them. Bison played a vital role in sustaining prairie ecosystems through selective grazing, wallowing behavior that disturbed soil and created new microhabitats (Ratajczak et al., 2022), and dung that attracted insects, providing a key food source for many grassland birds. Today, livestock have replaced bison as the dominant ecological force shaping the structure and composition of Minnesota's one million acres of pastures and hayfields (USDA, 2022).

Depending on how grazing is managed, it can have both negative and positive impacts on the abundance and diversity of grassland birds. Intensive grazing often has harmful consequences, including the reduction of vegetative cover needed by ground-nesting species, which increases the risk of nest predation; trampling of nests by livestock; and the spread of exotic plant species that degrade native vegetation used for feeding and nesting (Wuerthner, 2012). Bird abundance is especially sensitive to grazing in riparian habitats (Barzan et al., 2021). Grassland species known to respond negatively to intensive grazing include northern harriers (*Circus hudsonius*), short-eared owls, and [Henslow's sparrow \(*Centronyx henslowii*\)](#) (Bock et al., 1993). Some studies have also found that cattle grazing may be more detrimental to bird populations than grazing by sheep or mixed livestock herds (Barzan et al., 2021).

Livestock grazing can also benefit grassland conservation by creating favorable conditions for certain grassland bird SGCN species. For example, moderate to heavy grazing that produces short-grass prairies or grasslands can support species like the [chestnut-collared Longspur \(*Calcarius ornatus*\)](#) (Lebbin & Fenwick, 2010). Indeed, for many years, the only site in Minnesota where longspurs were regularly observed was the B-B Cattle Ranch in Clay County. However, very few grassland bird species in Minnesota's tallgrass prairie ecosystem are strictly dependent on short-grass conditions. In these landscapes, the absence of grazing can actually reduce habitat suitability for birds. Moderate grazing helps maintain the vegetative heterogeneity and structural diversity that many bird species, such as grasshopper sparrows and western meadowlarks (*Sturnella neglecta*), thrive upon (Barzan et al., 2021). Indeed, a recent study concluded that rotational cattle grazing provides similar benefits to breeding birds as bison grazing. The authors concluded that the rotational cattle or bison grazing system used was less important to grassland bird diversity and abundance than other factors such as stocking rate, soil

composition, precipitation, topography and woody vegetation (Boyce et al., 2021). Grazing on private "working lands" also helps to maintain grassland habitats and prevents conversion to other land uses.



Mining and Quarrying

The loss of habitat is the primary impact of mining and quarrying operations. This has been especially evident along the former Lake Agassiz beach ridges in northwestern Minnesota. The beach ridges not only support some of the best and largest remnants of native prairie in the state that provide habitat for many declining grassland birds, they also are a significant source of sand and gravel to growing communities in the Red River Valley (DNR, 1997). Potential impacts to birds in sensitive wetlands from proposed sulfide mines in northeastern Minnesota also have been raised as a concern (Breining, 2014).



Wind and Solar Energy Infrastructure

Renewable energy site development can negatively impact birds through habitat degradation and direct collision (also see discussion of window and tower collisions under the Development stressor). Loss and his colleagues (2013b) estimated between 140,000 and 328,000 birds are killed annually at wind turbines across the contiguous United States. Based on data collected at wind turbines supported by a single vertical tower they demonstrated that mortality increases with the height of the turbine hub. Mortality also differed geographically with the lowest risk in the Great Plains.



Roads, Trails, and Railroads

Bird species that depend on large tracts of forest that are not fragmented (e.g. [American goshawk](#) (*Accipiter atricapillus*) and [red-shouldered hawk](#) (*Buteo lineatus*)) can be impacted by creation of new roads which decrease the amount of unfragmented forest and increase the level of disturbance in nesting areas, especially during the breeding season. Ortega and Capen (2002) found that habitat quality for a suite of forest-interior songbirds may be higher when territories are located more than 150 m from a road.

Vehicle collisions pose another threat to birds that use roadside fence lines, utility lines, and poles as perches. Car collisions have been noted as a significant cause of mortality for species such as the [loggerhead shrike](#) (*Lanius ludovicianus*) and red-headed woodpecker (*Melanerpes erythrocephalus*) (Pfanmuller et al., 2024).



Utility Corridors

Grassland bird species and forest interior songbird species can be impacted by the creation and maintenance of utility and service corridors that traverse large tracts of habitat. Mortality can also result from collision with the lines. Powerlines, for example, present a risk to waterfowl when situated between feeding and roosting areas. This is especially true for lines near lakes, rivers and wetlands. Less agile birds, such as [American white pelicans](#) (*Pelecanus erythrorhynchos*), are particularly susceptible to collision with overhead lines located near waterbodies (APLIC, 2012). Mortality can result from the collision with wires or from electrocution from both wires and poles (Loss et al., 2015).

Pipelines used to transport oil and gas fragment habitats across the state and pose a risk of spills, which can lead to water-borne pollution.



Hunting and Collecting Animals

Lead poisoning in birds, resulting from the consumption of lead shot, is well-documented (Tranel & Kimmel, 2009; Haig et al., 2014; Pain et al., 2019). It can result from ingesting spent lead shot or from ingesting food that contains lead shot. Among bird species documented with lead poisoning in Minnesota are bald eagles and trumpeter swans. Data from the Raptor Rehabilitation Center at the University of Minnesota document that over 150 bald eagles are admitted each year. Between 80-90% of these birds have some level of lead in their blood; 25-30% have lead toxicity and either die or are euthanized. Many more bird species are susceptible to the impacts of lead poisoning by consuming lead shot, including many SGCN. A few examples include bufflehead (*Bucephala albeola*), canvasback (*Aythya valisineria*), Northern pintail (*Anas acuta*), Northern harrier (*Circus hudsonius*) and American goshawk (Kimmel & Tranel, 2007; Pain et al., 2009). Lead shot use has been banned in some areas in Minnesota, including federal Waterfowl Protection Areas, National Wildlife Refuges, state Scientific and Natural Areas (SNA) that regularly allow hunting, and in state parks and SNAs that hold special hunts. Lead shot may be used in all other areas, including most state Wildlife Management Areas.

Although falconry is also tightly regulated by Minnesota rules and federal regulations, concerns remain regarding the illegal take of young raptors from nests.



Timber harvest

Forest-dependent bird species are impacted by timber harvest that alters their habitat, including woodpeckers, owls and hawks, grouse, and warblers. Timber harvest alters habitat at a landscape-scale such as changing forest composition, age classes, patch sizes, and causing habitat fragmentation. At a site-level, timber harvest can change the plant species and structural composition of a stand. Changes that increase the amount of young forest (approximately 20 years old) benefit some species including song sparrows, indigo buntings, American goldfinches, chestnut-sided warblers, and veerys. Rarely are these Species of Greatest Conservation Need, although golden-winged warblers (*Vermivora chrysoptera*) may be found in young aspen stands. Changes that decrease the amount of older forests (i.e., stands that have more complexity in their structural layers and more diversity in the forest canopy) negatively impacts forest birds dependent on these forest components, including such SGCN as the American goshawk, boreal chickadee (*Poecile hudsonicus*), Swainson's thrush, and black-throated blue warbler.

Numerous studies document the relationship between forest cover and bird distribution and abundance in Minnesota forests. The most comprehensive studies have been conducted by the [University of Minnesota's Natural Resources Research Institute's Avian Ecology Lab](#) (e.g. Niemi et al., 2016). Fortunately, most logging occurs during the winter months to reduce impacts to roads and trails. Winter harvesting however, can impact resident species that often begin nesting in late winter, such as the American goshawk and great gray owl. When spring and summer timber harvests occur they can destroy nests, kill nestlings, and cause birds to abandon their territories. Although the amount of timber harvested during the winter season increased during the 1990s, in 1996 the total volume harvested in all other seasons of the year was 45%, including 21% in summer (Puettmann and Ek, 1999). Regardless of the time of year harvesting occurs, rare species need to be considered when silvicultural practices are implemented at either the site or landscape level.



Fishing

Waterbirds and fish-eating raptors can all be impacted by lead fishing tackle, and by fishing lines left behind by anglers. The impacts of lead sinkers and jigs have been most extensively studied in loons and swans but also documented in multiple waterfowl species, colonial waterbirds, bald eagles and osprey. In addition to common loons, other SGCN species that have been documented ingesting lead sinkers include the American black duck (*Anas rubripes*) and black-crowned night-heron (*Nycticorax nycticorax*) (Rattner et al., 2008; Grade et al., 2018; Slabe et al., 2019).

Fish consumption by some waterbirds is often perceived by the angling public to be a threat to local fish populations. Although double-crested cormorants (*Nannopterum auritum*) are the most commonly perceived threat, one SGCN, the American white pelican, is also frequently vilified. Field studies, however, have shown that pelicans forage primarily on rough forage fish that have little economic value.



Recreation

Nature-based recreational activities, such as cycling, hiking, and boating, can negatively impact birds. Reproductive success may decline, behaviors may change, and population numbers may decrease (Steven et al., 2011). Wave actions from recreational boats, for example, can disrupt nesting areas for overwater and shoreline nesting species while high levels of boating traffic also can disrupt waterbird and shorebird nesting activity. Recreational vehicles can cause nest abandonment, nest desertion, and reduced nest survival (Barton & Holmes, 2006; Spaul & Heath, 2016). Human activity near or within colonial waterbird nesting sites may result in colony abandonment or reduced nesting success, especially when events occur early in the nesting season (Vennesland, 2010). Ultimately, the level of impact that recreational activities cause depends on factors such as their proximity, duration, intensity, frequency, visual screening, and time of the year as well as individual species sensitivity and habituation.



Fire Management

Many forest birds utilize large open habitats created by recent fires; some would not be present in Minnesota's dense northern forests without the presence of fire or other large disturbances (such as logging and windstorms). A field study of the response of forest birds to the Little Sioux fire on the Superior National Forest reported a variety of species attracted to the 15,000 acre burn including the killdeer (*Charadrius vociferus*), Northern flicker (*Colaptes auratus*), Eastern bluebird (*Sialia sialis*), and indigo bunting (*Passerina cyanea*) and three SGCN including the black-backed woodpecker (*Picoides arcticus*), red-headed woodpecker and American kestrel (*Falco sparverius*), the latter two being more common in agricultural regions farther south in Minnesota (Niemi, 1978).

Disturbance (fire or grazing) is critical to maintaining suitable habitat for many grassland birds; without disturbance grasslands become shrublands and forests. Although grassland bird species each require vegetative cover that may differ in composition, density, and height at different times of year, they each depend on some level of active management, from [greater prairie-chickens](#) (*Tympanuchus cupido*) and upland sandpipers (*Bartramia longicauda*) to western meadowlarks and grasshopper sparrows (Wilsey et al., 2019).



Dams and Water Management

Wetland/marsh birds, shorebirds, and waterbirds are all impacted by changes in water management regimes. Dams, for example, will permanently flood seasonal wetlands that provide nesting, feeding and resting areas, eliminate the periodicity of floods and droughts that may provide critical signals to birds, and destroy the connectivity of associated wetland habitats in a large wetland complex (Abreu et al., 2020). Stream channelization projects can impact avian habitats by increasing downstream sedimentation and bank erosion and decreasing in-stream riffles and pools and stream-side vegetation. For example, habitat improvement projects that stabilize eroding stream banks, can eliminate microhabitat features that are critical to [Louisiana waterthrushes \(*Parkesia motacilla*\)](#) in southeastern Minnesota (Stucker & Cuthbert, 2000).

The installation of underground tiles and the construction of ditches to remove water from poorly drained soils provide many benefits to agricultural production but also cause significant environment impacts. With approximately 19,150 miles of drainage ditches and an extensive, untallied system of subsurface tiles ([Minnesota Board of Soil and Water Resources](#)), the hydrology, water quality, and abundance of wetland habitats across much of Minnesota's original prairie landscape have been negatively impacted ([University of Minnesota Extension](#)). Removing water from poorly drained soils not only results in the loss of wetland habitats, including wet meadows, that provide habitat for LeConte's sparrows (*Ammodramus leconteii*), [yellow rails \(*Coturnicops noveboracensis*\)](#), and sedge wrens, but can also cause higher flows into streams after high rainfall events. For example, decades ago, extensive tiling within the Heron Lake Watershed in southwest Minnesota caused untimely surges in Heron Lake's water level during the nesting season, ultimately leading to the demise of many waterbird nesting colonies and waterfowl nesting populations. Fortunately, many best management practices and water management options now are available to help mitigate many of these impacts.



Invasive Species (Problematic Non-native Species)

Grassland species are negatively impacted by the spread of non-native plant species in their habitat, including common buckthorn (*Rhamnus cathartica*), glossy buckthorn (*Frangula alnus*), and invasive honeysuckle species (*Lonicera spp.*). Although many studies investigating the impacts of common non-native shrubs such as buckthorn (*Rhamnus cathartica* and *Frangula alnus*), and non-native honeysuckles (e.g. *Lonicera tatarica*) have been equivocal, others have pointed to "altered nest site availability, habitat heterogeneity, and food supplies" (Nelson et al., 2017) and increased predation (Schmidt & Whelan, 1999). The increasing prevalence of garlic mustard (*Alliaria petiolata*) in riparian and floodplain forests decreases the diversity of native flora, instead creating a dense carpet of herbaceous vegetation and potentially impacting riparian species like the Louisiana waterthrush (Stucker pers. comm.). Thick stands of purple loosestrife (*Lythrum salicaria*) can similarly impact wetland SGCN such as black terns and least bitterns. Many of the most widespread species, such as smooth brome (*Bromus inermis*) and Kentucky blue grass (*Poa pratensis*), can outcompete native vegetation resulting in a decline in vegetative diversity. Although some grassland birds will use these habitats, studies have shown that "non-native vegetation cover is correlated with reduced nesting success and post-fledging survival" (Bernath-Plaisted et al., 2023). One hypothesis for the observed decline is that non-native plants attract fewer insects (Narango et al., 2018).

Ground nesting forest birds are negatively impacted by non-native, European earthworms used as bait for fishing (Loss & Blair, 2011). The worms' consumption of leaf litter and other organic materials reduces the amount of ground cover and plant diversity, increasing the vulnerability of ground-nesting species.

Several non-native invasive forest insects and diseases can cause defoliation and mortality. Currently the most significant threat is from the elm ash borer (*Agrilus planipennis*), which has impacted 6,100 acres in central Minnesota. Like many forest disturbances, some birds will benefit from the increase in dead trees (such as woodpeckers; Koenig et al., 2013) while others that depend on a healthy tree canopy may decline. Given the current distribution of [emerald ash borer](#) in Minnesota this could impact such SGCN as the cerulean warbler and Acadian flycatcher. Neither species is entirely dependent on ash but it is a canopy tree in many of the forests they inhabit. The fungal disease oak wilt (*Bretziella fagacearum*) is now common throughout southeastern and east-central Minnesota. Because oaks decay very slowly, the dead oaks will provide benefits to some birds for many years. Ultimately, however, the loss of oaks will result in the absence of acorns, an important food source for many birds, such as wood ducks, turkeys and blue jays and the SGCN, red-headed woodpeckers who are especially dependent on hard mast during the winter months (Frei et al., 2020).

Waterbird and marshbird nesting habitat also can be diminished by the spread of non-native vegetation. Non-native species such as common reed (*Phragmites australis* subsp. *australis*), can crowd out native vegetation and reduce accessibility and availability of open water habitat, especially for birds that utilize short wetland grasses (Benoit & Askins, 1999).

Two birds are officially classified as invasive species in Minnesota, the European starling (*Sturnus vulgaris*) and the Mute Swan (*Cygnus olor*). Starlings can cause damage to agricultural crops and aggressively compete with native birds for nesting cavities. Mute swans are also fiercely competitive with other waterbirds, including trumpeter swans, aggressive to people, and consume huge amounts of native aquatic vegetation daily (Pfannmuller et al., 2024).



Problematic Native Species

In Minnesota, open sand and gravel beaches are a rare resource so species that share that habitat are often in competition for nest sites. Common terns, a state threatened species, compete for these sites with ring-billed gulls (*Larus delawarensis*) and ground-nesting double-crested cormorants. Because adult gulls and cormorants arrive on the breeding grounds weeks before the terns, the prime nesting sites are already occupied. Intensive management efforts are underway at most Minnesota's tern breeding colonies to deter the gulls and cormorants (Pfannmuller et al., 2024).

Three native forest insects can impact large tracts of songbird breeding habitat in northern Minnesota. The eastern larch beetle (*Dendroctonus simplex*) has caused the death of mature tamaracks in more than 1.1 million acres since 2001. Many birds that are found in conifer dominated peatlands are likely impacted by this loss. The forest tent caterpillar (*Malacosoma disstria*) and spruce budworm (*Choristoneura fumiferana*) can also impact broad swaths of forest land, but their negative impacts are more short-lived. Some birds positively respond to this sudden increase in insect populations. Cuckoos are best known for their voracious appetite for tent caterpillars, but the list of avian predators includes at least two SGCN, evening grosbeaks (*Coccothraustes vespertinus*) and Philadelphia vireos (*Vireo philadelphicus*), while several warblers, including the SGCN warblers, bay-breasted (*Setophaga castanea*) and Cape May (*Setophaga tigrina*), are well-known consumers of spruce budworm.



Diseases and Pathogens

Birds are subject to a variety of diseases and pathogens that cause disease. Many birds naturally carry these diseases while showing few, if any, of the symptoms until they are stressed and lose their resilience. Many of the stressors listed above, such as problematic non-native species and habitat loss that causes overcrowding on migratory stopovers, can instigate an outbreak. Bird flu, caused by the H5N1 virus, has resulted in the death of at least 11,000 wild birds nationally since 2022 (Yates, 2025), with die-offs most pronounced in waterfowl. The virus's impact on Minnesota SGCN is unknown. It has generated widespread concern because of its impact on both high-density domestic poultry operations and backyard flocks. In just a two-month period, from December 2024 through January 2025, 41.4 million domesticated chickens were culled nationwide because of the flu.

West Nile Virus can affect a number of bird species but members of the Corvidae bird family are especially susceptible. Transmission was so high in the eastern United States in the early 2000s that some local populations of American crows declined nearly 70% (Reed et al., 2009); although not as severe, declines were also noted in Minnesota. West Nile Virus has been documented in 250 bird species including the SGCN American goshawk, Northern harrier and Cape May warbler (Taieb et al., 2020). The virulent Newcastle disease has been reported in a number of bird species, including the SGCN the American white pelican (Wisconsin DNR), but double-crested cormorants and pigeons appear to be most susceptible. Without precautions they can spread the disease to poultry farms where chickens are considered the most vulnerable. The most recent outbreak in Minnesota was in 1992 when more than 20,000 cormorants collectively died in North Dakota, South Dakota and Minnesota (DNR, 2025). Pelicans commonly co-nest with cormorants and were likely part of the loss documented in 1992. Another pathogen, avian botulism, occurs world-wide and affects many birds but predominantly waterfowl and other species common to wetland habitats. Shallow waters with low oxygen levels are most susceptible. It is not spread from bird to bird but by the consumption of aquatic invertebrates that carry the toxin or by birds that filter bottom sediments or organic materials where the toxin is embedded. Ultimately resulting in the bird's paralysis, thousands, often millions, of birds succumb to the disease each year (Gutierrez-Arnal & Marin, 2024). Botulism outbreaks are expected to only increase in frequency and magnitude with warming temperatures (Lafrancois et al., 2011).



Water-borne Pollution

Many water-borne pollutants that have been banned or strongly regulated for years, such as PCBs (polychlorinated biphenyls), dioxins and dibenzofurans, are persistent chemicals that have remained in the environment for decades. Their continued presence in fish-eating birds in the Great Lakes Basin was recently documented and included evidence of reduced reproductive success, physical deformities, and compromised immune function (Grasman et al., 2025). Other sources of water pollutants including "wastewater, heavy metals, chemicals, plastics, water from agricultural fields, and oil spills are considered among the most important factors responsible for mortality and population declines in freshwater birds (Shahid & Nadeem, 2021). One impactful water pollution event occurred in 2010 when the Deepwater Horizon oil rig exploded in the Gulf of Mexico, an area of coastal waters where at least one Minnesota SGCN, the common loon, winters (see Minnesota Common Loon Restoration Project Case Study in the Lakes Chapter). New concerns also are being raised about the impacts of PFAs (perfluoroalkyls), brominated flame retardants and neonicotinoids and their persistence in aquatic environments.



Air-borne Pollutants

Mercury's impact on fish-eating birds (waterfowl, shorebirds, loons, and wading birds) has been well-documented (Seewagen, 2010). Airborne mercury enters aquatic ecosystems primarily through atmospheric deposition, where it settles onto land and water bodies via rain, snow, or dry particles, and is carried into water bodies by rain events. Deposition of mercury on soil and foliage also impacts terrestrial species. Recent studies that examined approximately 50% of the songbirds in the U.S. and Canada documented reproductive problems attributed to mercury exposure in 41% of the species (Evers & Cristol, 2020). The risks are highest among birds that consume predatory invertebrates such as spiders.

Among grassland birds, the major population declines that began in the 1960s were coincident with a period of agricultural intensification across the Great Plains, including heavy pesticide use (Stanton et al., 2018). Studies have consistently demonstrated the negative impacts of aerially applied pesticides on birds, including mortality, reduced reproductive success, and disrupted migratory behavior (Bernath-Plaisted et al., 2023). Carbamates and organophosphates have been the most widely studied pesticides. As their use has decreased there is growing concern about the use of neonicotinoids (Stanton et al., 2018). Studies conducted to date point to both lethal and sublethal impacts (e.g. declines in reproductive success, growth, chick survival, etc.) from ingesting seeds treated with neonicotinoids (Wood & Goulson, 2017).



Light and Noise Pollution

Nearly half of the United States land surface has light-polluted skies (Falchi et al., 2016). Sources of light pollution at night include interior and exterior lights on homes, buildings, billboards, parking lots, and streets as well as lighted sports fields, industrial plants, and airports (DarkSky International, 2025). Studies also point to the ecological impacts created during the daytime by polarized light that reflects off surfaces such as asphalt roads, glass panes, and solar panels (Chock et al., 2020; Horvath et al., 2009). Because most of the nation's large urban areas (with multiple artificial light sources every night), are embedded in the Temperate Deciduous Forest biome, biota occupying temperate forests are the most vulnerable to these impacts (Aubrecht et al., 2010).

Impacts of light pollution include disrupting the behavior of amphibians whose breeding calls are delivered at night, drawing nocturnal insects to artificial lights that increase their predation risk, and disorienting birds that use environmental cues to guide their nighttime migration (DarkSky International, 2025). Bright day and dark night cycles synchronize the internal clocks of plants and animals, and alterations to these cycles can impact hormone production, reproductive behavior, neural activity, and metabolic functions (Bumgarner & Nelson, 2021).

Noise pollution from aircraft, vehicles, boats, and other human causes is pervasive. Anthropogenic noise was detected in 36% of national parks surveyed (Buxton et al., 2019), and 12% of wilderness areas had anthropogenic noise levels 3 decibels higher than levels predicted to occur naturally (Buxton et al., 2017). Exposure to increasing noise levels can change the spatial distribution and movement patterns of wildlife, cause avoidance of feeding and nesting areas, and interrupt sleeping patterns and communications (Kok et al., 2023).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce, 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al., 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

Many bird species will be affected by habitat shifts and alterations due to warming temperatures and the accompanying impacts. Warmer winters may change the diversity, distribution and abundance of birds that overwinter in the state, with fewer Canadian species moving south and more short-distance migrants remaining through the year.

Predictive models developed by the National Audubon Society (2019) forecast the most significant negative impacts to Minnesota's avifauna will occur to boreal forest species (see Climate Change Profile). More than 21 of the 27 species the models predict will be extirpated in the state should the temperature rise by 1.5 degrees Centigrade are restricted to Minnesota's northern forest (Pfannmuller et al., 2024). Even more species are lost should the temperatures rise even further. The breeding range of most of these species is predicted to move farther north across Canada, providing suitable habitat is available.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. Increased flooding and high winds will negatively impact ground-nesting waterfowl, waterbirds, and shorebirds by eroding nesting habitat and flooding nests. Nesting failures in canopy dwelling waterbirds (herons and egrets) could increase given the vulnerability of their highly

exposed nests. Flash floods, tornados and hail can cause adult and chick mortality, blow down nests in trees and shrubs, flood ground nests or nests embedded in wetland vegetation, delay migration, and alter habitat.

The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). Droughts will likely decrease insect abundance and increase the occurrence of fires, especially in northern counties. Grassland birds, wetland birds, and waterbirds may be negatively impacted if habitat quality degrades and food supplies decline. Drought conditions and low water levels also are predicted to increase the spread of infectious diseases and pathogens that impact birds. For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Case Study: Modeling Climate Impacts on Breeding Birds

In 2019 the National Audubon Society (NAS) published a report titled [Survival by Degrees: 389 Bird Species on the Brink](#). The report documented a comprehensive effort to model changes in the distribution of 640 North American birds that could result from warming temperatures. More than 70 data sources and over 140 million bird records were used as input, along with information on vegetative cover, land use, and climate. Using the most current climate modeling methods available at the time, the researchers assessed the distributional changes to birds under three global warming scenarios based on the 2014 Intergovernmental Panel on Climate Change. They include a 1.5°C warming above preindustrial levels (which is considered imminent between 2030 and 2052 if no climate change mitigation measures are taken); a 2° C warming (which could happen as soon as 2050); and a 3°C warming scenario (which could occur by 2080). These models are predictive models, meaning that they may or may not occur.

Distribution maps for each species for each of the three warming scenarios are available and can be used to assess impacts at the national, regional and state level. Pfannmuller et al. (2024) assessed the predicted results for Minnesota species. They designated a species as extirpated at a given warming scenario when the NAS models portrayed the absence of any current or newly available “thermal breeding range” for the species in Minnesota. At the lowest warming scenario (1.5°C), the models predicted the extirpation of 27 breeding birds from Minnesota. The vast majority of these species are dependent on the northern boreal forests of the Laurentian Mixed Forest Province. They include such SGCN as the spruce grouse (*Canachites canadensis*), great gray owl, olive-sided flycatcher (*Contopus cooperi*), and Cape May warbler. At the 2°C warming scenario Minnesota may lose a total of 41 breeding birds including the SGCN American goshawk, evening grosbeak, and LeConte’s sparrow and a total of 66 species at the 3°C warming scenario including the SGCN lesser scaup (*Aythya affinis*), red-necked grebe (*Podiceps grisegena*), and Franklin’s Gull (*Leucophaeus pipixcan*).

Priority Conservation Strategies

To implement the SGCN Goal of this Plan (*Conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions*), three strategies were identified:



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Examples of conservation actions are grouped below under these three strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Some actions that are focused on important population monitoring and public engagement needs might not always be tied to a specific stressor; these are labeled with ‘n/a’ in the stressor column.

Potential Conservation Actions for Birds



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.

Stressor	Action
	Increase monitoring of grassland birds to improve our understanding of statewide population levels. Conduct response/effectiveness monitoring to assess their response to different grassland management regimes, including prescribed burning, grazing, and prairie restoration efforts. Rigorous effectiveness monitoring is necessary to develop improved evidence-based management recommendations and avoidance measures for SGCN (Binley et al., 2025) as part of adaptive management. This information can then be used to help devise and guide new options for incentivizing bird-friendly management.

	When conflicts arise regarding the impacts of fishing-eating birds on local fish populations, apply science-based decision making to conflict resolution processes. Ensure mitigation strategies sustain regional populations and avoid impacts to non-target species. Monitor effects of any management actions. Provide educational opportunities related to "nuisance" species.
n/a	Promote participatory science opportunities such as the Minnesota Loon Monitoring Program , and eBird . See Chapter 4. Public Engagement and DNR's Wildlife Diversity and Community Science webpage for more bird-related opportunities.
n/a	Actively support and promote continued funding for long-term bird monitoring programs, including the Minnesota National Forest Breeding Bird Monitoring Program , the Hawk Ridge Bird Observatory , and monitoring surveys conducted by the DNR to track the status of colonial waterbirds, waterfowl, and upland game birds.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.

Stressor	Action
	To reduce window collisions, apply decals to windows or use UV paint visible to birds. Require or incentivize collision mitigation for new construction builds. Increase public outreach/education on collision prevention. The American Bird Conservancy and Audubon's Bird Friendly Buildings initiative provide more details.
	Protect and restore shoreline habitat. Advocate for stricter community planning and zoning. Increase public outreach about the value of natural shorelines and provide guidance for shoreline owners . Provide guidance regarding artificial nesting platforms, such as for loons.
	Encourage agricultural producers to enroll in conservation programs that incentivize integrating bird conservation efforts into their working farms or ranches such as by restoring and managing grasslands, improving water quality, reducing erosion and sedimentation, and conserving ground and surface water. Promote conservation grazing and prairie restorations with incentives to accommodate Best Management Practices proven to benefit native birds, especially SGCN. Add buffers or windbreaks around agricultural fields. Encourage cover crop and no-till farming practices. Support private lands biologists to assist landowners in enrolling in Farm Bill and other similar programs and learning about conservation easements. Encourage livestock producers to explore other creative options such as

	Audubon's Conservation Ranching Program . It is a collaborative effort between the National Audubon Society and ranchers to protect and restore grassland habitats for declining bird populations. It focuses on sustainable ranching practices, particularly rotational grazing, and provides a market connection for "bird-friendly" beef and bison. Additional opportunities to link agricultural production with bird conservation can be found at the Natural Resources Conservation Service and the Board of Soil and Water Resources.
	Integrate habitat requirements into permits for mining, quarrying and drilling. Work with appropriate government agencies to restrict drilling and quarrying in areas of known and historical leks (grouse/prairie chickens).
	Require native vegetation in energy development areas. Prohibit renewable energy development in remnant prairie areas. Influence wind farm siting to avoid flyways. Advocate for policies to improve reporting of mortality and implementation of mitigation strategies required under state and federal laws. Seek collaborative opportunities to work with wind energy companies to publicly acquire habitat adjacent to wind towers for conservation.
	Minimize new roads and trails and damaging recreational activities in important habitats, including Important Birds Areas delineated by Audubon Minnesota.
	Co-locate utilities when possible to reduce environmental impact and consider wildlife in siting plans. Mow utility areas outside the breeding and nesting season. Avoid routing lines near or across lakes where feasible. Provide technical guidance to utility companies related to reducing avian collisions with powerlines (such as by marking lines). Support research related to reducing avian collisions with powerlines
	Strategize acquisitions of forest lands to emphasize connectivity and corridor creation. Quantify spatial use needs and territory size requirements of birds to help guide management decisions. Use best management practices such as winter harvest and the maintenance of snags, downed logs, and live trees; increase rotation age; reduce clear-cutting; retain live trees in clumps; reduce cord targets when appropriate; and protect sensitive areas. Encourage harvests that better mimic natural disturbance regimes. Better incorporate ecological values into sustainable timber harvest models and evaluate proposed harvest rates for impacts on wildlife habitat. Monitor impacts of forest management on forest birds.
	When conflicts arise regarding the impacts of fishing-eating birds on local fish populations, apply science-based decision making to conflict resolution processes. Ensure mitigation strategies sustain regional populations and avoid impacts to non-target species. Monitor effects of any management actions. Provide educational opportunities related to "nuisance" species.
	Restrict recreational access of people to colonies during breeding season and peak migration. Seasonal beach restrictions protect nesting and foraging areas of birds such as piping plovers and common terns (Hillman et al., 2015). Wildlife sanctuary signage placed around breeding

	habitat reduces recreational disturbance to these sites. Local support from landowners, conservation partners and enforcement officers is important in supporting these closures.
	Establish and enforce no wake zones around key nesting areas, especially for colonial nesting waterbirds, and conduct public outreach to promote wildlife-friendly boating behavior.
	Integrate seasonal timing of water management, especially wetland drawdowns, relative to the bird nesting season. Provide sufficient water levels, especially in non-forested wetlands that support bird SGCN, from spring through July. Encourage the application of best management practices for installing new or upgrading old drainage systems (Board of Soil and Water Resources and the Natural Resources Conservation Service)
	Support Lights Out programs. Advocate for mandates and bylaws to change city lighting (downward facing and off after a certain time). Promote artificial light guidance from the International Dark Sky Association and implement their best practices for wildlife-friendly outdoor lighting.
	Leave some dense vegetation structure in conifer forests to provide thermal refugia for spruce grouse.
	Increase buffers around wetlands and increase the size of wetlands to mitigate wind and water level rise. Provide more upland nesting habitat on higher ground. Protect lake shoreline habitat. Create artificial nesting habitat. Stop island erosion and flooding.
	Continue and support efforts to provide suitable nesting habitats for common terns with actions that reduce the access of other waterbirds, such as ring-billed gulls and double-crested cormorants, that compete for the limited nesting habitat that is available.
	Increase the use of prescribed fire. Proactively apply prescribed fire and grazing in ways that sustain native ecosystems and support SGCN birds while also avoiding or minimizing negative impacts to them through factors such as timing, intensity, acreage, and leaving unburned areas for escape cover. When developing grazing and burn plans, consider the requirements of SGCN. Aim to provide diversity across the landscape and over time (see also Prairies and Other Grasslands sub-chapter).
	Support and promote efforts to control problematic non-native species, placing a special emphasis on restoration of native vegetation in grasslands and wetlands.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Stressor	Action
	Provide land managers (including private landowners) with information on bird-friendly landscape design; highlight bird-friendly yard certification and bird-friendly forest management materials. Provide tax relief/incentive payments for woodland owners. Audubon's " How to Make Your Yard Bird-Friendly " and the National Wildlife Federation's " Create a Bird-Friendly Habitat " offer suggestions. The Sustainable Forest Incentive Act provides payments to landowners to keep woodlands undeveloped.
	Share information with land managers, including private landowners, about habitat needs of rare forest species. Engage conservation stakeholders in forest management topics. Increase representation of all stakeholder groups in decisions.
	Promote the use of non-lead fishing tackle options to anglers and increase public outreach on negative impacts to wildlife. Encourage anglers to recycle lead tackle at hazardous waste facilities and/or encourage local communities to hold tackle collections days. The Minnesota Pollution Control Agency has additional information and suggestions.
	When conflicts arise regarding the impacts of fishing-eating birds on local fish populations, apply science-based decision making to conflict resolution processes. Ensure mitigation strategies sustain regional populations and avoid impacts to non-target species. Monitor effects of any management actions. Provide educational opportunities related to "nuisance" species.
	Promote non-toxic ammunition options to reduce exposure of wildlife and humans to lead. For example, the Non-Toxic Working Group of the Minnesota Chapter of the Wildlife Society , along with partners, distributes information and holds hands-on demonstration events to inform and encourage the use of copper ammunition for hunting.
	Initiate an information campaign to raise awareness of the impacts of seed coatings to the health and sustainability of grassland birds.
	Support Lights Out programs. Advocate to change city lighting practices (downward facing and off after a certain time). Promote artificial light guidance from the International Dark Sky Association and implement their best practices for wildlife-friendly outdoor lighting.
	Support and promote efforts to control problematic non-native species, placing a special emphasis on restoration of native vegetation in grasslands and wetlands.

Stressor	Action
	Promote programs such as Audubon's Bird City Minnesota which helps birds by restoring, protecting and improving bird habitat; reducing threats such as window collisions, toxins and harmful human activities, often due to a lack of understanding of birds' needs; and engaging people in learning about, enjoying and helping birds thrive.

Case Study: Linking Grassland Birds to Profitable Livestock Production

A monitoring team was launched by the Land Stewardship Project and the Minnesota Institute for Sustainable Agriculture in the 1990s to create a set of sustainability indicators that would link healthy grassland habitat to profitable livestock production. At the core of the team were six farm families in southeastern Minnesota who utilized rotational grazing and wanted to know what impact this system of dairy and beef production was having on the land and its wild inhabitants. Farmers worked with University of Minnesota environmental scientists, an economist, and a sociologist, as well as experts from the Minnesota DNR and the U.S. Fish and Wildlife Service, to develop indicators that could be easily implemented while carrying out livestock management chores. The farmers were taught the basics of monitoring soil health, water quality, forage viability, and frog and toad populations.

But by far the most popular, and effective, indicator for the farmers was the presence of grassland birds. It became a user-friendly method for the farmers to link the productivity of their pastures, and therefore their animals, with the presence of species that rely on healthy, heterogeneous, grassland habitat. Because it can be worked into chores like moving cattle and fixing fences, birding is a handy way to gauge some of the key impacts a farmer is having on the land. Indeed, by the third year of the monitoring initiative, bird sightings were among the first items farmers mentioned during their monthly reports. When they got together, they were not bragging about their corn yields, milk production, or improvements in soil structure. Instead, they were quick to share the number of successful bluebird or bobolink nestings they had observed.

As they became more aware of the birds on their farms, the farmers took the next step: they started wondering what impacts their livelihoods were having on the life cycles of their feathered neighbors. They began discussing with wildlife biologists concerns they had about nesting disruptions caused by haying, pasture clipping (a method for keeping the grass more palatable for livestock), and even grazing.

Although several farmers observed that cattle were sometimes able to graze a paddock with an active nest without destroying the eggs or nestlings, it became clear that the hay fields and managed pastures could become avian “population sinks,” or booby traps, rather than “population sources.” As a result, several farmers reduced or delayed pasture clipping to allow fledglings to become more mobile before the mower disrupted the nests. Dairy producer Art Thicke also eliminated clipping on some of his paddocks one year and by the fourth grazing, he says, “You couldn’t tell where you clipped and where you didn’t.” In other words, despite the lack of clipping, the cows still found the grass palatable.

One management technique that improved nesting success was the establishment of rest areas within their grazing paddocks. One year, each farm held one paddock out of grazing from the beginning of the season until at least the end of July. The densely vegetated rest areas provided a place for birds to nest undisturbed by cattle or machines. They also allowed birds disturbed in adjoining paddocks to retreat to the lush cover and re-nest. Farmers noticed greater concentrations of bobolinks and dickcissels in the rest areas during the first year of the experiment. As an added bonus, the rested paddocks gave the vegetation an opportunity to mature so it could re-seed directly or through the livestock’s digestive systems. Allowing the longer growing period also increased the plant’s root structure, thus improving soil structure, which is important when a farm is exposed to extreme drought or extreme precipitation events.

These are examples of management practices that benefit the farm financially, as well as improve the environment. In short, more bobolinks can equal more beef. If the farmers had been told right out of the gate that their farming methods were threatening grassland bird species, defenses would have gone up, reducing opportunities to implement even minor management changes. But monitoring took them through a process that started with newfound knowledge and ended with appreciation and action that improved grassland bird habitat without negatively impacting their profits (DeVore, 2003).

References for Birds

- Abreu, T.L.S., Berg, S.B., de Faria, L.P., Gomes, L.P., Marinho-Filho, J.S., & Colli, G.R. (2020). River dams and the stability of bird communities: A hierarchical Bayesian analysis in a tropical hydroelectric power plant. *Journal of Applied Ecology*, 57(6), 1124–1136. <https://doi.org/10.1111/1365-2664.13607>
- Adikwan, Ogimaagiizhig Odoodeman (Charles Grolla). (2019). Binesi-Dibaajimowinan: Ojibwe Bird Stories Unique to Northern Minnesota. Cass Lake, MN: Leech Lake Tribal College, LLTC Design and Print Shop.
- Arnold, T.W., & Zink, R.M. (2011). Collision Mortality Has No Discernible Effect on Population Trends of North American Birds. *PLoS ONE* 6(9): e24708. <https://doi.org/10.1371/journal.pone.0024708>
- Aubrecht, C., Jaiteh, M., & de Sherbinin, A. (2010). Global Assessment of Light Pollution Impact on Protected Areas. Technical Report by the Center for International Earth Science Information Network, The Earth Institute at Columbia University, and Austrian Institute of Technology.
- Avian Power Line Interaction Committee (APLIC). (2012). Reducing Avian Collisions with Power Lines: The State of the Art in 2012. Edison Electric Institute and APLIC. Washington, D.C.
- Barton, D. C., & Holmes, A. L. (2007). Off-Highway vehicle trail impacts on breeding songbirds in northeastern California. *Journal of Wildlife Management*, 71(5), 1617–1620. <https://doi.org/10.2193/2006-026>
- Barzan, F.R., Bellis, L.M., & Dardanelli, S. (2021). Livestock grazing constrains bird abundance and species richness: A global meta-analysis. *Basic and Applied Ecology*, 56, 289–298. <https://doi.org/10.1016/j.baae.2021.08.007>.
- Benoit, L.K., & Askins, R.A. (1999). Impact of the spread of *Phragmites* on the distribution of birds in Connecticut tidal marshes. *Wetlands* 19, 194–208. <https://doi.org/10.1007/BF03161749>.
- Bernath-Plaisted, J. S., Correll, M. D., Somershoe, S. G., Dwyer, A. M., Bankert, A., Beh, A., Berlanga, H., Boyle, W. A., Cruz-Romo, J. L., George, T. L., Herkert, J., Koper, N., Macías-Duarte, A., Panjabi, A. O., Ramírez-Flores, O. M., Robinson, B., Ruvalcaba-Ortega, I., Sibbing, J., Strasser, E. H., . . . VerCauteren, T. (2023). Review of conservation challenges and possible solutions for grassland birds of the North American Great Plains. *Rangeland Ecology & Management*, 90, 165–185. <https://doi.org/10.1016/j.rama.2023.07.002>
- Binley, A. D., Haddaway, L., Buxton, R., Lalla, K. M., Lesbarreres, D., Smith, P. A., & Bennett, J. R. (2025). Endangered species lack research on the outcomes of conservation action. *Conservation Science and Practice*. <https://doi.org/10.1111/csp2.13304>
- Bock, C.E., Saab, V.A., Rich, T.D., & Dobkin, D.S. (1993). Effects of Livestock Grazing on Neotropical Migratory Landbirds in Western North America. In: Finch & Stangel (Eds.) *Status and Management of Neotropical Migratory Birds*. Rocky Mountain Forest and Range Experiment Station.
- Boyce, A. J., Shamon, H., Kunkel, K. E., & McShea, W. J. (2021). Grassland bird diversity and abundance in the presence of native and non-native grazers. *Avian Conservation and Ecology*, 16(2). <https://doi.org/10.5751/ace-01944-160213>
- Breining, G. (2014). Mining Minnesota's Canoe Country. Audubon Magazine. March-April 2014. <https://www.audubon.org/magazine/mining-minnesotas-canoe-country>

- Bumgarner, J.R. & Nelson, R. (2021). Light at Night and Disrupted Circadian Rhythms Alter Physiology and Behavior. *Integrative and Comparative Biology*, 61(3), 1160-1169. <https://academic.oup.com/icb/article/61/3/1160/6206372>.
- Buxton, R. T., McKenna, M. F., Mennitt, D., Brown, E., Fristrup, K., Crooks, K. R., Angeloni, L. M., & Wittemyer, G. (2019). Anthropogenic noise in US national parks – sources and spatial extent. *Frontiers in Ecology and the Environment*, 17(10), 559–564. <https://doi.org/10.1002/fee.2112>
- Buxton, R. T., McKenna, M. F., Mennitt, D., Fristrup, K., Crooks, K., Angeloni, L., & Wittemyer, G. (2017). Noise pollution is pervasive in US protected areas. *Science*, 356(6337), 531-533.
- Chock, R. Y., Clucas, B., Peterson, E. K., Blackwell, B. F., Blumstein, D. T., Church, K., Fernández-Juricic, E., Francescoli, G., Greggor, A. L., Kemp, P., Pinho, G. M., Sanzenbacher, P. M., Schulte, B. A., & Toni, P. (2020). Evaluating potential effects of solar power facilities on wildlife from an animal behavior perspective. *Conservation Science and Practice*, 3(2). <https://doi.org/10.1111/csp2.319>
- DarkSky International. (2025). DarkSky International | Protecting the night skies for present and future generations. Accessed June 2025. <https://darksky.org>
- DeVore, B. (2003). Creating habitat on farms the Land Stewardship Project and monitoring on agricultural land. *Conservation in Practice*, 4(2), 28–35. <https://doi.org/10.1111/j.1526-4629.2003.tb00060.x>
- Evers, D.C., & Cristol, D.A. (2020.) The Impact of Mercury on North American Songbirds: Effects, Trends, and Predictive Factors. Biodiversity Research Institute. Portland, Maine, USA. BRI Science Communications Series 2020-29. 22 pp. https://briwildlife.org/wp-content/uploads/2021/06/Mercury_in_Songbirds_Booklet_2020.pdf
- Falchi, F., Cinzano, P., Duriscoe, D., Kyba, C. C. M., Elvidge, C. D., Baugh, K., Portnov, B. A., Rybnikova, N. A., & Furgoni, R. (2016). The new world atlas of artificial night sky brightness. *Science Advances*, 2(6). <https://doi.org/10.1126/sciadv.1600377>
- Finch, D. M., Butler, J. L., Runyon, J. B., Fetting, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Frei, B., Smith, K.G., Withgott, J.H., Rodewald, P.G., Pyle, P., & Patten, M.A. (2020). Red-headed Woodpecker (*Melanerpes erythrocephalus*), version 1.0. In Rodewald, P.G. (Ed.) *Birds of the World*. Lab of Ornithology, Ithaca, NY, USA. <https://doi.org/10.2173/bow.rehwoo.01>
- Gerald Niemi, Janet Green, Bonnie Sample, Nick Walton, Ed Zlonis, Tim Brown, Annie Bracey, George Host, Jane Reed, Kim Rewinkel, and Norm Will. 2017. The First Minnesota Breeding Bird Atlas (2009-2013). <https://mnbirdatlas.org/>
- Grade, T. J., Pokras, M. A., Laflamme, E. M., & Vogel, H. S. (2018). Population-level effects of lead fishing tackle on common loons. *Journal of Wildlife Management*, 82(1), 155–164. <https://doi.org/10.1002/jwmg.21348>

- Grasman, K. A., Annis, M., Eakin, C., Moore, J., & Williams, L. L. (2025). Monitoring and assessment of population, reproductive, and health effects in colonial waterbirds breeding at contaminated Great Lakes sites in Michigan. *Environmental Toxicology and Chemistry*, 44(1), 77–91. <https://doi.org/10.1093/etoinl/vgae001>
- Green, J.C. (1995). *Birds and Forests: A Management and Conservation Guide*. Minnesota Department of Natural Resources.
- Green, J.C., & Janssen, R.B. (1975). *Minnesota Birds: Where, When, and How Many*. Minneapolis: University of Minnesota Press.
- Gutiérrez-Arnal, J., & Marín, C. (2024). The Latent Threat in Wild Birds: *Clostridium botulinum*. *Veterinary Sciences*, 11(1), 36. <https://doi.org/10.3390/vetsci11010036>
- Haig, S. M., D’Elia, J., Eagles-Smith, C., Fair, J. M., Gervais, J., Herring, G., Rivers, J. W., & Schulz, J. H. (2014). The persistent problem of lead poisoning in birds from ammunition and fishing tackle. *Ornithological Applications*, 116(3), 408–428. <https://doi.org/10.1650/condor-14-36.1>
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Hillman, M.D., Karpanty, S.M., Fraser, J.D., & Derosé-Wilson, A. (2015). Effects of aircraft and recreation on colonial waterbird nesting behavior. *The Journal of Wildlife Management* 79(7), 1192-1198. <https://doi.org/10.1002/jwmg.925>.
- Horváth, G., Kriska, G., Malik, P., & Robertson, B. (2009). Polarized light pollution: a new kind of ecological photopollution. *Frontiers in Ecology and the Environment*, 7(6), 317–325. <https://doi.org/10.1890/080129>
- Janssen, R.B. (1987). *Birds in Minnesota*. Minneapolis: University of Minnesota Press.
- Janssen, R.B. (2019). *Birds in Minnesota*. Rev. ed. Minneapolis: University of Minnesota Press.
- Kimmel, R.O., & Tranel, M.A. (2007). Evidence of Lead Shot Problems for Wildlife, the Environment, and Human Health – Implications for Minnesota. In DonCarlos, M.W. et al. (Eds.) *Summaries of Wildlife Research Findings 2007*. Minnesota Department of Natural Resources.
- Koenig, W.D., Liebhold, A.M., Bonter, D.N., Hochachka, W.M., & Dickinson, J.L. (2013). Effects of the emerald ash borer invasion on four species of birds. *Biological Invasions* 15, 2095-2103. DOI 10.1007/s10530-013-0435-x
- Kok, A. C. M., Berkhout, B. W., Carlson, N. V., Evans, N. P., Khan, N., Potvin, D. A., Radford, A. N., Sebire, M., Sabet, S. S., Shannon, G., & Wascher, C. a. F. (2023). How chronic anthropogenic noise can affect wildlife communities. *Frontiers in Ecology and Evolution*, 11. <https://doi.org/10.3389/fevo.2023.1130075>
- Lafrancois, B. M., Riley, S. C., Blehert, D. S., & Ballmann, A. E. (2011). Links between type E botulism outbreaks, lake levels, and surface water temperatures in Lake Michigan, 1963–2008. *Journal of Great Lakes Research*, 37(1), 86–91. <https://doi.org/10.1016/j.jglr.2010.10.003>
- Lebbin, D.J. & Fenwick, G.H. (2010). *The American Bird Conservancy Guide to Bird Conservation*. American Bird Conservancy. 456 pages.

- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Loss, S. R., & Blair, R. B. (2011). Reduced density and nest survival of Ground-Nesting songbirds relative to earthworm invasions in northern hardwood forests. *Conservation Biology*, 25(5), 983–992. <https://doi.org/10.1111/j.1523-1739.2011.01719.x>
- Loss, S. R., Boughton, B., Cady, S. M., Londe, D. W., McKinney, C., O’Connell, T. J., Riggs, G. J., & Robertson, E. P. (2022). Review and synthesis of the global literature on domestic cat impacts on wildlife. *Journal of Animal Ecology*, 91(7), 1361–1372. <https://doi.org/10.1111/1365-2656.13745>
- Loss, S. R., Will, T., & Marra, P. P. (2013a). The impact of free-ranging domestic cats on wildlife of the United States. *Nature Communications*, 4(1). <https://doi.org/10.1038/ncomms2380>
- Loss, S. R., Will, T., & Marra, P. P. (2013b). Estimates of bird collision mortality at wind facilities in the contiguous United States. *Biological Conservation*, 168, 201–209. <https://doi.org/10.1016/j.biocon.2013.10.007>
- Loss, S. R., Will, T., & Marra, P. P. (2015). Direct Mortality of Birds from Anthropogenic Causes. *Annual Review of Ecology Evolution and Systematics*, 46(1), 99–120. <https://doi.org/10.1146/annurev-ecolsys-112414-054133>
- Loss, S. R., Will, T., Loss, S. S., & Marra, P. P. (2014). Bird–building collisions in the United States: Estimates of annual mortality and species vulnerability. *Ornithological Applications*, 116(1), 8–23. <https://doi.org/10.1650/condor-13-090.1>
- Machtans, C. S., Wedeles, C. H. R., & Bayne, E. M. (2013). A First Estimate for Canada of the Number of Birds Killed by Colliding with Building Windows. *Avian Conservation and Ecology*, 8(2). <https://doi.org/10.5751/ace-00568-080206>
- Minnesota Department of Natural Resources (DNR). (1997). Clay County Beach Ridges Forum for gravel mining and prairie protection. Final Report. <https://www.lrl.mn.gov/docs/2015/mandated/155087.pdf>
- Minnesota Department of Natural Resources (DNR). (2025). Double-crested Cormorant. <https://www.dnr.state.mn.us/birds/doublecrestedcormorant.html> [Accessed May 9, 2025].
- Minnesota Natural Shoreline Partnership. (2023). Minnesota’s Vanishing Natural Shorelines: A Loss that Contributes to Degraded Lake Quality. https://drive.google.com/file/d/1vU-nx9_8DHivVmwjHjL0nem9VihC_Ffz/view.
- Minnesota Ornithologists Union (MOU). (2024). Checklist of The Birds of Minnesota. Compiled and Annotated by the Minnesota Ornithologists’ Union Record Committee. <https://moumn.org/mourc/MOURC%20Checklist.pdf>
- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). 2025. Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state’s greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>

- Narango, D. L., Tallamy, D. W., & Marra, P. P. (2018). Nonnative plants reduce population growth of an insectivorous bird. *Proceedings of the National Academy of Sciences*, 115(45), 11549–11554. <https://doi.org/10.1073/pnas.1809259115>
- National Audubon Society. (2019). Survival by Degrees: 389 Bird Species on the Brink. <https://www.audubon.org/climate/survivalbydegrees>.
- Nelson, S. B., Coon, J. J., Duchardt, C. J., Fischer, J. D., Halsey, S. J., Kranz, A. J., Parker, C. M., Schneider, S. C., Swartz, T. M., & Miller, J. R. (2017). Patterns and mechanisms of invasive plant impacts on North American birds: a systematic review. *Biological Invasions*, 19(5), 1547–1563. <https://doi.org/10.1007/s10530-017-1377-5>
- Niemi, G.J. (1978). Breeding birds of burned and unburned areas in northern Minnesota. *Loon*, 50, 73-84.
- Niemi, G.J., Howe, R.W., Sturtevant, B.R., Parker, L.R., Grinde, A.R., Danz, N.P., Nelson, M.D., Zlonis, E.J., Walton, N.G., Giese, E.E.G., & Lietz, S.M. (2016). Analysis of Long-term Forest Bird Monitoring Data from National Forests of the Western Great Lakes Region. USDA, Forest Service. Northern Research Station, General Technical Report NRS-159.
- Ortega, Y. K., & Capen, D. E. (2002). Roads as Edges: Effects on birds in forested landscapes. *Forest Science*, 48(2), 381–390. <https://doi.org/10.1093/forestscience/48.2.381>
- Pain, D.J., Fisher, I.J., & Thomas, V.G. (2009). A global update of lead poisoning in terrestrial birds from ammunition sources. In: Watson, R.T., Fuller, M., Pokras, M., & Hunt, W.G. (Eds.) *Ingestion of Lead from Spent Ammunition: Implications for Wildlife and Humans*. The Peregrine Fund, Boise, Idaho, USA.
- Pain, R.J., Mateo, R., & Green, R.E. (2019). Effects of lead from ammunition on birds and other wildlife: A review and update. *Ambio*, 48(9), 935-953. <https://pubmed.ncbi.nlm.nih.gov/30879267/>
- Pfannmuller, L., Niemi, G., Green, J., Sample, B., Walton, N., Zlonis, E., Brown, T., Bracey, A., Host, G., Reed, J., Rewinkel, K., & Will, N. (2017).
- Pfannmuller, L.A., Niemi, G.J., & Green, J.C. (2024). The Breeding Birds of Minnesota: History, Ecology, and Conservation. University of Minnesota Press. 685 pages.
- Puettmann, K.J., & Ek, A.R. (1999). Status and Trends of Silvicultural Practices in Minnesota, *Northern Journal of Applied Forestry*, 16(4), 203–210. <https://doi.org/10.1093/njaf/16.4.203>
- Ratajczak, Z., Collins, S.L., Blair, J.M., Koerner, S.E., Louthan, A.M., Smith, M.D., Taylor, J.H., & Nippert, J.B. (2022). Reintroducing bison results in long-running and resilient increases in grassland diversity, *Proc. Natl. Acad. Sci. U.S.A.*, 119(36) e2210433119. <https://doi.org/10.1073/pnas.2210433119>.
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwccasc.umn.edu/node/821>
- Rattner, B.A., Franson, J.C., Sheffield, S.R., Goddard, C.I., Leonard, N.J., Stang, D., & Wingate, P.J. (2008). Sources and Implications of Lead Ammunition and Fishing Tackle on Natural Resources. Technical Review 08-01, The American Fisheries Society, The Wildlife Society, Bethesda, Maryland, USA.

- Reed, L.M., Johansson, M.A., Panella, N., McLean, R., Creekmore, T., Puella, R., & Komar, N. (2019). Declining Mortality in American Crow (*Corvus brachyrhynchos*) Following Natural West Nile Virus Infection. *Avian Diseases* 53, 458-461. <https://vectorbio.rutgers.edu/publications/Reed2009MortalityCrows.pdf>
- Roberts, T.S. (1932). *The Birds of Minnesota*. 2 vols. Minneapolis: University of Minnesota Press.
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Runwal, P. (2020). Building Collisions are a Greater Danger for Some Birds than Others. Audubon Magazine. July 9, 2020. <https://www.audubon.org/news/building-collisions-are-greater-danger-some-birds-others>
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>
- Seewagen, C. L. (2010). Threats of environmental mercury to birds: knowledge gaps and priorities for future research. *Bird Conservation International*, 20(2), 112–123. <https://doi.org/10.1017/s095927090999030x>
- Shahid, A., & Nadeem, A. (2021). Effects of Pollution on Freshwater birds: A review. *Journal of Chemical Biological and Physical Sciences*, 11(3). <https://doi.org/10.24214/jcbps.b.11.3.46474>
- Slabe, V.A., Anderson, J.T., Cooper, J., Brown, B., Ortiz, P., Buchweitz, J., McRuer, D., & Katzner, T. (2019). Lead in piscivorous raptors during breeding season in the Chesapeake Bay region of Maryland and Virginia, USA. *Environ. Toxicol. Chem.* 38, 862–871.
- Spaul, R. J., & Heath, J. A. (2016). Nonmotorized recreation and motorized recreation in shrub-steppe habitats affects behavior and reproduction of golden eagles (*Aquila chrysaetos*). *Ecology and Evolution*, 6(22), 8037–8049. <https://doi.org/10.1002/ece3.2540>
- Stanton, R.L., Morrissey, C.A., & Clark, R.G. (2018). Analysis of trends and agricultural drivers of farmland bird declines in North America. *Agriculture, Ecosystems and Environment* 254(2018), 244-254.
- Steven, R., Pickering, C., & Castley, J.G. (2011). A review of the impacts of nature based recreation on birds. *Journal of Environmental Management*, 92(10), 2287-2294.
- Stucker, J.H., & Cuthbert, F.J. (2000). Biodiversity of southeastern Minnesota forested streams: relationships between trout habitat improvement practices, riparian communities, and the Louisiana waterthrush. Final report submitted to the Natural Heritage and Nongame Research Program, Minnesota Department of Natural Resources. 147 pp.
- Taieb, L., Ludwig, A., Ogden, N. H., Lindsay, R. L., Iranpour, M., Gagnon, C. A., & Bicout, D. J. (2020). Bird species involved in West Nile virus epidemiological cycle in Southern Québec. *International Journal of Environmental Research and Public Health*, 17(12), 4517. <https://doi.org/10.3390/ijerph17124517>
- Tranel, M.A. & Kimmel, R.O. (2009). Impacts of lead ammunition on wildlife, the environment, and human health—a literature review and implications for Minnesota. The Peregrine Fund, Boise, Idaho, USA. <http://science.peregrinefund.org/legacy-sites/conference-lead/PDF/0307%20Tranel.pdf>

- United States Department of Agriculture (USDA). (2022). Census of Agriculture State Profile. National Agricultural Statistics Service. Minnesota.
https://www.nass.usda.gov/Publications/AgCensus/2022/Online_Resources/County_Profiles/Minnesota/cp99027.pdf
- Vennesland, R.G. (2010). Risk perception of nesting Great Blue Herons: Experimental evidence of habituation. *Canadian Journal of Zoology* 88(1), 81-89.
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report].
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>
- Wilsey, C.B., Grand, J., Wu, J., Michel, N., Grogan-Brown, J., & Trusty, B. (2019). North American Grasslands. National Audubon Society, New York, New York. USA. https://nas-national-prod.s3.amazonaws.com/audubon_north_american_grasslands_birds_report-final.pdf
- Wood, T. J., & Goulson, D. (2017). The environmental risks of neonicotinoid pesticides: a review of the evidence post 2013. *Environmental Science and Pollution Research*, 24(21), 17285–17325.
<https://doi.org/10.1007/s11356-017-9240-x>
- Wuerthner, G. (2012). Birds and Livestock Grazing – A Review. The Wildlife News. November 13, 2012.
<https://www.thewildlifeneeds.com/2012/11/13/birds-and-livestock-grazing-a-review/>
- Yates, D. (2025). How are migrating wild birds affected by H5N1 infection in the U.S.? University of Illinois Urbana-Champaign Strategic Communications and Marketing News Bureau. February 13, 2025.
<https://news.illinois.edu/how-are-migrating-wild-birds-affected-by-h5n1-infection-in-the-u-s>

Fish

Overview

Minnesota is well known for its excellent aquatic habitats and fisheries. Considered a top destination for fishing, the state provides quality fishing opportunities on 4,500 lakes managed for fishing and 16,000 miles of fishable rivers and streams. Fishing is a major recreational activity for adults and families throughout the state with approximately 26% of the Minnesota population age 16 and older participating in the year 2022. In addition to attracting Minnesota residents to quality lakes and rivers across the state, over 500,000 nonresidents also participated (University of Chicago, 2022). Yet, of the 165 fish species that are known to occur in the state, a relatively small percentage are targets of most angling expeditions.

This chapter focuses on Minnesota fishes that are Species in Greatest Conservation Need. These rare, declining, or vulnerable species are found throughout nearly every aquatic habitat in the state. These habitats and the fishes that depend on them experience cumulative effects of multiple stressors: loss of connectivity, changes in land use, altered surficial hydrology, invasive species, erosion and sedimentation, disease, climate change and degraded water quality (Aadland, 2015; Walsh et al., 2011). Many SGCN fish species are migratory, moving among large rivers, tributary streams and connected lakes and are impacted by alterations to multiple kinds of aquatic habitats. Large river species such as [lake sturgeon \(*Acipenser fulvescens*\)](#), [paddlefish \(*Polyodon spathula*\)](#) and [skipjack herring \(*Alosa chrysochloris*\)](#) have historically been impacted by loss of connectivity due to dams and their impoundments that prevent upstream movement. Stream dwelling SGCN such as [gravel chub \(*Erimystax x-punctatus*\)](#), [slender madtom \(*Noturus exilis*\)](#), and hornyhead chub (*Nocomis biguttatus*) are especially affected by land use and hydrologic alterations that increase sediment loads in streams and result in the burial of coarse gravel substrates that these fishes depend on for spawning habitat and the aquatic invertebrates that they feed on. Lake dwelling SGCN species such as [least darter \(*Etheostoma microperca*\)](#), [pugnose shiner \(*Notropis anogenus*\)](#) and [northern sunfish \(*Lepomis peltastes*\)](#) depend on native aquatic vegetation and are adversely impacted by alterations to natural shorelines, invasive plant species, and increased nutrient loads.

Overall, however, the loss of connectivity, much of it historic, may be the most significant stressor to fish communities across Minnesota. Aadland (2015) evaluated the effects of barriers to connectivity on fish distributions upstream and downstream of 32 dams in the Mississippi, Minnesota, St. Croix, St. Louis, Missouri, and Red River watersheds. “On average, species richness declined by 41% for complete barriers, 37% for near-complete barriers and 20% for barriers that are/were inundated at bankfull flows. A detailed assessment of the Cottonwood River Watershed indicated that a single barrier near the mouth of the river caused a watershed-wide loss of species richness.” Fish that were most vulnerable to the loss of connectivity were species that are imperiled or intolerant of the habitat conditions created by barriers, including 9 SGCN fish (see Dams and Water Management stressor).

Many of the stressors that negatively impact fish communities influence one or more aspects of water quality such as turbidity, oxygen levels, sediment loads, the presence of pollutants, invasive species, or increasing water temperatures. Minnesota’s list of [impaired waters](#) is a good resource regarding water quality throughout the state’s rivers, streams, and lakes. As of 2024, the total number of impaired water bodies was 2,798; impaired waters were most prevalent in the southern counties.

Summary of Fish SGCN and SNI

A total of 36 fish species have been designated Species in Greatest Conservation Need (SGCN); this represents nearly 22% of the 165 fish species that have been documented in Minnesota. One SGCN, the [Topeka shiner](#) (*Notropis topeka*), is listed as a federally endangered species. This small minnow is restricted to small prairie streams that are within the Missouri River watershed in southwestern Minnesota. Four fish SGCN have been classified as State Endangered Species (skipjack herring, [crystal darter](#) (*Crystallaria asprella*), [pallid shiner](#) (*Hybopsis amnis*), and [slender madtom](#) (*Noturus exilis*); five have been classified as State Threatened (gravel chub, [plains topminnow](#) (*Fundulus sciadicus*), [black buffalo](#) (*Ictiobus niger*), pugnose shiner, and paddlefish); and 21 species are state listed as Special Concern.

Five species that were listed as SGCN in the 2015-2025 SWAP were removed for the 2025-2035 plan update. The American eel (*Anguilla rostrata*) was deemed a temporary resident and, therefore, not a conservation target in the state. Most of the remaining four species ([kiyi](#) (*Coregonus kiyi*), [blue sucker](#) (*Cycleptus elongatus*), [yellow bass](#) (*Morone mississippiensis*) and pugnose minnow (*Opsopoeodus emiliae*)) were removed because their state populations are considered secure. Five fish were designated Species in Need of Information (SNI): Nipigon cisco (*Coregonus nipigon*), shortjaw cisco (*Coregonus zenithicus*), shoal chub (*Macrhybopsis hyostoma*); [pygmy whitefish](#) (*Prosopium coulterii*), and ninespine stickleback (*Pungitius pungitius*). A complete list of SGCN and SNI species can be found in Appendix B.

Habitat Associations

The 2025-2035 SWAP uses the Strahler Stream Order (SSO) method to classify streams and rivers. Streams without any tributaries are assigned an SSO of 1; then the stream order increases when tributaries combine into larger streams. As illustrated in Table 2.5, most fish SGCNs depend on small streams (SSO 1-4); these species also use a variety of secondary riverine habitats. Among the large river (SSO 7-8) SGCN, five species are entirely restricted to this habitat: skipjack herring, [bluntnose darter](#) (*Etheostoma chlorosoma*), pallid shiner, black buffalo, and paddlefish. Riverine cold-water specialists include three fish SGCN: longnose sucker (*Catostomus catostomus*), slimy sculpin (*Cottus cognatus*) and brook trout (*Salvelinus fontinalis*). Eight SGCN select shallow lakes as their primary habitat. Five are coldwater specialists in lakes: slimy sculpin, spoonhead sculpin (*Cottus ricei*), [lake chub](#) (*Couesius plumbeus*), deepwater sculpin (*Myoxocephalus thompsonii*), and brook trout.

Table 2.5. Habitat Associations for Fish SGCN

Habitats	Primary Habitat	Secondary Habitat	Total
Riverine Streams (SSO 1-4)	13	4	17
Riverine Rivers (SSO 5-6)	7	15	22
Riverine Large Rivers (SSO 7-8)	11	2	13
Riverine Large Rivers: Backwaters	3	0	3
Riverine Coldwater Specialist	3	0	3
Lakes Shallow	0	6	6
Lakes Not Shallow	8	3	11
Lakes Coldwater Specialist	5	0	5

Primary Stressors

Stressors are factors that pose direct or indirect challenges to vulnerable plant and wildlife species. Habitat loss and degradation are considered the primary threat for most SGCN, and some stressors reduce the quality of those habitats. Other stressors may operate more directly on the SGCN, such as disease and pathogens. In some cases, stressors may operate both directly and indirectly. For example, roads, trails, and railroads can directly harm SGCN through vehicular strikes, or operate indirectly through habitat fragmentation and reduce landscape connectivity. The stressor list below is adapted from an internationally recognized threats classification developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For more information, see the “*Stressors*” section of Chapter 1. Species in Greatest Conservation Need.

It is important to note that some factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts (e.g. catastrophic wildfire following a prolonged period of fire suppression) while serving as valuable conservation tools in others (e.g., appropriately planned and applied prescribed fire). Information about a subset of primary stressors specifically impacting this species group is included below, followed by a set of conservation actions addressing those stressors.



Development

Development around aquatic habitat for fish species can degrade habitat, reduce habitat connectivity, degrade water quality, and alter the local hydrology. Elimination or degradation of natural shorelines can alter runoff or discharge patterns that increase nutrients, sediments, and pollutants in waters. Some of these changes are prevalent when natural stream banks have been channelized or stabilized with rocks or cement. Impervious surfaces, such as parking lots, streets, and rooftops decrease the amount of rainwater that can replenish local soils, leading instead to increased runoff to aquatic habitats that carries sediment, pollutants, and debris. The loss of riparian vegetation can increase the amount of sunlight and water temperatures (USGS, 2012). Over time, urbanization can impact species richness and composition, decreasing the ratio of native to non-native species as some non-native species, such as common carp (*Cyprinus carpio*), are more tolerant of decreased water quality conditions (Pandit et al., 2024).



Crop Production

Many activities associated with intensive agriculture pose threats to aquatic systems and fish communities. Changes to sediment dynamics, water quality, nutrient or chemical runoff, and drainage practices are some of the potential stressors resulting from this activity (Schottler et al., 2014). Stream channelization, wetland drainage, and the loss of surrounding floodplain and riparian cover all increase inputs of pesticides, herbicides, sediment and nutrients into adjacent water bodies. This can lead to algal blooms that deplete oxygen in the water column and negatively impact sensitive fish that require well oxygenated water to survive. Drainage from fields reduces the natural absorption of the landscape for heavy rain events and increases sheet flow. Agricultural practices such as the installation of wells or drain tiles can also change the flow of groundwater to streams and lakes and are often associated with straightening of the channel (Blann et al., 2009). These changes result in flashy hydrology and incised channels that cause streams to function poorly (Blann et al., 2009). Smaller headwater streams are disproportionately affected by agricultural practices (Wohl, 2017). Fish SGCN that depend on cold, clear waters, such as brook trout (*Salvelinus fontinalis*) and the longnose sucker (*Catostomus catostomus*), as well as the federally endangered Topeka shiner (*Notropis topeka*), are impacted by row crop agriculture operations.



Livestock Management

Livestock operations have the potential to impact fish communities in several ways. Pastures that are intensively grazed can lead to erosion along the banks of rivers, streams, and lakes, particularly if livestock have unlimited access to these aquatic habitats. In addition, stream bottoms can become disturbed and lose shade-providing cover through herbivory and trampling (Storch, 1979). It may only take a few weeks of overgrazing to completely destroy years of effort to restore a stream and/or riparian habitat. Inappropriate grazing can cause water temperatures to rise, sediment loads to increase, and water quality to decline. If stockyards are not managed appropriately, runoff can impact water quality in ground water and adjacent wetlands and streams. Fish SGCN that respond positively to the implementation of best management practices include the Topeka shiner and plains topminnow (*Fundulus sciadicus*).



Mining and Quarrying

Sand mining operations conducted in rivers can negatively impact fish by either destroying the habitat, especially the bottom sediments, as well as increasing sedimentation turbidity. Two Minnesota fish SGCN that are especially vulnerable to such activities include the crystal darter and the Western sand darter (*Ammocrypta clara*). Both species inhabit the sandy bottoms of large rivers.

Taconite mining, a major industrial and economic activity throughout northeastern Minnesota for decades, is known to increase threats to fish populations by the discharge of waste materials into waters from tailing basins, as well as increasing sediments and pollutants such as sulfates and methylmercury. Federal studies associated with taconite operations at Silver Bay, along the north shore of Lake Superior, demonstrated impacts to fish populations largely caused by the discharge of tailings (over 60,000 tons per day for 25 years) that increased water turbidity and established an entire new delta of contaminated materials along the shoreline (WTIP, 2024). Although mining operators and permitting agencies have learned a great deal about how to prevent and/or mitigate these threats, concerns remain. Sulfates continue to leach into the waters of Birch Lake in northeastern Minnesota from the Dunka Pit, which was closed in 1994, despite several efforts to capture pollutants (Johnson, 2025) and they remain a high concern considering the prospects of establishing new copper-nickel mines in the same region. Unfortunately, the published literature regarding the impacts to fish from mining activities are sparse. Regardless, the biggest concern remains the role that high sulfide levels have in increasing the availability of methylmercury which easily accumulates in higher trophic levels, such as predatory fish.



Roads, Trails, and Railroads

Railroads can present an overlooked threat to aquatic organisms as derailments resulting in the release of toxic substances into lakes and rivers can cause fish kills. Although the vast majority of derailments happen in shipping yards, a relatively small number of derailments result in contaminant spills that can cause disproportionate environmental damage. In 2015, a train derailment spilled approximately 20,000 gallons of ethanol into the Mississippi River. Although no fish kills were observed, ethanol can deplete oxygen levels in the water (Pheifer, 2015). A train derailment near Superior, Wisconsin in 1992 spilled 22,000 gallons of chemicals including liquid benzene and toluene into the Nemadji River, killing thousands of fish (Wikipedia, 2025).

Roads, trails, and railroads not only are a source of erosion, sedimentation and pollution into lakes, they can reduce connectivity with crossing structures such as culverts that are insufficiently sized or placed and prevent regular access and movement for fish. Connectivity is especially important for providing fish access to important habitats such as thermal refugia and spawning habitat. Once restoration efforts have occurred in one part of a lake (e.g. along the shoreline or along an inflowing stream), recolonization of these improved habitats depends on the ability of fish to safely and easily access the restored areas. Finally, impervious surfaces, such as tarred roads, parking lots, and trails, adjacent to lakes can lead to increased turbidity and runoff, including the addition of road salts to de-ice surfaces that can increase a lake's salinity.



Shipping Lanes

Maintaining shipping lanes and navigation channels to accommodate commercial boat traffic is essential for economic trade. The Duluth harbor on Lake Superior handles approximately 35 million tons of cargo each year (including iron ore, grain, coal, and cement; [Duluth Seaway Port Authority](#)). Farther south, the Mississippi River moves approximately 589 million tons a year, which accounts for 92% of all U.S. agricultural exports (Waterways Council, 2023). Maintaining these shipping lanes and navigation channels for ships and barges is a priority both for the industries that ship cargo and for the U.S. Army Corps of Engineers who is responsible for dredging. Nevertheless, these activities may directly or indirectly lead to connectivity loss, habitat alteration, water quality changes, and altered hydrology, which all impact fish communities. Routine dredging of river channels, as well as near ports on large lakes, to accommodate commercial boat traffic creates disturbances that prevent natural channel formation, movement, and sediment transport. Field studies in Mississippi River pools that are dredged to maintain a navigation channel for barges and small ships demonstrated numerous impacts. Most important was the loss of habitat heterogeneity as large, deep pools replace shallower ripple and run stretches of river, and the loss of habitat structure as gravel, logs, and other natural debris is removed. Impacts to the fish community include a decrease in species richness and abundance, the replacement of fish dependent on flowing water with fish more suited to still water, and the replacement of habitat specialists, such as darters, with invasive habitat generalists, such as catfish and common carp (Freedman et al., 2012).

Nutrient cycling also is limited in dredged channels because the river has limited access to floodplain habitats that offer an exchange of terrestrial and aquatic nutrients (Schramm et al., 2015). Channel dredging creates constant disturbance that prevents natural channel formation, movement, and sediment transport. Impounded sections of river upstream of locks and dams trap sediment and create warmer water temperatures (Poff & Zimmerman, 2010). Wakes and turbulence from barges can cause excess sedimentation (Spear et al., 2024). All these changes influence the composition of the river's fish community.

Shipping in the Great Lakes does not pose a direct threat to Lake Superior's fish but the ships themselves are a vector for the introduction of invasive species when they release their ballast water. As of April 2010, 89 non-native aquatic species had already been introduced into Lake Superior. Among the long list of non-native species are the following fish: Eurasian ruffe (*Gymnocephalus cernuus*), round goby (*Apollonia melanostomus*), sea lamprey (*Petromyzon marinus*), alewife (*Alosa pseudoharengus*), and rainbow smelt (*Osmerus ordax*). Fortunately, changes to ballast water regulations have reduced the rate of invasive species introductions by an estimated 85% since 2006 (Ricciardi & MacIsaac, 2022). Many biologists consider aquatic invasive species the biggest threat to the Great Lakes (Lake Superior Binational Program, 2014; see also Non-native Invasive Species).



Timber Harvest

Timber harvest conducted in riparian forests along rivers or streams, can impact fish. In streams, for example, harvesting can increase stream discharge, inputs of fine sediments, and water temperatures while decreasing macroinvertebrate diversity and inputs of leaf litter and wood (Merten et al., 2010). This same study, which examined the response of fish to timber harvest in a northern hardwood forest in Minnesota, found that summer air temperatures near harvest sites have the largest impact on the fish community, illustrating the importance of the shade provided by tree and shrub canopies along riparian corridors. Partial harvesting in riparian zones, however, may not impact stream fish communities (Chizinski et al., 2010).



Fishing

Stocking fish for angling can impact non-stocked native fish communities already present (Eby et al., 2006). Species assemblages can change which, in turn, impact trophic levels of the food web as many of the fish introduced, such as northern pike and largemouth bass, are top predators. Some of these introductions may be intentional management actions by resource agencies, some may be “unauthorized” by anglers, or accidental introductions by emptying bait buckets (Vander Zanden et al., 2004). Natural dispersal also occurs. Whether these introduced fish replace native species depends on a variety of factors (Eby et al., 2006). Even smaller, nongame fish could be impacted by introductions, intentional or otherwise, as they may experience increased predation and/or increased harvest pressure for use as bait fish. Smallmouth bass are one example of an introduced fish that can outcompete or predate on native fishes (Loppnow et al., 2013). Native to the Mississippi River, they are commonly introduced to lakes and river systems beyond their natural range. Lake trout, for example, no longer show up in fisheries surveys in Little Trout Lake or Beast Lake in Voyageurs National Park, following unauthorized smallmouth bass introductions (R. Maki, pers. comm.) Stocked fish can also have impacts on water quality, such as the resuspension of sediments caused by common carp, which were intentionally stocked to provide angling opportunities for food fish (Moyle, 1986). Unless otherwise authorized, the release of live bait, bait water, and fish into any Minnesota waters is illegal ([MN Rules 6262](#)).



Recreation

A global review of all studies designed to assess the ecological impacts of water-based recreational activities on freshwater ecosystems found that the most consistently negative impacts of boating and shoreline use were on invertebrates and plants, both of which can indirectly impact fish (Schafft et al., 2021). What the study failed to address, however, was the significant impact that recreational boats can have on introducing invasive species such as Eurasian water milfoil (*Myriophyllum spicatum*) and zebra mussels (*Dreissena polymorpha*), both of which can negatively impact fish populations. Bait bucket releases can also be a risk for spreading invasive species or fish diseases.



Fire Management

Fire can be a stressor for fish populations in landscapes where fire is prevalent. Field studies, however, documenting the response of fish to fire are minimal (Dunham et al., 2003). The most significant stressor is the fire retardant that is aerially sprayed to suppress wildfires. Comprised of water and ammonium phosphate, when it is dropped in lakes and rivers it suffocates fish. Although the United States Forest Service is now required to protect a 300-foot buffer zone around waterways, it can be difficult: “[between](#) 2012 and

2019, the agency dropped 761,282 gallons into water, 95,708 gallons into buffer zones and 248,285 gallons into the habitats of endangered and threatened species. The majority of incidents were accidental, but some were sanctioned [exceptions](#); Forest Service policies consider it acceptable to drop retardant into waterways when life or property is threatened or damage to natural resources outweighs loss of aquatic life (Mohr, 2023).” Although these applications only represented 0.3% of the [279 million gallons](#) used in those years, this could be a significant stressor to local populations.



Dams and Water Management

Dams cause fundamental changes to the ecology of river systems. Aadland (2015) noted that a total of “1,078 dams are fragmenting Minnesota streams” in addition to countless smaller dams and impassable culverts and other barriers to fish passage. Among the changes are interrupted sediment transport, altered nutrient dynamics, decreased fish species diversity upstream of the dam, and the propagation of fish diseases and parasites (Aadland, 2015). Although freshwater fish migrations are generally shorter than those observed for anadromous fish such as salmon, migrations may be necessary for spawning, for reaching optimal foraging areas, to respond to seasonal changes in habitat needs, or to move from drought-stricken waters or waters where water quality has declined. Aadland (2015) noted migration may be particularly important in northern latitudes, such as Minnesota, where severe winters may cause low oxygen levels, extremely cold water conditions, and ice cover resulting in increased stress, disease, and mortality.

Lock and dam operations block the migration and movement of many large fish species, including the SGCN skipjack herring, lake sturgeon, and paddlefish. Many smaller SGCN also face barriers to migration and reduced habitat connectivity when dams and other barriers are installed including the Western sand darter, slimy sculpin, crystal darter, gravel chub, Mississippi silvery minnow (*Hybognathus nuchalis*), and gilt darter (*Percina evides*) (Aadland, 2015). Restricted fish movement further impacts native mussel populations by restricting the movement of host fish that carry and disperse the mussel’s larval glochidia. Dams along the Mississippi River caused the near extirpation of two mussel species from Minnesota ([ebonyshell \(*Fusconaia ebena*\)](#) and elephant ear (*Elliptio crassidens*)). The upstream transport of goods on barges also facilitates the movement of undesirable and deleterious species like invasive zebra mussels that can attach to barges. Impounded sections of river located upstream of locks and dams are often reservoirs for invasive species, allowing them to persist where they would otherwise not occur in naturally flowing river reaches (Johnson et al., 2008). This is especially true for zebra mussels (Stoeckel et al., 2004). Commercial traffic also has been shown to increase suspended sediments and turbidity (Smarts et al., 1985, Spear et al., 2024).

As noted under the stressor Crop Production, extensive efforts to improve drainage for agricultural purposes is a major stressor on fish communities. Straightening stream channels, establishing and deepening ditches, and the installation of tiling systems, have extensively altered the hydrology of Minnesota’s western and southern counties (Blann et al., 2009). Fish that depend on cold, clear waters, such as brook trout and the longnose sucker as well as the federally endangered Topeka shiner are impacted by these alterations.



Invasive Species (Problematic Non-native Species)

As noted under the Shipping Stressor, at least 89 non-native species have been introduced into Lake Superior, and an uncounted number in Minnesota’s other lakes and rivers. Among fish these include the common carp, ruffe, sea lamprey, round goby, tubenose goby (*Proterorhinus semilunaris*), white perch (*Morone americana*) as well as countless other invertebrates and plants. Invasive aquatic plants, for

example, such as Eurasian watermilfoil and curly-leaf pondweed (*Potamogeton crispus*), can grow into dense canopies of aquatic vegetation that block sunlight and reduce oxygen levels leading to fish kills. Dense beds of vegetation also provide shelter to smaller fish, reducing predation levels by larger fish and altering the balance of fish trophic levels; they also reduce the overall diversity of native plants, fish, and aquatic invertebrates ([Minnesota Aquatic Invasive Species Research Center](#) (MAISRC)). The movement of several invasive carp species (e.g. silver carp (*Hypophthalmichthys molitrix*), grass carp (*Ctenopharyngodon idella*), black carp (*Mylopharyngodon piceus*) and bighead carp (*Hypophthalmichthys nobilis*)) up the Mississippi River is especially concerning. Silver and bighead carp are filter feeders that feed on microscopic plants and animals that many native fish depend on for food; grass carp are voracious feeders of plants that provide habitat for many young fish; black carp (not yet observed in Minnesota) eat mollusks, competing with native invertivores and further threatening native mussels ([MAISRC](#)). Upstream movement of invasive carp is related to local hydrologic conditions and passable dams (Fritts et al., 2024). Invasive carp reproduction has not been detected in Minnesota to date but has been detected further downriver (up to Pool 16) where abundance is higher (DNR, 2025). Notably, native fish conditions have been impacted in the Illinois River where there is a very high abundance of silver carp (Irons et al., 2007). Recently, another invasive fish, goldfish (*Carassius auratus*), has become established in many metro area lakes and ponds. They are commonly introduced by pet owners. Their quick rate of reproduction can quickly result in thousands of goldfish disturbing sediments and competing with native fish for food and space (DNR, 2021).



Diseases and Pathogens

Diseases and pathogens that impact fish may be naturally occurring or novel; however, the severity of a disease and its population-level effects may be responsive to water quality changes, altered hydrology, and land use changes. The [Minnesota DNR](#) provides a complete description of 12 common diseases that impact Minnesota fish. Disease, low oxygen levels, and industrial spills are believed responsible for most fish kills that occur on lakes and ponds ([DNR](#)). There is considerable concern about a deadly invasive fish virus, viral hemorrhagic septicemia (VHS) which is present in Lake Superior but has not yet been detected inland.



Water-borne Pollution

The impacts of water-borne pollutants, especially agricultural pesticides, are well-known and widely investigated (Ali et al., 2020). Polychlorinated biphenyls (PCBs), dioxins, per- and polyfluoroalkyl substances (PFAs) as well as toxins from algae can also accumulate in fish and negatively impact their health (see [Minnesota Department of Health: Contaminants in Fish](#)). [Minnesota's Fish Consumption Guidance](#), which initially focused on mercury and PCB contamination, has now been updated to include guidance for waterbodies with high levels of PFAs. Even microplastics, which may be coated with toxic substances, are now showing up in fish in Lake Superior (Hemphill, 2021; Durenberger-Grunow, 2022). The impact of microplastics and other newly emerging contaminants on fish requires further study.

Nonpoint source pollution, including phosphorus, nitrogen, sediment, bacteria, and other contaminants generally results from land runoff, precipitation, atmospheric deposition, drainage, seepage or hydrologic modification. As runoff flows, it picks up and carries away natural and human-made pollutants, depositing them into lakes, rivers, wetlands, groundwater, and estuaries. ([EPA NPS Webpage](#)) These nonpoint source pollutants represent the largest combined contribution (an estimated 86%) to the state's water pollution. ([MPCA Water Quality Initiatives](#)) Water quality measured by nutrients and contaminants has been shown to decline with higher agricultural or urban development in the watershed. Some key impacts include excess algal

growth, reduced plant diversity, and reduced diversity in fish and invertebrate communities. ([DNR Water quality non-point sources](#))



Air-borne Pollution

Mercury may be the best-known contaminant in Minnesota fish populations due to the periodically updated publication of [Minnesota's Fish Consumption Guidance](#) by the Minnesota Department of Health. Its main source is industrial waste and coal-combustion for power generation. Once released, it is converted into methylmercury by microorganisms in soil and waters. Unfortunately, it can be extremely harmful to humans but also to wildlife and fish. It is bioaccumulated in higher trophic levels so predatory fish, such as lake sturgeon, carry much higher concentrations. The Department of Health's guidelines provide the angling public with information on fish mercury levels from waterbodies across the state.



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (MPCA & MDOC, 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al., 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

In the Western Great Lakes region, changing hydrology and air and water temperatures are observed to be some of the largest climate change impacts affecting inland fisheries (Paukert et al., 2021). In Minnesota, air temperatures are increasing, with the largest increases observed in winter and nighttime temperatures ([Climate Trends](#)). Among the documented or expected changes to fish habitat in Minnesota lakes include: 1) increasing water temperature; 2) changing duration and/or timing of ice cover; 3) an extension of the open-water season; 4) changes in water clarity; and 5) increased algal productivity. As a result, coldwater fish populations in many Minnesota lakes will be increasingly stressed. Cisco and brook trout populations will likely be lost in lakes where water temperatures increase while warm-water species increase in abundance. Natural reproduction, recruitment and growth rates of coolwater species may be negatively affected. Unfortunately, warmer temperatures can exacerbate anthropogenic stressors, such as land use and invasive species, resulting in even more complex effects on fish populations. In Minnesota, increased warmer temperatures already are

causing longer periods of lake stratification which, in turn, decrease the abundance of cisco populations (Jacobson et al. 2012, 2019).



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota’s average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. In response, water quality and habitat characteristics of lakes are changing (Jacobson et al., 2019; Jane et al., 2021). Climatologists predict changes in both the timing and intensity of precipitation events which can impact connectivity, lake size and depth, and habitat diversity; it can also disrupt spawning cues (Paukert et al., 2021). In the Midwest, decreased precipitation can increase the salinity of waters while decreasing water levels. Changes in water flow can impact upstream and downstream migrations of fish and decrease water quality as water levels lower causing more turbid conditions. Formerly shallow lakes may become more turbid as nutrient runoff increases with the increase of extreme precipitation events that oscillate with drought conditions (Paukert et al., 2021).

The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Case Study: Restoration of Lake Sturgeon to the Red River of the North

[Lake Sturgeon \(*Acipenser fulvescens*\)](#) were extirpated from the Red River Basin in the mid-1900s. Through partnerships with the Rainy River First Nations, USFWS, White Earth Band, Red Lake Band, North Dakota Game and Fish and South Dakota Fish and Parks, the Red River Basin Lake Sturgeon Restoration Plan was implemented. The plan includes targeted stocking efforts, fish passage barrier removal efforts and population monitoring. Within the basin, 48 of 72 major fish barriers have been modified since 2002, which allows lake sturgeon and 70 other fish species and mussels to move more freely through the connected waterways. In 2022, the first natural spawning event in over 100 years occurred in the system. To learn more, see: [Red River of the North sturgeon restoration](#)

Priority Conservation Strategies

To implement the SGCN Goal of this Plan, to conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions, three strategies were identified:



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Examples of conservation actions are grouped below under these three strategies and tagged with icons for the stressor(s) that they address. Some actions combine multiple strategies, in which case we present it under the one it fits best. Some actions such as those focused on monitoring might not always be tied to a specific stressor; these are represented with ‘n/a’ in the stressor column.

Potential Conservation Actions for Fish



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.

Stressor	Action
	Conduct stream surveys of fish populations and water quality.
	Monitor fish health and transmission pathways for diseases and pathogens. Examples requiring careful consideration and protective strategies include inter-basin water transfer and importation of bait.
n/a	Conduct rigorous effectiveness monitoring to assess responses of SGCN to specific management activities in order to develop improved evidence-based management recommendations and avoidance measures for SGCN (Binley et al., 2025) as part of adaptive management.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.

Stressor	Action
n/a	Treat and slow stormwater runoff. Examples of conservation actions include reducing impervious surfaces and creating rain gardens. Create appropriately sized culverts to mimic natural conveyance or meet management goals.
	Encourage or incentivize agricultural best management practices (BMPs) across Minnesota landscapes. Examples include cattle exclosures in prairie streams, protecting and restoring riparian corridors (especially floodplains and unique habitats thereupon), erosion control BMPs, nutrient reduction BMPs, and approaches to drainage that have reduced impact or habitat benefits.
	Restore stream habitat and connectivity. Examples may include floodplain restorations, including oxbow systems, and aquatic connectivity projects, such as dam removal or modifications and road crossing remediations. Other activities may include stream habitat restorations. Projects within this general activity class should consider other co-occurring aquatic SGCNs and their required habitat features. The potential for invasive species movement (e.g, invasive carp, sea lamprey) should be considered when siting and designing connectivity projects.
	Prevent introductions of invasive species through outreach, regulations, deterrents where appropriate, and watercraft inspection. Carefully consider impacts on native species before stocking nonnative species. Manage existing invasive species populations using best practices, mitigating where possible any impacts on native species. Support research into better tools for prevention and management.
	Create riparian buffers and filtration ponds and plant native species to reduce water-borne pollutants entering bodies of water. Incentivize agricultural practices that increase soil organic matter.
	Protect high quality intact fish habitats and restore watersheds. For coldwater streams, enhance connectivity and riparian buffers to slow the rate of temperature change in the water as ambient temperatures rise. Protect groundwater inputs. Protect coldwater springs and seeps.
	In anticipation of greater frequencies of both floods and droughts, replace undersized culverts and road-crossings with those that are designed to hold up under flooding events and will also optimize aquatic connectivity during low flow periods (see Case Study in Climate Section of Chapter 6. Implementation. Employ culvert design to help support safe, effective wildlife passage based on tested designs.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Stressor	Action
	Develop and disseminate educational materials on the significance of aquatic connectivity to fish populations
	Examine ways to reinforce to the angling public the impacts to native fish populations of releasing live bait fish and fish into any waters of the state.
	Continue to support work to educate the angling and boating public about the negative impact of problematic non-invasive aquatic species; continue boat checks at major public accesses.
	Increase public awareness regarding the presence and impacts of viral hemorrhagic septicemia (VSH) and the importance of removing all visible plants from boats before leaving a water access; draining water from the boat, livewell, motor and bait containers; disposing of unwater bait into the trash; and drying boats before moving them to other water bodies. Education at public accesses in Lake Superior are of primary importance.

Case Study: Brook Trout

Brook Trout are one of two salmonids native to Minnesota. Following European settlement, widespread land use changes degraded cold-water stream habitat, especially in southeast Minnesota and contributed to declining populations of native fishes such as Brook Trout and Slimy Sculpin (Thorn & Ebbers, 1997). Most populations of native Brook Trout were presumed extirpated. Poor land use practices also caused excessive flooding and erosion of important farmland and stimulated the implementation of many conservation practices beginning in the 1930s (Thorn et al., 1997). Expansion of conservation practices reduced flooding and erosion and improved instream habitat. Expanding conservation practices across several decades may have also increased infiltration as groundwater-fed baseflow in many streams and rivers has increased in the Midwest (Juckem et al., 2008; Ayers et al., 2019). In the hope that some native populations persisted through this time period, Hoxmeier et al. (2015) conducted the first expansive genetic survey of remaining Brook Trout populations and identified some populations that were not associated with known hatchery populations, suggesting that native “heritage” Brook Trout still persist in the region. Since about 2020, fisheries managers have been aggressively surveying and prioritizing candidate streams for restoration and reintroduction of native Brook Trout, and we now have robust populations of this SGCN in the state.

References for Fish

- Aadland, L. (2015). Barrier Effects on Native Fishes of Minnesota. Minnesota Department of Natural Resources, Division of Ecological and Water Resources. <https://files.dnr.state.mn.us/eco/streamhab/barrier-effects.pdf>
- Ali, S., Wali, A. F., Yattoo, A. M., Majid, S., Rasool, S., Khan, R., Ali, M. N., Wani, J. A., Farooq, S., Rasool, S., Wani, H. A., & Rehman, M. U. (2020). Effect of pesticides on fish fauna: Threats, challenges, and possible remedies. In *Springer eBooks* (pp. 27–54). https://doi.org/10.1007/978-3-030-35691-0_2
- Ayers, J. R., Villarini, G., Jones, C., & Schilling, K. (2019). Changes in monthly baseflow across the U.S. Midwest. *Hydrological Processes*, 33(5), 748–758. <https://doi.org/10.1002/hyp.13359>
- Binley, A. D., Haddaway, L., Buxton, R., Lalla, K. M., Lesbarreres, D., Smith, P. A., & Bennett, J. R. (2025). Endangered species lack research on the outcomes of conservation action. *Conservation Science and Practice*. <https://doi.org/10.1111/csp2.13304>
- Blann, K. L., Anderson, J. L., Sands, G. R., & Vondracek, B. (2009). Effects of Agricultural drainage on Aquatic ecosystems: a review. *Critical Reviews in Environmental Science and Technology*, 39(11), 909–1001. <https://doi.org/10.1080/10643380801977966>
- Chizinski, C. J., Vondracek, B., Blinn, C. R., Newman, R. M., Atuke, D. M., Fredricks, K., Hemstad, N. A., Merten, E., & Schlessner, N. (2010). The influence of partial timber harvesting in riparian buffers on macroinvertebrate and fish communities in small streams in Minnesota, USA. *Forest Ecology and Management*, 259(10), 1946–1958. <https://doi.org/10.1016/j.foreco.2010.02.006>
- Dunham, J., Young, M., Gresswell, R., & Rieman, B. (2003). Effects of fire on fish populations: landscape perspectives on persistence of native fishes and nonnative fish invasions. *Forest Ecology and Management*, 178(1–2), 183–196. [https://doi.org/10.1016/s0378-1127\(03\)00061-6](https://doi.org/10.1016/s0378-1127(03)00061-6)
- Durenberger-Grunow, L. (2022). Microplastics in Lake Superior are showing up in fish, humans, and everywhere in between. Boreal.Org. July 25, 2022. <https://www.boreal.org/2022/07/25/406621/microplastics-in-lake-superior-are-showing-up-in-fish-humans-and-everywhere-in-between>
- Eby, L., Roach, W., Crowder, L., & Stanford, J. (2006). Effects of stocking-up freshwater food webs. *Trends in Ecology & Evolution*, 21(10), 576–584. <https://doi.org/10.1016/j.tree.2006.06.016>
- Finch, D. M., Butler, J. L., Runyon, J. B., Fetting, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Freedman, J. A., Carline, R. F., & Stauffer, J. R. (2012). Gravel dredging alters diversity and structure of riverine fish assemblages. *Freshwater Biology*, 58(2), 261–274. <https://doi.org/10.1111/fwb.12056>
- Fritts, M., Gibson-Reinemer, D., Appel, D., Lieder, K., Henderson, C., Milde, A., Brey, M., Lamer, J., Turney, D., Witzel, Z., Szott, E., Loppnow, G., Stiras, J., Zankle, K., Oliver, D., Hoxmeier, R. J., & Fritts, A. (2024). Flooding

and dam operations facilitate rapid upstream migrations of native and invasive fish species on a regulated large river. *Scientific Reports*, 14(1). <https://doi.org/10.1038/s41598-024-70076-4>

He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>

Hemphill, Stephanie. (2021). Microplastics proliferate in Lake Superior. Agate. May 16, 2021. <https://www.agatemag.com/2021/05/microplastics-proliferate-in-lake-superior/>

Hoxmeier, R. J. H., Dieterman, D. J., & Miller, L. M. (2015). Brook trout distribution, genetics, and population characteristics in the Driftless area of Minnesota. *North American Journal of Fisheries Management*, 35(4), 632–648. <https://doi.org/10.1080/02755947.2015.1032451>

Irons, K. S., Sass, G. G., McClelland, M. A., & Stafford, J. D. (2007). Reduced condition factor of two native fish species coincident with invasion of non-native Asian carps in the Illinois River, U.S.A. Is this evidence for competition and reduced fitness? *Journal of Fish Biology*, 71(sd), 258–273. <https://doi.org/10.1111/j.1095-8649.2007.01670.x>

Jacobson, P., Cross, T., Zandlo, J., Carlson, B., & Pereira, D. (2012). The effects of climate change and eutrophication on cisco *Coregonus artedii* abundance in Minnesota lakes. *Advances in Limnology*, 63, 417–427. <https://doi.org/10.1127/advlim/63/2012/417>

Jacobson, P.C., Hansen, G.J.A., Olmanson, L.G., Wehrly, K.E., Hein, C.L., & Johnson, L.B. (2019). Loss of coldwater fish habitat in glaciated lakes of the Midwestern United States after a century of land use and climate change. *American Fisheries Society Symposium* 90, 141-157.

Jane, S. F., Hansen, G. J. A., Kraemer, B. M., Leavitt, P. R., Mincer, J. L., North, R. L., Pilla, R. M., Stetler, J. T., Williamson, C. E., Woolway, R. I., Arvola, L., Chandra, S., DeGasperi, C. L., Diemer, L., Dunalska, J., Erina, O., Flaim, G., Grossart, H., Hambright, K. D., . . . Rose, K. C. (2021). Widespread deoxygenation of temperate lakes. *Nature*, 594(7861), 66–70. <https://doi.org/10.1038/s41586-021-03550-y>

Johnson, Chloe. (2025). Old Mine left Toxic Legacy for BWCAW. The Minnesota Star Tribune. June 19, 2025.

Johnson, P. T., Olden, J. D., & Zanden, M. J. V. (2008). Dam invaders: impoundments facilitate biological invasions into freshwaters. *Frontiers in Ecology and the Environment*, 6(7), 357–363. <https://doi.org/10.1890/070156>

Juckem, P. F., Hunt, R. J., Anderson, M. P., & Robertson, D. M. (2008). Effects of climate and land management change on streamflow in the driftless area of Wisconsin. *Journal of Hydrology*, 355(1–4), 123–130. <https://doi.org/10.1016/j.jhydrol.2008.03.010>

Lake Superior Binational Program. (2014). Lake Superior Aquatic Invasive Species Complete Prevention Plan. <https://www.epa.gov/sites/default/files/2015-11/documents/lake-superior-aquatic-invasive-species-complete-prevention-plan-201401-92pp.pdf>

- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Loppnow, G., Vascotto, K., & Venturelli, P. (2013). Invasive smallmouth bass (*Micropterus dolomieu*): history, impacts, and control. *Management of Biological Invasions*, 4(3), 191–206. <https://doi.org/10.3391/mbi.2013.4.3.02>
- M. Jake Vander Zanden, Olden, J. D., James H. Thorne, & Mandrak, N. E. (2004). Predicting Occurrences and Impacts of Smallmouth Bass Introductions in North Temperate Lakes. *Ecological Applications*, 14(1), 132–148. <http://www.jstor.org/stable/4493526>
- Merten, E. C., Hemstad, N. A., Eggert, S. L., Johnson, L. B., Kolka, R. K., Newman, R. M., & Vondracek, B. (2010). Relations between fish abundances, summer temperatures, and forest harvest in a northern Minnesota stream system from 1997 to 2007. *Ecology of Freshwater Fish*, 19(1), 63–73. <https://doi.org/10.1111/j.1600-0633.2009.00389.x>
- Minnesota Department of Natural Resources (DNR). (2021). Goldfish Fact Sheet. (https://files.dnr.state.mn.us/natural_resources/invasives/aquaticanimals/goldfish/goldfish-fact-sheet.pdf)
- Minnesota Department of Natural Resources (DNR). (2025). 2024 Invasive Carp Sampling Report. St. Paul, MN. 32 pp.
- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). (2025). Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state’s greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>
- Mohr, K. (2023). Fire retardant kills fish. Is it worth the risk? High Country News. May 10, 2023.
- Moyle, P.B. (1986). Fish Introductions into North America: Patterns and Ecological Impact. In: Mooney, H.A., Drake, J.A. (Eds.) Ecology of Biological Invasions of North America and Hawaii. *Ecological Studies*, 58. Springer, New York, NY. https://doi.org/10.1007/978-1-4612-4988-7_2
- Pandit, S. N., Poesch, M. S., Kolasa, J., Pandit, L. K., Ruppert, J. L. W., & Enders, E. C. (2024). Long-term evaluation of the impact of urbanization on native and non-native fish assemblages. *Aquatic Invasions*, 19(3), 345–360. <https://doi.org/10.3391/ai.2024.19.3.125642>
- Paukert, C., Olden, J. D., Lynch, A. J., Breshears, D. D., Chambers, R. C., Chu, C., Daly, M., Dibble, K. L., Falke, J., Issak, D., Jacobson, P., Jensen, O. P., & Munroe, D. (2021). Climate change effects on North American fish and fisheries to inform adaptation strategies. *Fisheries*, 46(9), 449–464. <https://doi.org/10.1002/fsh.10668>
- Pheifer, Pat. (2015). More than 18,000 gallons of ethanol spilled into Mississippi River after Wisconsin train derailment. The Minnesota Star Tribune. November 8, 2015.

- Poff, N. L., & Zimmerman, J. K. H. (2010). Ecological responses to altered flow regimes: a literature review to inform the science and management of environmental flows. *Freshwater Biology*, 55(1), 194–205.
<https://doi.org/10.1111/j.1365-2427.2009.02272.x>
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. & LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center.
<https://mwasc.umn.edu/node/821>
- Ricciardi, A., & MacIsaac, H. J. (2022). Vector control reduces the rate of species invasion in the world’s largest freshwater ecosystem. *Conservation Letters*, 15(2). <https://doi.org/10.1111/conl.12866>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.
<https://statesummaries.ncics.org/chapter/mn/>
- Salafsky, N., Relton, C., Young, B.E., Lamarre, P., Bohm, M., et al. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*. 39(3). <https://doi.org/10.1111/cobi.14434>
- Schafft, M., Wegner, B., Meyer, N., Wolter, C., & Arlinghaus, R. (2021). Ecological impacts of water-based recreational activities on freshwater ecosystems: a global meta-analysis. *Proceedings of the Royal Society B Biological Sciences*, 288(1959), 20211623. <https://doi.org/10.1098/rspb.2021.1623>
- Schottler, S. P., Ulrich, J., Belmont, P., Moore, R., Lauer, J. W., Engstrom, D. R., & Almendinger, J. E. (2014). Twentieth century agricultural drainage creates more erosive rivers. *Hydrological Processes*, 28(4), 1951–1961.
<https://doi.org/10.1002/hyp.9738>
- Schramm, H. L., Richardson, W. B., & Knights, B. C. (2015). Managing the Mississippi River floodplain: Achieving ecological benefits requires more than hydrological connection to the river. In *Springer eBooks* (pp. 171–201).
https://doi.org/10.1007/978-1-4939-2380-9_8
- Smarts, M. M., Radar, R. R., Nielsen, D. N., & Claflin, T. O. (1985). The effect of commercial and recreational traffic on the resuspension of sediment in Navigation Pool 9 of the Upper Mississippi River. *Hydrobiologia*, 126(3), 263–274. <https://doi.org/10.1007/bf00007504>
- Spear, M. J., Harris, B. S., Bookout, T. A., Ickes, B. S., Jankowski, K. J., Solomon, L. E., Maxson, K. A., Harris, A. L. W., Mathis, A. T., Schaick, S. J., Williams, J. A., DeBoer, J. A., Lenaerts, A. W., Hine, E. C., Chick, J. H., & Lamer, J. T. (2024). Reduction of large vessel traffic improves water quality and alters fish habitat-use throughout a large river. *The Science of the Total Environment*, 946, 172705. <https://doi.org/10.1016/j.scitotenv.2024.172705>
- Stoeckel, J. A., Rehmann, C. R., Schneider, D. W., & Padilla, D. K. (2004). Retention and supply of zebra mussel larvae in a large river system: importance of an upstream lake. *Freshwater Biology*, 49(7), 919–930.
<https://doi.org/10.1111/j.1365-2427.2004.01237.x>
- Storch, R. L. (1979). Livestock/streamside management programs in eastern Oregon. In Cope, O.B. (Ed.) *Proceedings of the forum – grazing and riparian/stream ecosystems*. Denver, Colorado.

Thorn, W. C., Anderson, C.S., Lorenzen, W.E., Hendrickson, D.L., & Wagner, J.W. (1997). A review of trout management in southeast Minnesota streams. *North American Journal of Fisheries Management*, 17(4), 860-872.

Thorn, W. C., & Ebbers, M. (1997). Brook Trout restoration in southern Minnesota. In R. E. Gresswell, P. Dwyer, and R. H. Hamre (Eds.), *Proceedings of Wild Trout VI Symposium: Putting the native back in wild trout* (pp. 188-192). Montana State University, Bozeman, Montana.

United States Geological Survey (USGS). (2012). Stream Ecosystems Change with Urban Development. <https://pubs.usgs.gov/gip/143/pdf/GIP143.pdf>

University of Chicago, National Opinion Research Center. (2022). 2022 Minnesota Report from the National Survey of Fishing, Hunting, and Wildlife-Associated Recreation. https://www.fishwildlife.org/application/files/2617/3316/0893/Minnesota_FHWAR_Report_Final.pdf#:~:text=In%202022%2C%202.8%20million1%20Minnesota%20residents%20age,such%20as%20observing%2C%20feeding%2C%20and%20photographing%20wildlife.

Walsh, S. J., Jelks, H. L., & Burkhead, N. M. (2011). The decline of North American freshwater fishes. *American Currents*, 36(4), 10-17. <https://www.nanfa.org/ac/freshwater-fishes-decline.pdf>.

Waterways Council, Inc. (2023). Agriculture, shipping industries prepare for uncertainty on Mississippi River in the future. *Democrat Gazette*. November 9, 2023.

Wikipedia. (2025). 1992 Nemadji River Train Derailment. Accessed June 19, 2025. https://en.wikipedia.org/wiki/1992_Nemadji_River_train_derailment

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

WITP. (2024). Reserve Mining. https://wtip.org/wp-content/uploads/2024/05/LSP_Reserve-Mining-text.pdf

Wohl, E. (2017). The significance of small streams. *Frontiers of Earth Science*, 11(3), 447–456. <https://doi.org/10.1007/s11707-017-0647-y>

Mammals

Overview

Minnesota is home to 74 native mammal species with established breeding populations. This includes a recent immigrant to the state, the North American evening bat (*Nycticeius humeralis*). Three species were historically present but are now considered extirpated from Minnesota: free-ranging American bison (*Bos bison*); caribou (*Rangifer tarandus*); and [North American least shrew \(*Cryptotis parva*\)](#). Other mammalian wanderers from adjacent states make an occasional appearance in Minnesota and there are several mammal species whose distributions approach the state borders. Excluded from this list are humans, two non-native rodent species that have established populations in the wild, the brown rat (*Rattus norvegicus*) and house mouse (*Mus musculus*), and several domesticated mammals that have become feral.

Among the mammals identified as Species in Greatest Conservation Need, most Minnesotans will recognize the large and conspicuous [gray wolf \(*Canis lupus*\)](#) and [moose \(*Alces alces*\)](#). However, they may not know much about the smaller mammals on this list, such as [prairie vole \(*Microtus ochrogaster*\)](#), [Northern bog lemming \(*Mictomys borealis*\)](#), or [smoky shrew \(*Sorex fumeus*\)](#). Seven of the eight species of bats found in the state are included but go largely unnoticed due to their nocturnal habits. These range from the more numerous [big brown bat \(*Eptesicus fuscus*\)](#) that frequents buildings to more secretive species such as the silver-haired bat (*Lasionycteris noctivagans*) and [tricolored bat \(*Perimyotis subflavus*\)](#).

For thousands of years mammals in Minnesota provided food and materials for clothing, tools, and shelter to the Ojibwe and Dakota people. Tribes in Minnesota maintained important relationships with American bison, American black bear (*Ursus americanus*), wolf, [wapiti \(elk; *Cervus canadensis*\)](#), snowshoe hare (*Lepus americanus*), North American beaver (*Castor canadensis*), and common muskrat (*Ondatra zibethicus*) (Swanson, 1940).

The arrival and expansion of European settlers caused one of the major stressors to some Species in Greatest Conservation Need (SGCN) through their unregulated take and market hunting that fed rapidly growing urban populations. In addition to the decimation of bison, current SGCN that declined during this period include elk, [Canadian lynx \(*Lynx canadensis*\)](#) gray wolf, and American badger (*Taxidea taxus*) (Swanson, 1940). Although tightly regulated hunting and trapping seasons now help to maintain sustainable populations of these species, concern regarding their status remains. Like all SGCN taxa today, the loss and degradation of habitat is a primary threat that impedes the ability these species to recover from historic losses.

With exploration and settlement of the Minnesota Territory came accounts of the region's abundant wildlife. However, by the time scholarly publications first described the mammals present in this region, they recognized that losses were already occurring due to changing conditions associated with "encroachments of civilization" (Herrick, 1892). Subsequent accounts of the mammals of Minnesota included observations of the impact of human activities, such as tree harvest, pest removal, or fur-trapping, that benefited some mammals while decimating others (Surber, 1932; Gunderson & Beer, 1953; Hazard, 1982). Change has continued to occur since the last comprehensive account of Minnesota's mammals was published in 1982. Statewide surveys, beginning in the late 1980s by the Minnesota Biological Survey, have made significant contributions to our knowledge of the distribution and abundance of many mammal species found in the state.

Long an iconic symbol of Minnesota's Northwoods, one prominent SGCN is the moose. As the state population has dramatically declined, field research has suggested the role that increasing temperatures are likely contributing, and propelling, the species to be emblematic ambassador regarding climate change. The moose population has experienced a 60% population decline since 2006 in northeast Minnesota. The state population estimate in 2006 was 8,840 animals; by 2025 it had declined to 4,040. The latter number actually was an increase above the lowest population estimate in the year 2018 of 3,030 animals ([DNR Aerial Moose Survey](#)). Heat stress, increased winter ticks, and introduction of parasites carried by white-tailed deer (brainworms and liver flukes) whose ranges are increasingly overlapping are key impacts from a changing climate. During summer heat, moose experience elevated body temperatures that sometimes persist until the next day and reduce their ability to forage (Carstensen et al., 2024). Impacts of warming temperatures to other mammalian SGCN associated with Minnesota's boreal forests, such as the Canada lynx and snowshoe hare, have also been noted and are discussed in the following pages.

SGCN and SNI Summary Information

This chapter provides information on 32 mammals that were designated as Species in Greatest Conservation Need (SGCN). In addition, three species were designated as Species in Need of Information (SNI): prairie shrew (*Sorex haydeni*), North American evening bat (*Nycticeius humeralis*), and long-tailed weasel (*Neogale frenata*). None of the SGCN on the 2015 list were removed while five new species were added: gray wolf, northern flying squirrel (*Glaucomys sabrinus*), snowshoe hare, rock vole (*Microtus chrotorrhinus*), and Eastern woodland jumping mouse (*Napaeozapus insignis*).

Among the newly added SGCN, snowshoe hare was selected due to its cultural significance to Native Americans within Minnesota, declines in parts of its range, and anticipated impacts that climate change will have on the species through warming winters and reduced snow duration and depth. The range of the Northern flying squirrel appears to be retracting as Southern flying squirrels (*Glaucomys volans*) expand northward, possibly due to changes in climate and habitats that are bringing the two species into contact. The rock vole and Eastern woodland jumping mouse both have distributions limited to far northeastern Minnesota, where they appear to occur in small, isolated populations that could be vulnerable to local habitat alteration. Gray wolves are federally listed as threatened in Minnesota and endangered in the remaining contiguous 48 states where they are found. The Minnesota population remains vulnerable to the same threats as elsewhere in its range.

Federally listed SGCN include one species as endangered, [Northern long-eared bat \(*Myotis septentrionalis*\)](#), and two federally threatened, Canadian lynx and gray wolf. Two species are state threatened, plains spotted skunk (*Spilogale interrupta*) and [Northern pocket gopher \(*Thomomys talpoides*\)](#); 19 species are state listed as Special Concern. A complete list of both SGCN and SNI species can be found in Appendix B.

In light of recent field studies (see Case Study box) that suggest large regional population declines for both migratory and cave-using populations of bats (Boman, pers. comm.), seven of the eight bat species found in Minnesota are considered SGCN: [little brown bat \(*Myotis lucifugus*\)](#), Northern long-eared bat, tricolored bat, big brown bat, hoary bat (*Lasiurus cinereus*), silver-haired bat, and Eastern red bat (*Lasiurus borealis*). The evening bat, first confirmed in Minnesota in 2016, is a Species in Need of Information as few individuals have been confirmed in the state.

Case Study: Declines in Minnesota's Bats

Bats are a unique group of SGCN mammals in Minnesota, occupying both terrestrial and aerial habitat as the only mammals capable of true flight. Historically, bats have been understudied across the globe compared to other mammal groups, largely due to their cryptic nature, nocturnal activity, and limited technology for studying movements of such small animals. Since the early 2000s, research has expanded exponentially across the continent in response to a deadly bat-specific disease known as White-Nose Syndrome (WNS, Frick et al., 2010). The disease is caused by a non-native fungus, *Pseudogymnoascus destructans*, that grows in cold and humid environments (Gargas et al., 2009), such as the conditions found in caves and mines where bats hibernate. The fungus colonizes the tissues of hibernating bats leading to frequent arousals and energy expenditure, premature loss of critical fat reserves, and mortality rates exceeding 75% two years after infections are documented (Blehert et al., 2009).

In Minnesota, the disease was first confirmed at Soudan Underground Mine in the winter of 2016. Similar to trends throughout North America, severe declines for multiple hibernating bat species were documented throughout the state within five years of confirmation of the disease in hibernacula. By winter of 2019, the population of bats overwintering in Soudan had declined by over 90% just three years after WNS was documented in the mine. This population continued to decrease until reaching a 98% decline by 2024. At the southern border of Minnesota, Mystery Cave also observed declines of 94% a few years after the disease was confirmed, following a similar timeline as the Soudan Mine farther north. All known and accessible hibernacula have documented evidence of WN. As of 2025 Minnesota is considered the western border of the endemic zone (i.e. the zone where the fungus is consistently present which stretches east to the Atlantic coastline and south to Arkansas, Tennessee, and North Carolina ([USGS](#))).

Species in Greatest Conservation Need such as the little brown bat, Northern long-eared bat, and tricolored bat have been heavily impacted by the disease. The fourth cave-hibernating species in Minnesota, the big brown bat, appears to be less susceptible to mortality from the fungus (Frank et al., 2014), and their populations have experienced lower mortality. Little brown bats make up a large portion of the bat declines in hibernacula; they were once the most common bat species in the state and across their known range throughout North America. Northern long-eared bats and tricolored bats were often encountered in Minnesota caves and mines, though in lower numbers compared to little brown bats (Nordquist and Birney, 1985). After WNS infections, these species are encountered much less often during monitoring surveys, particularly the Northern long-eared bat. Due to these declines, the Northern long-eared bat was uplisted from federally threatened to endangered in 2022 (USFWS, 2022); the tri-colored bat and little brown bat are both also being considered for federal listing at the time of this publication.

The need to monitor bat populations through time in response to WNS has generated improved tools for bioacoustics, allowing biologists to record bat vocalizations, also known as bat calls, and to assess changing activity through time. This development has increased the ability to monitor migratory bat species that travel long distances longitudinally, are not known to aggregate in colonies, and are difficult to survey with traditional methods. As their name suggests, migratory bats do not overwinter or use caves in Minnesota, instead they move to areas with milder climates (Hayes et al., 2015). As a result, they do not enter hibernation for any prolonged period (which is when the fungus infests the host bat) and their immune system is able to combat the WNS fungus. However, a disparate threat has emerged for these species that also causes high mortality rates with negative impacts for migratory bat populations.

Migratory SGCN bat species, such as hoary bats, silver-haired bats, and Eastern red bats, are frequently found dead below wind turbines (Arnett et al., 2008). Until recently, it was not understood how the frequency of mortality was impacting regional populations given the limited data on demographics or population status for migratory bats, particularly the hoary bat (Lentini et al., 2015). As a long-distance migrating species, hoary bats comprise the largest portion of fatalities at turbines (Kunz et al., 2007). In 2017, Frick et al. predicted that the estimated mortality in North America per megawatt (installation capacity levels in 2014) for hoary bats could threaten population viability and pose risk of extinction for this species (Frick et al., 2017). With the use of bioacoustics and improved monitoring protocols, empirical evidence of

summertime region-wide decline for hoary bat populations was confirmed in the Pacific Northwest (Rodhouse et al., 2019).

In Minnesota, bioacoustic monitoring was conducted from 2010-2019 by the U.S. Forest Service, Camp Ripley, and Minnesota Biological Survey (DNR). Bioacoustic monitoring uses special detection devices to record and analyze the ultrasonic calls that bats emit to navigate and locate food, enabling researchers to identify which species are present and to assess their abundance. Call abundance for WNS-affected species such as little brown bats and Northern long-eared bats declined 84% after 2015, closely following trends observed in winter hibernaculum (Figure 2.1, Herberg et al., 2020 and Melissa Boman, DNR, pers. comm.). The data collected by these partners also documented a large decline (greater than 78%) in the abundance of migratory *Lasiurus*, the hoary bat and Eastern red bat in Minnesota in the years following 2014 (Figure 2.1, Herberg et al., 2020). Unexpectedly, these trends were similar to cave hibernating species experiencing high mortality from WNS. Statewide, total call abundance declined by 54% from a peak in 2015. Big brown bats and silver-haired bats were combined into one category due to overlapping call parameters. This group composed the largest portion of call files (23%) and did not follow overall trends for steep declines. Due to high levels of interactions between migratory bats and wind turbines, these acoustic data suggest additional monitoring is needed to track migratory bat population trends as wind energy development expands nationwide. For all bat species in Minnesota, more information is needed on summer roosting requirements, important gathering areas, migratory pathways, and management actions that can benefit populations.




Figure 2.1 Graphs indicating population levels of bats in Minnesota for the years 2010-2019, based on bioacoustics data (bat calls). The left panel shows *Myotis* bats (Northern long-eared bat and little brown bat) that overwinter in Minnesota and the onset of white-nose syndrome in the state. The right panel shows data for two migrating bats for the same timeframe, also showing strong declining trends (Herberg et al., 2020 and Melissa Boman, DNR, pers. comm.).

Habitat Associations

Mammal SGCN occupy diverse habitats including grasslands, wetlands, brushlands, peatlands, forested uplands and lowlands, and transitional habitats. Some species are found in a variety of habitats due to their mobility and life history traits. Large predators, such as puma (*Felis concolor*), range widely in search of prey. Bats' ability to fly considerable distances allows them to utilize multiple habitats for foraging, rearing pups, and hibernating. Other SGCN mammals are strongly associated with a narrow range of habitat conditions, including the rock vole and Northern bog lemming.

A summary of mammal SGCN associations with the 15 primary habitat types in the SWAP are portrayed in Table 2.6. Primary habitats are those where the species is found most consistently; the species is most reliant on the habitat and, should the habitat be degraded or lost, it would have the greatest effect on the species. Secondary habitats are those where a species can be found some of the time, but less frequently than a primary habitat. Some species, especially those with more general habitat requirements, may be assigned to multiple primary or secondary habitats. A complete list of all primary and secondary habitats for each SGCN can be found in Appendix C.

Table 2.6. Habitat Associations for Mammal SGCN (no mammals are associated with vernal pools).

Habitats	Primary Habitat	Secondary Habitat	Total
Prairie and Other Grasslands	12	1	13
Savanna	5	13	18
Upland Conifer Forest and Woodland	1	18	19
Upland Deciduous Forest and Woodland	2	20	22
Mesic Hardwood Forest	2	14	16
Deciduous Wet Forest	1	11	12
Coniferous Forest Wetland	1	6	7
Riparian and Floodplain Forest	1	9	10
Non-forested Wetlands	3	3	6
Rivers and Streams	0	9	9
Lakes	0	9	9
Cliff, Talus, and Rock Outcrops	2	12	14
Caves	3	1	4
Urban and Other Developed Lands	2	15	17

Prairies and other grasslands are the primary habitat for the largest number of SGCN mammals. The extensive loss and degradation of this habitat across western and southern Minnesota is responsible for the declining populations and vulnerability of so many species. Many of these SGCNs are small to mid-sized mammals that are limited to grassland habitats. These mammals include the white-tailed jack rabbit (*Lepus townsendii*),

American badger, prairie vole, [Northern grasshopper mouse \(*Onychomys leucogaster*\)](#), [plains pocket mouse \(*Peroznathus flavescens*\)](#), and [Richardson's ground squirrel \(*Uroditellus richardsonii*\)](#).

The important ecological roles that small and mid-sized mammals play in grassland systems were summarized by Ayodele (2025) and highlighted by the DNR [Prairie Wildlife](#) webpage. They include nutrient cycling, soil disturbance, grazing, dispersal of seeds and mycorrhizal fungi, and serving as both predators and prey for other animals. SGCN mammals can be vital to populations of SGCN from other groups. For example, Richardson's ground squirrels are widely cited as being strongly associated with the population health of key SGCN predators such as the [burrowing owl \(*Athene cunicularia*\)](#) and Swainson's hawk (*Buteo swainsoni*) (Houston & Bechard, 1984; Conway, 2018), while white-tailed jackrabbit's selective grazing activities play an important role in the distribution of grassland plants while their burrows can act as seed pits or refuges (Schlater et al., 2021). Both these SGCN species have experienced population declines. White-tailed jackrabbit populations, for example, have significantly declined over the past century, primarily from eradication efforts (e.g., hunting and poisoning) aimed at reducing crop damage in the early 1900s (Brown et al., 2018) and from the loss of the species' preferred habitats, including grasslands, pastures, hayfields, and small grains. Today, the species' population is only a fraction of what it was throughout its range compared to the previous century (Brown et al., 2020) and it is rarely seen in Minnesota ([DNR ARS 2024](#)). Despite this decline, little research on the species has been conducted (Schlater et al., 2021).

A variety of forests provide habitat for many SGCN mammals that have more general habitat requirements. These include many large SGCN mammal species such as gray wolves, moose, and elk, which have had a long history of importance to human communities as apex predators and game species. As a result, these mammals are well studied. Primarily found in the forested regions of Minnesota many of these species are considered boreal species, with the majority of their populations occurring to the north in Canada or to the west in montane regions. They survive, in part, because these regions have been relatively undisturbed compared to the Great Plains and Eastern Deciduous Forest Biomes.

Minnesota forests provide primary habitat for four mammal SGCN species: rock vole (upland conifer and upland deciduous forests); [woodland vole \(*Microtus pinetorum*\)](#); mesic, deciduous wet, and riparian forests); tri-colored bat (upland deciduous and mesic forests); and Eastern heather vole (coniferous forest wetlands). Several small SGCN mammals rely on peatland habitat or mature conifer habitats with rich, damp forest floors that are often covered in sphagnum and ericaceous plants including the [Northern bog lemming \(*Mictomys borealis*\)](#), smoky shrew, and [Eastern heather vole \(*Phenacomys ungava*\)](#).

Eight species of bats are known to occur in Minnesota, seven of which are SGCN, equally divided into two groups that use different strategies for surviving long winters without insects. Migratory bat SGCN (hoary bats, Eastern red bats, and silver-haired bats) leave the state in early fall and head towards regions with milder weather. Evening bats, the only bat species not designated a SGCN, are likely to migrate out of the state, but little is known about overwintering habits in the northern Midwest. Cave bats (little brown bats, big brown bats, tricolored bats, and Northern long-eared bats) hibernate underground for the duration of winter until warm spring weather brings reliable insect hatches. All of the species use forested habitats during the spring, summer, and fall months, many using trees for pup-rearing.

Primary Stressors for Mammals

Stressors are factors that pose direct or indirect challenges to vulnerable plant and wildlife species. Habitat loss and degradation are generally considered the primary threat for most SGCN, and some stressors reduce the quality of those habitats. Other stressors may operate more directly on the SGCN, such as disease and pathogens. In some cases, stressors may operate both directly and indirectly. For example, roads, trails, and railroads can directly harm SGCN through vehicular strikes, or operate indirectly through habitat fragmentation and reduced landscape connectivity. The stressor list below is adapted from an internationally recognized threats classification developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For more information, see the “*Stressors*” section of Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts (e.g. catastrophic wildfire following a prolonged period of fire suppression) while serving as valuable conservation tools in others (e.g., appropriately planned and applied prescribed fire).

Information about a subset of primary stressors specifically impacting this species group is included below, followed by a set of conservation actions addressing those stressors.



Development

Development causes habitat loss and fragmentation and makes traversing the landscape more difficult and dangerous for mammals of all sizes (see also Roads, Trails, and Railroads). This stressor is only projected to increase in the coming years with projections of urban land use increasing by a factor of 1.8 to 5.9 between the years 2000 and 2100 (Gao & O’Neil, 2020). Hanson et al. (2024) reviewed the impacts of urbanization on mammals including changes in the diversity and abundance of species and changes in daily activity patterns. In some urban areas, mammals may actually thrive while in others the impacts of urbanization are neutral or negative. Several factors may be involved, including city-wide landscape factors such as housing density and overall structure. One of the most common behavioral changes is increased activity during nighttime hours in order to avoid humans and/or predators. Overall, Hanson and his colleagues demonstrated the direct relationship between the density/intensity of urbanization and the impacts to mammal communities. Maintaining natural areas near development sites was key to preserving the local mammal diversity, especially for species that cannot prosper in densely populated areas.



Crop Production

Small grassland mammals are impacted by conversion of habitat to rowcrop agriculture. In general, studies have consistently demonstrated that more intensive agricultural practices, including more intensive use of herbicides and pesticides, and more narrow specialization of crop production, reduces the availability of terrestrial invertebrates and subsequently, those higher trophic levels that depend on their abundance, especially small mammals (Balestrieri et al., 2019).

Some agricultural producers view small burrowing animals, such as Richardson’s ground squirrels, Franklin’s ground squirrels (*Spermophilus franklinii*), American badgers, and Northern pocket gophers as pests. The burrowing nature of these species is concerning to some agricultural producers and results in persecution of these animals, including shooting, trapping, and poisoning.



Livestock Management

As Minnesota's wolf population has increased so have the conflicts with livestock operations.

Wolves do kill and/or injure livestock such as cattle, sheep, poultry and occasionally pets. Among livestock, the wolves primarily attack young animals. The number of animals taken is low and impacts less than 1% of all farms in the state, but the financial impact on individual operators can be substantial. Farmers do have the legal authority to kill wolves without a permit if they are "posing an immediate threat to livestock, a guard animal, or a domestic animal located on property owned, leased or occupied by the owner of the livestock, guard animal, or domestic animal" (M.S. 97B.645). During the past ten years the average number of complaints regarding by the USDA Wildlife Services Gray Wolf Depredation Management Program was 174 per year. In 2024 the number increased to 252. Biologists attributed the increase to the lack of snow cover, which made it more difficult for wolves to catch their primary prey, white-tailed deer (Stanley, 2025). The USDA's Wildlife Service's Program works closely with livestock owners to address concerns and to implement changes that help deter and prevent wolf access to young livestock. Landowners also can be compensated for their losses; from about 2010 to 2020 an average of \$135,000 per year was paid to farmers from Minnesota's Wolf Compensation Fund.

Following the reintroduction of elk to northwestern Minnesota in 1935, their populations also have increased, as have conflicts with livestock operations. Most of the damage is in the northwest counties but lone animals occasionally wander outside of the primary range and do damage elsewhere. As the USDA Wildlife Services office reports, "damage typically involves consumption of or damage to stored and stockpiled livestock forage" but can frequently include "damage to infrastructure such as livestock fences (Minnesota Department of Agriculture, 2025)". Like wolves, these conflicts are mitigated with a combination of compensation for fence and crop damage (managed by the MNDA) and abatement (managed by the DNR).

Smaller mammals, on the other hand, may respond differently to livestock grazing operations depending upon how it is managed. For example, SGCN mammals that require very short grasslands (e.g. Richardson's ground squirrel) may benefit from more intensive grazing whereas others (e.g. Franklin's ground squirrel) are associated with taller grasslands.



Mining and Quarrying

Sand and gravel quarries and taconite mining, all significant industrial activities in Minnesota, impact mammals not only by the loss of habitat but also by the increase in noise near these operations. The impacts of noise pollution on mammals have received increased attention in past years, interfering with intra and inter-specific communication and inducing stress responses (Arcangeli et al., 2022; refer to Light and Noise Pollution Stressor). One SGCN, the northern grasshopper mouse, is found on sites with coarse or gravelly soils which include both active and inactive quarry sites. Despite its affinity for this habitat, active quarrying can disturb or destroy their burrows.



Wind and Solar Energy Infrastructure

Migratory, foliage and tree-roosting bats, including SGCN bat species such as hoary bats and eastern red bats, are frequently found dead below wind turbines (Arnett et al., 2008). Although variable in timing, most mortality occurs during late summer and fall, corresponding with migration and mating seasons. Estimates place the loss at approximately 888,000 bats per year (Smallwood, 2013). To date,

lighting and noise appear to not influence the level of mortality. Although several other factors that have been investigated, including tree bats' attraction to large structures resembling trees as well as factors that influence their foraging and mating behavior (e.g. tall trees are an attractant for mating), many questions remain (Guest et al., 2022). Overall, collisions with wind turbines are considered a major, but not the only, factor contributing to the widespread decline of bat populations.



Roads, Trails, and Railroads

Mammals are susceptible to collisions with vehicles on roadways. Some studies have reported that carnivores are less impacted than omnivores and herbivores while nocturnal species, which are more difficult to see, are more seriously impacted. Among SGCN, moose are at some risk for vehicle collision as they use roads and trails for their own travel and as salt licks in the winter. Predators, such as wolves, do utilize secondary roadways and trails in efforts to hunt moose and deer in northeastern Minnesota. Indeed, research in Minnesota on Canada lynx found road and train collision-related mortality (Moen et al., 2005).

In addition to direct mortality, and the impact of roads, trails, and railroads in fragmenting habitats, these travel corridors can introduce other disturbances that also impact mammals (see Recreation and Light and Noise Stressors). Finally, construction and maintenance of bridges can impact bats that use bridge spans for roosting sites.



Utility Corridors

Corridors cleared for utility lines and pipelines may have different impacts on different mammals. Species that depend on forest habitats, such as squirrels and mustelids, usually avoid forest corridors. On the other, the absence of woody cover in many corridors allows predators to more efficiently hunt for prey. They also are attractive to species like the white-tailed deer who, despite being more visible to predators, often forage along these utility lines.

At the landscape scale, habitat fragmentation is a major concern, especially for species that depend on large habitat areas or species that can utilize smaller tracts if they are interconnected. Corridors also create edge habitats which can negatively impact small forest-dwelling mammals by increasing predation. A recent and detailed analysis of the impacts of linear infrastructure on vertebrates found the impacts to mammals were more nuanced than previous studies had suggested; they depend on both the biological traits of each species and the overall environmental context (de Jonge et al., 2022). Carnivorous mammals were more abundant near infrastructure while non-carnivorous species “consistently display avoidance responses.”



Hunting and Collecting Animals

The tradition of hunting large mammals such as deer, elk, and moose is sustainably managed through the establishment of strict regulations and seasons. Small mammals, however, do not receive the same focused management attention and can be impacted by deliberate trapping or shooting. For example, while Northern pocket gophers are rare, the Plains pocket gopher is very abundant and is unprotected, meaning that hunting and trapping is not limited. The collection of bounties for some animals, including gophers, ground squirrels, and woodchucks, may contribute to their decline. Most natural resource professionals, however, doubt their effectiveness in reducing populations. Although the state no longer pays bounties, counties and townships still maintain that authority ([Sec. 348.13 MN Statutes](#)). Payouts are usually

based on the number of feet or tails that are collected, creating a small financial incentive. Today, Northern pocket gophers are very rare or extirpated from Minnesota. Several other SGCN such as Richardson's ground squirrels, Franklin's ground squirrels and American badgers, are still considered pests. Their burrowing nature concerns some agricultural producers, resulting in their continued persecution.



Timber Harvest

Managed timber harvests on public and private lands provide both benefits and losses to Minnesota's mammalian SGCN. Indeed, a recent modeling exercise that examined 41 years of forest management on native wildlife habitat in Minnesota found that over that time span "35.5% of species experienced significant improvement in habitat, 29% significant reductions, and 35.5% nonsignificant change. The extent of habitat (acreage) increased for 100% of species, but the quality declined for 63% of species (Zobel et al., 2021).

Timber harvest activities benefit at least one SGCN mammal, the moose. The young regenerating forests that follow a stand's harvest, creates young, regenerating forests that serve as foraging habitat.

Bats, however, may not all benefit from timber harvest. Another modelling exercise in Indiana examined the impacts of different harvest scenarios on two bats, including the Minnesota SGCN, the Northern long-eared bat. Overall, suitable habitat for the species was negatively correlated with timber harvest intensity. Illustrating the importance of planning forest management activities to simulate the range of natural disturbance regimes, the models documented the bat's preference for forest edges for nocturnal foraging while preferring mature, intact forests for daytime roosting (Pauli et al., 2015).

The snowshoe hare also is impacted by forest management activities, particularly those that decrease structural diversity in the forest. A study by the Leech Lake Band of Ojibwe Division of Resource Management found that snowshoe hare populations were restricted to northern white cedar stands that maintained old growth characteristics. They could not expand much beyond these stands because much of the surrounding forest lacked horizontal structure that provides hiding cover from predators (Tanya Roerick, Leech Lake Band of Ojibwe, pers. comm.). Impacts to smaller mammals, such as shrews, and voles, are less well known. Nevertheless, species that prefer the structural complexity of mature forest stands, including two non-SGCN species, the fisher and marten, benefit from timber harvest strategies that strive to retain these features (Suffice et al., 2023).



Recreation

A recent, thorough review of the literature on the effects of recreational activities on mammals by Sganzerla and colleagues (2025) found that most studies have focused on large, charismatic mammals that rarely are of conservation concern. Only in recent years has this bias begun to be addressed. Although most of the 209 articles they reviewed analyzed the impacts of recreational activities on foot (i.e. hiking and skiing), 74% of those reported at least one negative effect, including avoidance behavior, changes in habitat use, and physiological responses such as stress. Motorized vehicles, aerial disturbances and water-based disturbances were more likely to cause a significant response than hiking.

Regarding Minnesota's mammalian SGCN, recreation is not considered to be a major concern for moose (Larson, pers. com.). Field studies in Colorado that focused on winter recreational activities found that Canada

lynx did not consistently avoid areas of low or moderate dispersed recreation but did avoid some areas with high levels of motorized recreation. When the level of recreational activity was high, lynx were more active during the nighttime hours (Miller, 2019). Finally, in a wilderness setting in Alaska, wolves were found to reduce their presence up to 6 km away from areas of high human use (Thompson et al., 2025). In Minnesota, the Voyageurs National Park Wolf Project has found that wolves are attracted to areas of human development because of the high concentration of white-tailed deer but still avoid any encounters with people while hunting (Johnson-Bice et al., 2023).

Bats can be negatively impacted by recreation in caves, both by physical disturbance during hibernation and by spread of the fungus (*Pseudogymnoascus destructans*) that causes White-nose syndrome (WNS). When people visit caves where bats roost their activities can increase the noise levels and cause local changes in air flow and temperature that wake the bats, causing them to disperse and deplete their energy reserves during a critical time. This is when cave bats are most susceptible to WNS. Even climbing activities outside of the cave, but near its opening, can disturb roosting bats inside the cave. Pounding or chiseling the rock, placement of climbing gear in rocks and crevices, and high noise levels all create disturbances that can negatively impact bats (Schorr et al., 2025).



Fire Management

Proper fire management in grasslands is especially important for mammalian SGCN. Nearly 38% (12 species) of the SGCN depend on prairie and other grasslands as their primary habitat. Species such as the [Western harvest mouse \(*Reithrodontomys megalotis*\)](#), plains pocket mouse, and prairie vole depend on upland prairies. When such sites become overgrown with exotic invasives and woody vegetation due to fire suppression they no longer provide suitable habitat. Fire also was, and continues to be, a natural disturbance in Minnesota's upland forests, helping to create a dynamic mosaic of forest ages and vegetative cover that is critical to many species such as moose. Fires create openings and young, regenerating forest stands that provide important habitat to SGCN that utilize fire-dependent upland conifer and upland deciduous forests in Minnesota. In the absence of fire, timber harvests can be planned to help create this habitat mosaic.



Dams and Water Management

Tilling to convert wet grasslands and pastures to farmland is the major impact of water use management on grassland-dependent mammalian SGCN. Loss of habitat remains the primary threat to all grassland species.



Diseases and Pathogens

Moose are severely impacted by brainworm (*Parelaphostrongylus tenuis*), causing abnormally high mortality in Minnesota populations. Brainworm is carried by white-tailed deer, who have increased their range north into what previously was solely moose territory. Deer are usually not affected by the worm. In moose, on the other hand, *P. tenuis* damages the brain and spinal cord, causing weakness and disorientation. Although less pervasive than brainworm, other parasites that impact moose, occasionally causing death, include liver flukes (*Fascioloides magna*) and hydatid cysts (*Echinococcus granulosus*) that are found in the animal's lungs or liver (Carstensen et al., 2017).

Bat populations of multiple species have been decimated by White-nose syndrome, caused by a fungus (*Pseudogymnoascus destructans*) that wakes bats up from hibernation and reduces their energy stores so they eventually die from starvation or dehydration. In Minnesota, the disease was confirmed at Soudan Underground Mine in the winter of 2016. Species such as the little brown bat, northern long-eared bat, and tricolored bat have been heavily impacted statewide with population declines greater than 90% (Bowman, pers. comm; see Case Study: Bat Declines).



Water-borne Pollution

Aerial applications of herbicides and pesticides that end up in local wetlands, streams, and rivers can potentially impact mammal SGCNs, especially in agricultural regions where their use may be intensive. There also is growing concern about the impacts of microplastics on wildlife. Although most studies have focused on aquatic species, microplastics have been found in some terrestrial mammals including voles and foxes (Jeong et al., 2024).



Air-borne Pollution

Of more concern to mammals are the impacts of aerially applied herbicides and pesticides, especially on grassland species. A study in Europe examined the presence of 140 currently used pesticides and banned or restricted pesticides (i.e. legacy pesticides) in the hair of small mammals and found 112 of the compounds in the hair samples with a total of 32 to 65 chemicals found per individual specimen. The detection and quantity of chemicals found in the specimens directly correlated with the amount of pesticide sold. Little is known about the relative contribution of different pathways of exposure (e.g. direct contact, consumption of treated seeds, or ingestion of soil invertebrates). Some of these chemicals have also been reported in the body tissues of bats. As the authors note, the ubiquity of exposure raises concern not only about acute exposure but about chronic exposure as well (Fritsch et al., 2022). Although this study did not assess the pesticides' impacts to small mammals, they are widely recognized as causing neurological symptoms such as tremors and convulsions; gastrointestinal symptoms such as vomiting; and death.



Light and Noise Pollution

Nearly half of the United States land surface has light-polluted skies (Falchi et al., 2016). Sources of light pollution at night include interior and exterior lights on homes, buildings, billboards, parking lots, and streets as well as lighted sports fields, industrial plants, and airports (DarkSky International, 2025). Studies also point to the ecological impacts created during the daytime by polarized light that reflects off surfaces such as asphalt roads, glass panes, and solar panels (Chock et al., 2020; Horvath et al., 2009).

Several studies have documented the impacts of anthropogenic light on mammals, the majority of which are nocturnal. Bats, for example, can delay their nighttime emergence from daytime roosts and caves, sometimes even abandoning the site (Jagerbrand & Spoelstra, 2023). Individual species, however, may respond differently to lighting conditions. Foraging activity by big brown bats and little brown bats was reduced by nighttime lights while biologists found that eastern red bats and hoary bats responded neither positively nor negatively (Seewagen et al., 2023). Many other mammals decrease their activity in the presence of light and/or change the spaces in which they move (Jagerbrand & Spoelstra, 2023). In addition to these observed impacts, biologists know that bright day and dark night cycles synchronize the internal clocks of animals. Alterations to

these cycles have been shown to impact hormone production, reproductive behavior, neural activity, and metabolic functions (Bumgarner & Nelson, 2021).

Noise pollution from aircraft, vehicles, boats, and other human causes is pervasive. Anthropogenic noise was detected in 36% of national parks surveyed (Buxton et al., 2019), and 12% of wilderness areas had anthropogenic noise levels 3 decibels higher than levels predicted to occur naturally (Buxton et al., 2017). Exposure to increasing noise levels can change the spatial distribution and movement patterns of wildlife, cause avoidance of feeding and nesting areas, and interrupt sleeping patterns and communications (Kok et al., 2023).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Increasing temperatures and prolonged 'hot' temperature events negatively impact moose (*Alces alces*) health in particular. As temperatures rise, moose experience elevated internal body temperatures and these "hot" moose are more likely to die than moose with fewer "hot" episodes (Carstensen et al., 2024). Also, Shorter winters with less snow cover may also impact [Canada lynx \(*Lynx canadensis*\)](#) and snowshoe hare (*Lepus americanus*). University Minnesota-Duluth mammal researcher, Ron Moen also predicts steep declines and even potentially extirpation from Minnesota over the coming 50 - 100 years for four species of shrews and moles, nine species of rodents, the snowshoe hare, moose and three carnivores dependent on the north woods (NRRI, 2018).

Milder winters with limited snow cover and earlier snow melt because of higher-than-normal temperatures create more hospitable environments for some parasites like winter tick (*Dermacentor albipictus*), which can lead to increased tick loads on moose and potentially to moose mortality. Milder winters also can promote the northward range expansion for other parasites and their hosts, as well as insect populations that may adversely impact moose health.

Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota’s average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

In the north woods, the lack of winter precipitation (i.e., snow) or an early snow melt creates a better environment for parasites such as winter tick, which can cause moose mortality. Drier conditions could potentially create a less hospitable environment for other parasites and their vectors, such as brainworm (*Parelaphostrongylus tenuis*) and gastropod hosts.

Priority Conservation Strategies

To implement the SGCN Goal of this Plan, to conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions, three strategies were identified:



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnerships with agencies, Tribes, non-governmental organizations, and private landowners.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management practices.

Examples of conservation actions are grouped below under these three strategies and tagged with icons for the stressor(s) that they address. Some actions combine multiple strategies, in which case we present it under the one it fits best. Some actions such as those focused on monitoring might not always be tied to a specific stressor; these are shown with an 'n/a' in the stressor column.

Potential Conservation Actions for Mammals



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.

Stressor	Action
	Conduct more studies to better understand the conditions that attract bats to wind turbine sites.
	Conduct more research on bat locations and response to timber management. Quantify spatial use needs and territory size requirements.
	Investigate options to reduce the impacts of aerially applied pesticides and herbicides on grassland dependent small mammal SGCN.
	Continue important research and collaborative studies to investigate the impacts of a changing climate on mammalian SGCN and to develop potential mitigation strategies, such as the Nature Conservancy's establishment of conifer strongholds that could benefit conifer-dependent SGCN, including moose.
n/a	Investigate the causes of declines in mammal SGCN, investigating threats, habitat characteristics and effectiveness of conservation measures. Conduct rigorous effectiveness monitoring (Ayodele, 2025) to assess responses of SGCN to specific management activities in priority habitat types to enable the development of improved, evidence-based management recommendations and avoidance measures for SGCN (Binley et al., 2025) as part of adaptive management.
n/a	Support and develop opportunities for community scientists to engage in bat monitoring. Examples would be acoustic monitoring such as has been done by the North American Bat Monitoring Program , and bat box monitoring.

n/a	Increase the number of bat caves that are monitored each winter to help ensure their protection and the sustainability of the wintering bats they provide habitat for each year.
n/a	Improve moose population surveys by exploring improved/new methods and tools (e.g., thermal plane) and by applying advanced analytical methods. Monitor calving and calf survival with new tools (e.g., thermal plane) to improve population estimates.
n/a	Assess fecundity and recruitment of young moose to improve population modeling and info management activities and conduct research to understand predator-prey interactions among moose, deer, wolves, bear, etc.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.

Stressor	Action
	Install bat boxes using recommended guidelines to provide suitable roosting habitat for bats in urban or rural areas, particularly when bat colonies are excluded from building roosts (Boman, 2023; Boman et al., 2024).
	Retain wild areas for mammal SGCN that cannot thrive in developed areas.
	Promote and support restoration of prairie and grassland acres through innovative financial incentives, technical assistance, and engagement.
	Encourage non-lethal techniques for livestock owners to manage wolves. The DNR Wolf Plan includes a wolf, livestock depredation toolbox which includes non-lethal techniques aimed at minimizing wolf depredation.
	Build better captive elk farm fences to keep wild elk from breaking in to help prevent the spread of diseases such as chronic wasting disease and bovine tuberculosis.
	Reduce overgrazing and implement conservation grazing methods that benefit the farmer and wildlife simultaneously. Flash grazing, where livestock graze intensively on small areas for a short period of time, patch burn grazing, the use of prescribed fire to attract cattle to highly palatable vegetation or other methods of adaptive grazing management that are suited to each farmer's conditions and that have been evaluated for their effectiveness.
	At active quarry sites that support populations of the northern grasshopper mouse, inform operators of the species' presence and work to find solutions that allow mining operations and species protection to continue simultaneously. Explore ways to restore expended mine pits to promote recolonization of impacted SGCN.
	Encourage best management practices that reduce mortality, such as reducing blade rotation when wind speeds are low and bats are more active (e.g. late summer and fall) and delaying turbine activation when bat activity is high (Clean Wisconsin, 2023).
	Promote wildlife safe road crossing programs. Advocate for safe crossings when providing technical guidance on roadway projects.

	Minimize new roads and trails and damaging recreational activities in important habitats for mammalian SGCN.
	Co-locate utilities when possible to reduce environmental impact and consider wildlife in siting plans. Avoid routing lines near or across lakes where feasible or along known mammalian travel corridors.
	Consider implementing habitat enhancements for snowshoe hare. This could include increasing structural diversity in the forest by creating slash piles during or after timber harvest, dropping trees to leave as coarse woody debris, planting conifers to increase trees with low hanging branches that offer cover, and protecting areas that have structural diversity and provide snowshoe hare habitat.
	Implement best management practices such as winter harvest and increasing rotational age to improve habitat for bats.
	To manage for the wide array of forest-dependent SGCN, as well as forest-specialist furbearers like marten and fisher, leave some old forest patches or corridors within cuts. Overall protection and proper management of old-growth forests in accordance with the DNR's old-growth forest policy is equally important.
	Support collaborative efforts to implement the large-scale habitat management necessary to sustain moose in Minnesota. Determine moose habitat use/response prior to, during, and after landscape-level habitat management activities. Assess juxtaposition and patch size of key habitat types to ensure adequate forage and thermal cover (winter and summer) are available to inform forest management and promote moose population growth.
	Increase the use of prescribed fire. Proactively apply prescribed fire and grazing in ways that sustain native ecosystems and support SGCN mammals while also avoiding or minimizing negative impacts to them through factors such as timing, intensity, acreage, and leaving unburned areas for escape cover. When developing grazing and burn plans, consider the requirements of SGCN. Aim to provide diversity across the landscape and over time (see also Prairies and Other Grasslands sub-chapter).
	Restrict entry to known hibernacula during the winter months and, where possible, restrict climbing and other recreational activities immediately outside of caves where bats are known to hibernate.
	Encourage the application of best management practices for installing new or upgrading old drainage systems to help reduce the conversion of wet prairies and grasslands to cropland (Board of Soil and Water Resources and the Natural Resources Conservation Service).
	Protect bats from human disturbance and spread of White-nose Syndrome by increasing the number of caves that are gated and monitored.
	Designate specific areas of northern Minnesota as core moose areas and consider how to reduce transmission of brainworm from white-tailed deer. Integrate deer management as part of a comprehensive moose management strategy as outlined in the Minnesota DNR Moose Research and Management Plan (2011) .
	Implement best practices for wildlife-friendly outdoor lighting, such as turning off lights when they are not needed, using timers or motion detectors, shielding lights pointed downward, and employing lights in the warmer part of the spectrum. For more detailed information,

	refer to solutions advised by conservation organizations such as DarkSky , the Xerces Society and National Audubon’s Lights Out Program .



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Stressor	Action
	Engage in education and outreach to raise awareness of persecuted SGCN mammals provide information on their ecological roles.
	Initiate a public information campaign to raise awareness of the impacts of aerial pesticide and herbicide applications on SGCN mammals.
n/a	Develop and promote accessible opportunities for the public to report observations of priority SGCN.
n/a	Design and implement education and outreach activities that raise awareness of bats and their ecological roles and unique adaptations.

Case Study: Minnesota Department of Transportation Develops a New Tool to Identify Bats in Culverts and on Bridges

Minnesota Department of Transportation (MNDOT) takes its responsibility to protect state endangered, threatened and special concern species seriously. Knowing that the Northern long-eared bat is already an endangered species at the federal and state level, and that two other bat species (little brown myotis and tri-colored bat) are under consideration for federal listing, staff began to investigate how their construction and maintenance activities on bridges and culverts could impact these species. These transportation structures have long been known to provide roosting habitat for bats. Given the devastating declines that bats have experienced due to white-nose syndrome, attention has focused on additional measures to protect their declining populations. For example, a recent study in Iowa (Bektas et al., 2018) examined factors that increase a bridge’s suitability as a bat roosting site. Among the factors that increased suitability were the bridge’s proximity to wetlands (important foraging habitat) and the bridge’s height and depth (higher and deeper were most suitable). When there is evidence that bats are using a bridge for roosting habitat (e.g. presence of guano), MNDOT staff must assume that state and federal listed species are present. Physical inspection of the site, however, can cause the very disturbance that staff are trying to avoid. Lacking any other evidence to identify the presence of state or federally listed species, engineers must implement costly measures to protect the bats if a new bridge is being constructed or an old bridge is being maintained or upgraded. A new study by MNDOT begun in 2024 is aimed at addressing this challenge. Researchers are investigating the utility of analyzing the DNA present in bat guano as a means of identifying whether state or federally listed species are present and if special provisions are necessary to protect these species (MnDOT, 2024).

References for Mammals

- Arcangeli, G., Lulli, L. G., Traversini, V., De Sio, S., Cannizzaro, E., Galea, R. P., & Mucci, N. (2022). Neurobehavioral alterations from noise exposure in animals: a systematic review. *International Journal of Environmental Research and Public Health*, 20(1), 591.
- Arnett, E. B., Brown, W. K., Erickson, W. P., Fiedler, J. K., Hamilton, B. L., Henry, T. H., Jain, A., Johnson, G. D., Kerns, J., Koford, R. R., Nicholson, C. P., O'connell, T. J., Piorkowski, M. D., & Tankersley, R. D. (2008). Patterns of bat fatalities at wind energy facilities in North America. *Journal of Wildlife Management*, 72(1), 61–78. <https://doi.org/10.2193/2007-221>
- Ayodele, N. A. (2025). Small mammal communities as indicators of habitat health: A review. *International Journal of Science and Research Archive*, 15(2), 576–594. <https://doi.org/10.30574/ijrsra.2025.15.2.1413>
- Balestrieri, A., Gazzola, A., Formenton, G., & Canova, L. (2019). Long-term impact of agricultural practices on the diversity of small mammal communities: a case study based on owl pellets. *Environmental Monitoring and Assessment*, 191(12). <https://doi.org/10.1007/s10661-019-7910-5>
- Bektas, B. A., Hans, Z., & Phares, B. (2018). Assessing Bridge Characteristics for Use and Importance as Roosting Habitats for Bats. A report from Bridge Engineering Center, Institute for Transportation, Iowa State University. https://publications.iowa.gov/28696/1/bridges_as_roosting_habitats_for_bats_w_cvr.pdf
- Binley, A. D., Haddaway, L., Buxton, R., Lalla, K. M., Lesbarreres, D., Smith, P. A., & Bennett, J. R. (2025). Endangered species lack research on the outcomes of conservation action. *Conservation Science and Practice*. <https://doi.org/10.1111/csp2.13304>
- Blehert, D. S., Hicks, A. C., Behr, M., Meteyer, C. U., Berlowski-Zier, B. M., Buckles, E. L., Coleman, J. T. H., Darling, S. R., Gargas, A., Niver, R., Okoniewski, J. C., Rudd, R. J., & Stone, W. B. (2009). BAT White-Nose syndrome: an emerging fungal pathogen? *Science*, 323(5911), 227. <https://doi.org/10.1126/science.1163874>
- Boman, M., Crawford, R., Pierce, E., Wolff, P., & O'Keefe, J. (2024). Recommendations for use of artificial bat roosts and acoustic lure technology. https://www.denix.osd.mil/ndcee/denix-files/sites/44/2024/04/Guidance_Artificial-Bat-Roost-Construction-Monitoring-Temperature-Acoustic-Lure.pdf
- Boman, M.S. (2023). Altering artificial abodes: Addressing two management concerns of artificial roosts for imperiled bat species. (Doctoral dissertation, University of Illinois at Urbana-Champaign).
- Brown, D. E., Beatty, G., Brown, J. E., & Smith, A. T. (2018). History, status, and population trends of cottontail rabbits and jackrabbits in the western United States. *Western Wildlife*, 5, 16-42.
- Brown, D. E., Smith, A. T., Frey, J. K., & Schweiger, B. R. (2020). A review of the ongoing decline of the white-tailed jackrabbit. *Journal of Fish and Wildlife Management*, 11(1), 341-352.
- Bumgarner, J.R. & Nelson, R. (2021). Light at Night and Disrupted Circadian Rhythms Alter Physiology and Behavior. *Integrative and Comparative Biology*, 61(3), 1160-1169. <https://academic.oup.com/icb/article/61/3/1160/6206372>.

- Buxton, R. T., McKenna, M. F., Mennitt, D., Brown, E., Fristrup, K., Crooks, K. R., Angeloni, L. M., & Wittemyer, G. (2019). Anthropogenic noise in US national parks – sources and spatial extent. *Frontiers in Ecology and the Environment*, 17(10), 559–564. <https://doi.org/10.1002/fee.2112>
- Buxton, R. T., McKenna, M. F., Mennitt, D., Fristrup, K., Crooks, K., Angeloni, L., & Wittemyer, G. (2017). Noise pollution is pervasive in US protected areas. *Science*, 356(6337), 531-533.
- Carstensen, M., Hildebrand, E.C., Plattner, D., Dexter, M., Jennelle, C., & Wright, R.G. (2017). Determining cause-specific mortality of adult moose in northeast Minnesota, February 2013–July 2016. In Cornicelli, L., Carstensen, M., D’Angelo, G., Larson, M.A., & Lawrence, J.S. (Eds.) *Summaries of wildlife research findings 2017*, pp. 188–197. Minnesota Department of Natural Resources, St. Paul, USA.
- Carstensen, M., St-Louis, V., & Tri, A. (2024). Relating Ambient and Body Temperature in Free-Ranging Moose: Implications to Heat Stress and Survival in Minnesota. *Alces*, 59, 111-134.
- Chock, R. Y., Clucas, B., Peterson, E. K., Blackwell, B. F., Blumstein, D. T., Church, K., Fernández-Juricic, E., Francescoli, G., Greggor, A. L., Kemp, P., Pinho, G. M., Sanzenbacher, P. M., Schulte, B. A., & Toni, P. (2020). Evaluating potential effects of solar power facilities on wildlife from an animal behavior perspective. *Conservation Science and Practice*, 3(2). <https://doi.org/10.1111/csp2.319>
- Clean Wisconsin. 2023. Under the Lens: Mitigation bird and bat mortality at wind farms. December 19, 2023. <https://www.cleanwisconsin.org/under-the-lens-mitigating-bird-and-bat-mortality-at-wind-farms/#:~:text=Comparing%20the%20impact%20of%20wind,mortalities%20related%20to%20climate%20change>
- Conway, C. J. (2018). Spatial and temporal patterns in population trends and burrow usage of burrowing owls in North America. *Journal of Raptor Research*, 52(2), 129–142. <https://doi.org/10.3356/jrr-16-109.1>
- DarkSky International. (2025). DarkSky International | Protecting the night skies for present and future generations. Accessed June 2025. <https://darksky.org>
- de Jonge, M. M., Gallego-Zamorano, J., Huijbregts, M. A., Schipper, A. M., & Benítez-López, A. (2022). The impacts of linear infrastructure on terrestrial vertebrate populations: A trait-based approach. *Global Change Biology*, 28(24), 7217-7233.
- Falchi, F., Cinzano, P., Duriscoe, D., Kyba, C. C. M., Elvidge, C. D., Baugh, K., Portnov, B. A., Rybnikova, N. A., & Furgoni, R. (2016). The new world atlas of artificial night sky brightness. *Science Advances*, 2(6). <https://doi.org/10.1126/sciadv.1600377>
- Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weyand, D. M. Finch, C. F. Miniat, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4

- Frank, C. L., Michalski, A., McDonough, A. A., Rahimian, M., Rudd, R. J., & Herzog, C. (2014). The Resistance of a North American Bat Species (*Eptesicus fuscus*) to White-Nose Syndrome (WNS). *PLoS ONE*, 9(12), e113958. <https://doi.org/10.1371/journal.pone.0113958>
- Frick, W. F., Pollock, J. F., Hicks, A. C., Langwig, K. E., Reynolds, D. S., Turner, G. G., Butchkoski, C. M., & Kunz, T. H. (2010). An emerging disease causes regional population collapse of a common North American bat species. *Science*, 329(5992), 679–682. <https://doi.org/10.1126/science.1188594>
- Frick, W., Baerwald, E., Pollock, J., Barclay, R., Szymanski, J., Weller, T., Russell, A., Loeb, S., Medellin, R., & McGuire, L. (2017). Fatalities at wind turbines may threaten population viability of a migratory bat. *Biological Conservation*, 209, 172–177. <https://doi.org/10.1016/j.biocon.2017.02.023>
- Fritsch, C., Appenzeller, B., Burkart, L., Coeurdassier, M., Scheifler, R., Raoul, F., Driget, V., Powolny, T., Gagnaison, C., Rieffel, D., Afonso, E., Goydadin, A., Hardy, E. M., Palazzi, P., Schaeffer, C., Gaba, S., Bretagnolle, V., Bertrand, C., & Pelosi, C. (2022). Pervasive exposure of wild small mammals to legacy and currently used pesticide mixtures in arable landscapes. *Scientific Reports*, 12(1). <https://doi.org/10.1038/s41598-022-19959-y>
- Gao, J., & O'Neill, B. C. (2020). Mapping global urban land for the 21st century with data-driven simulations and Shared Socioeconomic Pathways. *Nature Communications*, 11(1). <https://doi.org/10.1038/s41467-020-15788-7>
- Gargas, A., Trest, M., Christensen, M., Volk, T., & Blehert, D. (2009). *Geomyces destructans* sp. nov. associated with bat white-nose syndrome. *Mycotaxon*, 108(1), 147–154. <https://doi.org/10.5248/108.147>
- Guest, E. E., Stamps, B. F., Durish, N. D., Hale, A. M., Hein, C. D., Morton, B. P., Weaver, S. P., & Fritts, S. R. (2022). An updated review of hypotheses regarding bat attraction to wind turbines. *Animals*, 12(3), 343. <https://doi.org/10.3390/ani12030343>
- Gunderson, H.L., & Beer, J.R. (1953). The Mammals of Minnesota. Minnesota Museum of Natural History, Occasional Papers, 6, 1-190.
- Hansen, C. P., Kays, R., & Millspaugh, J. J. (2024). From backyard to backcountry: changes in mammal communities across an urbanization gradient. *Journal of Mammalogy*, 105(1), 175–191. <https://doi.org/10.1093/jmammal/gyad110>
- Hayes, M. A., Cryan, P. M., & Wunder, M. B. (2015). Seasonally-Dynamic Presence-Only species distribution models for a cryptic migratory bat impacted by wind energy development. *PLoS ONE*, 10(7), e0132599. <https://doi.org/10.1371/journal.pone.0132599>
- Hazard, E. B. (1982). The Mammals of Minnesota. University of Minnesota Press. 279 pp.
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>

- Herberg, A.M., Boman, M.S., Catton, T.J., Nordquist, G.E., Dietz, N.J., Dirks, B.J., & Berndt, D. (2020). Changes in Bat Call abundance pre- and post- white-nose syndrome using mobile acoustic surveys in Minnesota. Poster presented at Midwest Bat Working Group Annual Meeting 2019.
- Herrick, C. L. (1892). The mammals of Minnesota. Bulletin (Geological Natural History Survey of Minnesota), 7, 1-299.
- Horváth, G., Kriska, G., Malik, P., & Robertson, B. (2009). Polarized light pollution: a new kind of ecological photopollution. *Frontiers in Ecology and the Environment*, 7(6), 317–325. <https://doi.org/10.1890/080129>
- Houston C. S. & Bechard, M.J. (1984). Decline of the ferruginous hawk in Saskatchewan. *American Birds* 2, 166-170. <https://sora.unm.edu/sites/default/files/journals/nab/v038n02/p00166-p00170.pdf>
- Jägerbrand, A. K., & Spoelstra, K. (2023). Effects of anthropogenic light on species and ecosystems. *Science*, 380(6650), 1125–1130. <https://doi.org/10.1126/science.adg3173>
- Jeong, E., Lee, J. Y., & Redwan, M. (2024). Animal exposure to microplastics and health effects: A review. *Emerging Contaminants*, 10(4), 100369.
- Johnson-Bice, S. M., Gable, T. D., Homkes, A. T., Windels, S. K., Bump, J. K., & Bruggink, J. G. (2023). Logging, linear features, and human infrastructure shape the spatial dynamics of wolf predation on an ungulate neonate. *Ecological Applications*, 33(7), e2911.
- Kok, A. C. M., Berkhout, B. W., Carlson, N. V., Evans, N. P., Khan, N., Potvin, D. A., Radford, A. N., Sebire, M., Sabet, S. S., Shannon, G., & Wascher, C. a. F. (2023). How chronic anthropogenic noise can affect wildlife communities. *Frontiers in Ecology and Evolution*, 11. <https://doi.org/10.3389/fevo.2023.1130075>
- Kunz, T. H., Arnett, E. B., Erickson, W. P., Hoar, A. R., Johnson, G. D., Larkin, R. P., ... & Tuttle, M. D. (2007). Ecological impacts of wind energy development on bats: questions, research needs, and hypotheses. *Frontiers in Ecology and the Environment*, 5(6), 315-324.
- Lentini, P. E., Bird, T. J., Griffiths, S. R., Godinho, L. N., & Wintle, B. A. (2015). A global synthesis of survival estimates for microbats. *Biology Letters*, 11(8), 20150371. <https://doi.org/10.1098/rsbl.2015.0371>
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Miller, S. (2019). Winter Sports and Wildlife: Can Canada lynx and winter recreation share the same slope? Rocky Mountain Research Station Science You Can Use Bulletin, United States Department of Agriculture. January/February 2019, 33. https://www.fs.usda.gov/rm/pubs_journals/rmrs/sycu/2019/sycu_033_2019_01_winter_sports.pdf
- Minnesota Department of Agriculture (2025). Cervidae Depredation on Minnesota Farms: Impacts, Current Mitigation Strategies, and Recommendations. <https://www.lrl.mn.gov/docs/2025/mandated/250297.pdf>

Minnesota Department of Transportation (MnDOT). (2024). New Project: Species to Feces: A New Tool to Identify Bats in Culverts and on Bridges. MNDOT Crossroads: Minnesota's transportation research blog. June 27, 2024. <https://mntransportationresearch.org/2024/06/27/new-project-species-from-feces-a-new-tool-to-identify-bats-in-culverts-and-on-bridges/>

Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). 2025. Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state's greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/Iraq-3sy25.pdf>

Moen R., Niemi, G., Burdett, C.L., & Mech, L.D. (2005). Canada lynx in the Great Lakes Region: 2005 annual report to USDA Forest Service, MN Cooperative Fish and Wildlife Research Unit, and Minnesota Department of Natural Resources. NRRI Technical Report No. NRRI/TR-2006-16. Center for Water and Environment, University of Minnesota, Duluth. 31 pp.

Natural Resources Research Institute (NRRI). 2018. NRRI prepares state for changes to mammal populations. NRRI News. February 4, 2018. <https://nrri.umn.edu/news/climate-mammals>.

Nordquist, G.E. & Birney, E.C. (1985). Distribution and status of bats in Minnesota. Final report submitted to the Nongame Wildlife Program, Minnesota Department of Natural Resources. 64 pp.

Pauli, B. P., Zollner, P. A., Haulton, G. S., Shao, G., & Shao, G. (2015). The simulated effects of timber harvest on suitable habitat for Indiana and northern long-eared bats. *Ecosphere*, 6(4), 1-24.

Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwcas.umn.edu/node/821>

Rodhouse, T. J., Rodriguez, R. M., Banner, K. M., Ormsbee, P. C., Barnett, J., & Irvine, K. M. (2019). Evidence of region-wide bat population decline from long-term monitoring and Bayesian occupancy models with empirically informed priors. *Ecology and evolution*, 9(19), 11078-11088.

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>

Schlater, S. M., Ringenberg, J. M., Bickford, N., & Ranglack, D. H. (2021). White-tailed jackrabbits: a review and call for research. *The Southwestern Naturalist*, 65(2), 161-172.

- Schorr, R. A., Warren, Z. A., Goodwin, K., Murdock, E., & Neubaum, D. J. (2025). Collaborative conservation of cave-roosting bats: guidance on managing rock climbing near caves. *Frontiers in Conservation Science*, 6, 1411427.
- Seewagen, C. L., Nadeau-Gneckow, J., & Adams, A. M. (2023). Far-reaching displacement effects of artificial light at night in a North American bat community. *Global Ecology and Conservation*, 48, e02729.
<https://doi.org/10.1016/j.gecco.2023.e02729>
- Sganzerla, F., Scillitani, L., Brivio, F., & Grignolio, S. (2025). The effects of recreational activities on wild mammals. *Global Ecology and Conservation*, e03485.
- Smallwood, K. S. (2013). Comparing bird and bat fatality-rate estimates among North American wind-energy projects. *Wildlife Society Bulletin* 37, 19-33.
- Stanley, G. Reports of wolves killing livestock in Minnesota reach a record high. The Minnesota Star Tribune. February 26, 2025. <https://www.startribune.com/reports-of-wolves-killing-livestock-in-minnesota-reach-a-record-high/601228271>
- Suffice, P., Mazerolle, M. J., Imbeau, L., Cheveau, M., Asselin, H., & Drapeau, P. (2023). Site occupancy by American martens and fishers in temperate deciduous forests of Québec. *Journal of Mammalogy*, 104(1), 159-170.
- Surber, T. (1932). The Mammals of Minnesota. Minnesota Game and Fish Department. 84 pp.
- Swanson, E.B. (1940). The Use and Conservation of Minnesota Wildlife 1850–1900. [Master’s thesis, University of Minnesota.] <https://www.lrl.mn.gov/docs/2007/other/070554.pdf>.
- Thompson, P. R., Paczkowski, J., Whittington, J., & St. Clair, C. C. (2025). Integrating human trail use in montane landscapes reveals larger zones of human influence for wary carnivores. *Journal of Applied Ecology*, 62(2), 344-359.
- U.S. Fish and Wildlife Service (USFWS). (2022). Endangered and Threatened Wildlife and Plants; Endangered Species Status for Northern Long-Eared Bat. 50 CFR Part 17. Department of the Interior.
<https://www.federalregister.gov/documents/2022/11/30/2022-25998/endangered-and-threatened-wildlife-and-plants-endangered-species-status-for-northern-long-eared-bat>
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report].
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>
- Zobel, J. M., Ek, A. R., & Edgar, C. B. (2021). Assessing the impact of 41 years of forest management on native wildlife habitat in Minnesota, USA. *Journal of Forestry*, 119(2), 164-176.

Reptiles

Overview

Despite its cold climate, Minnesota is home to a diverse assemblage of reptile species – 9 turtles, 16 snakes, and 3 lizards. In 1944 Walter Breckenridge published *Reptiles and Amphibians of Minnesota*, marking the state's beginning of modern herpetology. Breckenridge's publication was considered Minnesota's standard herpetological reference until *Amphibians and Reptiles Native to Minnesota* was published by Oldfield and Moriarty in 1994, followed by an update in 2014 (Moriarty and Hall). In the late 1970s, the Minnesota DNR's Nongame Wildlife Program was established, and has since funded studies on [wood turtles \(*Glyptemys insculpta*\)](#), [Blanding's turtles \(*Emydoidea blandingii*\)](#), [timber rattlesnakes \(*Crotalus horridus*\)](#), [gophersnakes \(*Pituophis catenifer*\)](#), [five-lined skinks \(*Plestiodon fasciatus*\)](#), and more. The Minnesota Biological Survey, initiated in 1987, surveyed counties across the state targeting rare wildlife and collecting records on common species. Their work has helped establish Minnesota's list of reptile SGCNs and has added thousands of records of reptiles to the DNR's Natural Heritage database.

Reptiles play an important role as both predator and prey in their ecosystems. Snakes eat many mammal species that humans consider pests and also reduce numbers of disease-carrying ticks that are attached to those mammals (Kabay et al. 2013). Turtles create and modify habitats as they dig burrows or nests, and function as seed dispersers (Lovich et al. 2018). Conservation and research initiatives targeting reptile species have historically been underfunded and underrepresented despite the ecological importance of these species and particular life history requirements that make them vulnerable to decline, such as low reproductive rate, reliance on rare habitat types, and sensitivity to changes in temperature (Gibbons, 1988; Todd et al., 2010; Christoffel & Lepczyk, 2012; Tingley et al., 2016). In part, this may be due to traits that make them more difficult to study, including the rarity and cryptic nature of many species (Crane et al., 2021) and the fear and disdain that members of the public often feel towards snakes, lizards, and turtles relative to charismatic bird and mammal species (Olson & Pilliod, 2022). SGCN reptiles need focused attention to ensure their long-term conservation. The [Minnesota Herpetological Society](#) is an important venue for public education and promoting the study and conservation of both reptiles and amphibians. [Herpmapper](#), a community science platform designed to gather reptile and amphibian observations worldwide, hosts the [Amphibian & Reptile Survey of Minnesota](#), which provides information on Minnesota's herpetofauna and collects observations of Minnesota's herpetofauna, including turtle road crossing and mortality hotspots through the Minnesota Turtle Crossing Tally and Count Project. On a larger scale, [Midwest Partners in Amphibian and Reptile Conservation](#) (PARC) is an important network dedicated to the conservation of reptiles and amphibians that provides an avenue for collaborative conservation initiatives, outreach and education resources, and more.

Case Study: Wood Turtle Nest Success

As part of a USFWS Competitive State Wildlife Grant, the Minnesota DNR Nongame Wildlife Program undertook a project in northeastern Minnesota to identify threats to wood turtles and effective conservation actions to address those threats. One area of focus was nest success. Biologists monitored wood turtle nests and found that 5% of nests were successful, with predation the primary cause of nest failure. Badgers were responsible for 85% of predated nests, followed by raccoons, ravens, skunks, and foxes. Different nest protection methods were trialed, including an electric fence around a nest site, nest boxes which completely enclose a nest site with a small opening at the bottom for turtles, and deploying nest cages over individual nests. The different nest protection methods have pros and cons in terms of amount of effort and logistics of working in remote areas. Overall, nest success increased to 50% with nest protection efforts. Nest protection has been a successful conservation action, and these efforts have led to a minimum of 630 wood turtle hatchlings.

The real measure of success will be finding out if these conservation actions are having an impact on the overall wood turtle population. To answer that question, DNR biologists have established long-term monitoring sites with a mark-recapture methodology. The goal is to monitor the sites every 5 years to obtain information on abundance, age-class structure, and sex ratio. Some long-term monitoring sites are controls, where no conservation actions are occurring. Other long-term monitoring sites are treatments, with conservation actions like nest protection and enhancing nesting habitat by removing encroaching vegetation and creating nest sites that are more flood-safe. If these conservation efforts are successful, biologists expect to see an increase in turtles in the younger age classes in the coming years in the treatment monitoring sites. See Brown et al., 2025 for more information.

SGCN and SNI Summary Information

The 2025 SGCN list includes 13 reptiles (46%) of the 28 reptiles in the state and one Species in Need of Information (SNI), the Southern map turtle (*Graptemys ouachitensis*) (See Appendix B). The SGCN include 8 snakes, three turtles, and two lizards. Four species are state-listed as Threatened, and six as Special Concern species. Two species, the Blanding's turtle and wood turtle, are presently under consideration for federal listing. Two species that were on the 2015 list were not included on this update, the Northern ring-necked snake (*Diadophis punctatus*), which is now thought to have a stable population in Minnesota, and [*massasauga* \(*Sistrurus catenatus*\)](#), due to extirpation.

Habitat Associations

Reptile SGCN utilize a variety of habitats, but the highest number of species were associated with savanna, prairie and other grasslands, upland deciduous forest and woodland, and cliff, talus, and rock outcrops (see Table 2.7). Cliff, talus, and rock outcrops provide rock surfaces essential for basking and thermoregulation in snakes and lizards, while rock and talus crevices provide important shelter and nesting sites.

Table 2.7 summarizes the association of reptile SGCN with the habitat types identified in the 2025-2035 SWAP. Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently. Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C.

Table 2.7. Numbers of species of reptile SGCN associated with each of the 15 main habitat types (only showing habitats with reptile SGCN associations).

Habitats	Primary Habitat	Secondary Habitat	Total
Prairie and Other Grasslands	11	0	11
Savanna	12	0	12
Upland Conifer Forest and Woodland	2	0	2
Upland Deciduous Forest and Woodland	9	0	9
Mesic Hardwood Forest	5	0	5
Deciduous Wet Forest	3	0	3
Riparian and Floodplain Forest	2	0	2
Non-forested Wetlands	1	5	6
Rivers and Streams	3	0	3
Lakes	1	0	1
Vernal Pools	0	1	1
Cliff, Talus, and Rock Outcrops	8	0	8
Urban and Other Developed Lands	0	3	3

Across their entire range, common five-lined skinks are typically found in woodlands. However, in Minnesota they are primarily associated with limestone and granite rock outcrops and bluff prairies. Another lizard SGCN, the six-lined racerunner (*Aspidoscelis sexlineatus*), is also found in prairie, savanna, and rocky habitats, usually in areas with sand or gravel.

The [North American racer \(*Coluber constrictor*\)](#) utilizes dry habitats including edges of deciduous woodlands, prairies, rock outcrops, and old crop fields. [Plains hog-nosed snakes \(*Heterodon nasicus*\)](#) are a prairie species that prefer sandy or gravelly areas to burrow in. The closely related Eastern hog-nosed snake (*Heterodon platirhinos*) also prefers sandy or loamy soil, but occupies a wider range of habitats including grasslands, open woodlands and forest edges, and river shorelines and floodplains. The smooth greensnake (*Opheodrys vernalis*) is associated with prairies and meadows, edges of mixed forests, and the edges of marshes and bogs. The gophersnake prefers open landscapes like prairies, savannas, pastures, and old cropland, as well as rocky bluff habitats. The [lined snake \(*Tropidoclonion lineatum*\)](#) can be found in open prairies with large amounts of exposed rock. The timber rattlesnake requires hillsides with bluff prairies and rock outcrops, where they breed and overwinter in dens. Nearby forests, forest edges, and open fields are used as feeding grounds in the summer. The [western ratsnake \(*Pantherophis obsoletus*\)](#) is a woodland species and prefers forested uplands and valleys with rivers and rock outcrops.

Blanding's turtles prefer open areas with shallow, slow-moving water, like marshes, ponds, and bogs. During the winter, they overwinter at the bottom of these bodies of water. Wood turtles utilize rivers and adjacent floodplains and uplands, primarily in forested areas. They nest on sandbars and riverbanks. The smooth softshell is found in large rivers with moderate to fast currents that have sandy or muddy bottoms, which they burrow in during the winter.

Primary Stressors for Reptiles



Development

Development causes habitat loss and fragmentation and makes traversing the landscape more difficult and dangerous for reptiles (see also Roads, Trails, and Railroads). Development also can increase the abundance of mesopredators, such as raccoons, badgers, foxes and skunks, which prey on reptiles (Gelvin-Innvaer et al., 2025). Erosion control materials including rip rap and mesh, such as those used in construction of roads and to stabilize stream banks, can pose hazards as well, ensnaring snakes, lizards and turtles. (Stuart et al., 2001; Barton & Kinkead, 2005; Walley et al., 2005; Kingsbury & Gibson, 2012).



Crop Production

Row crop agriculture has converted native prairie and grasslands that reptiles like Blanding's turtles, gophersnakes, and plains hog-nosed snakes depend on (Congdon et al., 2008; Moriarty & Hall, 2014). Suitable nesting habitat may be a limiting factor for some SGCN reptiles (MWPARC, 2010). Cultivated fields often become the only remaining option for nesting, yet these open, disturbed soils can also be "ecological sinks." For example, female Blanding's turtles have been found to preferentially select bare crop fields for nesting over naturally occurring nest sites (Mui et al., 2016). As crops grow, they shade nests from the sun's warmth which impacts egg development, can lead to skewed sex ratios and reduce nest success (Freedberg & Pappas, 2011). Reptiles, such as snakes and turtles, are also injured and killed by farming equipment (Kapfer et al., 2008; Saumure & Bider, 1998; Saumure et al., 2007). Nests and eggs may be subject to destruction from mid-season tillage, and soil compaction from heavy equipment that could make it difficult for hatchlings to emerge from the nest (Dowling et al., 2010; Henning & Hinz, 2016; Chessman, 2017; King et al., 2025.) Agricultural chemical application also poses threats to turtles and their unhatched eggs. Turtles are long-lived and therefore may also be prone to bioaccumulation of toxins that could have lethal and/or sublethal effects (DNR, 2020). Lastly, row crop agriculture can negatively impact water quality, hydrology, and plant communities of wetlands, streams, and shallow lakes that SGCN turtles rely on and some SGCN snakes like gophersnakes use in parts of Minnesota for thermoregulation, foraging, and travel (DNR unpubl. data.)

SGCN reptiles use haylands and other managed grassy areas to meet a variety of needs, including for nesting and foraging (Knoot & Best, 2011; Dowling et al., 2010; Wittenberg, 2012; DNR, 2020). Haying for livestock forage and mowing, such as for roadside maintenance, filtration and buffer strips, or firebreaks, pose significant risks of injury and mortality to rare reptiles, especially turtles and snakes (Dalrymple & Reichenbach, 1984; Durbian, 2006; Wiese, 2016). The nesting period is the most vulnerable period for Blanding's turtles (Henning & Hinz, 2016; King et al., 2025). Wood turtles may also be at risk when foraging in uplands (Saumure et al., 2007; DNR, 2020).



Mining and Quarrying

Hard rock mining operations may remove or degrade rock outcrop habitats and put reptiles at direct risk of harm by equipment. Sand and gravel quarries can attract nesting turtles, increasing the risk of turtles being injured or killed and nests being destroyed by heavy equipment and blasting as part of mining operations (Zagorski et al., 2019; DNR, 2020).



Wind and Solar Energy Infrastructure

Solar farms shade nesting and basking areas for Blanding's turtles, including agricultural lands which may be the only available nesting areas in the vicinity of the turtle's aquatic habitat.

Additionally, unless fence design specifically incorporates periodic openings to accommodate turtles, fencing around solar facilities can block turtle movements and habitat use (MWPARC, 2010).

Easements for wind energy facilities often preclude acquisition and easement protections of adjacent open lands for conservation which in turn affects reptiles dependent on those habitats. (See also Prairies and Other Grasslands chapter).



Roads, Trails, and Railroads

Roads fragment reptile habitat and create a hazard for species that need to cross this barrier to reach habitats necessary for their survival and reproduction. Transportation infrastructure and vehicles affect reptiles in many ways (Andrews et al., 2006). Road mortality poses a major threat to snakes and turtles including Blanding's turtles (King et al., 2025; MWPARC, 2010) and gophersnakes (Kapfer et al., 2008). Railroads have also been documented as a mortality risk (King et al., 2025). Roads can attract cold-blooded reptiles seeking a warm surface to raise their body temperature. Gravel roads and sparsely vegetated road shoulders with disturbed soils may attract nesting turtles which increases risks to breeding females, their nests, and hatchlings from traffic and roadway maintenance (Jones et al., 2024). The turtles' slow movements and habit of withdrawing into their shells when perceiving a threat, such as an oncoming vehicle, further adds to the risks of crossing a road (Gooley, 2010).



Utility Corridors

Directional boring and trenching for utilities (cable and pipelines) can impact reptiles directly and indirectly. Extensive excavation may destroy reptile nests and hibernation dens. Availability of suitable hibernacula can be a limiting factor for snakes in northern latitudes including gophersnakes ("bullsnakes") (Johnson et al., 2022; Kapfer et al 2008; Martino et al., 2012) that overwinter in deep burrows. The DNR has documented Blanding's turtles overwintering in stream banks, including concentrations of many turtles in burrows that extend deep into the banks at least 10 feet (DNR unpubl. data). Such sites could be subject to utility boring (e.g. at stream crossings). Unfortunately, overwintering locations are not universally documented for all Blanding's turtle populations which hinders environmental review.



Hunting and Collecting Animals

In Minnesota, taking, importing, transporting or selling any state endangered or threatened reptile is prohibited without a state permit (M.S. 84.0895) (e.g. timber rattlesnake, Blanding's turtle, wood turtle, and western ratsnake; MN Rules chapter 6134). Native turtles also may not be sold as pets in the state. Although state residents with a fishing license and recreational turtle license may harvest common snapping turtles and western painted turtles for their personal use, commercial harvesting of these species was prohibited beginning in January 2024. The statute has exemptions that provide for the take of painted and snapping turtles by minors to continue the long-standing tradition of summer turtle races in small towns in Minnesota (M.S 97C.605). Once collected, and the races are over, the turtles cannot be released to the wild; their long-term care is the responsibility of the collector.

These rules and regulations provide a strong network of protection for Minnesota's native reptile species although illegal collection still occurs and poses a risk factor especially for imperiled species in Minnesota (DNR, 2020; Christman et al., 2024). Occasionally, a Threatened turtle species gets entered illegally in a race (DNR unpubl. data).

Concerns also remain regarding the persecution and killing of species people perceive to be harmful to themselves, their livestock, or their pets. The timber rattlesnake is considered especially vulnerable. Because of the species' low reproductive rate, irregular reproduction, and high juvenile mortality (DNR, 2009), the loss of even a single adult can significantly impact a local population (Brown, 1993). Another concern is when citizens are fearful of a snake, remove it from their home site and release it elsewhere, it may decrease its long-term survival.



Timber Harvest

The wood turtle is a forest-dependent SGCN reptile that is affected by a variety of factors related to forest management including timber harvest, road development, forest type conversion, and herbicide application which may harm reptiles directly and degrade their habitats (Tingley & Herman, 2008; DNR, 2020).



Fishing

Turtle by-catch and ingestion of fishing tackle can have individual and population level impacts (Borkowski, 1997; Barko et al., 2010; Midwood et al., 2015; Steen & Robinson, 2017). Anglers may unintentionally attract and concentrate predators like raccoons by leaving discarded fish cleaning remains and live bait, which can pose a significant threat to SGCN turtles, particularly hatchlings (see Problematic Native Species; Recreation).



Recreation

Recreation can have numerous direct and indirect impacts on rare, sensitive reptiles and their habitats ([Northeast Blanding's Turtle Working Group, 2022](#)). Snakes, for example, can be physically harmed by people when they are encountered during outdoor activities, due to widespread fear of snakes. Rare turtle species, like the wood turtle and Blanding' turtle, are vulnerable to a range of recreational activities, including ATV trails and canoe routes located near nesting locations. OHV parks established on sandy areas pose risks to snakes and turtles that depend on these areas. Bonfires, party sites, camping or fishing spots, and picnic areas located on or near sandy shorelines where turtles nest can create a range of disturbances. Garbage left behind at these locations usually attracts mesopredators to nearby nest sites while motorized vehicles along trails can cause direct mortality to the slow-moving turtles and snakes and destroy nests. An increase in human traffic on land also increases the chances turtles are picked up and removed from the local population. A 20-year field study in Connecticut documented a decline in two wilderness wood turtle populations after recreational use of the two sites was allowed (Garber & Burger, 1995).

Larger loose rock is an important habitat component for some SGCN reptiles, like five-lined skinks (Lang, 1982; Quirt et al., 2006). Rock collecting and recreational cairn building increases disturbance and impacts reptile habitat (Schlesinger & Shine, 1994; Hecnar & M'Closkey, 1998; Goldingay & Newell, 2006).

In the water, motorized watercraft can impact individual turtles and turtle communities, causing either direct injury, mortality, disturbance and habitat degradation (Lester et al., 2013, Selman et al., 2013; Smith et al., 2018).

Summer turtle races are local community events that result in the permanent removal of turtles from their populations, as required by state law. Despite providing recreational opportunities, turtles collected for the races are likely stressed, placed on hot pavement, and prone to disease and injury (Heeb et al., 2024).



Fire Management

Fire suppression increases the growth and spread of woody vegetation in open upland habitats that are critical for many reptiles. Encroachment of trees and shrubs impacts reptile species adapted behaviorally and physiologically to open habitats such as prairies and grasslands, including prairie rock outcrops. For example, woody species reduce suitability for Blanding's turtle nesting and egg development, primarily due to shading (MWPARC, 2010). It may also reduce the ability of hatchling Blanding's turtles to orient towards water after they emerge from the nest (Pappas et al., 2009). Common five-lined skink populations of the Upper Minnesota River Valley have been impacted by woody encroachment on prairie outcrops, a required habitat (Gelvin-Innvaer et al., 2025). Trees like red cedar can shade basking areas for timber rattlesnakes, reducing reproductive success by limiting the number of days with high ground temperatures that females need to incubate young (DNR, 2009). (See also Chapters: Prairies and Other Grasslands; Cliff, Talus, Rock Outcrops, and Dunes).

Prescribed fire is a valuable tool to maintain open upland habitats for nesting and basking to benefit SGCN such as five-lined skink, Blanding's turtle, and many SGCN snakes. Prescribed fire also may help create quality foraging habitat (i.e., flush of herbaceous vegetation and berries post-fire). However, many prairies and grasslands now exist as small and isolated patches which make managing both for vegetation and sensitive wildlife a challenge. Fire can cause injury or mortality when it is not applied with the life histories and risk factors of rare, sensitive reptiles in mind. For example, adult female Blanding's turtles have been injured and killed by prescribed burning conducted during their nesting periods in May and June when they were trapped away from aquatic refuge. Spring burning in native prairies and wet meadows have resulted in injury and mortality of Blanding's turtles (Lang, 2004). Fire can also cause direct mortality of prairie snakes and lizards (Setser & Cavitt, 2003, Schultz & Caven, 2021). During early spring emergence, snakes typically remain near their hibernation dens, and when temperatures are cool, they are less able to escape quickly if threatened by fire.



Dams and Water Management

Turtles can be impacted by a wide variety of water management when that occurs without considering turtle requirements and vulnerabilities. Examples include dams, water drainage and drawdowns, stream channelization, excavation/dredging culvert installation and even construction activities associated with stream restoration. The creation of physical barriers and changes to water flow that result in loss of feeding and nesting habitat, and changes to water quality are among the documented impacts (Barcenas-Garcia et al., 2022). Water appropriations and drawdowns that reduce water levels during critical low water periods pose another threat especially when there are no accessible alternative habitats or when turtles are inactive during overwintering (Hall & Cuthbert, 2000; MWPARC, 2010; DNR 2020; NatureServe, 2025).

Turtles face direct risk of injury and mortality from heavy equipment operation in waterbodies. Currently, there is no corroborated evidence that proposed actions to mitigate the impacts of dredging and other excavation to turtles effectively results in avoidance of turtle injury or mortality (Barcenas-Garcia et al., 2022). Dredging during the turtles' active season and posting spotters during dredging are not proven to be reliably effective measures due to the turtles' cryptic coloring, insufficient water clarity that obscures vegetation, and the tendency of many freshwater turtles to "hide" in place within their shells rather than to evade direct threats.



Invasive Species (Problematic Non-native Species)

Invasive, non-native species can alter open habitats that are critical for reptiles, including rare species (Gibbons et al., 2000). Prairies and grasslands dominated by dense stands and thick thatch layers of smooth brome (*Bromus inermis*) are unsuitable for nesting and successful hatching by Blanding's turtles (L. Gelvin-Innvaer, pers. comm.). Bluff prairies overgrown with buckthorn (*Rhamnus cathartica* and *Frangula alnus*) negatively impact timber rattlesnakes in southeastern Minnesota (DNR, 2009) and five-lined skinks on rock outcrops along the Upper Minnesota River Valley (Gelvin-Innvaer et al., 2025). (See chapter on Shore, Cliff, Talus, Rock Outcrops and Dunes). Dense stands of reed canary grass (*Phalaris arundinacea*) in floodplains reduce the availability of sandy nesting areas used by wood turtles (DNR, 2020). Free-roaming cats are non-native predators on snakes and lizards including the common five-lined skink (Gelvin-Innvaer et al., 2025). That impact has been often overlooked and warrants further attention as an additive stressor (Lloyd, 2012; Hall, 2019).



Problematic Native Species

Reptiles are impacted by nest predation from mesopredators like raccoons, badgers, foxes, and skunks. Mesopredators also injure and kill juveniles and adults. Although an important component of native ecosystems, the populations of these native predators can be artificially elevated and concentrated through factors such as habitat fragmentation, constructed trails, and attractants such as garbage and discarded bait. Human-subsidized nest predators have been identified as major factor in Blanding's and wood turtle declines (DNR, 2020).



Diseases and Pathogens

North American reptiles face a variety of risks from diseases, pathogens, and parasites (Gibbons et al., 2000). For example, snake fungal disease has been found in several species that are known to occur in Minnesota, including milk snakes, black rat snakes, and timber rattlesnakes. Caused by a fungus (*Ophidiomyces ophiodiicola*) it can spread internally, causing eye infections and pneumonia, and externally, causing lumps or ulcerations. Mortality can occur (Cornell College of Veterinary Medicine, 2021). Turtles are also carriers of various pathogens and can be impacted by a variety of diseases, including metabolic bone syndrome. Emerging diseases such as *Emydomyces testavorans* and *Emydoidea herpesvirus 2* are of particular concern for turtles, including Blanding's turtles (Woodburn et al., 2019, Andersson et al., 2021). Turtles also are well-known carriers of *Salmonella* bacteria which is easily transmitted to people who handle them. Overall, there is a lack of focus on and awareness of releasing and transporting captively held species without rigorous precautions and criteria.



Water-borne Pollution

Freshwater turtles are bioindicators of environmental pollution due to their longevity, ecological diversity, and trophic positioning (higher in food web). Mercury contamination has been studied the most extensively (Dias et al., 2025) although other environmental pollutants also can have direct and indirect effects on reptiles (Gibbons et al., 2000). In general, insufficient research on the direct, indirect, and synergistic effects of pesticides and herbicides on reptiles hampers informed proposal review and technical guidance.



Air-borne Pollution

Airborne mercury enters aquatic ecosystems primarily through atmospheric deposition, where it settles onto land and water bodies via rain, snow, or dry particles, and then can be washed into water bodies. Otherwise, chemical application on uplands often is via aerial spraying. As noted above, several life traits of freshwater turtles make them suitable bioindicators of environmental pollution (Thompson et al., 2018; Dias et al., 2025) and effects on reptiles are vastly under researched.



Light and Noise Pollution

With the exception of sea turtles, where light pollution has shown to negatively impact hatchling movements, reptiles have rarely been the subject of field research addressing the impacts of light or noise pollution. However, some studies suggest that freshwater turtles like the Blanding's turtle orient toward dark horizons upon hatching to help them disperse from nests into wetlands (Pappas et al., 2009).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (MPCA & DOC, 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

More specifically, reptiles depend on external conditions to regulate their body temperatures, which influences their activity and metabolism. Substantially warmer winters have implications for Minnesota's reptiles which have evolved long dormancy periods for winter survival. Although reptiles must have overwintering sites below the frostline to avoid freezing, those sites must also be sufficiently cool for them to

maintain a reduced metabolic state during dormancy. Later freezes and earlier winter thaws affect reptile activity periods and have implications for technical guidance to help ensure their conservation. Warmer temperatures during their nesting season may cause skewed sex ratios for Blanding's turtles and Southern map turtles due to temperature dependent sex determination (Gutzke & Packard, 1987; Vogt, 2018) which will impact population structure and reproductive output.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. Flooding can be detrimental to species like turtles, which may be killed, washed downstream, buried under sand, or experience nest failure ([Climate trends affecting lakes and rivers | Minnesota DNR](#)). Flooding also can cause erosion and incising of stream banks that degrade a stream's natural connections to floodplains and cause the loss of meanders and other natural features favored by Blanding's and wood turtles. Steep, high stream banks also hamper turtle access to terrestrial habitats. Increased flooding often results in more proposals for riprap, other hardened infrastructure, and drainage systems which can pose risks to turtles and otherwise alter their native habitats.

The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). Droughts may especially impact reptile species associated with water, like Blanding's (Anthonysamy et al., 2013) and wood turtles. Longer term, drought can also alter terrestrial habitats and food chains for all reptiles. For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the SGCN Goal of this Plan, to conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions, three strategies were identified:

- 

Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.
- 

Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnerships with agencies, Tribes, non-governmental organizations, and private landowners.
- 

Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management practices.

Examples of conservation actions are grouped below under these three strategies and tagged with icons for the stressor(s) that they address. Some actions combine multiple strategies, in which case we present it under the one it fits best. Some actions such as those focused on monitoring might not always be tied to a specific stressor; these are marked with ‘n/a’ in the stressor column.

Potential Conservation Actions for Reptiles



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.

Stressor	Action
	Research responses of reptiles to solar facilities and measures to make solar facilities more reptile-friendly (Hernandez et al., 2014).
	To address road mortality, develop or collaborate with existing citizen science programs to identify stretches of road that are mortality hotspots (e.g. the Minnesota Turtle Crossing Tally and Count .) Once identified, use these data to work with MnDOT to develop and apply effective mitigation strategies such as safe under-road passages to help reduce collisions, which benefits wildlife and motorists.
	Conduct rigorous effectiveness monitoring to assess responses of SGCN to specific management activities such as prescribed fire, conservation grazing, forestry practices, water management, etc. in order to develop improved evidence-based management recommendations and avoidance measures for SGCN (Binley et al., 2025) as part of adaptive management. Graduate research projects would help address needs for intensive research.
	Research and apply reptile-friendly solutions for controlling invasive vegetation like smooth brome, reed canary grass, and buckthorn.
	Prioritize research and monitoring on snake fungal disease as well as diseases putting Blanding’s turtles and other turtle species at risk such as adenoviruses and <i>Emydomyces testavorans</i> (Fredrickson et al., 2024).

n/a	Resurvey/monitor reptile SGCN populations whose last observations in the NHIS database are over 10 years old to confirm continued presence of the species, assess population status, and identify specific conservation needs.
n/a	To help address data gaps which hinder conservation, support continued research on efficacy of tools like environmental DNA (eDNA) and camera traps to augment (not replace) standard survey and monitoring techniques for rare SGCN reptiles in various environments, such as Blanding's and wood turtles (Kranz, 2023; Davis et al., 2025).
n/a	Support more in-depth research of Blanding's turtle populations to help answer key conservation questions such as: • Habitat use and movements of hatchlings • Response to nesting habitat creation and management • Stream restoration (including construction phase) • Different grassland and forest management regimes



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.

Stressor	Action
	To address habitat fragmentation from development, protect key areas through land acquisition, conservation easements, and other protected designations. Conserve, reconstruct and manage suitable nesting habitats for SGCN reptiles.
	Reducing landscape fragmentation by increasing buffering, connectivity, and conditions similar to native remnant habitats can help reduce impacts of native and non-native predators on rare sensitive prey species (Schneider, 2001).
	To help avoid entanglement by reptiles and other wildlife and impacts to their habitats, consider these practices (DNR, 2004): A) Use biodegradable netting, preferably natural materials with short degradation periods. Erosion control blankets should be limited to bio-netting or natural netting types due to the risk of entanglement and death of small animals. Avoid photodegradable materials, as they do not degrade properly when covered/shaded, and products containing plastic mesh netting or other plastic components. B) Avoid mulch products that contain synthetic (plastic) fiber additives near waterbodies. Identify acceptable materials in Category 3N or 4N mulches. See Standard Specifications for Construction - MnDOT (state.mn.us) .
	When possible, restore agricultural lands to high quality reconstructed prairie.
	When haying and mowing, promote and apply measures such as found in the Habitat Management Guidelines for Amphibians and Reptiles of the Midwestern United States (Kingsbury & Gibson, 2012) and Mowing Guidelines in Rare Turtle Habitat (MANHESP, 2009) which can be adapted to accommodate larger turtles like the Blanding's turtle. These include mowing when reptiles are less active (November to February) and mowing at high blade settings.
	Apply current guidelines for solar facility fencing to allow access for SGCN turtles (DNR, 2023).

	Work with wind energy companies and conservation partners to reduce restrictions of wind easements on land acquisition and conservation of adjacent lands that reptiles and other wildlife depend upon.
	Using the best available information, establish safe under-road passages and surmountable curbs to aid safe wildlife crossings for the benefit of wildlife and motorists (Markle & Stapleton, 2022, Markle et al., 2025). Avoid creating new roads in important areas for reptiles such as adjacent to water bodies as well as identified crossing hotspots for turtles (Congdon & Keinath, 2006) and snakes.
	Increase representation of all stakeholder groups in forest management decisions. Incorporate ecological values into sustainable timber harvest model and evaluate proposed harvest rates for impacts on wildlife habitat.
	Temporarily close off key nesting areas for reptile SGCN during critical periods (e.g. nesting, emergence). Consult the Northeast Blanding's Turtle Working Group Guidelines for Recreational Area Development and Maintenance (2022).
	When planning prescribed burns, research whether any SGCN reptiles are known or potentially occur on the parcel. Burns should be conducted at times and under conditions when specific rare reptiles are less likely to be on land and access to water refuges remain available (Congdon & Keinath, 2006). Consult guidelines regarding prescribed fire and reptiles (MWPARC, 2008b) with appropriate modifications for target species of concern. Also consider how annual weather variation and climate change affect reptile activity, phenology, and access to safe refuge. Develop more comprehensive prescribed burn guidance based on scientific literature and expertise that considers factors such as fire frequency, intensity, timing, acreage, configuration of treated area, buffers around hibernacula, and access to fire refuge and escape routes in a landscape context (Kapfer et al., 2008). Where prescribed fire poses a significant risk, consider other forms of beneficial habitat disturbance such as conservation grazing. Managing for a mosaic of grassland structure and leaving unburned refugia may also benefit SGCN turtles and snakes in those habitats (Setser & Cavitt, 2003).
	Replace dams and restore streams in a way that minimizes negative impacts and increases benefits to turtles during demolition and construction as well as final restoration design. Consider turtle phenology for the given area and life history characteristics. Time water drawdowns and ditch cleaning at a time of year less likely to impact turtles. Employ measures such as temporary barriers to keep turtles out or temporary drainage prior to excavation to encourage turtles to move to safety or to be more visible for active rescue. Avoid dredging where Endangered or Threatened turtles occur unless or until rigorous empirical research clearly identifies measures that will result in effective avoidance of injury or mortality.
	Manage water levels and conduct riverine management and restorations with SGCN turtles to address their requirements and include measures to avoid take (Congdon & Keinath, 2006).
	Protect nests of rare reptile species from native and non-native predators. Manage human-subsidized predators, like raccoons, especially near nesting areas (MWPARC, 2008a; Golba, 2024).
	Disinfect shoes and gear before traveling to a new outdoor site, whether you are a field biologist, land manager, or outdoor enthusiast. Use scrub brushes and immerse items in disinfectant such as 1% solution of Virkon Aquatic, 3% solution of bleach, or 70% solution of ethanol (Olson et al., 2021; Rzakowska et al., 2016; Bletz et al., 2023.)

	Create buffer zones around bodies of water to reduce water-borne pollutants. Increase native plantings and add filtration ponds.
	Create buffer zones along rivers and streams to reduce the impact of flooding on shoreline habitat. Increase the availability of flood-safe nesting habitat.
n/a	Headstarting of Blanding's turtles and wood turtles in suitable habitats with compatible management of habitat and mesopredators to boost recruitment in populations with age ratios heavily skewed to older individuals (Golba, 2024).



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Stressor	Action
	Increase public education and outreach about how releases of captive reptiles (including wild-caught individuals) can spread disease and introduce non-native species.
n/a	Public outreach about turtle by-catch and what to do if you accidentally hook a turtle .
n/a	Increase enforcement of nongame illegal take issues; educate conservation officers on the issue and monitor key areas for poaching. Develop effective solutions for masking rare reptile locations on websites like iNaturalist. Collaborate with initiatives like Collaborative to Combat the Illegal Trade in Turtles (CCITT) for effective approaches (Christman et al., 2024).
n/a	Increase public awareness of state rules and regulations around take, possession, purchasing, and selling of native reptile species and the reasons for these laws. Promote alternative ways to appreciate Minnesota's reptiles in their native habitats.
n/a	Outreach/education to reduce persecution of venomous and non-venomous snakes via interpretive programs, handouts, social media posts, and other communications. Increase public understanding and appreciation of snakes and how to co-exist with them.
n/a	Educate land managers, including private landowners, about habitat needs of rare forest species.
n/a	Develop and implement a state conservation plan for Blanding's turtles.
n/a	Revise and develop comprehensive best management practices (BMPs), including but not limited to required avoidance measures, for listed reptiles. Prioritize Endangered and Threatened species that are often flagged in internal or external project reviews. These BMPs should be based on the best available updated scientific/technical information, vetted by species experts; be date versioned and in digital format; address the range of practices/actions that affect the species and that commonly arise in reviews; address relevant variation in habitat use and phenology across that species' Minnesota range as it relates to technical guidance for the species (e.g. critical periods, avoidance periods, activity); and be linked from the species' Rare Species Guide account on the DNR website

Case Study: Listed Turtle Environmental DNA Project

Rare reptiles can be particularly difficult to detect, which can hinder conservation and environmental review. Although conventional survey methods like visual encounter surveys or trapping are the “gold standard”, they can be costly and time-consuming. Therefore, scientists from the DNR Nongame Wildlife Program and MnDOT Office of Environmental Stewardship identified a need for additional tools to augment (not replace) standard monitoring methods. Environmental DNA (eDNA) is an emerging technology that holds promise as a rapid, coarse survey tool to help detect rare wildlife. Water samples can be analyzed for DNA shed by animals into their environment to detect their presence, without needing to directly observe or capture them. These scientists championed a proposal to assess the efficacy of eDNA for detecting Blanding’s turtles and wood turtles across their ranges in Minnesota. MnDOT accepted and funded the proposal which led to contracting an expert research team from the University of Illinois. A Minnesota Technical Advisory Panel comprised of representatives from several DNR agencies, MnDOT, the Mille Lacs Band of Ojibwe, and the Minnesota Zoo provided additional guidance and input. This project, which is wrapping up in 2025, has demonstrated that environmental DNA is a rapid, sensitive, and cost-effective tool for monitoring these species under certain conditions. Their final report will provide guidance on methodology and recommendations for next steps to further refine its use (Kranz 2023; Davis et al. 2025).

References for Reptiles

- Andersson, K. E., Adamovicz, L., Mumm, L. E., Winter, J. M., Glowacki, G., et al. (2021). Detection of a novel herpesvirus associated with squamous cell carcinoma in a free-ranging Blanding’s turtle. *Journal of Veterinary Diagnostic Investigation*, 33(2), 348–351. <https://doi.org/10.1177/1040638721989302>
- Andrews, K.M., Gibbons, J.W., & Jochimsen, D.M. (2006). *Literature Synthesis of the Effects of Roads and Vehicles on Amphibians and Reptiles*. Federal Highway Administration Publication. FHWA-HEP-08-005 https://www.researchgate.net/publication/228731156_Literature_Synthesis_of_the_Effects_of_Roads_and_Vehicles_on_Amphibians_and_Reptiles
- Anthonyamy, W. J. B., Dreslik, M. J., & Phillips, C. A. (2013). Disruptive influences of drought on the activity of a freshwater turtle. *The American Midland Naturalist*, 169(2), 322–335. <https://doi.org/10.1674/0003-0031-169.2.322>
- Bárcenas-García, A., Michalski, F., Morgan, W. H., Smith, R. K., Sutherland, W. J., Gibbs, J. P., & Norris, D. (2022). Impacts of dams on freshwater turtles: A global review to identify conservation solutions. *Tropical Conservation Science*, 15. <https://doi.org/10.1177/19400829221103709>
- Barko, V. A., Briggler, J. T., & Ostendorf, D. E. (2004). Passive fishing techniques: a cause of turtle mortality in the Mississippi River. *Journal of Wildlife Management*, 68(4), 1145–1150. [https://doi.org/10.2193/0022-541x\(2004\)068](https://doi.org/10.2193/0022-541x(2004)068)
- Barton, C., & Kinkad, K. (2005). Do erosion control and snakes mesh? *Journal of Soil and Water Conservation*, 60(2). <https://doi.org/10.1080/00224561.2005.12435774>

- Binley, A. D., Haddaway, L., Buxton, R., Lalla, K. M., Lesbarreres, D., Smith, P. A., & Bennett, J. R. (2025b). Endangered species lack research on the outcomes of conservation action. *Conservation Science and Practice*. <https://doi.org/10.1111/csp2.13304>
- Bletz, M., et al. (2023). Disinfection Protocols for Herpetofaunal Pathogens. *Herpetological Review*, 54, 200-203.
- Borkowski, R. (1997). Lead Poisoning and Intestinal Perforations in a Snapping Turtle (*Chelydra serpentina*) Due to Fishing Gear Ingestion. *Journal of Zoo and Wildlife Medicine* 28, 109-113.
- Breckenridge, W. (1944). *Reptiles and Amphibians of Minnesota*. University of Minnesota Press.
- Brown, D., et al. (2025). Turtles All the Way Down. *The Wildlife Professional* 2025(2), 50-54. https://www.fs.usda.gov/pnw/pubs/journals/pnw_2025_brown001.pdf
- Brown, W. S. (1993). *Biology, Status, and Management of the Timber Rattlesnake (Crotalus Horridus)*. Society for the Study of Amphibians & Reptiles. *Herpetological Circular*, 22.
- Chessman, B. C. (2017). Freshwater turtle hatchlings that stay in the nest: strategists or prisoners? *Australian Journal of Zoology*, 66(1), 34. <https://doi.org/10.1071/zo17054>
- Christman, M., Wixted, K., Buchanan, S. W., Boratto, R., Karraker, N., et al. (2024). The Collaborative to Combat the Illegal Trade in Turtles: Addressing Illegal Wildlife Trade with an Adaptive Socio-Ecological Approach. *Chelonian Conservation and Biology*, 23(2). <https://doi.org/10.2744/ccb-1634>
- Christoffel, R. A., & Lepczyk, C. A. (2012). Representation of herpetofauna in wildlife research journals. *Journal of Wildlife Management*, 76(4), 661–669. <https://doi.org/10.1002/jwmg.321>
- Congdon, J.D., & Keinath, D.A. (2006). Blanding’s Turtle (*Emydoidea blandingii*): A Technical Conservation Assessment. USDA Forest Service, Rocky Mountain Region.
- Congdon, J.D., Graham, T.E., Herman, T.B., Lang, J.W., Pappas, M.J., & Brecke, B.J. (2008). *Emydoidea blandingii* (Holbrook 1838) - Blanding’s turtle. In: Rhodin, A. G. J., Pritchard, P.H.C., van Dijk, P.P., Saumure, R.A., Bulmann, K.A., & Iverson, J.B. (Eds.) *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs* 5, 015.1-12.
- Cornell College of Veterinary Medicine. (2021). *Snake Fungal Disease*. <https://cwhl.vet.cornell.edu/disease/snake-fungal-disease>
- Crane, M., Silva, I., Marshall, B. M., & Strine, C. T. (2021). Lots of movement, little progress: a review of reptile home range literature. *PeerJ*, 9, e11742. <https://doi.org/10.7717/peerj.11742>
- Dalrymple, G. H., & Reichenbach, N. G. (1984). Management of an endangered species of snake in Ohio, USA. *Biological Conservation*, 30(3), 195–200. [https://doi.org/10.1016/0006-3207\(84\)90083-1](https://doi.org/10.1016/0006-3207(84)90083-1)

- Davis, M., Costantini, M., Katz, A., Sperry, J., & Larson, E. (2025). *Assessing a New Tool for Early Detection of Endangered Turtles on Proposed Transportation Projects*. Final Technical Report submitted to MN Department of Transportation.
- Dias, F. V. N., Garcês, L. M., & De Oliveira, A. T. (2025). Contaminant levels and their effects on the American continent chelonian: A systematic review. *Acta Zoologica*. <https://doi.org/10.1111/azo.12550>
- Dowling, Z., Hartwig, T., Kiviat, E., & Keesing, F. (2010). Experimental Management of Nesting Habitat for the Blanding's Turtle (*Emydoidea blandingii*). *Ecological Restoration* 28, 154-159.
- Durbian, F. E. (2006). Effects of Mowing and Summer Burning on the Massasauga (*Sistrurus catenatus*). *The American Midland Naturalist*, 155(2), 329–334. [https://doi.org/10.1674/0003-0031\(2006\)155](https://doi.org/10.1674/0003-0031(2006)155)
- Finch, D. M., et al. (2021). *Effects of Climate Change on Invasive Species*. In Poland, T.M., Patel-Weyand, T., Finch, D.M., Miniati, C.F., Hayes, D.C., & Lopez, V.M. (Eds.). *Invasive Species in Forests and Rangelands of the United States*, 57–83. Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Fredrickson, K., Adamovicz, L., Terio, K., Davidson, A., Ryan, M., et al. (2024). *Emydomyces testavorans* surveillance in multiple free-ranging terrestrial and aquatic chelonian species in Illinois, USA. *Journal of Wildlife Diseases*, 60(4). <https://doi.org/10.7589/jwd-d-23-00164>
- Freedberg, S., Lee, C., & Pappas, M. (2011). Agricultural practices alter sex ratios in a reptile with environmental sex determination. *Biological Conservation*, 144(3), 1159–1166. <https://doi.org/10.1016/j.biocon.2011.01.001>
- Garber, S. D., & Burger, J. (1995). A 20-YR study documenting the relationship between turtle decline and human recreation. *Ecological Applications*, 5(4), 1151–1162. <https://doi.org/10.2307/2269362>
- Gelvin-Innvaer, L. (2018). Reptile Renaissance. *Minnesota Conservation Volunteer*, July-August 2018, 9–19. <https://www.dnr.state.mn.us/mcvmagazine/issues/2018/jul-aug/reptile-science.html>
- Gelvin-Innvaer, L., Leclerc, J. B., Worland, M., & Hall, C. D. (2025). Microhabitat Use by Disjunct Common Five-Lined Skink (*Plestiodon fasciatus*) Populations at Their Northwestern Range Extent. *Herpetologica*, 81(2). <https://doi.org/10.1655/herpetologica-d-24-00016>
- Gibbon, J. W., Scott, D. E., Ryan, T. J., Buhlmann, K. A., Tuberville, T. D., Metts, B. S., Greene, J. L., Mills, T., Leiden, Y., Poppy, S., & Winne, C. T. (2000). The global decline of reptiles, déjà vu amphibians. *BioScience*, 50(8), 653. [https://doi.org/10.1641/0006-3568\(2000\)050](https://doi.org/10.1641/0006-3568(2000)050)
- Gibbons, J.W. (1988). The management of amphibians, reptiles, and small mammals in North America: The need for an environmental attitude. In: Szaro, R.C., Severson, K.E., & Patton, D.R. (Eds.). *Management of Amphibians, Reptiles, and Small Mammals in North America: Proceedings of the Symposium*, Flagstaff, Arizona. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station. General Technical Report RM-166. https://www.fs.usda.gov/rm/pubs_rm/rm_gtr166/rm_gtr166_004_010.pdf.
- Golba, C. B. (2024). *Evaluation of Conservation Strategies for Blanding's turtle (Emydoidea blandingii)* [PhD. Dissertation Northern Illinois University].

- Goldingay, R. L., & Newell, D. A. (2006). A preliminary assessment of disturbance to rock outcrops in Gibraltar Range National Park. *Proceedings of the Linnean Society of New South Wales*, 127(1), 75–81.
http://epubs.scu.edu.au/esm_pubs/1091/
- Gooley, A. C. (2010). *Testing the Behavioral Responses of West Virginia Turtles to Roads and Vehicles* [M.S. Thesis Marshall University].
- Gutzke, W. H. N., & Packard, G. C. (1987). The Influence of Temperature on Eggs and Hatchlings of Blanding's Turtles, *Emydoidea blandingii*. *Journal of Herpetology*, 21(2), 161. <https://doi.org/10.2307/1564476>
- Hall, C.D. & Cuthbert, F.J. (2000). Impact of a controlled wetland drawdown on Blanding's turtles in Minnesota. *Chelonian Conservation and Biology* 3, 643-649.
- Hall, J. (2019). *North Carolina Wildlife Profiles: five-lined skink*. North Carolina Wildlife Resources Commission. <https://www.ncwildlife.gov/five-lined-skink-profile/open>
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges*, 3(9).
<https://doi.org/10.1002/gch2.201800095>
- Hecnar, S., & M'Closkey, R. (1998). Effects of human disturbance on five-lined skink, *Eumeces fasciatus*, abundance and distribution. *Biological Conservation*, 85(3), 213–222. [https://doi.org/10.1016/s0006-3207\(98\)00014-7](https://doi.org/10.1016/s0006-3207(98)00014-7)
- Heeb, A. B. et al. (2024). Turtle Races: A threat to Wild Turtle Populations? *Herpetological Review*. 55(1), 11-18.
- Henning, B.M., & Hinz, L.C. Jr. (2016). Conservation guidance for Blanding's Turtle (*Emydoidea blandingii*). INHS Technical Report 2016 (57) Prepared for Illinois Department of Natural Resources.
- Hernandez, R., Easter, S., Murphy-Mariscal, M., Maestre, F., Tavassoli, M., Allen, E., Barrows, C., Belnap, J., Ochoa-Hueso, R., Ravi, S., & Allen, M. (2014). Environmental impacts of utility-scale solar energy. *Renewable and Sustainable Energy Reviews*, 29, 766–779. <https://doi.org/10.1016/j.rser.2013.08.041>
- Johnson, N. B., Poulin, R. G., & Somers, C. M. (2022). Thermoregulation by Bullsnares (*Pituophis catenifer sayi*): do burrows make life easier on the prairies? *Canadian Journal of Zoology*, 100(5), 303–314.
<https://doi.org/10.1139/cjz-2021-0191>
- Jones, J. D., Urquhart, O., Garrah, E., Eberhardt, E., & Danby, R. K. (2024). Patterns and drivers of amphibian and reptile road mortality vary among species and across scales: Evidence from eastern Ontario, Canada. *Global Ecology and Conservation*, 50, e02855. <https://doi.org/10.1016/j.gecco.2024.e02855>
- Jones, M. T., & Sievert, P. R. (2009). Effects of Stochastic Flood Disturbance on Adult Wood Turtles, *Glyptemys insculpta*, in Massachusetts. *The Canadian Field-Naturalist*, 123(4), 313.
<https://doi.org/10.22621/cfn.v123i4.1000>

Kabay, E. (2013). Timber Rattlesnakes may reduce incidence of Lyme disease in the Northeastern United States. *98th ESA Annual Meeting (August 4 -- 9, 2013)*.

<https://eco.confex.com/eco/2013/webprogram/Paper44305.html>

Kapfer, J. M., Coggins, J. R., & Hay, R. (2008). Spatial Ecology and Habitat Selection of Bullsnares (*Pituophis catenifer sayi*) at the Northern Periphery of Their Geographic Range. *Copeia*, 2008(4), 815–826.

<https://doi.org/10.1643/ce-07-104>

King, R. et al. (2025). Archival Data Reveals Human Impacts on Blanding's Turtle Population Persistence.

<http://dx.doi.org/10.2139/ssrn.5144875>

Kingsbury, B.A. & Gibson, J. (2012). *Habitat Management Guidelines for Amphibians and Reptiles of the Midwestern United States*. Partners in Amphibian and Reptile Conservation Technical Publication HMG-1, 2nd Edition.

Knoet, T.G., & Best, L.B. (2011). A multiscale approach to understanding snake use of conservation buffer strips in an agricultural landscape. *Herpetological Conservation and Biology*, 6(2), 191-201.

Kranz, M. (2023). New Project: Assessing a New Tool for Early Detection of Endangered Turtles on Proposed Transportation Projects. <https://mntransportationresearch.org/2023/07/13/new-project-assessing-a-new-tool-for-early-detection-of-endangered-turtles-on-proposed-transportation-projects/>

Lang, J. W. (2004). *Synopsis of Blanding's Turtle Occurrences at or near Altona WMA, Pipestone-Lincoln Co*, Research on State Threatened Blanding's Turtle (*Emydoidea blandingii*) in SW Minnesota 1987-2004. 23 Aug 2024 interim report to Nongame Wildlife Program, Minnesota Dept. of Natural Resources.

Lang, J.W. (1982). Distribution and abundance of the five-lined skink (*Eumeces fasciatus*) in Minnesota. Final report submitted to Nongame Wildlife Program, Minnesota Department of Natural Resources.

Lester, L. A., Avery, H. W., Harrison, A. S., & Standora, E. A. (2013). Recreational boats and turtles: Behavioral mismatches result in high rates of injury. *PLoS ONE*, 8(12), e82370.

<https://doi.org/10.1371/journal.pone.0082370>

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution climate Projections over Minnesota for the 21st century. *Earth and Space Science*, 9(3).

<https://doi.org/10.1029/2021ea001893>

Lloyd, K.A.T. (2012). *Sociopolitical, ecological, and behavioral aspects of free-roaming cats* [PhD. Dissertation University of Georgia].

Lovich, J. E., Ennen, J. R., Agha, M., & Gibbons, J. W. (2018). Where have all the turtles gone, and why does it matter? *BioScience*, 68(10), 771–781. <https://doi.org/10.1093/biosci/biy095>

Markle, T. & Stapleton, S. (2022). *Reduce vehicle-animal collisions with installation of small animal exclusion fencing* (No. MN 2022-19). Minnesota Department of Transportation, Office of Research & Innovation.

https://rosap.nhtl.bts.gov/view/dot/63370/dot_63370_DS1.pdf

Markle, T., Smith, C.E., & Stapleton, S. (2025). Turtles and Turnarounds: Small Animal Exclusion Fencing Effectively Reduces Turtle Road Mortality. <http://dx.doi.org/10.2139/ssrn.5215062>

Martino, J. A., Poulin, R. G., Parker, D. L., & Somers, C. M. (2011). Habitat selection by grassland snakes at northern range limits: Implications for conservation. *Journal of Wildlife Management*, 76(4), 759–767. <https://doi.org/10.1002/jwmg.313>

Massachusetts Natural Heritage and Endangered Species Program (MANHESP). (2009). *Mowing advisory guidelines in rare turtle habitat: pastures, successional fields, and hayfields*. Massachusetts Division of Fisheries and Wildlife.

Midwest Partners in Amphibian and Reptile Conservation (MWPARC). (2010). Blanding’s turtle (*Emydoidea blandingii*) Conservation Assessment Survey.

Midwest Partners in Amphibian and Reptile Conservation (MWPARC). (2008a). Raccoons and turtle conservation. https://www.mwparc.org/files/ugd/fdd25d_25eaeca060aa462c82488d20902f60ce.pdf

Midwest Partners in Amphibian and Reptile Conservation (MWPARC). (2008b). Prescribed Fire Use and Important Management Considerations for Amphibians and Reptiles within the Midwest. https://www.mwparc.org/files/ugd/fdd25d_8a68f84aa9934d60853cba3c24b8d0dc.pdf

Midwood, J. D., Cairns, N. A., Stoot, L. J., Cooke, S. J., & Blouin-Demers, G. (2015). Bycatch mortality can cause extirpation in four freshwater turtle species. *Aquatic Conservation Marine and Freshwater Ecosystems*, 25(1), 71–80. <https://doi.org/10.1002/aqc.2475>

Minnesota Department of Natural Resources (DNR). (2004). Best Practices for Meeting DNR General Public Waters Work Permit GP 2004-0001. https://www.dnr.state.mn.us/waters/watermgmt_section/pwpermits/gp_2004_0001_manual.html

Minnesota Department of Natural Resources (DNR). (2009). Timber Rattlesnake recovery plan (*Crotalus horridus*). Division of Ecological Resources, Minnesota Department of Natural Resources. https://files.dnr.state.mn.us/eco/nongame/projects/tr_recovery_plan_final2009.pdf

Minnesota Department of Natural Resources (DNR). (2020). Wood Turtle (*Glyptemys insculpta*) Conservation Plan for Minnesota. <https://www.lrl.mn.gov/docs/2024/other/240264.pdf>

Minnesota Department of Natural Resources (DNR). (2023). Commercial Solar Siting Guidance. https://files.dnr.state.mn.us/publications/ewr/commercial_solar_siting_guidance.pdf

Minnesota Pollution Control Agency (MPCA) & Department of Commerce (DOC). (2025). Greenhouse gas emissions in Minnesota 2005-2022. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>

Moriarty, J.J., & Hall, C.D. (2014). *Amphibians and reptiles in Minnesota*. University of Minnesota Press.

Mui, A., Edge, C., Paterson, J., Caverhill, B., Johnson, B., Litzgus, J., & He, Y. (2016). Nesting sites in agricultural landscapes may reduce the reproductive success of populations of Blanding’s Turtles (*Emydoidea blandingii*). *Canadian Journal of Zoology*, 94(1), 61–67. <https://doi.org/10.1139/cjz-2015-0154>

- NatureServe. (2025). NatureServe Network Biodiversity Location Data accessed through NatureServe Explorer [web application]. NatureServe, Arlington, Virginia. Available <https://explorer.natureserve.org/>. (Accessed: May 14, 2025).
- Northeast Blanding's Turtle Working Group (2022). A Guide of the Best Management Practices for the Blanding's Turtle (*Emydoidea blandingii*) in the Northeastern U.S.
- Oldfield, B. & Moriarty, J.J. (1994). *Amphibians and Reptiles Native to Minnesota*. University of Minnesota Press.
- Olson, D. H., & Pilliod, D. S. (2022). Elevating human dimensions of amphibian and reptile conservation, a USA perspective. *Conservation Science and Practice*, 4(6). <https://doi.org/10.1111/csp2.12685>
- Olson, D. H., Haman, K. H., Gray, M. J., Harris, R. N., Thompson, T., Iredale, M., Christman, M., Williams, J. M., Adams, M. J., & Ballard, J. R. (2021). Enhanced between-site biosecurity to minimize herpetofaunal disease-causing pathogen transmission. *Herpetological Review*, 52(1), 29–39.
<https://pubs.er.usgs.gov/publication/70222102>
- Pappas, M. J., Congdon, J. D., Brecke, B. J., & Capps, J. D. (2009). Orientation and dispersal of hatchling Blanding's turtles (*Emydoidea blandingii*) from experimental nests. *Canadian Journal of Zoology*, 87(9), 755–766. <https://doi.org/10.1139/z09-065>
- Quirt, K. C., Blouin-Demers, G., Howes, B. J., & Lougheed, S. C. (2006). Microhabitat selection of Five-Lined skinks in northern peripheral populations. *Journal of Herpetology*, 40(3), 335–342.
[https://doi.org/10.1670/0022-1511\(2006\)40](https://doi.org/10.1670/0022-1511(2006)40)
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center.
<https://mwcasc.umn.edu/node/821>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.
<https://statesummaries.ncics.org/chapter/mn/>
- Rzadkowska, M., Allender, M. C., O'Dell, M., & Maddox, C. (2016). Evaluation of Common Disinfectants Effective against *Ophidiomyces ophiodiicola*, the Causative Agent of Snake Fungal Disease. *Journal of Wildlife Diseases*, 52(3), 759–762. <https://doi.org/10.7589/2016-01-012>
- Saumure, R. A., Herman, T. B., & Titman, R. D. (2007). Effects of haying and agricultural practices on a declining species: The North American wood turtle, *Glyptemys insculpta*. *Biological Conservation*, 135(4), 565–575.
<https://doi.org/10.1016/j.biocon.2006.11.003>
- Saumure, R.A., & Bider, J.R. (1998). Impact of agricultural development on a population of Wood Turtles (*Clemmys insculpta*) in southern Québec, Canada. *Chelonian Conservation and Biology* 3, 37-45.
- Schlesinger, C. A., & Shine, R. (1994). Choosing a rock: Perspectives of a bush-rock collector and a saxicolous lizard. *Biological Conservation*, 67(1), 49–56. [https://doi.org/10.1016/0006-3207\(94\)90008-6](https://doi.org/10.1016/0006-3207(94)90008-6)

- Schneider, M. F. (2001). Habitat loss, fragmentation and predator impact: spatial implications for prey conservation. *Journal of Applied Ecology*, 38(4), 720–735. <https://doi.org/10.1046/j.1365-2664.2001.00642.x>
- Schultz, C., & Caven, A. (2021). Lined Snake (*Tropidoclonion lineatum*) Prescribed Fire Mortality. *Transactions of the Nebraska Academy of Sciences and Affiliated Societies*, 42–45. <https://doi.org/10.32873/unl.dc.tnas.41.3>
- Selman, W., Qualls, C., & Owen, J. C. (2013). Effects of human disturbance on the behavior and physiology of an imperiled freshwater turtle. *Journal of Wildlife Management*, 77(5), 877–885. <https://doi.org/10.1002/jwmg.538>
- Setser, K., & Cavitt, J.F. (2003). Effects of burning on snakes in Kansas, USA, Tallgrass prairie. *Natural Areas Journal* 23, 315-319. http://www.naturalareas.org/docs/v23_4_03_pp315_319.pdf
- Smith, G. R., Iverson J.B., and Rettig, J.E. (2018). Frequency of propeller damage in a turtle community in a northern Indiana, USA, lake: a long-Term study. *Herpetological Conservation and Biology* 13, 691–699.
- Steen, D. A., & Robinson, O. J. (2017). Estimating freshwater turtle mortality rates and population declines following hook ingestion. *Conservation Biology*, 31(6), 1333–1339. <https://doi.org/10.1111/cobi.12926>
- Stuart, J.N., Watson, M.L., Brown, T.L., & Eustice, C. (2001). Plastic netting: an entanglement hazard to snakes and other wildlife. *Herpetological Review* 32, 162-164.
- Thompson, M., Coe, B. H., Andrews, R. M., Cristol, D. A., Crossley, D. A., & Hopkins, W. A. (2018). Agricultural land use creates evolutionary traps for nesting turtles and is exacerbated by mercury pollution. *Journal of Experimental Zoology Part a Ecological and Integrative Physiology*, 329(4–5), 230–243. <https://doi.org/10.1002/jez.2198>
- Tingley, R., & Herman, T.B. (2008). *The effects of agriculture and forestry on the distribution, movements and survival of wood turtles in an intensively managed landscape*. Report prepared for the Nova Scotia Habitat Conservation Fund.
- Tingley, R., Meiri, S., & Chapple, D. G. (2016). Addressing knowledge gaps in reptile conservation. *Biological Conservation*, 204, 1–5. <https://doi.org/10.1016/j.biocon.2016.07.021>
- Todd, B., Willson, J., & Gibbons, J.W. (2010). The global status of reptiles and causes of their decline. In: Sparling, D.W., Linder, G., Bishop, C.A. (Eds.) *Ecotoxicology of Amphibians and Reptiles*, 47–67. CRC Pres, Boca Raton.
- Vogt, R.C. (2018). *Graptemys ouachitensis* Cagle 1953 – Ouachita Map Turtle. In: Rhodin, A.G.J., Iverson, J.B., van Dijk, P.P., Buhlmann, K.A., Pritchard, P.C.H., & Mittermeier, R.A. (Eds.) *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs* 5(11), 103.1–13. doi: 0.3854/crm.5.103.ouachitensis.v1.2018; www.iucn-tftsg.org/cbftt/.
- Walley, H.D., King, R.B., Ray, J.M., and Robinson, J. (2005). What should be done about erosion mesh netting and its destruction of herpetofauna? *Journal of Kansas Herpetology* 16, 26-28.

Wiese, J.D. (2016). Common Gartersnake (*Thamnophis sirtalis*) Mortality due to Haying Practices in South-central Nebraska. *Journal of Kansas Herpetology* 5, 15-16.

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020*. Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin, chapter E. U.S. Geological Survey Scientific Investigations Report 2023–5064.
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Wittenberg, R.D. (2012). Foraging ecology of the timber rattlesnake (*Crotalus horridus*) in a fragmented agricultural landscape. *Herpetological Conservation and Biology*, 7(3), 449-461.

Woodburn, D. B., Miller, A. N., Allender, M. C., Maddox, C. W., & Terio, K. A. (2018). *Emydomyces testavorans*, a New Genus and Species of Onygenalean Fungus Isolated from Shell Lesions of Freshwater Aquatic Turtles. *Journal of Clinical Microbiology*, 57(2). <https://doi.org/10.1128/jcm.00628-18>

Zagorski, G. M., Boreham, D. R., & Litzgus, J. D. (2019). Endangered species protection and evidence-based decision-making: Case study of a quarry proposal in endangered turtle habitat. *Global Ecology and Conservation*, 20, e00751. <https://doi.org/10.1016/j.gecco.2019.e00751>

Aquatic Invertebrates

Overview

This sub-chapter highlights different groups of aquatic invertebrates, including mussels and aquatic insects, a few groups of which have been well-surveyed in Minnesota, but the majority of which have not. Mussels have been well-investigated, primarily by DNR's Center for Aquatic Mollusk Programs (see [Mussels of Minnesota](#)). Some groups of aquatic insects have been surveyed by researchers at several universities, the Minnesota Pollution Control Agency, and the Minnesota Dragonfly Society, including dragonflies (Anisoptera) and damselflies (Zygoptera), caddisflies (Trichoptera; Houghton et al., 2022), mayflies (Ephemeroptera), and stoneflies (Plecoptera).

Aquatic invertebrates play an important role in freshwater ecosystems by acting as intermediaries between primary producers (algae, plants, fungi, bacteria) and secondary consumers (fish). It is estimated that aquatic invertebrates are responsible for processing 20-73% of the leaf litter that enters a stream (Covich et al., 1999). Aquatic macroinvertebrates also increase the amount of bioavailable nutrients dissolved in the water column through their feeding activities, burrowing in sediment, and excretion, which in turn accelerates the growth of benthic microbes, algae, and rooted macrophytes (Covich et al., 1999).

Preferred habitats vary by species. Some inhabit only water bodies with specific conditions (water depth, temperature, trophic state, flow rate, surrounding habitat type), while others inhabit a wider variety of water body types. The vast diversity of aquatic macroinvertebrate species and their different tolerance levels to biotic and abiotic environmental factors can serve as an important indicator of the biological condition of a water body.

Historical data on aquatic invertebrate populations is often absent or lacking in quality, which poses a challenge when trying to determine how aquatic invertebrate communities have shifted over time and what trends could be observed. Additionally, aquatic invertebrate assemblages vary greatly across different aquatic habitat types and seasonal periods, making it difficult to determine overall trends on a large spatial or temporal scale. Some studies for particular taxa that are considered less tolerant of lower water quality, such as caddisflies and mayflies, have shown overall losses in biodiversity in association with anthropogenic disturbances (Houghton & Dewalt, 2023, Stepanian et al., 2020). The paragraphs below provide some insights into the diversity of invertebrates in our state.

Mussels

Mussels are relatively well-studied, and have been the focus of survey, research, and conservation work for many years in Minnesota. Efforts by DNR malacologists to conserve freshwater mussels began with statewide inventory surveys from 1999-2010, completing surveys at more than 2700 sites. During this time, 16 quantitative, long term monitoring stations were established in rivers across the state to detect long-term trends in mussel populations. Those monitoring sites have been sampled at least three times as of this writing. Since 2004, DNR malacologists also studied the early life histories of freshwater mussels, determining host fish requirements, timing of larval brooding, and host-infection strategies for 20 species. This included several species whose hosts were poorly or entirely unknown. For example, DNR malacologists were the first to identify the host of the endangered [spectaclecase \(*Cumberlandia monoonta*\)](#) despite nearly 100 years of study

(see case study). Life history work was expanded to in situ culturing of native species for reintroduction to suitable sites. In situ culture work has evolved, starting with culture cages for [Higgins' Eye \(*Lampsilis higginsii*\)](#) in the Mississippi River to having dedicated staff and full functioning hatchery space devoted to culturing mussels since 2016. Since 2016, DNR has successfully reintroduced over 25,000 cultured juvenile mussels representing six species ([black sandshell \(*Ligumia recta*\)](#), plain pocketbook (*Lampsilis cardium*), [mucket \(*Actinonaias ligamentina*\)](#), [snuffbox \(*Epioblasma triquetra*\)](#), Higgins' Eye, and spectaclecase into the Cannon, Cedar, and Mississippi watersheds. Efforts are currently underway to evaluate survival and growth of the reintroduced mussels and to look for signs of recruitment. Some of the priority mussel conservation goals for the DNR from 2025-2035 include a second round of statewide surveys to begin elucidating trends, examining mussel response to stream restoration efforts, and continuing to address mussel culture needs for state and federally listed mussels. The depth of DNR's mussel conservation efforts would not be possible without the continuous support and collaboration from a wide variety of federal, state, tribal, local and private partners (U.S. Army Corps of Engineers, U.S. Fish and Wildlife Service, National Park Service, Illinois DNR and Natural History Survey, Iowa DNR, Wisconsin DNR, Minnesota Department of Transportation, Minnesota Zoo, Prairie Island Indian Community, Minneapolis Park and Recreation Board, Macalester College, University of Minnesota, Shedd Aquarium, James Ford Bell Museum of Natural History, Friends of the Hormel Nature Center, Mower County Soil and Water Conservation District, Daguna Consulting), and many volunteers.

Collectively, DNR malacologists have dozens of scientific publications and have participated in national-scale research projects examining causes of mussel declines and development of a mussel assemblage health index. DNR malacologists have also developed numerous outreach products including a mussel guidebook (updated version expected summer 2025), a Mussels of Minnesota poster, and a mussel life cycle poster, all of which can be found on the DNR's [Mussels of Minnesota website](#). DNR's Clam Counter Minnesota (Google [link](#); [Apple link](#)) mussel identification app became available in spring 2025 and is modeled after the Clam Counter app originally developed by the Toronto Zoo and Fisheries and Oceans Canada. DNR's partners are also actively contributing to mussel conservation outreach, such as the Minnesota Zoo's "[Show Us Your Mussel's Challenge](#)" for high school students. These products, along with a variety of in-person outreach, [press and news releases](#), [informational videos](#), and social media [efforts](#) around the state, have helped Minnesotans grow to love mussels and appreciate the role they play in healthy aquatic ecosystems.

Native freshwater mussels are a diverse group of animals with over 300 species in the US and 51 species in Minnesota representing families Unionidae and Margartiferidae. Of Minnesota's species, 33 are considered at risk to some degree with 6 extant species needing federal protection as endangered species: spectacle case; snuffbox; Higgins' Eye; [sheepnose \(*Plethobasus cyphus*\)](#); [winged mapleleaf \(*Quadrula fragosa*\)](#); and [salamander mussel \(*Simpsonia ambigua*; currently proposed for federal endangered status\)](#). Often the reasons for imperilment can be tied to damming large rivers for navigation and hydropower that affect all aspects of the mussel life cycle and biology. General habitat and water quality degradation associated with industry and agriculture are linked to mussel declines (Bakshi et al., 2023). Despite intensive study, precise explanations for most mussel declines are unclear and declines are often described as enigmatic (Haag, 2012; 2019). Disease ecology of freshwater mussels remains understudied and effects of certain invasive species like Basket Clams (*Corbicula* spp.) need further evaluation (Haag, 2019; Richard et al., 2022). Fortunately in Minnesota, *Corbicula* species are relatively uncommon when compared to other regions of the U.S. Many of the widespread mussel declines documented in Minnesota during the last 100 years seem to have stabilized in recent years as water quality standards have been implemented and large dams are no longer being built. However, many populations linger as shells of their former selves (Hornbach et al., 2019) while other

populations in select high quality and connected habitats may be rebounding (DNR unpublished data). The last decade has seen some success in DNR hatchery rearing of extirpated species to re-establish populations in restored or recovered stream segments that historically supported those species. In cases where certain mussel species are extirpated upstream of dams, properly executed dam removals have benefitted those species by allowing downstream populations to naturally recolonize habitats upstream of the former dam (DNR unpublished data).

Mussels are effective filter feeders that occupy bottom sediments of lakes and rivers, often occurring in dense and diverse aggregations referred to as mussel beds. Mussel beds can be very productive at filtering nutrients, algae, and certain bacteria from the water. Mussel beds can also stabilize substrates and serve as habitats for other organisms like macroinvertebrates. Ecosystem services provided by freshwater mussels cannot be understated (Vaughn, 2018). Freshwater mussels are mostly sessile and may live in the same patch of sediment for their entire lives. Because of their limited mobility and tendency to live in flowing water, limiting any movement to a downstream direction, mussels rely on one or more host species, usually fish, to disperse their larvae to new areas. Host fish are often attracted by and enticed into eating a prey-like lure made by a female mussel. Upon attempting to eat the lure, fish are instead parasitized by a cloud of mussel larvae that develop and fall off weeks later, wherever the fish happens to end up. Due to this complicated life cycle and being immobile residents of benthic sediments, mussels are particularly sensitive to stressors that reduce substrate stability, highly polluted water, and absence of host species. Taken together these attributes can make mussels indicators of overall stream and lake health in some systems.

Case Study: Spectaclecase, an Example of Full-circle Conservation

Through decades of collaboration with many partners and funding from the State and Tribal Wildlife Grants Program, the spectaclecase is just one example of freshwater mussel conservation coming full circle in Minnesota. The spectaclecase is a long-lived (over 100 years) mussel found in cavities within piles of large boulders and slab rocks in a few locations in the Mississippi and St. Croix rivers in Minnesota. Given this unique and difficult-to-survey habitat, it is a challenging species to study, but survey data collected by the DNR and other agencies suggest that the species has declined. Spectaclecase currently occupies four HUC8 (a hydrologic unit code that typically describes a medium-sized river) basins in Minnesota (extirpated from 1). The Lower St. Croix basin is one of only a few strongholds currently remaining for this species across its range. Spectaclecase received protection under the Endangered Species Act (ESA) in 2012, and since then DNR and its partners have been working to restore the spectaclecase populations in the Upper Mississippi River basin.

There are a number of reasons why mussel species are in decline, but given freshwater mussels' need for host-fish to complete their life cycle, understanding specific host needs for a freshwater mussel species is critical to its conservation. Since the hosts for spectaclecase were unknown at the time of its listing under the ESA, host-fish identification was a high priority conservation action. The DNR and University of Minnesota (UM) spent many years conducting laboratory trials to identify the host for spectaclecase without any success. Then, after realizing that spectaclecase found upstream of the St. Croix Falls Dam were showing no signs of reproduction, the DNR and UM deduced that the absence of host-fish upstream of the dam could be the reason why spectaclecase were unable to reproduce. A search began for fishes that were missing upstream of the dam, and Mooneye, Goldeye, and American Eel were identified. Host-trials for these three species determined that Mooneye and Goldeye are indeed hosts for spectaclecase, and finding glochidia on the gills of Mooneye in the wild corroborated the laboratory results.

Ever since appropriate hosts were identified in 2017, DNR has been culturing spectaclecase, with the plan to reintroduce them into the Mississippi and St. Croix rivers to bolster populations. At the DNR's mussel facility in Lake City, the first few years of spectaclecase culture begins in a laboratory setting. Optimal flow rate, sediment type, sediment size, water temperature, and feeding regimes have been experimentally determined to maximize their survival and growth. The mussels continue growing in storage totes in the river for a few more years until they are big enough to tag and are more resistant to predation. The DNR's first spectaclecase cohort was produced in 2017 (130 individuals) and is planned to be reintroduced in 2025. In the coming years, they will be monitored for signs of growth, survival, and recruitment – which are indicators of a self-sustaining population and the ultimate sign of conservation success.

Next steps for the spectaclecase include continued culturing of additional cohorts to support genetic robustness. Additionally, DNR plans to collaborate with partners to better understand their unique habitat requirements and incorporate creation of additional spectaclecase habitat into future stream restoration projects.

Snails

Aquatic snails have been studied in Minnesota for over 100 years and surveys have identified around 60 species. Most of these surveys occurred well over 20 years ago, and a concerted statewide aquatic snail survey has not been completed in decades. Because of the lack of recent survey effort, the distribution and abundance of most species is not well known, making assessment of conservation statuses and implementation of subsequent conservation actions difficult. This is further complicated by strong roots in morphological species concepts and a high degree of intraspecific shell variation having potentially led to an over description of some genera. Apparent difficulties with mitochondrial genetic testing has complicated modern understanding of taxonomic relationships for many aquatic snails, but genomic sequencing is

promising. To begin addressing the above conservation issues during 2025-2035, the DNR is incorporating snails into statewide mollusk survey efforts, compiling existing snail survey data and specimens, and identifying partners to help address taxonomic uncertainties.

Native snails can range from a few millimeters to around 2.5 cm and occur in almost all aquatic habitats in Minnesota from small springs to large rivers to deep lakes. These snails graze algae, detritus, and other organic matter and are food items for other organisms like fish and some mammals. Historical threats were likely relegated to habitat loss or degradation from draining of wetlands for agriculture and subsequent sedimentation from fields in the prairie regions of Minnesota. Current threats faced by snails include invasive mollusk species, pollutants, and ongoing habitat degradation. Much of the snail toxicity literature often studies widespread and relatively tolerant snail species, although more recent studies have begun focusing on rarer and more sensitive species as culture techniques are developed (Barrick et al., 2024).

Aquatic Insects

The diversity of insects in Minnesota is likely greater than the combined diversity of all other animals, but insects are among the least well understood. The Minnesota Pollution Control Agency, researchers at several universities, and the Minnesota Dragonfly Society have surveyed the state for some aquatic insect groups including dragonflies (Anisoptera) and damselflies (Zygoptera), caddisflies (Trichoptera; Houghton et al., 2022), mayflies (Ephemeroptera), and stoneflies (Plecoptera).

In the conservation status assessment process, we evaluated 826 species of aquatic invertebrates, but the true diversity could be orders of magnitude more (Jessica Petersen, DNR, personal communication). Other orders of insects that were not even evaluated during our conservation status assessment are hyper-diverse and we know essentially nothing about their diversity, including aquatic beetles (Coleoptera) and dobsonflies, alderflies and fishflies (Megaloptera).

Lacking basic biological information about so many species, each with a unique role in sustaining the health of natural communities, compromises the ability to address their conservation needs. There is an immediate need to document the diversity of insects in Minnesota before further losses are sustained and therefore go undocumented (Cardoso et al., 2020).

Dragonflies and Damselflies

Despite being tropical in origin, over the past 300 million years dragonflies and damselflies (Odonata) have spread through our temperate climate all the way to the arctic. Minnesota's location in the center of the continent, our four recognized biomes, and our diversity of aquatic breeding habitats all lead to an interesting and diverse mix of Odonata species in the state. Despite being readily recognizable as dragonflies, even by lay-people, we do not have a deep historical record of Odonata species distribution in Minnesota. Prior to the formation of the Minnesota Odonata Survey Project (Minnesota State Wildlife Program Grant T-36-R-1, 2005-2012) there were counties in Minnesota which had zero records of dragonflies and damselflies, despite widespread aquatic breeding habitat. The MOSP, with the help of many volunteers, added thousands of new county records and nearly two dozen new state records for the Odonata in Minnesota. Not only were surveys of adult Odonata conducted in the field, every known museum collection in Minnesota was also investigated and catalogued. [The Minnesota Dragonfly Society](#) (MDS), a 501c3 tax exempt organization, was formed in 2013

as a volunteer organization to ensure the conservation of Minnesota's dragonflies and damselflies through education and research.

Very little formal work on the Odonata distribution in Minnesota has been done since the ending of the MOSP. Despite that, volunteers and enthusiasts continue to contribute to our Odonata knowledge in Minnesota, with several state records and many new county records added in the past years. Given Minnesota's geographical location and our abundance and diversity of habitats, it is highly probable that there are small populations of other Odonata species yet to be found in Minnesota. [Odonata Central](#) and [iNaturalist](#) are tools readily used by citizen scientists to report new findings.

Minnesota has 154 Odonata species on its list, with 112 species of dragonflies (suborder Epiprocta) and 42 species of damselflies (suborder Zygoptera). Several of these records are represented by only one individual which could mean that those species are strays or that more survey work is needed. With a warming and changing climate, there is anecdotal evidence that we are seeing the expansion of more-southerly species into southern Minnesota.

The two general habitat types that hold the most Odonata SGCN species are boggy, northern waters and larger, clear rivers. The biggest threats, statewide, to all Odonata are human disturbances within sensitive watersheds or to the habitat, itself.

Mayflies

Mayflies (Ephemeroptera) are an order of hemimetabolous insects (i.e. they have no pupal stage in the transition from larva to adults) which spend most of their lives in the water before emerging briefly as subimago (dun) or winged, sexually immature adults. Within hours of emerging from the water, the subimago molts to reach full reproductive maturity. In total, this winged adult phase of their lives can last anywhere from a couple of hours to a few days depending on the species and the sex of the individual. After mating, female mayflies disperse their eggs on the surface of the water or, in some cases, on substrates just beneath the surface of the water.

As aquatic nymphs, mayfly larvae can inhabit a wide variety of habitats, from fast flowing rocky-bottom streams to lakes and wetlands. While most mayfly species are collectors and/or scrapers of benthic algae and detritus for the duration of their development, mayflies have evolved a wide variety of feeding strategies and habitat preferences. Numerous mayfly genera burrow in fine sediments, while others are predators, shredders of organic material, or adept swimmers, able to respond quickly to changing stream conditions.

Mayflies play a vital role in maintaining healthy aquatic ecosystems. As aquatic nymphs, they help break down aquatic vegetation and detritus, contributing significantly to the recycling of nutrients. Mayfly nymphs are also a key food source for fish, other macroinvertebrates, and amphibians. As adults, mayflies provide nourishment for fish, birds, and many terrestrial insects, serving as an essential link in the transfer of nutrients between aquatic and terrestrial ecosystems.

Mayflies are familiar to many fishermen who target their emergence events, utilizing artificial flies to "match the hatch". More broadly, mayflies play a critical role in sustaining productive fisheries and healthy ecosystems, which are essential not only for biodiversity, but also for Minnesota's outdoor recreation economy. Recreational fishing and other aquatic activities supported by these ecosystems contribute

significantly to the state's economy through tourism, equipment sales, licenses, and local businesses. Protecting mayfly populations helps ensure the long-term viability of these economic and recreational opportunities for communities across Minnesota.

Mayflies in Minnesota face a wide range of threats. Human development has altered much of the state's natural landscape, disrupting the specific habitat and chemical conditions which many mayfly species require to survive. Major threats include agriculture, urban development, mining, invasive species, and climate change. Of the 658 currently known and taxonomically valid mayfly species in North America, 159 have been documented in Minnesota. Among these, 14 species are considered vulnerable (S3), three are imperiled (S2), and two are critically imperiled (S1), indicating conservation concern for these populations. Eight species are designated SGCN and 83 as SNI. Further work is needed to better understand the health of many of Minnesota's mayfly species.

Stoneflies

Stoneflies (Plecoptera) are an order of aquatic insects strongly associated with running, well-oxygenated freshwater during most of their life cycle. Their presence is often an indicator of good water quality, as many species have specific habitat requirements such as cold temperatures, stable stream substrates, and low pollution levels.

Stoneflies begin their lives as eggs which develop into aquatic nymphs, often after an extended period of limited development, or diapause. As nymphs, stoneflies adopt a wide variety of feeding strategies depending on the species, and stage of development. For example, stoneflies in the family Pteronarcyidae, shred and eat detritus and aquatic vegetation throughout their development, while many stoneflies in the family Perlidae transition from a diet of algae/detritus early in their development, to a diet of aquatic macroinvertebrates as larger and more mature instars. The winged adult phase of the stonefly life cycle is relatively brief, lasting only a few days in most cases and up to a few weeks in a few species.

Stoneflies play a vital role in maintaining healthy aquatic ecosystems. As aquatic nymphs, stoneflies help break down aquatic vegetation and detritus, while contributing significantly to the recycling of nutrients. Predatory nymphs help to regulate the productivity of some other aquatic macroinvertebrates and help to maintain the delicate balance of these aquatic ecosystems. Stoneflies, both as adults and nymphs, are a key food source for fish, other macroinvertebrates, amphibians and birds. Like other aquatic insects they often serve as an essential link in the transfer of nutrients between aquatic and terrestrial ecosystems.

Due to their stringent habitat requirements, stoneflies are often some of the most pollution sensitive organisms found in each stream or river. Across Minnesota, stoneflies face a wide range of threats. Human development has altered much of the state's natural landscape, disrupting the specific habitat and chemical conditions which many mayfly species require to survive. Major threats include agriculture, urban development, mining, invasive species, and climate change. Of the 70 stonefly species documented in Minnesota, 12 are considered vulnerable (S3), four are imperiled (S2), and one is considered critically imperiled (S1), indicating conservation concern for these populations. Further work is needed to better understand the health of many of Minnesota's mayfly species. Five are designated SGCN and 34 as SNI.

Caddisflies

Caddisflies (Trichoptera) are a diverse order of insects that utilize aquatic habitats for much of their life cycle. The name trichoptera is derived from the Greek *trichos* (hair) and *pteron* (wing) which is in reference to the hair that covers the wings on adult caddisflies.

Larval caddisflies have a major role in aquatic ecosystems including rivers, streams, marshes, and lakes. While some individual species are uniquely adapted for specific habitat types, others are considered generalists and are found across a wide array of aquatic ecosystems. Caddisflies are major components of aquatic ecosystems and contribute towards the transfer of energy and nutrients throughout trophic levels within freshwater systems worldwide. Caddisflies also display a wide array of life history characteristics. Many caddisfly taxa produce shelters or cases to protect and disguise themselves from predators. These cases are constructed using materials from their surroundings that are fastened together by silk, and some species even produce nets that they use to filter flowing water as a passive source of food. While many species use silk to protect themselves or as a food source, there are also 'freeliving' caddisflies, such as the *Rhyacophila* genera which do not build cases, and act as predators towards smaller aquatic fauna. Many taxa have very specific habitat preferences or tolerance/intolerance to specific pollutants and water chemistry parameters making them excellent indicators of watershed disturbance. The diversity of caddisflies as a whole, and their ingenious use of silk to fully optimize their unique life history strategies makes caddisflies an invaluable group for water quality monitoring.

Out of more than 300 caddisfly species that were reviewed in this process, 48 were designated SGCN. Four have been given a rank of S1 (critically imperiled), 20 have been given a rank of S2 (imperiled), and 70 have been given a rank of S3 (vulnerable). These species have been highlighted in these S ranks for a variety of reasons, but most commonly due to rarity, sensitivity, or unique habitat preferences. In addition to the S1-S3 ranks more than 130 caddisfly species were designated an S4 (apparently secure) or S5 (secure) ranking due to generally stable, well-established populations throughout Minnesota. Lastly, ranks of SH (possibly extirpated), SU (unrankable), and SNA (not applicable) were assigned to 23, 56, and 6 species respectively. SH taxa are typically species with exclusively old records, where efforts have been made to search for these species and they have not been found. SU species lack sufficient information to rank (very low numbers of occurrences), while SNA species are reserved for non-native species, vagrants, or species that are unlikely to have ever existed in Minnesota.

There is a high level of variability in information for any given species of caddisfly within Minnesota. For some species, there may be very few occurrences from a localized area, whereas other species may have thousands of records across the state. In general, the larger bodied caddisflies are typically easier to identify, and more correlations can be drawn about their habitat preferences, range, and population stability. Conversely, smaller caddisflies such as the Microcaddisfly family Hydroptilidae have many species, and are difficult, if not impossible to identify depending on their life stages without the aid of genetic analysis. Short of specific studies designed to identify adults or fully mature larva of certain groups such as Hydroptilidae, little is known about these taxa as many of the broad scale sampling efforts that occur in the state do not capture the necessary details to identify some groups down to species.

SGCN and SNI Summary Information

There are 123 SGCN aquatic invertebrates and 186 SNI aquatic invertebrates (see Table 2.8). See Appendix B for full list of species. This includes some state-listed species (24 caddisflies, 8 dragonflies and damselflies, 28 mussels) and federally listed species (5 mussels, with 1 additional proposed for federal listing).

Table 2.8. A summary of Aquatic Invertebrate numbers of Species in Greatest Conservation Need and Species in Need of Information by species group.

Taxa	# SGCN	# SNI	*Total Assessed	# SGCN removed from 2015 List	# SGCN added to 2015 List	# 2015 SGCN demoted to SNI	# SNI added
Beetles (Class Insecta; Order Coleoptera)	1	0	1	0	1	0	0
Caddisflies (Class Insecta; Order Trichoptera)	50	51	314	0	29	3	48
Dragonflies and Damselflies (Class Insecta; Order Odonata)	25	8	145	14	5	6	2
Mayflies (Class Insecta; Order Ephemeroptera)	8	83	163	0	8	0	83
Mussels (Class Bivalvia)	33	1	51	0	3	0	1
Shrimp (Class Branchiopoda)	0	1	9	0	0	0	1
Snails - Aquatic (Class Gastropoda)	1	8	67	0	1	0	8
Stoneflies (Class Insecta; Order Plecoptera)	5	34	76	0	5	0	34
TOTAL	123	186	862	14	52	9	177

*Many of these groups are very under-surveyed.

Habitat Associations

The range of wetlands, rivers, and streams that aquatic invertebrate SGCN inhabit is found in Table 2.9. The totals for each taxa listed below include both primary and secondary habitats for mussels (Bivalvia), caddisflies (Trichoptera), and mayflies (Ephemeroptera). For the other taxa (stoneflies (Plecoptera), aquatic beetles (Coleoptera), aquatic snails (Gastropoda) and dragonflies (Odonata), primary and secondary habitats were not delineated, only the habitats where the species occurred without ranking. Primary habitats are those where the species is found most consistently; the species is most reliant on the habitat and, should the habitat be degraded or lost, it would have the greatest effect on the species. Secondary habitats are those where a species can be found some of the time, but less frequently than a primary habitat. Some species, especially those with more general habitat requirements, may be assigned to multiple primary or secondary habitats. Table 2.9 does not include habitat associations for the one taxa that is only represented by Species in Need of

Information, shrimp (Branchiopoda). Appendix C provides a comprehensive list of the primary and secondary habitats utilized by each SGCN.

Table 2.9 Numbers of species of aquatic invertebrates associated with major aquatic habitats and wet forests.

Habitats	Beetles (aquatic)	Caddisflies	Dragonflies and Damselflies	Mayflies	Mussels	Snails (aquatic)	Stoneflies	Total
Deciduous Wet Forest	0	0	1	0	0	0	0	1
Coniferous Forest Wetland	0	0	1	0	0	0	0	1
Non-forested Wetlands	0	0	7	0	1	0	0	8
Rivers and Streams (Total)	1	43	12	8	32	0	5	101
Streams	0	23	10	7	10	0	5	55
Rivers	0	14	3	4	15	0	4	40
Large Rivers	0	5	1	0	30	0	0	36
Lakes (Total)	2	21	13	2	1	1	0	40
Lakes Not Shallow	0	15	10	0	1	1	0	27
Lakes Shallow	0	5	3	0	1	1	0	10
Vernal Pools	0	0	6	0	0	0	0	6

Rather than broadly characterizing aquatic invertebrate SGCN as simply river species, the 2025-2035 SWAP uses the Strahler Stream Order (SSO) method to classify streams and rivers. Streams without any tributaries are assigned an SSO of 1; then the stream order increases when tributaries combine into larger streams. As illustrated in Table 2.9, the majority of aquatic invertebrate SGCN occur in small streams (SSO 1-4). The mussels are primarily found in riverine waters where their diversity is highest (see the Rivers and Streams Chapter) with 10 species found in smaller streams (SSO 1-4), 15 species in medium rivers (SSO 5-6), and 30 species occurring in large rivers (SSO 7-8) (note: many species occur in more than one riverine habitat) (see Appendix C. SGCN Habitat Associations).

Minnesota is also home to the only known lake endemic mussel species, the lake floater (*Pyganodon lacustris*). It occurs primarily in silt or soft mud dominated substrates of backwater lakes that have little flow and are deep enough not to completely freeze over during the winter months. Some of the mussel SGCN are habitat specialists and examples include spectaclecase and salamander mussels which occupy crevices under boulders and slab rocks.

Primary Stressors for Aquatic Invertebrates

Stressors are factors that pose direct or indirect challenges to vulnerable plant and wildlife species. Habitat loss and degradation are generally considered the primary threat for most SGCN, and some stressors reduce the quality of those habitats. Other stressors may operate more directly on the SGCN, such as disease and pathogens. In some cases, stressors may operate both directly and indirectly. For example, roads, trails, and railroads can directly harm SGCN through vehicular strikes, or operate indirectly through habitat fragmentation and reduced landscape connectivity. The stressor list below is adapted from an internationally recognized threats classification developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For more information, see the “*Stressors*” section of Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts (e.g. catastrophic wildfire following a prolonged period of fire suppression) while serving as valuable conservation tools in others (e.g., appropriately planned and applied prescribed fire).

Information about a subset of primary stressors specifically impacting this species group is included below, followed by a set of conservation actions addressing those stressors. Also please reference the Rivers and Streams sub-chapter for more about stressors and actions in that habitat.



Development

Residential and commercial development can fragment and reduce suitable habitat available for insects and are among the most prevalent threats to aquatic invertebrates currently listed under the federal Endangered Species Act (Shirey et al., 2025).



Crop Production

Many activities associated with intensive agriculture pose threats to aquatic macroinvertebrates. Some examples include stream channelization, draining of wetlands, loss of surrounding floodplain and riparian cover, and increased inputs of pesticides, herbicides, sediments, and nutrients (Houghton & Dewalt, 2023). Over application of fertilizer results in increased nutrient runoff into adjacent water bodies. This can lead to algal blooms that deplete oxygen in the water column and negatively impact sensitive aquatic invertebrates that require well oxygenated water to survive.

Flashy hydrology caused by tiled agricultural fields can cause streams to scour which limits freshwater mussel presence to only the most stable stream reaches. This is compounded when stream channels are straightened, facilitating water conveyance “too efficiently” (Blann et al., 2009). Small streams are affected to a greater degree than larger streams by agricultural practices (Wohl, 2017), which in turn affects the mussels in those systems. However, in large rivers wide scale conversion to intensive row crop agriculture has been shown to reduce mussel diversity when compared to natural land uses in Minnesota (Hornbach et al., 2019).



Livestock Management

Direct links between cattle and freshwater mussel impacts are limited but effects of cattle on stream habitat degradation are well documented. Cattle cause stream bank erosion, sedimentation, increased total suspended solids, increased nutrient loading, and increased coliform bacteria (O’Callaghan et al., 2018).



Mining and Quarrying

Release of heavy metals into water bodies due to mining activities results in bioaccumulation of heavy metals throughout the aquatic ecosystem and toxicity to sensitive species of aquatic invertebrates. Indirect effects may also occur through the decline in algal food sources in response to heavy metal exposure (Mebane et al., 2020).

In-river gravel mines have been proposed in the Mississippi River in Pool 2. Dredging materials to be mined can directly remove freshwater mussels.



Roads, Trails, and Railroads

Railroads can present an overlooked threat to aquatic organisms as derailments resulting in release of toxic substances into lakes and rivers can cause mussel kills. Although most derailments happen in shipping yards, a relatively small number of derailments result in spills of contaminants (less than one percent; AAR, 2024) and can cause disproportionate environmental damage. Several mussel species in Minnesota that are only found in a single river could be devastated by a train derailment.

Freshwater mussels are also very sensitive to salt, and heavily salted roadways are considered a threat in some instances (Bakshi et al., 2023).



Shipping Lanes

Routine dredging of river channels to accommodate commercial boat traffic creates disturbances that prevent natural channel formation, movement, and sediment transport. Because freshwater mussels require stable river bottoms to burrow into, dredging can remove them from the sediments, bury them, or destabilize the sediments that remain. Increased water turbidity can also make it more difficult for these filter feeders to eat.

Nutrient cycling is limited in dredged channels because the river has limited access to floodplain habitats that offer an exchange of terrestrial and aquatic nutrients (Schramm et al., 2015). Lock and dam operations block the migration and movement of many fish species, especially those that migrate seasonally (e.g. [skipjack herring, *Alosa chrysochloris*](#)). Restricted fish movement further impacts native mussel species that require those fish species to complete mussel life cycles and this has led to functional extirpation of two mussel

species from Minnesota ([elephant-ear \(*Elliptio crassidens*\)](#) and [ebonyshell \(*Reginaia ebenus*\)](#)). The upstream transport of goods on barges facilitates the movement of undesirable and deleterious species like invasive zebra mussels (*Dreissena polymorpha*) that can attach to barges. Impounded sections of river upstream of locks and dams are often reservoirs for invasive species, allowing them to persist where they would otherwise not occur in naturally flowing river reaches (Johnson et al., 2008). This is especially true for zebra mussels (Stoeckel et al., 2004). Commercial traffic has been shown to increase suspended sediments and turbidity (Smarts et al., 1985; Spear et al., 2024). Elevated turbidity has been shown to negatively impact mussel reproduction (Gascho Landis et al., 2013, Gascho Landis & Stoeckel, 2015).



Many in Minnesota's large and diverse fishing community may know very little about the importance of aquatic invertebrates to the health and sustainability of good water quality and healthy fisheries. An engaged public would be a strong voice for water quality improvements.



Resuspension of sediments caused by recreational motorboat wakes from as few as 11 boats making a single pass per day has been shown to reduce filtration capacity of native freshwater mussels by nearly 7% in a river in Germany (Lorenz et al., 2013). Another study in the Mississippi River found recreational boats to have a measurable albeit small impact to mussel behavior in a common species (Miller et al., 1999). However, extrapolating these results to high traffic lakes and rivers in Minnesota during peak seasons could reduce mussel filtration capacity significantly. These effects were highly depth dependent and boat operation in water greater than 1.8 meters depth had lower effects (Lorenz et al., 2013). Though uncommon in most rivers, it is possible that modern wake boats designed to throw larger wakes could further impact native mussels due to the larger waves and associated increase sediment resuspension (Marr et al., 2022). This might be a significant concern on the Mississippi River where recreational boating with exceptionally large boats and yachts capable of generating large wakes is a common practice. Elevated turbidity has been shown to negatively impact mussel reproduction (Gascho Landis et al., 2013, Gascho Landis & Stoeckel, 2015).



Dams block the migration and movement of many fish species, especially those that migrate seasonally (e.g. skipjack herring). Restricted fish movement further impacts native mussel species that require those fish species to complete mussel life cycles and has caused functional extirpation of two species in Minnesota (Kelner & Sietman, 2000; DNR unpublished data). Since mussels are sessile and live in lake and river sediments, they rely on fish to transport their larvae upstream. Thus, dams severely limit upstream mussel dispersal. The population fragmentation associated with these barriers increases the chances of local mussel extirpations from stochastic events (Haag, 2012). Impounded sections of river upstream of dams are often reservoirs for invasive species, allowing them to persist where they would otherwise not occur in naturally flowing river

reaches (Johnson et al., 2008). Of specific importance to native mussels, impoundments and lakes upstream of flowing reaches often act as a continual source of zebra mussels that colonize downstream reaches (Stoeckel et al., 2004). zebra mussel biofouling effectively cements native mussel shells to close causing suffocation and reduced host infestation success.

Flashy hydrology caused by impervious surfaces or tiled agricultural fields can cause streams to scour which limits freshwater mussel presence to only the most stable stream reaches. This is compounded when stream channels are straightened, facilitating water conveyance “too efficiently” (Blann et al., 2009).



Invasive Species (Problematic Non-native Species)

Problematic non-native snail species such as the Chinese mystery snail (*Cipangopaludina chinensis* and *C. japonica*), banded mystery snail (*Viviparus georgianus*), faucet snail (*Bithynia tentaculata*), and New Zealand mudsnail (*Potamopyrgus antipodarum*) outcompete native macroinvertebrate species for food and habitat.

Zebra mussels endanger native mussel species and riverine infrastructure by densely encrusting to hard surfaces. Native mussel species can become encrusted to the point of suffocation by zebra mussels but also may lose the ability to close which is a primary defense against predators and temporarily unfavorable water quality conditions. Since many mussel species use lures to attract host fishes, complete encrustation by zebra mussels will prevent deployment of those lures. Because both native mussels and zebra mussels are filter feeders, high density zebra mussel populations can outcompete native species for food.



Diseases and Pathogens

While mussel diseases have not been identified specifically in Minnesota, rivers in the eastern US (including in Wisconsin; Knowles et al., 2024) have documented various mussel specific pathogens that can negatively affect mussels in various ways (Richard et al., 2022).



Water-borne Pollution

Some surfactants common in household products, like dish soap, are toxic to many aquatic mollusks (Gibson et al., 2016). Hydrobiid snails may be particularly sensitive to pollutants and at least one study has documented negative effects of copper, ammonia, and pentachlorophenol (Besser et al., 2016).

Ammonia is very toxic to freshwater mussels, especially at the larval and juvenile life stages (Wang et al., 2007). Another study found that mussels are broadly sensitive to total nitrate, total phosphorus, and chloride and that current water quality standards may not sufficiently protect mussels from these pollutants in Minnesota (Bakshi et al., 2023).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). More frequent and prolonged droughts are expected to reduce stream flows and water levels. Some stream reaches may dry up entirely, resulting in the loss of aquatic invertebrates due to desiccation. Warmer summer air temperatures will further raise stream temperatures, especially during low-flow periods. Many mussel species are particularly vulnerable, with thermal limits generally ranging from 25–35°C depending on tolerance, rate of warming, and duration of exposure (Pandolfo et al., 2012). Notably, many Minnesota streams already reach or exceed 30°C, and impounded reaches near dams may experience even greater temperature stress.

Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. Higher precipitation and extreme weather events can also severely impact mussel populations by increasing the sediment load to rivers and streams, suffocating mussels and burying mussel beds. An increase in nutrients and pollutants from surface runoff will decrease water quality and, depending on concentrations, kill or damage mussels. Extreme flooding can also scour the

riverbed displacing mussels and transporting them to less suitable habitat, damage mussel shells by impacts with swiftly moving debris, and cause mortality (Tarter et al., 2022).

The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the SGCN Goal of this Plan (Conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions), three strategies were identified:



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Examples of conservation actions are grouped below under these three strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Some actions that are focused on important population monitoring and public engagement needs might not always be tied to a specific stressor; these are shown as 'n/a' in stressors.

Potential Conservation Actions for Aquatic Invertebrates



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.

Stressor	Action
	Monitor mussel populations including focus on species at risk for disease. Take samples and analyze mussels for pathogens. Document disease and pathogen spread and implement prevention measures where possible.
	Monitor population distribution of sensitive aquatic macroinvertebrate species in impaired water bodies.
	Gather information on aquatic invertebrate assemblages across different seasons in key water bodies to use as comparative benchmarks for future studies quantifying the impact of the changing climate.
n/a	Conduct rigorous effectiveness monitoring to assess responses of SGCN and priority sensitive taxa to specific management activities to develop improved evidence-based

Stressor	Action
	management recommendations and avoidance measures for SGCN (Binley et al., 2025) as part of adaptive management.

 Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.	
Stressor	Action
	Support implementation of actions that improve overall watershed health (see Watershed Health Assessment Framework). Improve water quality through excellent stewardship of the land upslope of streams to reduce sedimentation and point and non-point pollution.
	Work with farmers to implement practical solutions to reduce sedimentation and nutrient loading into streams (riparian buffers, cover crops, conserve wetlands, etc.). Implement regenerative and precision agriculture practices to minimize the use of pesticides, herbicides, and fertilizer, preventing subsequent runoff into adjacent water bodies. Develop best management practices and communicate them to farmers, landowners, and public.
	Help develop incentives and alternative practices to reduce impacts of drain tiles on streams and find practical yet more environmentally sound solutions for water storage and conveyance.
	Working with permitting agencies, ensure that mining operations that impact downstream waters or mining operations that occur within waters properly mitigate for impacts to freshwater mussels (e.g. reducing impacts to established mussel beds). Ensure proper mining waste disposal standards are adhered to minimize risk of releasing heavy metals into adjacent water bodies. Perform proper remediation activities on affected water bodies to restore population of sensitive aquatic macroinvertebrate species.
	Work with the Army Corps of Engineers and others dedicated to problematic non-native species to reduce opportunities for invasive species to enter Minnesota waters and to mitigate the impacts problematic species already present.
	Support projects that focus on stream restoration, dam removal and replacement with rock arch rapids, restoration of natural water connections, the installation of effective fish passages, and mussel species reintroductions when and where appropriate.
	Promote the establishment of riparian buffers, native plantings, filtration ponds, and wetland basins. Insure the proper treatment of wastewater and stormwater. Mitigate ammonia runoff and point sources, especially during spring and summer when juvenile

	mussels are most vulnerable. Investigate surfactant levels in lakes with high residential land use. Bolster water quality standards sufficiently to minimize impacts to native mussels.
	Improve aquatic biodiversity and create a robust native community by stocking mussels when and where appropriate.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Stressor	Action
	Develop best management practices and communicate them to farmers, landowners, and public regarding solutions to reduce sedimentation and nutrient loading into streams (e.g., riparian buffers, cover crops, conserve wetlands, regenerative and precision agriculture practices, reducing impacts of drain tiles) to minimize impacts of runoff into adjacent water bodies.
	Engage with community members who enjoy water bodies recreationally to increase awareness around the importance of healthy aquatic invertebrate communities to support the overall health of the aquatic ecosystem.
	Increase education and enforcement efforts for decontamination of aquatic vessels to prevent transport of invasive macroinvertebrate species to new water bodies.
	Continue to implement proven aquatic invasive species detection and prevention methods (e.g. Clean, Drain, Dry educational initiatives for boaters).

References for Aquatic Invertebrates

Association of American Railroads (AAR). (2024). Freight Rail: Train Derailments Fact Sheet. <https://www.aar.org/issue/derailments/>. Accessed 5 June 2025.

Bakshi, B., Bouchard, R. W., Dietz, R., Hornbach, D., Monson, P., Sietman, B., & Wasley, D. (2023). Freshwater Mussels, Ecosystem Services, and Clean Water Regulation in Minnesota: Formulating an Effective Conservation Strategy. *Water*, 15(14), 2560. <https://doi.org/10.3390/w15142560>

- Barrick, A., Parham, S., Johnson, P., Brewer, S., & Hoang, T. (2024). Sensitivity of Alabama freshwater gastropod species to nickel exposure. *Environmental Toxicology and Chemistry*, 43(12), 2578–2588. <https://doi.org/10.1002/etc.5985>
- Besser, J. M., Dorman, R. A., Hardesty, D. L., & Ingersoll, C. G. (2016). Survival and growth of freshwater pulmonate and nonpulmonate snails in 28-Day exposures to copper, ammonia, and pentachlorophenol. *Archives of Environmental Contamination and Toxicology*, 70(2), 321–331. <https://doi.org/10.1007/s00244-015-0255-3>
- Binley, A. D., Haddaway, L., Buxton, R., Lalla, K. M., Lesbarreres, D., Smith, P. A., & Bennett, J. R. (2025). Endangered species lack research on the outcomes of conservation action. *Conservation Science and Practice*. <https://doi.org/10.1111/csp2.13304>
- Blann, K.L., Anderson, J.L., Sands, G.R., & Vondracek, B. (2009). Effects of agricultural drainage on aquatic ecosystems: a review. *Critical Reviews in Environmental Science and Technology*, 39, 909–1001.
- Cardoso, P., Barton, P. S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., Fukushima, C. S., Gaigher, R., Habel, J. C., Hallmann, C. A., Hill, M. J., Hochkirch, A., Kwak, M. L., Mammola, S., Noriega, J. A., Orfinger, A. B., Pedraza, F., Pryke, J. S., Roque, F. O., . . . Samways, M. J. (2020). Scientists’ warning to humanity on insect extinctions. *Biological Conservation*, 242, 108426. <https://doi.org/10.1016/j.biocon.2020.108426>
- Covich, A. P., Palmer, M. A., & Crowl, T. A. (1999). The role of benthic invertebrate species in freshwater ecosystems. *BioScience*, 49(2), 119. <https://doi.org/10.2307/1313537>
- Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weyand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Gascho Landis, A. M., & Stoeckel, J. A. (2015). Multi-stage disruption of freshwater mussel reproduction by high suspended solids in short- and long-term brooders. *Freshwater Biology*, 61(2), 229–238. <https://doi.org/10.1111/fwb.12696>
- Gascho Landis, A. M., Haag, W. R., & Stoeckel, J. A. (2013). High suspended solids as a factor in reproductive failure of a freshwater mussel. *Freshwater Science*, 32(1), 70–81. <https://doi.org/10.1899/12-093.1>
- Gibson, K. J., Miller, J. M., Johnson, P. D., & Stewart, P. M. (2016). Toxicity of sodium dodecyl sulfate TO federally threatened and petitioned freshwater mollusk species. *Freshwater Mollusk Biology and Conservation*, 19(1), 29. <https://doi.org/10.31931/fmbc.v19i1.2016.29-35>
- Haag, W. R. (2019). Reassessing enigmatic Mussel declines in the United States. *Freshwater Mollusk Biology and Conservation*, 22(2), 43. <https://doi.org/10.31931/fmbc.v22i2.2019.43-60>
- Haag, W.R. (2012). North American freshwater mussels: natural history, ecology, and conservation. Cambridge University Press, New York, NY.

He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019c). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges*, 3(9).

<https://doi.org/10.1002/gch2.201800095>

Hornbach, D. J., Hove, M. C., MacGregor, K. R., Kozarek, J. L., Sietman, B. E., & Davis, M. (2019). A comparison of freshwater mussel assemblages along a land-use gradient in Minnesota. *Aquatic Conservation Marine and Freshwater Ecosystems*, 29(11), 1826–1838. <https://doi.org/10.1002/aqc.3167>

Houghton, D. C., & DeWalt, R. E. (2023). The caddis aren't alright: modeling Trichoptera richness in streams of the northcentral United States reveals substantial species losses. *Frontiers in Ecology and Evolution*, 11.

<https://doi.org/10.3389/fevo.2023.1163922>

Houghton, D. C., DeWalt, R. E., Hubbard, T., Schmude, K. L., Dimick, J. J., Holzenthal, R. W., Blahnik, R. J., & Snitgen, J. L. (2022). Checklist of the caddisflies (Insecta, Trichoptera) of the Upper Midwest region of the United States. *ZooKeys*, 1111, 287–300. <https://doi.org/10.3897/zookeys.1111.72345>

Johnson, P.T., Olden, J.D., & Vander Zanden, M.J. (2008). Dam invaders: impoundments facilitate biological invasions into freshwaters. *Frontiers in Ecology and the Environment* 6(7), 357–363.

Kelner, D. E., & Sietman, B. E. (2000). Relic Populations of the Ebony Shell, *Fusconaia ebena* (Bivalvia: Unionidae), in the Upper Mississippi River Drainage. *Journal of Freshwater Ecology*, 15(3), 371–377.

<https://doi.org/10.1080/02705060.2000.9663755>

Knowles, S., Leis, E. M., Richard, J. C., Standish, I. F., Bojko, J., Weinzing, J., & Waller, D. L. (2024). *Hirsutonosema embarrassi* n. gen. n. sp. (Phylum Microsporidia) in the Ovary of Mucket (*Actinonaias ligamentina*), Plain Pocketbook (*Lampsilis cardium*), and Fatmucket (*Lampsilis siliquoidea*) (Unionidae) from the Embarrass River, Wisconsin, USA. *Parasitologia*, 4(2), 184–198. <https://doi.org/10.3390/parasitologia4020016>

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>

Lorenz, S., Gabel, F., Dobra, N., & Pusch, M. T. (2013). Modelling the effects of recreational boating on self-purification activity provided by bivalve mollusks in a lowland river. *Freshwater Science*, 32(1), 82–93.

<https://doi.org/10.1899/12-054.1>

Marr, J., Riesgraf, A., Herb, W., Lueker, M., Kozarek, J., & Hill, K. (2022). A field study of maximum wave height, total wave energy, and maximum wave power produced by four recreational boats on a freshwater lake.

Retrieved from the University Digital Conservancy, <https://hdl.handle.net/11299/226190>

Mebane, C. A., Schmidt, T. S., Miller, J. L., & Balistrieri, L. S. (2020). Bioaccumulation and toxicity of cadmium, copper, nickel, and zinc and their mixtures to aquatic insect communities. *Environmental Toxicology and Chemistry*, 39(4), 812–833. <https://doi.org/10.1002/etc.4663>

Miller, A. C., Payne, B. S., & Shaffer, L. R. (1999). A shell gape monitor to study effects of physical disturbance on freshwater mussels. *Journal of Freshwater Ecology*, 14(2), 241–247.

<https://doi.org/10.1080/02705060.1999.9663675>

- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). 2025. Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state's greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce.
<https://www.pca.state.mn.us/sites/default/files/Iraq-3sy25.pdf>
- O'Callaghan, P., Kelly-Quinn, M., Jennings, E., Antunes, P., O'Sullivan, M., Fenton, O., & hUallacháin, D. Ó. (2018). The Environmental Impact of cattle access to Watercourses: A review. *Journal of Environmental Quality*, 48(2), 340–351. <https://doi.org/10.2134/jeq2018.04.0167>
- Pandolfo, T. J., Kwak, T. J., & Cope, W. G. (2012). Thermal Tolerances of Freshwater Mussels and their Host Fishes: Species Interactions in a Changing Climate. *Freshwater Mollusk Biology and Conservation*, 15(1), 69. <https://doi.org/10.31931/fmbc.v15i1.2012.69-82>
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwccasc.umn.edu/node/821>
- Richard, J. C., Leis, E. M., Dunn, C. D., Harris, C., Agbalog, R. E., Campbell, L. J., Knowles, S., Waller, D. L., Putnam, J. G., & Goldberg, T. L. (2022). Freshwater Mussels Show Elevated Viral Richness and Intensity during a Mortality Event. *Viruses*, 14(12), 2603. <https://doi.org/10.3390/v14122603>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.
<https://statesummaries.ncics.org/chapter/mn/>
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>
- Schramm, H.L., Richardson, W.B., & Knights, B.C. (2015). Managing the Mississippi River floodplain: Achieving ecological benefits requires more than hydrological connection to the river. In: Hudson, P., & Middlekoop, H. (Eds.) *Geomorphic Approaches to Integrated Floodplain Management of Lowland Fluvial Systems in North America and Europe*. Springer, New York, NY. https://doi.org/10.1007/978-1-4939-2380-9_8
- Shirey, V. M., Lewthwaite, J. M., Gawel, A. M., & Guzman, L. M. (2025). 50 years of invertebrate conservation under the United States Endangered Species Act—history and threats to species. *Frontiers in Conservation Science*, 6. <https://doi.org/10.3389/fcosc.2025.1505451>
- Smarts, M. M., Radar, R. R., Nielsen, D. N., & Claflin, T. O. (1985). The effect of commercial and recreational traffic on the resuspension of sediment in Navigation Pool 9 of the Upper Mississippi River. *Hydrobiologia*, 126(3), 263–274. <https://doi.org/10.1007/bf00007504>
- Spear, M. J., Harris, B. S., Bookout, T. A., Ickes, B. S., Jankowski, K. J., Solomon, L. E., Maxson, K. A., Harris, A. L. W., Mathis, A. T., Schaick, S. J., Williams, J. A., DeBoer, J. A., Lenaerts, A. W., Hine, E. C., Chick, J. H., & Lamer, J.

- T. (2024). Reduction of large vessel traffic improves water quality and alters fish habitat-use throughout a large river. *The Science of the Total Environment*, 946, 172705. <https://doi.org/10.1016/j.scitotenv.2024.172705>
- Stepanian, P. M., Entrekin, S. A., Wainwright, C. E., Mirkovic, D., Tank, J. L., & Kelly, J. F. (2020). Declines in an abundant aquatic insect, the burrowing mayfly, across major North American waterways. *Proceedings of the National Academy of Sciences*, 117(6), 2987–2992. <https://doi.org/10.1073/pnas.1913598117>
- Stoeckel, J. A., Rehmann, C. R., Schneider, D. W., & Padilla, D. K. (2004). Retention and supply of zebra mussel larvae in a large river system: importance of an upstream lake. *Freshwater Biology*, 49(7), 919–930. <https://doi.org/10.1111/j.1365-2427.2004.01237.x>
- Tarter, A. A., Ford, D. F., Symonds, D. E., Ford, N. B., & Schwalb, A. N. (2022). Impact of extreme climatic events on unionid mussels in a subtropical river basin. *Hydrobiologia*, 850(6), 1427–1442. <https://doi.org/10.1007/s10750-022-04819-7>
- Vaughn, C. C. (2018). Ecosystem services provided by freshwater mussels. *Hydrobiologia*, 810(1), 15–27. <https://doi.org/10.1007/s10750-017-3139-x>
- Wang, N., Ingersoll, C. G., Hardesty, D. K., Ivey, C. D., Kunz, J. L., May, T. W., Dwyer, F. J., Roberts, A. D., Augspurger, T., Kane, C. M., Neves, R. J., & Barnhart, M. C. (2007). Acute toxicity of copper, ammonia, and chlorine to glochidia and juveniles of freshwater mussels (unionidae). *Environmental Toxicology and Chemistry*, 26(10), 2036–2047. <https://doi.org/10.1897/06-523r.1>
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>
- Wohl, E. (2017). The significance of small streams. *Frontiers of Earth Science*, 11(3), 447–456. <https://doi.org/10.1007/s11707-017-0647-y>

Terrestrial Invertebrates

Overview

This sub-chapter highlights different groups of terrestrial invertebrates, including insects, spiders, and snails. Only a few of these species groups have been well-surveyed in Minnesota. For aquatic invertebrates, including mussels and insects such as dragonflies and caddisflies, please see the Aquatic Invertebrates sub-chapter. The paragraphs below provide insights into the diversity of invertebrates in our state.

Insects

The diversity of insects in Minnesota is likely greater than the combined diversity of all other animals, yet we know very little about most insect species. Our conservation status assessment includes approximately 3,500 insect species, but the true richness is likely 30,000 or more species (Jessica Petersen, DNR, pers. comm). Insect groups for which we have the best statewide information include grasshoppers (Acrididae; Haarstad 1990), stink bugs (Pentatomidae; Koch et al. 2014), ground beetles (Carabidae; Gandhi et al. 2005), bees (Anthophila; Portman et al., 2023), butterflies and moths (Lepidoptera), and tiger beetles (Cicindelidae). These groups represent a tiny fraction of the full suite of insect species in Minnesota, leaving much yet to be discovered.

There are general concerns about declines in insects (Seibold et al., 2019), but long-term datasets to assess insect population trends are lacking. One exception is a recent analysis of butterfly monitoring projects, based on repeated transects in Midwest states (Illinois, Iowa, Michigan, Missouri, and Ohio), and North American Butterfly Association Fourth of July circle counts. Data from these and other programs were compiled into a study that estimated butterfly abundances among Midwestern species declined 17% from 2000 to 2020 (Edwards et al., 2025; [State of the Butterflies Report](#)). Nationally, 62 of 119 butterfly species (52%) showed declines, while only 25 (21%) showed increases (Edwards et al., 2025). Because population trend data are limited for most insects, other measures are sometimes used to indicate trends, such as [range extent](#) of the [rusty-patched bumble bee \(*Bombus affinis*\)](#) and [extent of occupied overwintering area](#) of the eastern migratory monarch butterfly (*Danaus plexippus*).

Acknowledging that it is not feasible to survey all groups of insects for which we lack information, the following groups could be prioritized for future baseline surveys:

- Fireflies (Lampyridae) are completely missing from the Minnesota Species of Greatest Conservation Need list due to lack of preliminary surveys and understanding of their biodiversity.
- Hover flies (Syrphidae), to better understand biodiversity and contributions to pollination.
- Bee flies (Bombyliidae), to better understand biodiversity and contributions to pollination.
- Beetles, such as leaf beetles (Chrysomelidae) that require specific plant species for food or other critical resources, may be of conservation concern if their required plants are under threat.
- Robber flies (Asilidae) could be of conservation concern given that most are predators and some may be habitat specialists.
- Many bush crickets (Tettigoniidae) are likely habitat specialists and prairie endemics, but their diversity is poorly understood.

- A statewide survey of ants (Formicidae) is warranted, as ants are known to be indicators of functioning ecosystems (Underwood & Fisher, 2006).

Lacking basic biological information about so many species, each with a unique role in sustaining the health of natural communities, compromises our ability to address their conservation needs. There is an immediate need to document the diversity of insects in Minnesota before further losses are sustained and therefore go undocumented (Cardoso et al., 2020).

Bees

Thanks to a statewide bee survey completed in 2022, we now know that Minnesota hosts over 500 native bee species, and around a dozen non-native species (Portman et al., 2023; see the [Minnesota Statewide Bee Survey](#) report). This [preliminary list](#) provides a baseline for conservation. This list is rapidly evolving, and updated to accommodate new findings, most recently to add two species that had not been documented in Minnesota in over 50 years. In addition, we are continuing to add more information about the 30% of bee species that are considered oligolectic, or specialists on a particular species or genus of plant pollen that they provide to their brood. These specialist species were designated as SGCN due to their being environmentally constrained to a limited resource, making them vulnerable to threats such as climate change, development, and natural system modifications. We know far more than we did 10 years ago about the bees of Minnesota, but we still have a lot to learn about population locations and their trends. See [Native Bees of Minnesota](#) for more information.

Because bees rely on nectar and pollen from flowers, and most bee species gather plant pollen to feed their broods, bees are often considered the most important animal pollinators. Global threats to bees include loss or degradation of habitat, climate change, phenological mismatches between flower bloom time and bee flight period, insecticide exposure, parasites, pathogens and disease, and low genetic diversity (Lima et al., 2022). Pathogens, disease, and low genetic diversity are thought to be the primary threats to bumble bees (Cameron et al., 2011). There are several boreal species (e.g., *Osmia neactica*) on the southern edge of their range in Minnesota that are threatened by climate change and habitat modifications including timber harvest. The rusty patched bumble bee is now found in only a fraction of its historical range and was listed as federally endangered in 2017 ([Rusty patched bumble bee status review](#)). It has a relatively robust population in Minnesota, particularly in comparison to many parts of its former and current range. Rusty patched bumble bee is one of the preferred hosts of Ashton's cuckoo bumble bee (*Bombus ashtoni*), a social parasite that invades the nests of other bumble bee species. Ashton's cuckoo bumble bee has not been detected in Minnesota since 1998 and may be extirpated from the state.

Conservation actions for bees include reconstructing prairie habitats, protecting remnant prairie habitats that have never been converted to row-crop agriculture, managing for floral diversity across all habitat types, conserving nesting habitat, and minimizing insecticide use. Reconstructed prairies can support robust bee communities and even some [rarer species](#), though some at-risk species are preferentially found in remnant habitats (Lane et al., 2023). Generalists like bumble bees are attracted to habitats of many types and sizes, including urban gardens.

Butterflies and Moths

Lepidoptera (moths and butterflies) stand out as the dominant order of herbivores in terrestrial ecosystems, both in terms of species richness and biomass (Grimaldi & Engel, 2005). Despite this importance, most families have received little study in Minnesota historically, with most attention on butterflies (Macy & Sheppard, 1941). Recent intensive surveys – primarily by the Minnesota Biological Survey – are striving to change this. As of this writing, over 2100 species have been documented in Minnesota and more continue to be discovered on a regular basis. A conservative estimate of Minnesota Lepidoptera richness is 2500 species, but 3000 would not be surprising given many species remain undescribed due to taxonomic difficulty and/or lack of survey effort (Kyle Johnson, DNR, pers. comm.). Of the 2100+ known species in the state, the majority are poorly understood in terms of their distributions, habitat preferences, life histories, and conservation concerns. Therefore, the current SGCN list underrepresents Lepidoptera. Despite these challenges, our collective knowledge of habitat specialists and species of conservation concerns has improved drastically over the past two decades and continues to improve at a rapid pace.

Butterfly species richness and abundance have both declined in Minnesota. Eight species of butterflies are now considered extirpated or likely extirpated: [Arogos skipper \(*Atrytone arogos*\)](#), [Assiniboia skipper \(*Hesperia assiniboia*\)](#), [Ottoe skipper \(*Hesperia ottoe*\)](#), [Uncas skipper \(*Hesperia uncas*\)](#), [Garita skipperling \(*Oarisma garita*\)](#), [Poweshiek skipperling \(*Oarisma poweshiek*\)](#), [Uhler's arctic \(*Oeneis uhleri varuna*\)](#), and [Karner blue \(*Plebejus samuelis*\)](#). Likewise, only a single population of [Dakota skippers \(*Hesperia dacotae*\)](#) remains despite extensive searching of historically occupied sites. Abundance has also declined among even previously common species (Jessica Petersen, DNR, pers. comm.). Similarly, there have been range contractions for species such as the [regal fritillary \(*Argynnis idalia*\)](#) and dusted skipper (*Atrytonopsis hianna*). Once common species like eastern tailed-blues (*Cupido comyntas*) and pearl crescents (*Phyciodes tharos*) are less abundant than in the past. There is a conservation need for butterfly monitoring of common species in Minnesota to document changes in abundance.

Prairies, savannas, barrens, and peatlands are particularly rich in habitat specialists and species of conservation concern. Habitat loss and degradation are likely the greatest threats to Lepidoptera overall (as in the case of most invertebrates) but other important issues such as pesticides and climate change likely threaten them as well. The essential character of prairies, savannas, and barrens depends on periodic disturbances, notably fire and grazing. Providing these disturbances in isolated habitat remnants can have deleterious effects if not implemented carefully. Timing and intensity, both of fire and grazing, are important considerations, and leaving untreated refugia to assure adequate survival for population recovery is essential. Studies on how different habitat management regimes affect Lepidoptera are lacking and would be quite helpful in guiding future conservation efforts.

Beetles

Of the 18 beetles designated as SGCN or SNI, 11 are rare tiger beetles of the genus *Cicindela*. The [Laurentian tiger beetle \(*Cicindela denikei*\)](#), threatened in Minnesota and globally vulnerable (G3), is limited to rock outcrops and sandy openings in hardwood forests in the Border Lakes Subsection. The [sandy tiger beetle \(*C. limbata*\)](#) is known from only one location in the state, sand dunes in Polk County, and may be extirpated now from the state. The [sandy stream tiger beetle \(*C. macra*\)](#) is known from several locations along moist, sandy stream shores in southeastern Minnesota. The [splendid tiger beetle \(*C. splendida*\)](#) requires steep clay embankments and is known from several locations in the Blufflands Subsection. The [crimson saltflat tiger](#)

[beetle \(*C. fulgida*\)](#) is restricted to alkaline shorelines and mudflats in western Minnesota. Very few re-surveys of these species have occurred with any regularity over the past 10 years.

Other Invertebrates

Spiders

Despite their abundance and diversity in most ecosystems, spiders are largely understudied as a group. Yet they serve as important predators of other invertebrates and in turn serve as prey for birds and other organisms. Minnesota first listed eight jumping spiders as species of special concern in 1995. The Minnesota County Biological Survey then incorporated jumping spider surveys into their work around the state and did so through 2013. The list of spiders in Minnesota's State Wildlife Action Plan has seen some revision as new data points have been established; the most recent revision takes into consideration recent collection work and citizen-science data from websites such as iNaturalist (www.inaturalist.org) and BugGuide (www.bugguide.net).

The greatest threat to Minnesota's rare spiders is habitat loss: both quality and connectivity. Thankfully, their conservation can be tied to conservation of other species through habitat preservation efforts that benefit rare birds, butterflies, plant communities, etc. For example, one [jumping spider \(*Tutelina formicaria*\)](#), listed as threatened in Minnesota is only known from protected sand prairies in Anoka County (although also distributed in other states). A couple of other species of special concern are denizens of rare plant communities that in Minnesota are limited to goat prairies of southeastern Minnesota. The updated list has more diversity than in the past with four jumping spiders, an orbweaver, a cobweb spider and a purse-web spider. For some, extensive surveys in appropriate habitat have indicated their rarity. Other species occur on the updated list because of their limited dispersal methods or their medical significance to humans. Spider distributions are poorly understood in Minnesota and additional survey work could reveal their true abundance and improve plans for their conservation.

While some spider collection has occurred historically in Minnesota, most notably by Willis J. Gertsch (1930s), Daniel T. Jennings (1960s), Bruce Cutler (1970-90s), and more recently William J. Ehmann (1996-2015), such collections were often focused on one taxon (Richman & Cutler 1978; Cutler & Jennings 1992; Ehmann 2002, 2011; Ehman & Boyd 1997), limited in their geographic scope (Cutler et al. 1975; Heimer et al. 1984) or both (Jennings & Cutler 1996). As a result, significant gaps remain in the knowledge of not only which species occur in Minnesota, but how they are distributed within the state.

Recent work has been done to improve our understanding of spider distribution in Minnesota by reviewing the taxonomic literature, finding museum records, collecting and identifying spiders throughout the state and monitoring citizen-science websites such as BugGuide and iNaturalist. Nearly 2,000 county records have been documented by photographs taken by Minnesotans in [an iNaturalist project](#) established specifically for Minnesota spiders (Chad Heins, Bethany Lutheran College, pers. comm.). The collection housed at Bethany Lutheran College currently has 3,087 catalogued vials representing 34 families collected in 70 of Minnesota's counties. These include several first state records as well as species awaiting formal description, with a formal checklist in preparation (Heins, in prep).

Presently in Minnesota we have confirmed the presence of 543 species of spiders, while there may be as many as 600-800 species (Chad Heins, pers. comm.). We still know very little about spider distributions in Minnesota,

let alone North America. In 2024, Chad Heins found three new species for Minnesota's list. The closest literature records for those species were Colorado, eastern Illinois, and Michigan.

Snails

Minnesota is home to approximately 80 species of native terrestrial snails and slugs (gastropods). Terrestrial snails can be found in most habitats in Minnesota ranging from semi-aquatic marsh edges to mesic bluffs. Terrestrial snails are grazers of organic material including detritus, rotting wood, and algae; they also serve as food items for other organisms. Despite nearly ubiquitous distributions, there have been no recent concerted statewide surveys for terrestrial gastropods in Minnesota. Several important studies in the northwestern and northeastern portions of Minnesota have documented 60 species (Nekola et al., 1999; Nekola, 2002), however these data are over 20 years old.

The lack of recent, targeted surveys in southern Minnesota are concerning given the presumed endemism within the region. For instance, in southeastern Minnesota the [bluff vertigo \(*Vertigo meramecensis*\)](#) is a rare species classified as threatened in Minnesota. Along with other species of land snails it persisted in unglaciated Pleistocene refugia found on steep, moist, shaded, cool north-facing slopes and cliff faces in the Blufflands Subsection. Bluff vertigo also occurs on algific slopes and moderate cliffs but generally avoids areas with continuous cooling from cold air or water discharge. It is known to occur at only six sites in Minnesota in Olmsted, Winona, and Wabasha counties, not all of which are protected.

Further conservation issues arise with snail taxonomy due to strong roots in morphological species concepts and a high potential for ecophenotypic plasticity. This has led to lingering questions about the taxonomic distinctness and overall distribution of some of the SGCN land snails.

Summary of Terrestrial Invertebrate SGCN and SNI

There are 267 terrestrial invertebrate Species in Greatest Need of Conservation (SGCN): 117 bees, 122 butterflies and moths, 11 beetles, 1 cicada, 1 leafhopper, 8 snails, and 7 spiders (see Table 2.10).

There are 430 terrestrial invertebrate Species in Need of Information (SNI): 125 bees, 10 beetles, 257 butterflies and moths, 5 flies, 6 grasshoppers, 1 millipede, 3 springtails, 15 snails, and 8 true bugs (see Table 2.10). For the full list of SGCN and SNI invertebrate species, see Appendix B.

Table 2.10. Summary of number of species of terrestrial invertebrate SGCN and SNI.

Species Group	# SGCN	# SNI	Total Assessed*	# SGCN removed from 2015 List	# SGCN added to 2015 List	# 2015 SGCN demoted to SNI	# SNI added
Bees (Class Insecta; Order Hymenoptera)	117	125	515	0	112	0	125
Beetles (Class Insecta; Order Coleoptera)	11	10	89	0	3	1	9
Butterflies, Moths (Class Insecta; Order Lepidoptera)	122	257	2087	2	95	4	253
Flies (Class Insecta; Order Diptera)	0	5	67	0	0	0	5
Grasshoppers, Crickets (Class Insecta; Order Orthoptera)	0	6	11	0	0	0	6
Millipedes (Class Diplopoda)	0	1	1	0	0	0	1
Snails (Class Gastropoda) - Terrestrial	8	15	81	0	3	0	15
Spiders (Class Arachnida; Order Araneae)	7	0	528	6	3	0	0
Springtails (Class Collembola)	0	3	3	0	0	0	3
True Bugs, Cicadas, Leafhoppers (Class Insecta; Order Hemiptera)	2	8	12	1	1	1	7
TOTAL	267	430	3394	9	217	6	424

* Only bees, butterflies and moths, and spiders are considered to have been well-surveyed in Minnesota, such that the number of species assessed is somewhat close to the number of species likely present in the state, while other groups are much more under-surveyed.

Habitat Associations

Terrestrial invertebrate Species in Greatest Conservation Need use a broad range of habitats (see Table 2.11). Some species, especially those with more general habitat requirements, were assigned to multiple habitats.

Appendix C provides a comprehensive list of the SGCN and into which habitats they associate. Spiders' habitat use was further distinguished into primary or secondary. Primary habitats are those that a species uses most consistently and is most reliant on; if it were gone or degraded, it would have the greatest effect on the species. Secondary habitats are those where a species can be found some of the time, but less frequently than a primary habitat.

Overall, prairies and grassland provide habitat for the greatest number of terrestrial invertebrate SGCN. Savanna, cliff, talus, and rock outcrops, and non-forested wetlands also provide habitat to a significant number of SGCN.

Table 2.11. Habitat Associations of Terrestrial Invertebrate SGCN.

Habitats	Bees	Beetles (terrestrial)	Butterflies	Moths	Snails (terrestrial)	Spiders	True Bugs	Total
Prairie and Other Grasslands	83	4	15	56	0	6	2	166
Savanna	4	6	8	40	0	5	0	63
Upland Conifer Forest and Woodland	5	2	5	7	5	0	0	24
Upland Deciduous Forest and Woodland	9	1	0	4	6	2	0	22
Mesic Hardwood Forest	8	1	0	0	8	2	0	19
Deciduous Wet Forest	0	0	0	1	1	2	0	4
Coniferous Forest Wetland	0	0	3	12	2	0	0	17
Riparian and Floodplain Forest	0	2	0	1	0	2	0	5
Non-forested Wetlands	20	1	6	14	0	1	0	42
Rivers and Streams	1	0	0	0	0	0	0	1
Lakes	4	0	0	0	0	1	0	5
Cliff, Talus, and Rock Outcrops	15	4	8	14	8	1	0	50
Urban and Other Developed Lands	2	3	1	0	0	0	0	6

Primary Stressors for Terrestrial Invertebrates



Development

Habitat loss and fragmentation is among the most prevalent threats to terrestrial invertebrates currently listed under the federal Endangered Species Act (Shirey et al., 2025). Residential and commercial development is a driver of both habitat reduction and fragmentation. Indeed, the [hairy-necked tiger beetle \(*Cicindela hirticollis rhodensis*\)](#), a state-listed endangered species in Minnesota, is restricted to a very small area along the Lake Superior shoreline. Habitat loss and shoreline development, including breakwalls affecting natural shoreline processes, are ongoing concerns (DNR, 2022). Species of butterflies that are negatively impacted by loss of habitat due to development in the Twin Cities Metro include [Leonard's skipper \(*Hesperia leonardus leonardus*\)](#) and Olympia marble (*Euchloe olympia*).



Crop Production

Prairie species such as the Dakota skipper, Poweshiek skipperling, and regal fritillary are impacted by conversion of native prairie to row crop agriculture. This loss of prairie habitat is both historical (Samson & Knopf, 1994), and ongoing in Minnesota (Lark et al., 2018).

Insecticides used to control crop pests (especially those applied towards the end of the growing season to control soybean aphids) have been correlated with the recent decline of listed grassland specialist butterflies like Poweshiek skipperling and Dakota skipper, even in protected prairies that are adjacent to row crop fields (Runquist et al., 2024).



Livestock Management Chan

Grazing livestock can potentially have positive impacts on terrestrial invertebrates and serve as an ecological replacement for the loss of disturbance from wild ranging ungulates (i.e., bison; reducing duff accumulation and promoting growth of native vegetation). The Poweskiek skipperling, for example, appears to benefit from light, or “conservative” grazing regimes (Knight et al., 2024). However, herds that are too large or left too long in sensitive management units may have a destructive impact on native insect communities and their habitats. The term “conservation grazing” is used broadly and rigorous assessment of responses of target species and their habitats is warranted.

Care must be taken when treating livestock for pests and parasites, as treatments often contain insecticides. Some insecticides persist in the environment weeks after treatment and can harm beneficial insects and contaminate soil and water via the animals’ feces.



Mining and Quarrying

Mining for sand and gravel in Minnesota is common throughout the grassland region of Minnesota and can disturb and destroyed prairies, which could have widespread effects for many insect species, including prairie obligate species such as the Dakota skipper. Mining and quarrying operations can also cause loss of habitat for cave invertebrate species. Some species that specialize on sandy habitats, such as the [ghost tiger beetle](#) (*Cicindela lepida*) may be particularly affected by sand and gravel mining and quarrying activity.



Roads, Trails, and Railroads

Roads replace and fragment habitat for insects. In places where little native vegetation remains, such as in highly agricultural or developed areas, when managed well, roadsides can facilitate insect dispersal and provide important nectar resources (Ries et al., 2001; Darst et al. 2024). Roadside rights-of-ways can provide native vegetation that provides valuable resources for foraging and reproducing insects, such as monarchs using milkweed (Kasten et al., 2016; Cariveau et al., 2024); indeed, roadsides and other rights-of-way have been proposed as important components of monarch conservation efforts (Thogmartin et al., 2017). When road rights-of-way are planted in nonnative plants, there is a missed opportunity to provide native plants on the landscape, which can be used by a wide variety of wildlife, particularly insects (Mitchell et al. 2024). However, roads also present risks, including collision with vehicles, which vary significantly across space and species (Kantola et al., 2019; Keilsohn et al., 2018; McKenna et al., 2001). In addition, roads introduce pollutants from de-icing salts, heavy metals from vehicle wear and tear, and the legacy of leaded gasoline in Minnesota roadside soils and plants (Mitchell et al 2020, Shephard et al. 2022).

Road maintenance activities are necessary for maintaining sight lines for safety. Broadcast herbicide use can diminish nectar resources and have negative non-target effects. Frequent mowing of roadsides removes nectar resources and reduces the ability for monarch butterflies to successfully reproduce on that milkweed (Hopwood et al. 2015), although mowing earlier in the season, may be beneficial in stimulating new growth in milkweed where monarchs prefer to oviposit (Cariveau et al. 2024, Knight et al. 2024).



Hunting and Collecting Animals

Harvest of monarch butterflies for unregulated captive rearing programs (for educational use or release) is not a recommended action due to the risk of pathogen spread and fitness consequences for wild populations (Pelton, 2018).



Gathering Plants and Fungi

Illegal harvest of specific host or nectar plant species can negatively impact terrestrial insects that depend on them.



Timber Harvest

Timber harvest of mesic hardwood forests where the fragile spring ephemeral understory may not recover from that disturbance could have impacts to the various bee species that are specialists of spring ephemerals. For example, the spring beauty mining bee (*Andrena erigeniae*) exclusively feeds its brood pollen from spring beauty (*Claytonia* sp.), the trout lily miner bee (*Andrena erythronii*) feeds its brood pollen from trout lily (*Erythronium* sp.), *Andrena polemonii* feeds its brood pollen from Jacob's ladder (*Polemonium* sp.), and *Andrena uvulariae* feeds its brood pollen from bellwort (*Uvularia* sp.). Similarly, many of the bumble bee queens that emerge in the early spring use pollen and nectar from spring ephemerals to begin to build their colonies, including rusty patched bumble bee. Timber harvest in peatlands paints a mixed picture. Recent surveys of Lepidoptera in Minnesota peatlands show a diverse fauna in sparsely or semi- treed peatlands approximately 2-20 years after timber harvest. But latter successional stages have consistently had poor results, and even 70-year+ stands have been strikingly depauperate, relative to their old-growth counterparts. [Taiga Alpine \(*Erebia mancinus*\)](#) is a SGCN which has yet to be found in logging regenerated peatland habitats, even those with similar floral composition. One hypothesis is that the tree structure in regenerated stands blocks more sunlight and offers less favorable flyways relative to old-growth stands.



Recreation

In some locations, hairy-necked tiger beetles have been negatively impacted by beach use.



Fire Management

Prairie dependent insects such as regal fritillary and Dakota skippers have been negatively impacted by the alteration of fire regimes that have resulted from decades of fire suppression. Prescribed fire has been found to be beneficial for some wild bees including ground-nesting bees (Brokaw et al., 2023; Galbraith et al., 2019). However, some SGCN insects such as the crossline skipper (*Polites origenes*) and the yellow bumble bee (*Bombus fervidus*) that overwinter as larvae or nest above ground (Scott, 1979; Williams et al., 2014) many be vulnerable to spring fires.



Invasive Species (Problematic Non-native Species)

Invasive, non-native plant species can negatively impact terrestrial insect species by outcompeting and displacing species-specific host and nectar plants. Additionally, some invasive plant species may become populations 'sinks' by triggering oviposition on unsuitable hosts that would not have been present naturally. This has been found to be the case for monarchs ovipositing on swallow-worts (Casagrande & Dacey, 2007) and Dakota skippers on invasive cool-season grasses (Nordmeyer et al., 2021).

Some non-native forbs provide nectar, pollen, and can serve as host plants for some insects. For example, Melissa blue (*Plebejus melissa*) a species in need of information, uses crown vetch as a host plant. Native plants that Melissa blue larvae use as a host plant include Astragalus and associated peas. Spot spraying crown vetch has perhaps negatively impacted Melissa blue. Spotted knapweed provides nectar for several species of insects including Leonard's skipper and Rusty patched bumble bee, at times when no other nectar resources are available.

Some non-native bee species are thought to be contributing to the decline of native species (LeCroy et al., 2020). *Osmia taurus* was collected by the Minnesota Biological Survey at Carley State Park, near the city of

Plainview in Wabasha County on May 10, 2023. At the time of the publication of the bees of Minnesota (Portman et al., 2023), there were 11 non-native species of bees documented in Minnesota. Another mason bee, *Osmia cornifrons* was documented in 2025 in the Twin Cities by community members.



Problematic Native Species

Pickerelweed (*Pontederia cordata*) is a native, aquatic plant that is thought by some to be outcompeting wild rice in northern Minnesota lakes. Some managers are attempting to remove and control pickerelweed on lakes where wild rice is thought to have declined due to increases in abundance of pickerelweed. There are two SGCN bees, the pickerelweed long-horned bee (*Melissodes apicatus*) and the pickerelweed bee (*Dufourea novaeangliae*) that exclusively feed their brood pollen from pickerelweed and would likely suffer if populations of these species occur at or near lakes where pickerelweed is being actively removed.



Diseases and Pathogens

Bees are affected by a number of pathogens. Pathogens spread from commercially reared bumble bee species may contribute to some of the declines in native bumble bee species (Colla et al., 2006; Szabo et al., 2012).



Air-borne Pollution

Pesticide drift is a major issue for terrestrial insects in Minnesota. Approximately 25 % of pesticides end up drifting to non-target areas, and drift can occur over 100s of miles from where pesticides were released (Albaseer et al., 2025). Prairies in agricultural areas are subject to insecticide exposure from off-site aerial spraying (Goebel et al., 2022). Prairie reconstructions on land formerly used for agriculture may also be exposed to systemic insecticides from treated seed. Systemic insecticides eventually degrade in soil, but this can take up to several years (Goulson, 2013).

Aerially applied insecticides intended to control biting insects (e.g., mosquitos; both nuisance and disease transmitting) can impact non-target species (Runquist et al., 2024). Pesticide use has been identified as one of the leading drivers of insect declines in the upper-Midwest (Van Deynze et al., 2024). Depending on the chemicals used, these can cause direct mortality upon contact/ingestion, or act as growth inhibitors causing other arthropod species to develop to maturity.

Biological insecticides (such as Btk, *Bacillus thuringiensis kurstaki*) used to control invasive Lepidoptera such as spongy moth, *Lymantria dispar*, can cause significant declines in native Lepidoptera species (Strazanac et al., 2007).



Light and Noise Pollution

Nearly half of the United States land surface has light-polluted skies (Falchi et al., 2016). Sources of light pollution at night include interior and exterior lights on homes, buildings, billboards, parking lots, and streets as well as lighted sports fields, industrial plants, and airports (DarkSky International, 2025a). Studies also point to the ecological impacts created during the daytime by polarized light that reflects off surfaces such as asphalt roads, glass panes, and solar panels (Chock et al., 2021; Horvath et al., 2009). Because most of the nation's large urban areas (with multiple artificial light sources every night), are embedded in the Temperate Deciduous

Forest biome, biota occupying temperate forests are the most vulnerable to these impacts (Aubrecht et al., 2010).

Impacts of light pollution include disrupting the behavior of amphibians whose breeding calls are delivered at night, drawing nocturnal insects to artificial lights that increase their predation risk, and disorienting birds that use environmental cues to guide their nighttime migration (DarkSky International, 2025a; 2025b). In addition, the phenology, growth rates and biomass of plants are impacted (Bucher et al., 2023). Bright day and dark night cycles synchronize the internal clocks of plants and animals, and alterations to these cycles can impact hormone production, reproductive behavior, neural activity, and metabolic functions (Bumgarner & Nelson, 2021).

Noise pollution from aircraft, vehicles, boats, and other human causes is pervasive. Anthropogenic noise was detected in 36% of national parks surveyed (Buxton et al., 2019), and 12% of wilderness areas had anthropogenic noise levels 3 decibels higher than levels predicted to occur naturally (Buxton et al., 2017). Exposure to increasing noise levels can change the spatial distribution and movement patterns of wildlife, cause avoidance of feeding and nesting areas, and interrupt sleeping patterns and communications (Kok et al., 2023). For plants, noise can reduce seedling recruitment (Phillips et al., 2021).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

Changes in temperature associated with climate change can have significant impacts on ectothermic organisms, like terrestrial insects, whose development and reproduction are closely tied to ambient temperatures. Weather patterns have been shown to influence insect abundance (Muller et al., 2024). Furthermore, climate warming is driving poleward and elevational shifts in many insect species. Cold-adapted specialists may experience population declines, while warm-tolerant species expand their ranges, reshaping community structure (Parmesan, 2006; Robinson et al., 2021). Shifts in annual temperature patterns can also alter the timing and phenology of insect life cycles. In extreme cases, these changes result in mismatches between co-evolved species during critical life history stages or lead to increased voltinism—the number of insect generations per year—which can, paradoxically, cause population declines due to incomplete or poorly timed generations (Altermatt, 2010; Van Dyck et al., 2015).

Climate warming is a great threat to boreal insects near their southern range limit in northern Minnesota. Some peatland butterflies and moths are already showing worrying signs, with recent surveys failing to find them at the southern range extremes. There are also concerns that habitat shifting caused by changes in temperature and hydrology may decrease the amount of suitable habitat available for prairie butterflies and pollinators.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Extreme precipitation fluctuations are likely to stress populations of wetland specialist invertebrates, both directly (mortality) and indirectly (habitat change). Recent surveys (2024-2025) in peatlands following unusual drought and flood cycles yielded unprecedented low numbers for some specialists (e.g. *Boloria freija*, *B. frigga*). It is plausible that such extreme years might cause extirpation of populations, especially at small and/or vulnerable sites. Likewise, for species with extremely small or localized populations, the impacts of severe weather conditions can be devastating. Taxa that historically had larger ranges may have had sufficient population redundancy to endure local extirpation caused by severe weather or other catastrophic events.

Priority Conservation Strategies

To implement the SGCN Goal of this Plan (*Conserve rare, declining, and vulnerable wildlife and plant SGCN through targeted actions*), three strategies were identified:



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Examples of conservation actions are grouped below under these three strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Some actions that are focused on important population monitoring and public engagement needs might not always be tied to a specific stressor; these are labeled 'n/a' in the stressor column.

Potential Conservation Actions for Terrestrial Invertebrates



Strategy 1. Survey, monitor, and research to document the distribution and trends of SGCN animals and plants, threats they experience, and conservation actions that support resilient populations.

Stressor	Action
	Conduct population and habitat sampling for pathogens and pesticides.
	Research how native terrestrial invertebrates respond to different management regimes, including prescribed fire and grazing, with a focus on SGCN.
n/a	Expand survey programs to monitor SGCN species and other under-studied terrestrial invertebrates.
n/a	Encourage public participation in observational surveys through available platforms like Bumble Bee Watch and iNaturalist, and in-person events like BioBlitzes.



Strategy 2. Collaborate to deliver conservation actions that support resilient populations of SGCN and their habitats in partnership with agencies, Tribes, non-governmental organizations, and private landowners.

Stressor	Action
	Prevent conversion and fragmentation of remnant habitats by pursuing protection for them, especially prairies. Many prairie-dependent SGCN insects require remnant prairies and to date have not been shown to utilize planted or reconstructed prairies or degraded grasslands.
	Improve planted habitats for the various species of SGCN insects that rely on specific plant species to complete their life cycle by including those plants in reconstructions, such as loosestrife (<i>Lysimachia</i>) for oil collecting bees (<i>Macropis</i>), prairie violets (<i>Viola pedatifida</i>) for regal fritillaries (<i>Argynnis idalia</i>), milkweeds (<i>Asclepias sp.</i>) for monarchs (<i>Danaus plexippus</i>).
	Partner with private landowners to conserve prairie and provide technical guidance. Incentivize/subsidize grassland habitat. Reconstruct native prairies. Increase crop diversity and promote perennial crops.
	Maintain or enhance the integrity of native plant communities by maximizing native plant diversity and reducing invasive plant species whenever possible.
	Follow best practices for treatment of livestock with systemic insecticides for pest or parasite control (such as ivermectin, doramectin, or eprinomectin), not to be released on conservation lands for at least 30 days following treatment. If parasite treatment is needed during the grazing period, the best practice is to remove affected animals for treatment and return them to the site after the recommended amount of time (DNR, 2025).
	Support and promote use of native blooming plants including host plants such as milkweed in rights-of-way and energy development projects that may benefit monarch butterflies and bumble bees.
	Create pollinator-friendly energy and transportation rights of way following best management practices such as Minnesota Department of Agriculture's Pollinator Best Management Practices for Roadsides and Other Rights-of-Way and resources found among the University of Illinois-Chicago Rights of Way Working Group's compilation of Best Management Practices . Example practices include seeding with native plants, mowing only the width of the area for safety or needed to protect sight lines, mowing only once per growing season, and minimizing broadcast spraying of herbicides.
	Identify and protect important areas for SGCN from overuse. For instance, block off important areas of beach from public access when sensitive species are present.
	Apply prescribed fire strategically to help manage prairies and other grasslands while also taking into consideration vulnerabilities of rare SGCN. Manage diversity of native vegetative structure and composition across the landscape and over time. Apply fire only to some areas while leaving aside skipped refuge areas to provide for recolonization of burned sites.
	Advocate for pesticide drift buffers. Before installing habitat on previously agricultural land treated with herbicides or insecticides, allow the land to rest before planting. A temporary cover crop may be planted during this time. Refer to herbicide rotation restrictions for forage and cover cropping systems for wait times prior to planting. This document can be used as a surrogate for native species wait times.

	Implement best practices for wildlife-friendly outdoor lighting, such as turning off lights when they are not needed, using timers or motion detectors, shielding lights pointed downward, and employing lights in the warmer part of the spectrum. For more detailed information, refer to solutions advised by conservation organizations such as DarkSky , the Xerces Society , and National Audubon's Lights Out Program .
	Increase seed collection and restoration to head-start new pockets of habitat. Consider sourcing seeds from areas that are more similar climatically to future site conditions, or assisted migration.



Strategy 3. Develop and disseminate informational material to guide conservation actions for SGCN wildlife, such as species accounts, threat assessments, recovery plans, relevant regulations, avoidance measures, and beneficial habitat management strategies.

Stressor	Action
n/a	Develop and disseminate education materials that promote the value of terrestrial invertebrates, including Minnesota's Pollinators - DNR webpage .
n/a	Encourage the use of the DNR Pollinator Best Management Practices and Habitat Restoration Guidelines .
n/a	Encourage public participation in promoting habitat for pollinators including monarchs and bumble bees (see also Chapter 4. Public Engagement).
n/a	Advise to limit commercial and recreational rearing of at-risk or declining species, particularly the monarch butterfly.

References for Terrestrial Invertebrates

Albaseer, S. S., Jaspers, V. L. B., Orsini, L., Vlahos, P., Al-Hazmi, H. E., & Hollert, H. (2025). Beyond the field: How pesticide drift endangers biodiversity. *Environmental Pollution*, 366, 125526. <https://doi.org/10.1016/j.envpol.2024.125526>

Altermatt, F. (2010). Climatic warming increases voltinism in European butterflies and moths. *Proceedings of the Royal Society B Biological Sciences*, 277(1685), 1281–1287. <https://doi.org/10.1098/rspb.2009.1910>

Aubrecht, C., Jaiteh, M.S., & de Sherbinin, A. (2010). Global Assessment of Light Pollution Impact on Protected Areas. Center for International Earth Science Information Network, The Earth Institute at Colombia University.

Brokaw, J., Portman, Z. M., Bruninga-Socular, B., & Cariveau, D. P. (2023). Prescribed fire increases the number of ground-nesting bee nests in tallgrass prairie remnants. *Insect Conservation and Diversity*, 16(3), 355–367. <https://doi.org/10.1111/icad.12628>

Bucher, S. F., Uhde, L., Weigelt, A., Cesarz, S., Eisenhauer, N., Gebler, A., Kyba, C., Römermann, C., Shatwell, T., & Hines, J. (2023). Artificial light at night decreases plant diversity and performance in experimental grassland communities. *Philosophical Transactions of the Royal Society B Biological Sciences*, 378(1892). <https://doi.org/10.1098/rstb.2022.0358>

- Bumgarner, J.R. & Nelson, R. (2021). Light at Night and Disrupted Circadian Rhythms Alter Physiology and Behavior. *Integrative and Comparative Biology*, 61(3), 1160–1169.
- Buxton, R. T., McKenna, M. F., Mennitt, D., Brown, E., Fristrup, K., Crooks, K. R., Angeloni, L. M., & Wittemyer, G. (2019). Anthropogenic noise in US national parks – sources and spatial extent. *Frontiers in Ecology and the Environment*, 17(10), 559–564. <https://doi.org/10.1002/fee.2112>
- Buxton, R.T., McKenna, M.F., Mennitt, D.J., et al. (2017). Noise pollution is pervasive in US protected areas. *Science* 356, 531–33.
- Cameron, S. A., Lozier, J. D., Strange, J. P., Koch, J. B., Cordes, N., Solter, L. F., & Griswold, T. L. (2011). Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences*, 108(2), 662–667. <https://doi.org/10.1073/pnas.1014743108>
- Cardoso, P., Barton, P. S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., Fukushima, C. S., Gaigher, R., Habel, J. C., Hallmann, C. A., Hill, M. J., Hochkirch, A., Kwak, M. L., Mammola, S., Noriega, J. A., Orfinger, A. B., Pedraza, F., Pryke, J. S., Roque, F. O., . . . Samways, M. J. (2020). Scientists’ warning to humanity on insect extinctions. *Biological Conservation*, 242, 108426. <https://doi.org/10.1016/j.biocon.2020.108426>
- Cariveau, A. J. B., Haynes, G. A., Perish, P., Tuerk, K., Nootenboom, C., Mitchell, T., Lonsdorf, E., Oberhauser, K., & Snell-Rood, E. C. (2024). Host plants and landscape predict use of roadside habitat by breeding monarchs. *Insect Conservation and Diversity*, 17(5), 894–909. <https://doi.org/10.1111/icad.12758>
- Casagrande, R. A., & Dacey, J. E. (2007). Monarch Butterfly Oviposition on Swallow-Worts (*Vincetoxicum* spp.). *Environmental Entomology*, 36(3), 631–636. [https://doi.org/10.1603/0046-225x\(2007\)36](https://doi.org/10.1603/0046-225x(2007)36)
- Chock, R. Y., Clucas, B., Peterson, E. K., Blackwell, B. F., Blumstein, D. T., Church, K., Fernández-Juricic, E., Francescoli, G., Greggor, A. L., Kemp, P., Pinho, G. M., Sanzenbacher, P. M., Schulte, B. A., & Toni, P. (2021). Evaluating potential effects of solar power facilities on wildlife from an animal behavior perspective. *Conservation Science and Practice*, 3(2). <https://doi.org/10.1111/csp2.319>
- Colla, S. R., Otterstatter, M. C., Gegear, R. J., & Thomson, J. D. (2006). Plight of the bumble bee: Pathogen spillover from commercial to wild populations. *Biological Conservation*, 129(4), 461–467. <https://doi.org/10.1016/j.biocon.2005.11.013>
- Cutler, B., & Jennings, D.T. (1992). Habitat Segregation by Species of *Metaphidippus* (Araneae: Salticidae) in Minnesota. *The Journal of Arachnology*, 20(2), 88–93.
- Cutler, B., Grim, L. H., & Kulman, H. M. (1975). A Study in the Summer Phenology of Dionychious Spiders from Northern Minnesota Forests. *The Great Lakes Entomologist*, 8(3). <https://doi.org/10.22543/0090-0222.1250>
- DarkSky International. 2025a. DarkSky International | Protecting the night skies for present and future generations. Accessed June 2025. <https://darksky.org>
- DarkSky International. 2025b. Artificial Light at Night: State of the Science 2025. DOI:10.5281/zeno.

- Darst, A. L., Mitchell, T. S., Verhoeven, M. R., Evans, E., Tonsfeldt, L., Kjaer, S., & Snell-Rood, E. C. (2024). Diversity of bumble bees and butterflies in Minnesota roadsides depends on floral diversity and abundance but not floral native status. *Insect Conservation and Diversity*, 17(4), 690–701. <https://doi.org/10.1111/icad.12739>
- Edwards, C. B., Zipkin, E. F., Henry, E. H., Haddad, N. M., Forister, M. L., Burls, K. J., Campbell, S. P., Crone, E. E., Diffendorfer, J., Douglas, M. R., Drum, R. G., Fallon, C. E., Glassberg, J., Grames, E. M., Hatfield, R., Hershcovich, S., Black, S. H., Larsen, E. A., Leuenberger, W., . . . Schultz, C. B. (2025). Rapid butterfly declines across the United States during the 21st century. *Science*, 387(6738), 1090–1094. <https://doi.org/10.1126/science.adp4671>
- Ehmann, W. J. (2002). Conservation biology of special concern jumping spiders (Araneae: Salticidae) of Minnesota. Final Report submitted to the Natural Heritage and Nongame Research Program, Minnesota Department of Natural Resources. 11 pp.
- Ehmann, W. J. (2011). Reconnaissance for State-Listed Jumping Spiders (Araneae: Salticidae) in St. Louis and Lake Counties, Minnesota. Final Report submitted to the Minnesota County Biological Survey, Minnesota Department of Natural Resources. 11 pp.
- Ehmann, W. J., and B. E. Boyd. (1997). Surveys for proposed special concern jumping spiders of Minnesota. Final report submitted to the Minnesota Department of Natural Resources.
- Falchi, F., Cinzano, P., Duriscoe, D., Kyba, C. C. M., Elvidge, C. D., Baugh, K., Portnov, B. A., Rybnikova, N. A., & Furgoni, R. (2016). The new world atlas of artificial night sky brightness. *Science Advances*, 2(6). <https://doi.org/10.1126/sciadv.1600377>
- Finch, D. M., Butler, J. L., Runyon, J. B., Fetting, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weyand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Galbraith, S. M., Cane, J. H., Moldenke, A. R., & Rivers, J. W. (2019). Wild bee diversity increases with local fire severity in a fire-prone landscape. *Ecosphere*, 10(4). <https://doi.org/10.1002/ecs2.2668>
- Gandhi, K.J.K., Gilmore, D.W., Ball, G.E., Holzenthal, R.W., Katovich, S.A., et al. (2005). A review of ground beetle species (Coleoptera: Carabidae) of Minnesota, United States: new records and range extensions. *Proceedings of the Entomological Society of Washington*, 107(4), 917-940.
- Goebel, K. M., Davros, N. M., Andersen, D. E., & Rice, P. J. (2022). Tallgrass prairie wildlife exposure to spray drift from commonly used soybean insecticides in Midwestern USA. *The Science of the Total Environment*, 818, 151745. <https://doi.org/10.1016/j.scitotenv.2021.151745>
- Goulson, D. (2013). REVIEW: An overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology*, 50(4), 977–987. <https://doi.org/10.1111/1365-2664.12111>

Grimaldi, D., & Engel, M. S. (2005). *Evolution of the Insects*. Cambridge University Press.

Haarstad, J. 1990. The Acrididae of Minnesota. Final report submitted to the Minnesota Department of Natural Resources. 28 pp.

https://www.dnr.state.mn.us/eco/nongame/projects/research_reports/abstracts/insects/haarstad1990.html

He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>

Heimer, S., Nentwig, W., & Cutler, B. (1984). The spider fauna of the Itasca State Park (Minnesota, USA). *Faunistische Abhandlungen, Staatliches Museum für Tierkunde in Dresden*, 11, 119-124.

Hopwood, J., Black, S. H., Lee-Mader, E., Charlap, A., Preston, R., Mozumder, K., and Fleury, S. (2015). Literature Review: Pollinator Habitat Enhancement and Best Management Practices in Highway Rights-of-Way. Prepared for: The Federal Highway Administration by The Xerces Society for Invertebrate Conservation in collaboration with ICF International. https://rightofway.erc.uic.edu/wp-content/uploads/2018/05/1A11-pollinators_BMPs_in_highway_ROW-1.pdf

Horváth, G., Kriska, G., Malik, P., & Robertson, B. (2009). Polarized light pollution: a new kind of ecological photopollution. *Frontiers in Ecology and the Environment*, 7(6), 317–325. <https://doi.org/10.1890/080129>

Jennings, D.T. & Cutler, B. (1996). Crab Spiders (Araneae: Philodroidae, Thomisidae) of Ramsey County, Minnesota. United States Department of Agriculture, Report NC-185. 42 pp.

Kantola, T. Tracy, J.L., Baum, K. A., Quinn, M. A., and Coulson, R. N. (2019). Spatial risk assessment of eastern monarch butterfly road mortality during autumn migration within the southern corridor, *Biological Conservation*, Volume 231, <https://doi.org/10.1016/j.biocon.2019.01.008>.

Kasten, K., Stenoien, C., Caldwell, W., & Oberhauser, K. S. (2016). Can roadside habitat lead monarchs on a route to recovery? *Journal of Insect Conservation*, 20(6), 1047–1057. <https://doi.org/10.1007/s10841-016-9938-y>

Keilsohn, W., Narango, D. L., & Tallamy, D. W. (2018). Roadside habitat impacts insect traffic mortality. *Journal of Insect Conservation*, 22(2), 183–188. <https://doi.org/10.1007/s10841-018-0051-2>

Knight, S. M., Bleho, B. I., Grantham, M. A., Westwood, R., Koper, N., & Hamel, C. D. (2024). Management practices benefit endangered Poweshiek skipperling (*Oarisma poweshiek*) in Manitoba tall grass prairie. *FACETS*, 9, 1–12. <https://doi.org/10.1139/facets-2023-0049>

Knight, S. M., Norris, D. R., Derbyshire, R., & Flockhart, D. T. T. (2019). Strategic mowing of roadside milkweeds increases monarch butterfly oviposition. *Global Ecology and Conservation*, 19, e00678. <https://doi.org/10.1016/j.gecco.2019.e00678>

Koch, R. L., Rider, D. A., Tinerella, P. P., & Rich, W. A. (2014). Stink bugs (Hemiptera: Heteroptera: pentatomidae) of Minnesota: an annotated checklist and new state records. *The Great Lakes Entomologist*, 47(2). <https://doi.org/10.22543/0090-0222.2305>

- Kok, A. C. M., Berkhout, B. W., Carlson, N. V., Evans, N. P., Khan, N., Potvin, D. A., Radford, A. N., Sebire, M., Sabet, S. S., Shannon, G., & Wascher, C. a. F. (2023). How chronic anthropogenic noise can affect wildlife communities. *Frontiers in Ecology and Evolution*, 11. <https://doi.org/10.3389/fevo.2023.1130075>
- Lane, I. G., Portman, Z. M., Herron-Sweet, C. R., Petersen, J. D., Bruninga-Socolar, B., & Cariveau, D. P. (2023). Higher floral richness promotes rarer bee communities across remnant and reconstructed tallgrass prairies, though remnants contain higher abundances of a threatened bumble bee (*Bombus Latreille*). *Biological Conservation*, 279, 109862. <https://doi.org/10.1016/j.biocon.2022.109862>
- Lark, T. J., Larson, B., Schelly, I., Batish, S., & Gibbs, H. K. (2018). Accelerated conversion of native prairie to cropland in Minnesota. *Environmental Conservation*, 46(2), 155–162. <https://doi.org/10.1017/s0376892918000437>
- LeCroy, K. A., Savoy-Burke, G., Carr, D. E., Delaney, D. A., & Roulston, T. H. (2020). Decline of six native mason bee species following the arrival of an exotic congener. *Scientific Reports*, 10(1). <https://doi.org/10.1038/s41598-020-75566-9>
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution climate Projections over Minnesota for the 21st century. *Earth and Space Science*, 9(3). <https://doi.org/10.1029/2021ea001893>
- Lima, M.A.P., Cutler, G.C., Mazzeo, G. & Hrncir, M. (2022). The decline of wild bees: Causes and consequences. *Frontiers in Ecology and Evolution*, 10.
- Macy, R. W., & Shepard, H. H. (1941). *Butterflies: A Handbook of the Butterflies of the United States, Complete for the Region North of the Potomac and Ohio Rivers and East of the Dakotas*. University of Minnesota Press.
- McKenna, D. D., McKenna, K. M., Malcom, S. B., & Bebenbaum, M. R. (2001). Mortality of Lepidoptera along roadways in central Illinois. *Journal-lepidopterists society*, 55(2), 63-68.
- Minnesota Department of Natural Resources (DNR). (2022). Rare Species Guide: an online encyclopedia of Minnesota’s rare native plants and animals [web application]. Division of Ecological and Water Resources.
- Minnesota Department of Natural Resources (DNR). 2025. DNR Pollinator Best Management Practices and Habitat Restoration Guidelines. V2.
- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). (2025). Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state’s greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/Iraq-3sy25.pdf>
- Mitchell, T. S., Agnew, L., Meyer, R., Sikkink, K. L., Oberhauser, K. S., Borer, E. T., & Snell-Rood, E. C. (2020). Traffic influences nutritional quality of roadside plants for monarch caterpillars. *Science of The Total Environment*, 724, 138045. <https://doi.org/10.1016/j.scitotenv.2020.138045>

- Mitchell, T. S., Verhoeven, M. R., Darst, A. L., Patterson, C. H., & Snell-Rood, E. C. (2024). Seeding roadsides is necessary but not sufficient for restoring native floral communities. *Ecological Solutions and Evidence*, 5(3). <https://doi.org/10.1002/2688-8319.12364>
- Müller, J., Hothorn, T., Yuan, Y., Seibold, S., Mitesser, O., Rothacher, J., Freund, J., Wild, C., Wolz, M., & Menzel, A. (2024). Weather explains the decline and rise of insect biomass over 34 years. *Nature*, 628(8007), 349–354. <https://doi.org/10.1038/s41586-023-06402-z>
- Nekola, J.C. (2002). Distribution and ecology of terrestrial gastropods in northwestern Minnesota. Final report to Minnesota DNR Natural Heritage and Nongame Research Program. https://files.dnr.state.mn.us/eco/nongame/projects/consgrant_reports/2002/2002_nekola.pdf.
- Nekola, J.C., Barthel, M., Massart, P., & North, E. (1999). Terrestrial gastropod inventory of igneous outcrops in northeastern Minnesota. Final report to Minnesota DNR Natural Heritage and Nongame Research Program. https://files.dnr.state.mn.us/eco/nongame/projects/consgrant_reports/1999/1999_nekola.pdf
- Nordmeyer, C., Runquist, E., & Stapleton, S. (2021). Invasive grass negatively affects growth and survival of an imperiled butterfly. *Endangered Species Research*, 45, 301–314. <https://doi.org/10.3354/esr01136>
- Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology Evolution and Systematics*, 37(1), 637–669. <https://doi.org/10.1146/annurev.ecolsys.37.091305.110100>
- Pelton, E. (2018). Keep Monarchs Wild: Why Captive Rearing Isn't the Way to Help Monarchs. Xerces Society. <https://xerces.org/blog/keep-monarchs-wild>. Accessed 21 Feb 2025.
- Phillips, J. N., Termondt, S. E., & Francis, C. D. (2021). Long-term noise pollution affects seedling recruitment and community composition, with negative effects persisting after removal. *Proceedings of the Royal Society B Biological Sciences*, 288(1948). <https://doi.org/10.1098/rspb.2020.2906>
- Portman, Z. M., Gardner, J., Lane, I. G., Gerjets, N., Petersen, J. D., Ascher, J. S., Arduser, M., Evans, E. C., Boyd, C., Thomson, R., & Cariveau, D. P. (2023). A checklist of the bees (Hymenoptera: Apoidea) of Minnesota. *Zootaxa*, 5304(1), 1–95. <https://doi.org/10.11646/zootaxa.5304.1.1>
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. & LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. <https://mwccasc.umn.edu/node/821>
- Richman, D. B., & Cutler, B. (1978). A list of the jumping spiders (Araneae: Salticidae) of the United States and Canada. *Peckhamia*, 1(5), 82-110.
- Ries, L., Debinski, D.M., & Wieland, M. L. (2001) Conservation Value of Roadside Prairie Restoration to Butterfly Communities. *Conservation Biology* 15(2). <https://conbio.onlinelibrary.wiley.com/doi/10.1046/j.1523-1739.2001.015002401.x>

Robinson, S. I., Mikola, J., Ovaskainen, O., & O’Gorman, E. J. (2021). Temperature effects on the temporal dynamics of a subarctic invertebrate community. *Journal of Animal Ecology*, 90(5), 1217–1227.

<https://doi.org/10.1111/1365-2656.13448>

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.

<https://statesummaries.ncics.org/chapter/mn/>

Runquist, E., Nordmeyer, C., & Stapleton, S. (2024). Widespread annual occurrence of pesticides within designated critical habitats for endangered prairie butterflies. *Journal of Insect Conservation*, 28(3), 539–552.

<https://doi.org/10.1007/s10841-024-00572-5>

Samson, F., & Knopf, F. (1994). Prairie conservation in North America. *BioScience*, 44(6), 418–421.

<https://doi.org/10.2307/1312365>

Scott, J. A. (1979). Hibernial diapause of North American Papilionoidea and Hesperioidea. *Journal of Research on the Lepidoptera*, 18(3), 171–200. <https://doi.org/10.5962/p.266881>

Seibold, S., Gossner, M. M., Simons, N. K., Blüthgen, N., Müller, J., Ambarlı, D., Ammer, C., Bauhus, J., Fischer, M., Habel, J. C., Linsenmair, K. E., Nauss, T., Penone, C., Prati, D., Schall, P., Schulze, E., Vogt, J., Wöllauer, S., & Weisser, W. W. (2019). Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature*, 574(7780), 671–674. <https://doi.org/10.1038/s41586-019-1684-3>

Shephard, A. M., Agnew, L., Herdtle, A., Mitchell, T. S., Borer, E. T., & Snell-Rood, E. C. (2022). Traffic patterns, more than adjacent land use, influence element content of roadside forbs for insect pollinators. *Ecological Solutions and Evidence*, 3(4), e12195. <https://doi.org/10.1002/2688-8319.12195>

Shirey, V. M., Lewthwaite, J. M., Gawel, A. M., & Guzman, L. M. (2025). 50 years of invertebrate conservation under the United States Endangered Species Act—history and threats to species. *Frontiers in Conservation Science*, 6. <https://doi.org/10.3389/fcosc.2025.1505451>

Strazanac, J.S., Seidel, G.E., Kondo, V., Fritzler, C.J., & Butler, L. (2007). Nontarget impact of *Bacillus thuringiensis kurstaki* in central Appalachian mixed broadleaf-pine forests: long-term evaluation of arthropods. In: Gottschalk, K. W. (Ed.) *Proceedings, 17th U.S. Department of Agriculture interagency research forum on gypsy moth and other invasive species 2006*; Gen. Tech. Rep. NRS-P-10. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station: 94.

Szabo, N. D., Colla, S. R., Wagner, D. L., Gall, L. F., & Kerr, J. T. (2012). Do pathogen spillover, pesticide use, or habitat loss explain recent North American bumblebee declines? *Conservation Letters*, 5(3), 232–239.

<https://doi.org/10.1111/j.1755-263x.2012.00234.x>

Thogmartin, W. E., López-Hoffman, L., Rohweder, J., Diffendorfer, J., Drum, R., Semmens, D., Black, S., Caldwell, I., Cotter, D., Drobney, P., Jackson, L. L., Gale, M., Helmers, D., Hilburger, S., Howard, E., Oberhauser, K., Pleasants, J., Semmens, B., Taylor, O., ... Wiederholt, R. (2017). Restoring monarch butterfly habitat in the Midwestern US: ‘All hands on deck.’ *Environmental Research Letters*, 12(7), 074005.

<https://doi.org/10.1088/1748-9326/aa7637>

Underwood, E. C., & Fisher, B. L. (2006). The role of ants in conservation monitoring: If, when, and how. *Biological Conservation*, 132(2), 166–182. <https://doi.org/10.1016/j.biocon.2006.03.022>

Van Deynze, B., Swinton, S. M., Hennessy, D. A., Haddad, N. M., & Ries, L. (2024). Insecticides, more than herbicides, land use, and climate, are associated with declines in butterfly species richness and abundance in the American Midwest. *PLoS ONE*, 19(6), e0304319. <https://doi.org/10.1371/journal.pone.0304319>

Van Dyck, H., Bonte, D., Puls, R., Gotthard, K., & Maes, D. (2015). The lost generation hypothesis: could climate change drive ectotherms into a developmental trap? *Oikos*, 124(1), 54–61. <https://doi.org/10.1111/oik.02066>

Williams, P., Thorp, R., Richardson, L. & Colla, S. (2014). Bumble bees of North America: an identification guide. Princeton University Press.

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Plants and Lichens

Overview

This section provides information for Minnesota's plant and lichen species in greatest conservation need (SGCN) and species in need of information (SNI). 2,468 species of native vascular plant species (hereafter, 'plants') were assessed comprehensively for inclusion in this document using the species conservation assessment process described in the beginning of Chapter 2 (S-ranking). Non-vascular plants and lichens (1,087 species) were assessed more opportunistically and included where information was sufficient to suggest their inclusion as SGCN or SNI.

The impacts of stressors and actions to take to mitigate those stressors for plant and lichen SGCN are presented in this chapter through example species or species groups. However, it should be noted that many of the stressors for plants are also described in detail within the Habitat Sub-chapters in Chapter 3. Actions to mitigate many threats to plant and lichen SGCN are mediated by habitat and so addressed in the appropriate habitat chapters.

This sub-chapter does not present actions in the same way as was done for animals. Instead, the authors preferred to highlight information needs and several case studies illustrating the application of conservation actions benefiting plant SGCN. Since State Wildlife Grant funds are not intended for direct application to plants or lichens in the same way as they are intended for animals, it also makes sense to stop short of action planning. We believe the inclusion of plants and lichen, however, will be a good resource for ecologists and land managers to learn more about the imperiled and vulnerable plants in our state so that we can all be implementing conservation actions together.

Plants

Minnesota contains four distinct ecological biomes. This results in Minnesota having a large and variable flora, and a proportionally large number of rare plant species across all habitats found in the state. Statewide surveys for terrestrial vascular plant species have been a priority in Minnesota for over 30 years as part of the Minnesota County Biological Survey, as well as other efforts. Continued surveys, monitoring and research for all vascular and non-vascular plant species and within aquatic systems continue to be a strategic priority for the [Minnesota Biological Survey](#) (MBS) Unit with the DNR. Past efforts produced a variety of datasets and information that contributed to the conservation assessments underlying the plant SGCN and SNI lists.

Lichens

Lichen "species" are actually a species complex consisting of two or more, typically obligate, symbionts. The dominant partner is a fungus, which creates the main body of the organism. The endosymbiont is either a green alga or a cyanobacteria that lives within the fungal tissue. One fungal species can contain more than one species of endosymbiont. Identification to a single species name is based on the fungal component. Historically, the study of fungi has been included within the broadest definition of the term "botany," thus, they are included with plants here. However, fungi are an entirely distinct form of life at the Kingdom level.

No statewide survey effort for fungi has taken place in Minnesota. However, there is a history of lichenology in Minnesota (e.g., Wetmore, 1978; Brako, 1981), and there are lichen experts currently active in the state (John

Thayer and Otto Gockman; Gockman et al., 2020). Two experts assisted with a modified conservation status assessment process for this SWAP revision using the best available data. They are also working with the DNR's MBS Unit to conduct a statewide survey of lichen species for conservation purposes. The SNI designation will be helpful in guiding that effort.

SGCN and SNI Summary Information

Plant and lichen species were assigned to SGCN or SNI categories following the flowchart illustrated in Appendix A and summarized in the overview to Chapter 2 with one exception. For designation of SNI species, we also used a previously created [Watchlist of plant and fungi](#) that recognizes species that may be in greater conservation need, but for which insufficient information is available to inform conservation and protection decisions. In most cases, a species on the Watchlist was designated as SNI, unless it was already designated a SGCN by another factor in the flowchart.

Plants

Of the 2468 plant species evaluated, 448 plant species were designated as SGCN and 87 designated as SNI (Appendix B). Four SGCN are federally listed. These include the [dwarf trout lily \(*Erythronium propullans*\)](#), [Leedy's roseroot \(*Rhodiola integrifolia* ssp. *leedyi*\)](#), [prairie bush clover \(*Lespedeza leptostachya*\)](#), and [Western prairie fringed orchid \(*Platanthera praeclara*\)](#). A total of 301 are state listed, and another 73 had been previously recognized on the DNR's Watchlist of plant and fungi species. Almost 70% of plant SGCN are categorized as Critically Imperiled (S1; 131) or Imperiled (S1S2 & S2; 174). Another 27% are Vulnerable (S2S3 & S3; 124). The remaining 19 plant SGCN were recognized as at risk through their Global Conservation Status (G-rank), state listing status, or inclusion on the Watchlist despite being categorized at the state level as Apparently Secure (S3S4 & S4; 9), Possibly Extirpated (SH; 6) or Unrankable (SU; 4). One species of plant, wild rice (*Zizania palustris*), was tribally nominated and included for having cultural significance and being of conservation concern to Indian Tribes in Minnesota.

The 448 plant SGCN represent 95 phylogenetic families. Most plant SGCN are forbs (67.8%; 304). Of those, 27 are floating or submerged aquatic plant species. Graminoids represent the next largest proportion (22.5%; 101 species) with the majority in the Cyperaceae (59) and Poaceae (35) families. Of the remaining, 39 are trees or shrubs, two are epiphytes, and two are succulents.

Lichens

There are 151 lichen species designated as SGCN and 130 designated as SNI (Appendix B). Thirty-two lichen SGCN are state listed, and another 67 had been previously recognized on the DNR's Watchlist of plant and fungi species. Almost 90% of lichen SGCN are categorized as Critical Imperiled (S1; 86) or Imperiled (S2; 49). Another 5% are Vulnerable (S3; 8). The remaining eight lichen SGCN were recognized as at risk through their Global Conservation Status (G-rank), state listing status, or by expert opinion despite being categorized at the state level as Apparently Secure (S4; 1), Possibly Extirpated (SH; 6) or Unrankable (SU; 1). Most lichen SNI are Possibly Extirpated or Unrankable but due to insufficient information.

Habitat Associations

Plant SGCN are geographically distributed across all parts of the state and found in 15 different Ecological Systems (classified by the [DNR's native plant community classification system](#)). To associated plants with the 15 habitat types in this Plan, first plant SGCN were categorized by the primary Ecological System in which they are found. This was used as a representation of the habitat in which they occur (with a few species categorized in two main Ecological Systems) and crosswalked to the habitat types in this Plan. The primary Ecological System was assigned as the most common association via a review of occurrence data (Relevés or Natural Heritage Information System observations), and by expert opinion where those data were insufficient. Note, there are plant SGCN that occur in more than one Ecological System. Plant SGCN are organized in Appendix D by the SWAP habitat chapter that describes the species' primary Ecological System. A summary of their habitat distribution is shown in Table 2.12. Lichens were not evaluated for habitat associations.

Table 2.12. Habitat Associations for Plant SGCN.

Primary Habitat Type (Chapter)	Number of Plant SGCNs
Prairie and Other Grasslands	119
Savanna	8
Upland Conifer Forest and Woodland	35
Upland Deciduous Forest and Woodland	33
Mesic Hardwood Forest	52
Deciduous Wet Forest	7
Coniferous Forest Wetland	21
Riparian and Floodplain Forest	22
Non-forested Wetlands	52
Rivers and Streams	11
Lakes	36
Cliff, Talus, and Rock Outcrops	69

There are 114 plant SGCN with Upland Prairie (UP) as the primary habitat. These species are represented in either the Prairie and Other Grasslands chapter or the Savanna chapter. There are 14 with Wet Prairie (WP) as the primary habitat, represented in the Prairie and Other Grasslands chapter. Together, prairie and savanna habitats support the largest proportion of plant SGCN statewide (28.5%).

There are 52 plant SGCN with Fire-Dependent Forest (FD) as the primary habitat. These species are represented in either the Upland Deciduous Forests and Woodlands chapter or the Upland Conifer Forests and Woodlands chapter. Of the 52 total species, 16 are present in both habitats; 19 are restricted to the Upland Conifer Forest and Woodlands and 17 to the Upland Deciduous Forest and Woodlands. There are also 52 plant SGCN with Mesic Hardwood (MH) as the primary habitat, represented in the Mesic Hardwood Forest chapter. Together, these two Ecological Systems house 23% of Minnesota's plant SGCN.

There are 48 plant SGCN with Rock Outcrop (RO) as the primary habitat and 21 with Cliff/Talus (CT) as the primary habitat. All are represented in the Cliff, Talus and Rock Outcrop chapter. These unique systems house 15% of Minnesota's plant SGCN.

There are 36 plant SGCN with Lakes (LK) as the primary habitat.

There are 29 plant SGCN with Open Rich Peatland (OP) as the primary habitat, represented in the Open Wetlands chapter and 19 with Wet Meadow/Carr (MW) as the primary habitat. Another three species have Marsh (MR) as the primary habitat and one species is found in Acid Peatlands (Apn9),

There are 22 plant SGCN with Floodplain Forest (FF) as the primary habitat, represented in the Riparian and Floodplain Forest chapter.

There are 17 plant SGCN with Forested Rich Peatlands (FP) as the primary habitat and two with the Acid Peatland as the primary habitat in the Coniferous Forest Wetland chapter. Eight plant SGCN species have Wet Forest (WF) as their primary habitat; two are found in the Coniferous Forest Wetland and 7 in the Deciduous Wet Forest; one species (*Ranunculus macounii*) can be found in both.

Lastly, there are 11 species with Riverine (RV) as the primary habitat, represented in the Rivers chapter.

Information Needs for Plants

Despite decades of surveys documenting native plant species in Minnesota, rare plants and lichens often lack basic information on which to make sound conservation decisions. Thus, there are un-met information needs associated with most actions taken to conserve plant and lichen SGCN. This is especially true for efforts targeting individual species but can impact habitat mediated actions as well, as described in the habitat chapters.

One of the most basic needs can be uncertainty regarding species or variety distinctions. That can lead to uncertainty related to the appropriate unit of conservation, for example an inability to distinguish a rare species or variety from a closely related common one. For example, there are 14 SGCN and 7 SNI in the genus *Rubus* in Minnesota. While the SGCN species are well defined as units of conservation, there is general uncertainty in species identification for many members of the genus. The uncertainty has many causes, from lack of basic phylogenetic research needed to distinguish genetic units, all the way to a lack of botanical knowledge, experience, and tools for in-field identification of known, well-defined species. A variety of taxonomic research and applications are needed to address gaps in this type of information. The [Biodiversity Atlas](#), through the Bell Museum, is an excellent source of this type of botanical information where it is available.

There is also a widespread lack of information on rare plant species' population genetics. These data are needed to understand elements such as whether a population is experiencing in-breeding, or to identify past reductions in dispersal and genetic diversity due to threats like habitat fragmentation or loss of pollination. This sort of information is crucial to conservation actions involving population relocations, augmentations or re-introductions of rare species. For example, Leedy's rose root is stressed by a genetic bottleneck, and so population augmentation via 'headstarting' or re-introductions are being considered. Another example where relocating populations should be considered is Western prairie fringed orchid. Moving individuals to locations

less vulnerable to displacement during road and trail development may help preserve genetic diversity in this species. Rare plant rescue efforts led by the [Anoka Sand Plains Partnership](#) are being conducted for a handful of other species in MN. Because re-introductions, relocations or other population augmentations carry genetic risks as well as the potential for genetic benefits, these efforts should be informed by a good understanding of a species' population genetic structure and history. They also require dedicated resources for monitoring the long-term success and impacts of these actions.

Another frequent gap in information about rare plants is comprehensive assessments of species' distributions. Distribution is a key biological measure of rarity used to inform a species conservation status or need for legal protection. In Minnesota, there are several taxonomic groups that, in large part, lack the basic survey data needed to fully document species' distributions. Particularly under surveyed taxa are non-vascular plants (algae and mosses), lichens and most other types of fungi. These groups need comprehensive, taxa-wide survey efforts to:

- Improve the total list of species that occur within Minnesota,
- Document species' distributions sufficient to identify rare versus common species, and
- Gather information to improve species' conservation status assessments.

The DNR's Minnesota Biological Survey began a conservation effort focused lichen species survey in 2024 to meet those objectives. Survey efforts target habitats known to contain high lichen species diversity, an abundance of rare, SGCN, or SNI lichen species, or habitats that have a unique conservation concern related to lichen SGCN.

Even when a rare plant species' distribution has been relatively well documented, further ecological data are often needed for conversation planning. Detailed population assessments, including population size, viability and geographic extent inform decisions such as sustainable foraging limits and avoiding negative impacts during management or other activities in their habitats. Because these population attributes change over time, those data need to be regularly updated. To address this need, the MN [PlantWatch](#) program was launched in 2023. See the Case Study section for more information on MN PlantWatch.

Regular population assessments to update status, while valuable in many ways, are typically not sufficient to quantify trends in population viability. Detailed demographic monitoring is the best way to predict future trends. These studies can be done to determine viability alone or designed to assess the impact of disturbance or management actions on population viability. Trend information is one component of conservation status assessments and is useful in many aspects of conservation planning. See the Case Studies section for an example of long-term demographic monitoring for Western prairie fringed orchid.

Like monitoring a rare species population, monitoring a rare species' habitat is crucial to documenting and predicting trends in habitat attributes. Many aspects of rare plant conversation are mediated by elements of the habitat, such as habitat size, connectedness, quality, or disturbance history. See the Case Studies section for an example of long-term SGCN habitat monitoring through the Ecological Monitoring Network.

Primary Stressors for Plants

Stressors are factors that pose direct or indirect challenges to vulnerable plant and wildlife species. Habitat loss and degradation are generally considered the primary threat for most SGCN, and some stressors reduce

the quality of those habitats. Other stressors may operate more directly on the SGCN, such as disease and pathogens. In some cases, stressors may operate both directly and indirectly. For example, roads, trails, and railroads can directly harm SGCN through vehicular strikes, or operate indirectly through habitat fragmentation and reduced landscape connectivity. The stressor list below is adapted from an internationally recognized threats classification developed by the International Union for the Conservation of Nature (Salafsky et al. 2024). For more information, see the “*Stressors*” section of Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts (e.g. catastrophic wildfire following a prolonged period of fire suppression) while serving as valuable conservation tools in others (e.g., appropriately planned and applied prescribed fire).

Information about a subset of primary stressors specifically impacting this species group is included below.



Development

The development of land for housing and other urban land uses encroaches on or destroys habitat for many plant SGCN. Development pressure varies across the state and by habitat or substrate type and so actions will be addressed in habitat chapters. One area in the state under high development pressure is the Anoka Sandplains. Development in this area also makes it more difficult to achieve prehistoric fire frequencies, reducing the quality of habitat for fire-dependent plants. And, development can fragment and isolate populations of SGCN whose state-wide population core resides mostly in this area, such as [bristle-berry \(*Rubus stipulatus*\)](#).



Crop Production

The conversion of natural habitat to rowcrop agriculture, livestock operations and similar uses encroaches on, degrades or destroys habitat for many plant and lichen SGCN. These pressures vary across the state and by habitat, although they are especially impactful in grasslands. Actions will be addressed in habitat chapters.



Mining and Quarrying

Mining and quarrying impact plant and lichen SGCN primarily through direct, often complete, loss of habitat. Habitats with extensive exposed bedrock are most at risk, thus actions to this threat will be addressed in Cliff, Talus, and Rock Outcrop chapter. Generally, prevention is key to conserving SGCN threatened by mining and quarrying.

An example of a plant SGCN threatened by mining and quarrying is [wolf’s spikerush \(*Eleocharis wolfii*\)](#). The known habitats of this species in Minnesota are associated with level outcrops of [bedrock](#) in the southern part of the state ([Minnesota River Prairie](#) and [Inner Coteau](#) subsections). Specifically, it resides in the margins of shallow rainwater pools that form in depressions on the bedrock or in thin sediments over the [bedrock](#) within a [prairie or savanna](#) setting.

Arguably one of the rarest plant species in Minnesota, [ball cactus \(*Coryphantha vivipara*\)](#) occurs in crevices on granite outcrops. Historically, it also occurred in thin-soiled prairies which were largely converted to

agriculture. Today, its remaining populations are threatened by granite quarrying. Much of its potential habitat in Minnesota has already been destroyed by mining.

Wild rice (*Zizania palustris*) could be impacted by mining and quarrying development near shallow lakes.

Many lichen SGCN are also vulnerable to impacts from mining and quarrying operations given their high diversity in rock-dominated habitats. For example, [Bloody Beard Lichen \(*Usnea mutabilis*\)](#) is a Minnesota State Threatened species that occurs primarily on sandstone outcrops near cool rivers associated with a variety of other lichen and moss species. It has been documented living on wood in other parts of its global range. We know very little about why it is restricted to outcrops in Minnesota.



Roads, Trails, and Railroads

SGCN species that can persist in marginal habitats are particularly at risk from maintenance of roads, trails and railroads. [Lance-leafed violet \(*Viola lanceolata*\)](#) is a prime example of a species that is frequently found persisting in this type of altered or fragmented habitat. It is a pioneer species that does not compete well with larger, later successional species. It requires some disturbance to its habitat to be successful. In some cases, anthropogenic disturbance that removes competing vegetation and releases the seed bank can be beneficial to the species. But these activities can also degrade or destroy the habitat beyond what lance-leaf violet can tolerate.

As detailed in the Information Needs section, Western prairie fringed orchid is another example of a species highly vulnerable to road and trail development.



Gathering Plants and Fungi

Example of species threatened by gathering pressure in Minnesota are ball cactus, many species of orchids for horticultural purposes, and [American ginseng \(*Panax quinquefolius*\)](#) for its medicinal value.

American ginseng is a perennial herb native to hardwood forests in eastern North America. The roots are commercially harvested, with interstate and international trade regulated by the [Convention on International Trade in Endangered Species of Wild Fauna and Flora \(CITES\)](#), and harvest further regulated in Minnesota under [Minnesota Rules, Chapter 6282](#). It is listed as a Species of Special Concern in Minnesota. Early harvest was unsustainable across the Eastern United States, resulting in wild ginseng being quite uncommon in Minnesota today. Under the CITES treaty, the export of wild ginseng root can only be allowed if a state can demonstrate that continued export will not cause ginseng populations within that state to decline any further. At this time, Minnesota is considered in compliance. However, populations are currently so depleted and fragmented that adequate recovery is unlikely without further restrictions on harvest. The DNR hosts a [webpage](#) to help collectors assure the future of wild ginseng in Minnesota.

As the example of American ginseng illustrates, protecting plant species at risk from gathering requires a multifaceted approach. First, species-specific research and monitoring are needed to determine the level and methods for sustainable harvest. Then outreach and education on those sustainable harvesting practices will need to be ongoing. Designating species with the appropriate legal protection to stop gathering when sustainable harvesting cannot be achieved is critical. Lastly, the threat of gathering cannot be mitigated

without adequate resources toward the enforcement of either a species' legal protection status or sustainable harvest regulations.



Timber Harvest

Timber Harvest has a variety of impacts to plant and lichen SGCN. Impacts can be direct, such as removal of their microhabitat, or indirect, such as changes to the forest dynamics or structure.

Actions to mitigate those impacts will be addressed in appropriate habitat chapters. Briefly, there are several broad considerations to mitigate the impacts of timber harvest on plant and lichen SGCN. First, consider increasing SGCN legal protection through the state listing process or through state policies, such as the DNR's forthcoming Listed Species Operational Order. Consider relocating SGCNs when it is known that a population will be impacted by a planned timber harvest. Incorporate ecological values into sustainable timber harvest practices and planning, including evaluating proposed harvest rates for impacts to both the SGCN and its habitat.



Recreation

Recreational activities have a variety of impacts to plant and lichen SGCN, including fragmentation by trails, soil compaction by trampling, and degradation through more frequent introductions of non-native or invasive species. These, again, are often habitat specific and will be addressed in appropriate habitat chapters.



Fire Management

The impacts of loss or alteration of natural ecological dynamics through fire suppression, dams in streams and rivers, and loss of wild-ranging ungulates to plant and lichen SGCN are variable, widespread, and habitat specific. They will be addressed in appropriate habitat chapters. But generally, increasing the use of management to mimic natural dynamics, such as increasing the frequency of fire in fire-dependent communities, will require building awareness, knowledge, and support among private landowners. Especially when using fire, considering how to improve safety, reduce liability and direct more resources for training and organizing fire crews are also needed.

For example, many prairie plant SGCN of the great plains have their most eastern populations in Western Minnesota's dry prairies. In the past, many short statured plants like [cross-leaved milkwort \(*Polygala cruciata*\)](#) or [prairie shooting star \(*Dodecatheon meadia*\)](#) were maintained by fire and grazing in prairie systems, which would have created light openings for these plants. Other species, like [Kinnickinnick Dewberry \(*Rubus multiflorus*\)](#) or [kitten-tails \(*Syntherisma bullii*\)](#), occur in the transitional communities between prairies and forests that were also maintained in greater abundance by past fire and grazing regimes.



Invasive Species (Problematic Non-native Species)

The impacts of invasive species to plant and lichen SGCN cannot be overstated. Here also, impacts can be direct, species-to-species competition for space or resources, or indirect, such as degradation due to the release of allelopathic chemicals into a SGCN's habitat. Negative impacts are variable, widespread, and habitat-specific. They will be addressed in appropriate habitat chapters.



Problematic Native Species

Native wild rice is at risk of genetic modification due to the introduction of cultivars for commercial harvest into wild lakes, or due to the close proximity of commercial rice fields to natural wild rice beds. There are also no requirements to use wild type seed for wild rice restorations.



Diseases and Pathogens

The threat of disease and pathogens cannot be understated, as they have the capacity to cause large-scale, rapid declines to even seemingly secure species. For example, [butternut \(*Juglans cinerea*\)](#) was once common throughout the Eastern United States, including Minnesota. A lethal fungal disease, [butternut canker \(*Ophiognomonia clavigignenti-juglandacearum*\)](#), reached Minnesota in the 1970's, and as of today, nearly all butternut trees in the state are dead or dying. Active research into treatment options and the potential for population augmentations using naturally occurring resistant genotypes is on-going.

Diseases and pathogens can also have indirect impacts to plant and lichen SGCN. [Caloplaca parvula](#) is a Minnesota State Endangered lichen species endemic to the Great Lakes region. It has been documented in two counties in Northeast Minnesota and is found almost exclusively on black ash trees. This species is at risk of being lost from the state as Emerald Ash Borer continues to spread throughout Minnesota, effectively destroying this species' habitat.



Air-borne Pollution

Plant and lichen SGCN are most affected by water-borne and air-borne pollutants. For example, many lichen species are sensitive to air pollution. [Salted Shell Lichen \(*Coccocarpia palmicola*\)](#) is a Minnesota State Threatened species only documented from two counties in Minnesota. This species is thought to be particularly sensitive, with air pollution believed to be the cause of its decline throughout the Midwest.



Water-borne Pollution

Wild rice, like many aquatic plant SGCN, can be impacted by polluted runoff into lakes. Water-borne pollutants can be introduced from a variety of sources, for example from mining and quarrying and industrial effluent, and will be addressed in the appropriate habitat chapters. Broadly, increased resources for monitoring and sampling of water quality are needed to detect and correct polluting activities that threaten aquatic SGCN habitat. These data will also aid in planning, helping to, for example, prevent problematic practices in watersheds containing natural wild rice lakes.



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (MPCA & MDOC, 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow

temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al., 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

Minnesota supports disjunct populations of several species that are more commonly found elsewhere, often in arctic or alpine regions. These populations may possess unique genetic or ecological traits that enable them to persist in atypical habitats; however, they also tend to be small and geographically restricted, making them more vulnerable to decline. A notable concentration of such species occurs in Rock Outcrop and Lake Shore habitats along Lake Superior, where the lake's climate-modifying influence creates localized conditions resembling those of colder, high-elevation environments. Examples include [butterwort \(*Pinquicula vulgaris*\)](#), [northern paintbrush \(*Castilleja septentrionalis*\)](#), [small false asphodel \(*Tofieldia pusilla*\)](#), [alpine bistort \(*Bistorta vivipara*\)](#), and alpine bilberry (*Vaccinium uliginosum*). These specialized habitats and the rare species they support are increasingly threatened by rising temperatures associated with climate change.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding and landslide risk on steep slopes. The state-endangered [Eastern hemlock \(*Tsuga canadensis*\)](#) faces significant threats in Minnesota. The largest known population occurs on steep slopes within a landscape highly susceptible to landslides. Groundwater seepage has likely supported the persistence of this population, but a major rain event in June 2012 triggered slope failures throughout the area. Water flowing through sand layers confined between clay layers appears to have raised artesian pressure to a level that caused multiple landslides. As a result, four of the 17 mature hemlocks (24%) and numerous seedlings were washed away. While seedling numbers have since rebounded, most are located near the edges of landslides, leaving them vulnerable to future extreme rain events. Efforts are underway to propagate trees from local genetic stock to support population augmentation. More broadly, strategies that reduce the "flashiness" of streamflow during heavy rainfall, such as stream restoration or the installation of water control structures, can help mitigate these risks and support long-term population stability.

The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on

emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation. 

Case Study: Minnesota PlantWatch Population Assessments

 The MN PlantWatch program is a collaboration between the [University of Minnesota Landscape Arboretum Plant Conservation Program](#) and the DNR's Minnesota Biological Survey, with additional support from [The Bell Museum of Natural History](#), [The Minnesota Native Plant Society](#), and [The Minnesota Master Naturalist Program](#). As of 2025, the program is funded by the Minnesota Environment and Natural Resources Trust Fund (ENTRF) as recommended by the Legislative-Citizen Commission on Minnesota Resources (LCCMR).

MN PlantWatch is a community science initiative that conducts detailed population assessments for rare plants in their natural habitats. Volunteers collect data on population size, phenology and extent. They also provide information on the rare species' habitat type, the quality of the site and the presence of threats. These data are used to keep the State's Natural Heritage Information System up to date and are provided to landowners and administrators to support on the ground conservation efforts.

MN PlantWatch volunteer's also collect seed for conservation seed banking at the UMLA's Rare Plant Seed Bank. Seed banking is done to help preserve the genetic diversity of a species across its range, provide material for re-introductions or augmentations, and allow for research opportunities to better understand the species.

In its first two years, MN PlantWatch engaged 69 volunteers, logged over 2,500 volunteer hours, completed 141 surveys, within which 63 populations of rare plant species were re-located and assessed, and collected seed from 33 populations.

Case Study: Monitoring Populations of Federally-listed Plants

The Minnesota Department of Natural Resources has been monitoring federally listed plant species since the mid-1980s. The level and intensity of monitoring has varied for each species and is dependent on the life history and the informational needs for each species. The simplest level of monitoring has been done at two levels that has consisted of updating the NHIS database with more accurate locational information using GPS technology and a census count of the individuals found at a given location each year. This information has been critical in guiding conservation actions of the species, be it habitat management or environmental review projects.

Demographic monitoring has been directed toward the western prairie fringed orchid and prairie bush clover. This monitoring is more substantive and involves returning to marked plots on the landscape and tracking individual plants from year to year, as well as any new recruits that establish within the transect. In the case of western prairie fringed orchids, we have pursued an even more intensive phenological monitoring protocol. Under this monitoring, individuals are tracked multiple times within the same year to document their emergence, growth, capsule development and seed dehiscence. This combination of nested monitoring has led to the publication of several peer-reviewed papers as well as habitat management guidelines that assist land managers in making decisions that further enhance the conservation of these species.

Case Study: Western Prairie Fringed Orchid

The western prairie fringed orchid is found throughout the prairie region of Minnesota. Despite this wide geographic distribution, we revisit and complete a flowering census of approximately 80% of the state's populations each season. This effort is only possible because of a collaborative effort between the DNR, The Nature Conservancy, The US Fish and Wildlife Service and the National Park Service. This collaborative partnership also means that information about the species' life history and management needs are shared across the entities responsible for the conservation of this rare prairie orchid.

Capturing data on plant emergence of an herbaceous species in a grassland can be challenging. The first step requires locating plants during the growing season in the preceding season. These areas are then marked and visited in early spring. We attempted to find and measure the heights of these individuals to gauge when they emerged. After several years of data collection and anticipating the appropriate timing, we were able to identify a window of typical emergence using a growing degree day model. This understanding was critical to guide managers with the information necessary to minimize impacting this imperiled species while still carrying out needed fire management on the land, as a poorly timed fire could be fatal (Biederman et al., 2018). Additionally, by following the phenological development of individual plants, we discovered that the first plants to emerge are those that would flower in July. These plants also tend to grow more quickly, making them even more susceptible to damage and mortality with a fire too late into the growing season.

Case Study: Habitat Monitoring by the Ecological Monitoring Network

The Minnesota Department of Natural Resources established the [Ecological Monitoring Network](#) (EMN) in 2017 to track ecological change throughout the state. The network's establishment was funded by the Minnesota Environment and Natural Resources Trust Fund (ENTRF) as recommended by the Legislative-Citizen Commission on Minnesota Resources (LCCMR). The EMN provides data on the status of SGCN habitats and how those habitats are changing in a changing world. This effort is being led by the Minnesota Biological Survey, in collaboration with other DNR divisions and partners.

Before EMN, there was no comprehensive statewide monitoring network that consistently measures and evaluates status and trends in native grassland, wetland, and forest habitats. Without such information, it will be increasingly difficult for professionals and policy makers to make informed land management and land use decisions for the benefit of plant and animal SGCNs.

The EMN employs a nested plot design to sample all major habitat types across Minnesota, ensuring a statewide distribution that is independent of habitat quality or land ownership. Our data collection focuses on ground layer vegetation and trees, alongside assessments of ecological factors such as deer browse, earthworms, water chemistry, prairie structure, and coarse woody debris. Early findings have provided valuable insights into plant community variation across the state (see [EMN website](#)). The EMN will complete establishment in 2025 and begin the first resampling cycle in 2028. Resampling will directly assess how SGCN habitats have changed over the past decade.

From data collected through 2023, a total of 387 plots have been established, 45% in upland forests, 23% in open wetlands, 15% in forested wetlands, 12% in upland prairies and 5% in wetland prairies. Eighty-nine percent of plots are located on government owned land (20% Federal, 42% State and 17% local governments). A few notable findings are native plant communities in northern latitudes appear to be generally higher quality than their southern counterparts. Low quality (D-ranked) prairies are more abundant than high quality (A-ranked) prairies. Wet meadows are typically either high or low quality with very few examples in the middle. Acid peatlands are almost always A-ranked. This is likely due to the sensitivity of these systems. Any disturbance likely increases pH causing the system to rapidly transition toward a wet meadow, fen or marsh-like community.

References for Plants

Biederman, L. A., Anderson, D., Sather, N., Pearson, J., Beckman, J., & Prekker, J. (2018). Using phenological monitoring *in situ* and historical records to determine environmental triggers for emergence and anthesis in the rare orchid *Platanthera praeclara* Sheviak & Bowles. *Global Ecology and Conservation* 16, e00461.

Brako, L. (1981). The lichens of Wabasha and Winona counties, Minnesota. [Master's Thesis, University of Minnesota]. 278 pp.

Finch, D. M., Butler, J. L., Runyon, J. B., Fetting, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing.

https://doi.org/10.1007/978-3-030-45367-1_4

- Gockman, O., Thayer, J., & Henderson, B. (2020). Contributions to the Lichen Flora of Minnesota: New Records for Lichens and Lichenicolous Fungi. *Opuscula Philolichenum* 19, 14-35. <https://doi.org/10.5962/p.386187>
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). (2025). Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state's greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M., & LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwccasc.umn.edu/node/821>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>
- Wetmore, C.M. (1978). Bruce Fink's lichen collections in Minnesota. *Michigan Botanist* 17, 103-112.
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Chapter 3. Habitats

Overview

This Chapter explains the second goal of the Plan, to protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for Species in Greatest Conservation Need (SGCN). This chapter emphasizes the importance of habitat for SGCN and outlines five strategies to guide habitat conservation. We explain the context of how habitats were depicted across the three Plan revisions and our approach to creating an updated habitat spatial layer. A set of 15 detailed habitat sub-chapters, one for each habitat, follows.

Significance

Habitat conservation is arguably the most central pillar in recovering vulnerable wildlife. Habitat loss and degradation are repeatedly cited as the top reason for wildlife population declines. In the 2015 SWAP, 70% of the SGCN were classified as being impacted by habitat-related stressors as a predominant factor (DNR, 2015). This also led the 2015 SWAP to be organized with the ‘habitat approach’ as the primary strategy for the Plan, a priority that is still relevant today. This makes habitat, and its conservation, the ‘heart’ of the Plan.

Historical Loss

Habitats are lost from an accumulation of past and present land use changes such as conversions to development or agriculture.

Many of the natural ecosystems in our state have sustained great losses. A compelling source of information regarding historic changes in Minnesota’s habitats come from a comparison of present day landcover to mapped native vegetation present during the Public Land Survey (1847 to 1907) (Marschner, 1974). A whole series of maps illustrate dramatic changes, especially in southern and central Minnesota (Marschner, 1974; Wovcha et al., 1995). In the St. Croix River Valley and Anoka Sandplain, for example, less than 90,000 acres, (6% of the total area) remained in native habitat types when the survey of these two regions was completed in 1990 (Wovcha et al., 1995; Figure 3.1). In the 35 years since that survey, additional changes have likely occurred as well. Land use change in Minnesota can also be visually depicted through historical aerial photography (see [Minnesota Historical Aerial Photographs Online](#)).

For example, the historic loss of prairie throughout North America is well-recognized; indeed, tallgrass prairies are classified as critically endangered (the highest level of imperilment short of collapse), according to a recent publication by the International Union for the Conservation of Nature (IUCN) (Comer et al., 2022 2024). The prevalence of native, remnant prairies in Minnesota has dramatically declined from once covering over a third of the state (about 18 million acres; Marschner, 1974) to about 250,000 acres today, or about 1–2 percent of the historical area (DNR, 2018).




Figure 3.1. A comparison of landcover in the St. Croix – Anoka Sandplain region from the mid 1800’s to 1990 (Wovcha et al., 1995). Map A depicts pre-settlement vegetation types was compiled by FJ Marschner in 1929-30 (Marschner, 1974) using notes from the General Land Office surveys of the mid-1800’s in constructing this map. Map B depicts the remaining native vegetation, as recorded by the Minnesota County Biological Survey between 1987 and 1990.

Ongoing Stress

Remaining natural lands are pressured by ongoing economic demands as well as impacts such as pollution, invasive species, and changes in climate. These stressors are described carefully and methodically through the use of a threats lexicon (Salafsky et al., 2024). A structured lexicon better specifies the proximate causes of current-day ecological degradation and enables the targeting of conservation actions to ameliorate these threats. We call these ‘stressors’ in this Plan and outline them in relation to SGCN and habitats (see stressor section of Chapter 2. Species in Greatest Conservation Need). It should be noted that some identified stressors might also be applied for conservation purposes, such as in ‘fire management.’ We take care to explain these nuances in the context of each habitat type. Climate change is pervasive throughout all habitat types and interacts with the other stressors. We discuss changes in temperature, precipitation, and hydrology within the stressors section for each habitat sub-chapter. Further background information and strategies for climate-smart management are presented in the Climate Adaptation section of Chapter 6. Implementation.

Building on the two Prior Plans

2005-2015 Tomorrow's Habitat for the Wild and Rare

Minnesota's original State Wildlife Action Plan, *Tomorrow's Habitat for the Wild and Rare*, identified significant loss and degradation of habitat as the principal management challenge for achieving its goal to stabilize and increase SGCN populations (DNR, 2005). The principal strategy for addressing this management challenge was the need to identify key SGCN habitats and focus management efforts on them. Habitat descriptions were subsequently provided for 16 key habitats in profiles of all 25 ecological subsections in Minnesota ([Ecological Classification System](#)). Each subsection profile included a matrix of SGCN use for all habitats that occurred in the subsection along with an assessment of which habitats were the most important to the greatest number of SGCN. Conservation actions to maintain and enhance the key habitats in each subsection were delineated.

Minnesota's Wildlife Action Plan 2015-2025

Minnesota's Wildlife Action Plan 2015-2025 identified a three-pronged approach to ensuring the long-term health and viability of Minnesota's wildlife, the first and most comprehensive being the habitat approach (DNR 2015). This Plan also presented a Wildlife Action Network to represent high-quality habitats for terrestrial and aquatic SGCN as well as movement corridors among them. To prioritize the implementation of on-the-ground conservation work with our partners, sites within the Wildlife Action Network were assigned priority rankings using five metrics including SGCN population viability and SGCN richness. These rankings were used in combination with a set of Conservation Focus Areas where resources, such as partners and/or funding, were available to address conservation needs through on the ground habitat restoration and management.

Habitat Conservation

In this chapter we outline five strategies to implement the Habitat Goal of the SWAP (see below). Habitat conservation that is meaningful at a landscape scale requires partnerships between land managers and collaborations across land ownership boundaries. Here is a current breakdown of MN land uses according to the [Minnesota state land portfolio](#) | Minnesota DNR:

- 5.6 million acres (11%) are state-owned, DNR managed lands
- 3.8 million acres (7%) are federally owned lands
- 2.8 million acres (6%) are state-owned, county-administered tax-forfeited lands
- 0.7 million acres (2%) are tribal owned lands
- 0.3 million acres (1%) are county owned lands
- 25.4 million acres (45%) are in agricultural use (USDA, 2024)
- 42 million acres (75%) privately owned ([Private landowner information](#) | Minnesota DNR)

SWAP Habitat Goal: Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN.

Habitat Strategies

- 1) **Protect**, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.
- 2) **Restore**, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience
- 3) **Collaborate** with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas
- 4) **Monitor** SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to stressors, natural disturbances, conservation actions, and climatic conditions.
- 5) **Connect** to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

This strategy articulates the need to protect remaining areas of quality habitat, such as remnant prairie, old growth forest, or lakes of biological significance. In addition, areas around those core areas also need safeguarding to provide SGCN access to all habitat needs throughout their annual cycle and allow for range shifts in response to potential environmental or climate related changes, thus creating a more ecologically resilient landscape. The Conservation Action Network (CAN) is a mapped resource representing important areas to the SWAP, including core areas of high biodiversity (high numbers of SGCN), highest quality habitats, and buffers and corridors to further protect and connect them. The CAN is intended to guide and prioritize conservation actions given limited resources (see Conservation Action Network section in Chapter 6. Implementation).



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

This strategy emphasizes the need to actively manage lands, often layered upon protected lands, but also can be implemented anywhere, by any land manager or private landowner who seeks to improve the ecological function of their property. Restoration projects can be dramatic, such as a prairie reconstruction where a crop field is planted and tended into prairie over time, or they can be more subtle, such as by the dedicated application of conservation practices like prescribed fire or conservation grazing. Enhancement is done by augmenting plant communities that are already present, such as by re-seeding, planting plugs, or removing invasive species. Maintenance management is critical for many habitats, particularly those dependent on disturbance, such as fire-dependent forests, woodlands, savanna, and prairie. Invasive species management

can also be an ongoing maintenance need vitally important to the function of many of our habitats, both terrestrial and aquatic.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

This strategy calls attention to our need to work together, in broad partnerships, for ecosystem change. As detailed in the stressor sections of the various habitat sub-chapters, the extent to which our environments are impacted by human activities, both past and present, require special attention to protecting, restoring, enhancing, and maintaining the habitats we have left, and optimizing their conditions to support species in need. The implementation of habitat conservation practices with a full set of partners, including private landowners, protected area managers, working land managers, and Tribes, enables us to collectively address the scale of stressors on the landscape.

This Plan relies on actions undertaken every year by numerous entities throughout the state. The role of the DNR is to facilitate partnerships with a broad array of entities so that we are working in concert. Collective actions when coordinated across entities such as municipalities, soil and watershed conservation districts, private landowners, Tribes, other state agencies, national agencies, educational institutions, and non-profit organizations will have the most impact. Most of work we do is in partnerships.

The Conservation Action Network can be used in a planning context to focus attention on core areas of high biodiversity, the highest quality habitats, and buffers and corridors protect and connect them. You can find more about the CAN in the Conservation Action Network section of Chapter 6. Implementation.

Conservation Opportunity Areas (COAs) are places where on-the-ground conservation actions are concentrated to benefit SGCN and their habitats. COAs build on the Conservation Action Network. The COAs are prioritized areas delineated specifically for partnerships to focus efforts together on habitat conservation projects designed to benefit SGCN. For more information, please see the Conservation Opportunity Areas section in Chapter 6. Implementation.

Actions that are specifically related to developing partnerships will be tagged with a ‘handshake’ icon, displayed at the beginning of this section. However, collaboration as an approach truly should apply to all actions in the Plan, as all conservation actions would benefit by broader implementation by a greater number of entities in a more locations.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions

This strategy highlights the importance of scientific information to inform our conservation actions. We need baseline information about our natural communities to learn about changes that may be happening in the systems such as in response to changing climatic factors. Well-designed monitoring projects provide insights about the effects of conservation actions as distinguished from natural changes that are happening in the

system. Layered on this we need a sound understanding of whether the conservation actions create persistent change (effectiveness monitoring) and whether SGCN are responding positively to those changes as desired (response monitoring). Greater research into the threats that cause ecosystem change can also help to direct conservation efforts.

Actions related to this strategy are tagged with a 'magnifying glass' icon, as presented at the beginning of this section. Please refer to Chapter 5. Monitoring and Research for a comprehensive assessment of monitoring and research needs and approaches.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

The SWAP puts science into action. However, sometimes there is a deficit in knowing what to do to help particular SGCN on their road to recovery. And, as much as we know about managing healthy ecosystems, we are still learning more each year about practices that are most effective and how in turn these conservation practices may help SGCN. Indeed, there is still a great unfulfilled demand for information to guide conservation efforts by habitat managers and landowners throughout the state.

In some cases, research or investigations are needed to develop the management guidelines, avoidance measures, or habitat best management practices. More studies about the effectiveness of our conservation practices, in learning whether or how species are reacting to our management actions, are essential (see Chapter 5. Monitoring and Research).

In other cases, there may be good knowledge of conservation actions coming out of the research community or become known through applied conservation practice that requires translation into easily applied materials and resources. There is a huge opportunity to investigate, collate, and write up habitat management guidance for more SGCN.

Sometimes existing and emerging resources just need better communications to put them into the hands of more practitioners. Examples include the Landowner Guides to Woodlands, Native Plant Classification guides, and climate adaptation resources. Workshops, field trips, and engaging webinars may work well for engaging managers, with one-pagers and webpages to provide the materials. Sometimes it is a matter of spreading the word, such as when new research or threats assessments or best management practices become available. This is a parallel strategy to Strategy 3 in Chapter 2. Species in Greatest Conservation Need and overlaps also with some of the strategies and actions in Chapter 4. Public Engagement.

Actions that are specifically related to developing and disseminating habitat management practices will be tagged with a 'one-pager' icon, displayed at the beginning of this section.

Conservation Actions

Strategies are implemented by conservation actions, which are described in the habitat sub-chapters. Conservation actions are related to stressors whenever possible (see stressors section in Chapter 2. Species in Greatest Conservation Need) through a set of icons.

Habitat Definition

The term ‘habitat’ can be used in several ways and warrants clarification (Krausman and Morrison 2016; Hall et al. 1997). Specifically, habitat may be defined as, “the resources and conditions present in an area that produce occupancy, which may include survival and reproduction, by a given organism” (Krausman and Morrison 2016). In contrast to this definition that is organism-specific, ‘habitat type,’ can be defined to mean, “type of vegetation association in an area or to the potential for vegetation to reach a specified climate stage” (Krausman and Morrison 2016 based on Daubenmire 1968).

In this Plan, we use the term ‘habitat’ broadly, like the ‘habitat types’ defined above. We take this approach because with over 1,000 Species in Greatest Conservation Need, it is not practical to discuss the specific habitat requirements of each species. Indeed, by adding plants as SGCN, we emphasize the importance of biodiversity and a broad conservation approach. And, when we manage natural communities for values such as ecosystem health or function, including the support of biodiversity, we are also providing habitat for animals that live in that community.

To analyze conservation needs and actions for the different types of habitat types present in our state, we created a set of 15 habitat types (see list below). For most terrestrial types we used ecological systems to group native plant communities into habitat types such as ‘prairies and other grasslands.’ For aquatic systems we followed hydrological classifications to define types such as ‘rivers and streams.’ Other habitats were inherently more specific, such as ‘caves’ and ‘cliffs, talus, and rock outcrops.’ In another case, we created a grouping based on non-natural landcover types, classified as ‘urban and other developed lands.’

These groupings are only meant for organizational purposes and do not replace the specificity of native plant community descriptions. In addition, the full array of specific habitat requirements, particular conditions, or micro-habitats needed by particular SGCN are not easily depicted at a statewide scale or in these sub-chapters. Rather, we provide a list of animal SGCN that use the habitat in Appendix C and a list of plant SGCN that use the habitat in Appendix D and briefly highlight a subset to illustrate use of the habitats in the sub-chapters. For more detailed information about species’ habitat requirements, we recommend referencing DNR’s webpages for [Animals](#), [Plants](#), and the [Rare Species Guide](#) or other external resources cited in the SGCN species group sub-chapters in Chapter 2. Species in Greatest Conservation Need.

Primary Habitat Types

1. Prairie and Other Grasslands
2. Savanna
3. Upland Conifer Forest and Woodlands
4. Upland Deciduous Forests and Woodlands
5. Mesic Hardwood Forest
6. Deciduous Wet Forest
7. Coniferous Forest Wetland
8. Riparian and Floodplain Forest
9. Non-forested Wetlands
10. Rivers and Streams
11. Lakes

12. Vernal Pools
13. Cliff, Talus, and Rock Outcrop
14. Caves
15. Urban and Other Developed Lands

Habitat Map Methods

Overview

For the 2025-2035 SWAP we recognized the need for an updated habitat map of Minnesota that was accurate at a site-level scale and provided full coverage of the state. There was also a need to provide accurate representation of anthropogenically modified habitats such as agriculture, and developed areas as SGCN are sometimes found in these locations. There are examples of layers that cover the entire state including the Crop Data Layer and Midwest Terrestrial Habitat layer. These layers are remotely sensed and often lose accuracy as map resolution is increased. Because they cover such broad areas, it is infeasible to ground-truth every cover-type. Other layers, such as the Native Plant Communities, National Wetland Inventory, and Lake Morphology layers are more accurate but lack comprehensive coverage. Due to these inherent strengths and weaknesses, it became apparent that the best course of action was to combine several layers into one, leveraging the accuracy from some and the coverage of others. Thus, the SWAP Habitat Map (Figure 3.2) was created by combining six different GIS layers. These layers consisted of the DNR Native Plant Communities (NPC), Stream Routes with Strahler Stream Order, Shallow Lakes Identified by DNR Wildlife, Lake Basin Morphology, National Wetland Inventory for Minnesota (NWI), and Midwest Terrestrial Habitat System (MWTSH) layers.




Figure 3.2. Map depicting Minnesota’s 2025-2035 Wildlife Action Plan Habitat Layer, depicting each of the primary habitat types described by sub-chapters, except for caves which are not depicted here. Also, Agriculture is not one of the SWAP Habitats but is included in this map as a prominent landcover type.

Component descriptions

Minnesota Ecological Classification System

There are many layers to the SWAP geographic scope which covers the state of Minnesota. Within that large frame the Ecological Classification System (ECS) of Minnesota delineates 4 provinces, 13 sections, 25 subsections, and many smaller land-type associations (see Figure 3.3).

While not used directly in the making of the habitat layer, we use the [Minnesota's Ecological Classification System \(ECS\)](#) to describe the geographic area of where habitats occur and because the NPC classification system, which is used in the habitat layer, is nested within each ecological sub-section. This system was developed by the Minnesota Department of Natural Resources and the U.S. Forest Service for ecological mapping and landscape classification. This ecological land classification hierarchy is used to identify, describe, and map progressively smaller areas of land with increasingly uniform ecological features. The system uses associations of biotic and environmental factors, including climate, geology, topography, soils, hydrology, and vegetation. There are eight levels of ECS units in the United States. Map units for six of these levels occur in Minnesota: Provinces, Sections, Subsections, Land Type Associations, Land Types, and Land Type Phases.

The three levels that are relevant to this SWAP are Provinces, Sections, and Subsections. Provinces are units of land defined using major climate zones, native vegetation, and biomes such as prairies, deciduous forests, or boreal forests. There are four Provinces in Minnesota. Sections are units within Provinces that are defined by origin of glacial deposits, regional elevation, distribution of plants, and regional climate. Minnesota has 10 sections. Subsections are units within the Sections that are defined using glacial deposition processes, surface bedrock formations, local climate, topographic relief and the distribution of plants, especially trees. Minnesota has 25 subsections.




Figure 3.3. Minnesota Ecological Provinces, Sections, and Subsections.

Native Plant Communities

DNR's [Native Plant Community Classification](#) hierarchy (Table 3.1) further divides plant communities into systems, classes, and types. The DNR's classification is intended to provide a framework and common language for improving our ability to manage vegetation, survey natural areas for biodiversity conservation, identify research needs, and promote study and appreciation of native vegetation in Minnesota. Three levels of this classification system can be found in the final habitat layer: System Groups, Native Plant Community Classes, and Native Plant Community Types.

Table 3.1 Minnesota's Native Plant Community Classification Hierarchy

Classification Level	Dominant Factors	Example
System Group	Vegetation structure & hydrology	Upland Forest & Woodland Systems
Ecological System	Ecological Processes	Fire-Dependent Forest/Woodland
Floristic Region	Climate & paleohistory	Central
NPC Class	Local environmental conditions	Central Dry Pine Woodland
NPC Type	Canopy dominants, substrate, or finer environmental conditions	Jack Pine-(Yarrow) Woodland
NPC Subtype	Finer distinctions in canopy dominants, substrate, or environmental conditions	Ericaceous Shrub

System Groups, the highest level, were created to allow development of manageable field keys for lower levels of the classification. System groups were formed by combining lower levels of the classification along major physiognomic and hydrologic splits in vegetation. The system level most closely matches the Habitat Chapters within the SWAP, and the Classes and Types are then further used to describe the habitats where SGCN are found. Native Plant Community Classes are units of vegetation that generally have uniform soil texture, soil moisture, soil nutrients, topography, and disturbance regimes. For wooded vegetation, native plant community classes were developed by emphasizing understory vegetation more than canopy trees, under the hypothesis that in much of Minnesota understory plants are often more strongly tied to local habitat conditions (such as levels of nutrients and moisture) than are canopy trees. These classes have informative descriptions that can be found [online](#). Native Plant Community Types are defined by dominant canopy trees, variation in substrate, or fine-scale differences in environmental factors such as moisture or nutrients. Type distinctions were also made to describe geographic patterns within a class.

The [DNR Native Plant Communities layer](#) was desirable for a few reasons: 1) The NPC hierarchy is a system widely used across the state by both internal and external partners, and 2) the geographic extent of NPC's are highly accurate because many of them have been identified and mapped on site. Due to these strengths, it was ranked as the highest importance for the final layer; however, it does have a few caveats: 1) It only covers around 10% of the total land area of the state, 2) it only includes terrestrial or emergent wetland plant communities, and 3) it does not classify human-modified plant communities or land cover.

Stream Routes with Strahler Stream Order

To incorporate rivers and streams into the habitat layer, components from two layers were combined into one. The attributes associated with the [Stream Routes with Strahler Stream Order layer](#) were desirable because streams and rivers could be categorized into different sizes. This layer also included the Kittle number designated by DNR Fisheries Stream Survey database meaning future attributes could be added based on this unique identifier. While this layer had the desirable attributes, it could not be used alone due to it being a line feature which meant it did not capture the spatial extent of the rivers and streams. For this reason, another layer was needed that represented the shape and size of rivers and streams as closely as possible. The NWI layer meets this need as its polygons outline the shapes of rivers and streams but lacked attributes that classified them into different sizes. To merge the two layers into one, all polygons categorized as “Riverine Systems” in the Circ39_Class attribute of the NWI layer were selected and exported as their own layer. Then a 500-foot buffer was added around the Stream Route layer and then clipped using the exported Riverine layer. In this way, a new feature class was created that followed the shape of the riverine polygons from the NWI layer while maintaining the attributes of the Stream Routes. Finally, a new attribute field was created to categorize the stream order sizes. Streams with stream orders from 1-4 were classified as “small”, 4-6 as “medium-large”, and 7-8 as “large”.

Shallow Lakes

For shallow lakes, the [Shallow Lakes Identified by DNR Wildlife layer](#) was used. The layer was developed to help natural resource managers identify these bodies of water for habitat restoration and enhancement. To qualify as a shallow lake, the body of water must be 50 acres or greater in size and have a maximum depth of 15 feet. Some lakes have an “undefined” depth and may be deeper than 15 feet but were still included.

Lake Basin Morphology

To capture deeper lakes, the [Lake Basin Morphology layer](#) was used. This layer describes the physical form and morphology of lakes in Minnesota. Like the Shallow Lakes layer, it contains an attribute called “DOWLKNUM” which is a unique identifier for each lake in Minnesota. This number enables the sharing of data and attributes from different layers that also have the “DOWLKNUM” attribute field. Any lakes that were not captured by the Shallow Lakes or Lake Basin Morphology layer were captured by the National Wetland Inventory Layer.

National Wetland Inventory for Minnesota

The [NWI Layer](#) was used to capture any lakes, rivers, or wetlands in Minnesota not mapped by the previous layers. Administered by the United States Fish and Wildlife Service, the [National Wetland Inventory data](#) provides detailed information on abundance, characteristics, and distribution of American wetland and deepwater habitats, as well as changes in these habitats through time. This information is provided through two resources: the Wetland Geospatial Dataset and Wetlands Status and Trends report. Within the NWI layer, several attribute fields were used to crosswalk wetland types with habitats: the Cowardin codes, the Simplified Plant Community Classification (SPCC), and the Circular 39 fields (similar but slightly different than the SPCC; see Table 3.2). For more detail, please refer to the [Minnesota Wetland Inventory: User Guide and Summary Statistics](#) (Kloiber et al. 2019).

Table 3.2. Table cross-walking National Wetland Inventory (NWI) types to Plan habitat sub-chapters, the simplified plant community classification (SPCC) and Cowardin codes (based on table in Kloiber et al. 2019).*

Chapter	SPCC	Cowardin code
Conifer Forest Wetland	Coniferous Bog	PFO2Dq, PFO4Dq
Conifer Forest Wetland	Coniferous Wetland	PFO2D, PFO4D
Non-forested Wetland	Deep Marsh	PEM2G, L2EM2G, L2EM2H, PEM2H, R2EM2G, R2EM2H
Deciduous Wet Forest	Hardwood Wetland	PFO1A, PFO1C, PFO1D
Lakes	Non-Vegetated Aquatic Community	L1UBH, L2RSC, L2USA, L2USC, PUSA, PUSC, R2UBF, R2UBG, R2UBH, R2USA, R2USC, R3UBG, R4SBA, R4SBC, L2UBF, L2UBG, L2UBH, PUBF, PUBG, PUBH
Non-forested Wetland	Open Bog	PEM1Dq, PSS1Dq, PSS2Dq, PSS3Dq, PSS4Dq
Non-forested Wetland	Seasonally Flooded/Saturated Emergent Wetland	PEM1A, PEM1D
Non-forested Wetland	Shallow Marsh	L2EM2F, PEM1C, PEM1F, PEM2F, R2EM2F
Non-forested Wetland	Shallow Open Water Community	L1ABH, L2ABF, L2ABG, L2ABH, PABH, R2ABG, R2ABH, PABF, PABG
Non-forested Wetland	Shrub Wetland	PSS1A, PSS1C, PSS1D, PSS1F, PSS2D, PSS3D, PSS4D
Deciduous Wet Forest	Artificially Flooded Hardwood Wetland	PFO1K
Lakes	Artificially Flooded Non-Vegetated Aquatic Community	L1UBK, PUSK, L2UBK, PUBK
Non-forested Wetland	Artificially Flooded Shallow Marsh	PEM1K
Non-forested Wetland	Artificially Flooded Shallow Open Water Community	L2ABK, PABK
Non-forested Wetland	Artificially Flooded Deep Marsh	L2EM2K, PEM2K
Non-forested Wetland	Artificially Flooded Shrub Wetland	PSS1K, PSS3K

*The relationship between these two classification systems is an approximation. In addition, Riverine types (R2ABH, R2EM2F, R2EM2H, R2UBF, R2UBH, R2USA, R2USC, R3UBH, R4SBA, R4SBC) were related to the Rivers habitat.

Midwest Landscape Initiative (MLI) Midwest Terrestrial Habitat System

The Midwest Association of Fish and Wildlife Agencies (MAFWA) passed a resolution to endorse the [Midwest Landscape Initiative](#) to help meet the need for landscape-level conservation planning in the Midwest. MLI created a regional terrestrial habitat system that provided common descriptions for natural habitats within the Midwest, thus providing a common naming convention that could be used to help coordinate conservation planning across state lines. The data used to create this map was initially modelled from the [LANDFIRE](#) Consortium using artificial intelligence and remote sensing. NatureServe then assigned the natural systems to their appropriate National Vegetation Classification (NVC). The [Midwest Terrestrial Habitat System 2025 Update](#) (MWTHS) incorporates three levels of classification; groups, macrogroups, and divisions. It also crosswalks the NVC groups with each state's unique heritage vegetation classes. This layer is a large component of the SWAP Habitat layer because it fills many of the spatial gaps left by the previous layers.

Habitat Map Compilation

To merge the individual component layers, the Union tool was utilized in ArcGIS Pro. However, before the individual component layers could be combined, each layer needed to be prepared. For the Union to work, all layers needed to be vectors. Fortunately, the MWTHS was the only layer that needed to be converted from a raster to a vector. The next step was to reduce the amount of attribute fields in each layer keeping only those relevant to habitats. Taking this step reduced the overall data size of the final layer which increased processing speeds. An additional field was created for each layer prior to the Union that assigned each instance in the attribute table to one of the SWAP habitat chapters.

Once all the layers were prepared, they were merged via the Union tool and ranked based on their importance to the final layer. Ranking the layers ensured that the more accurate ones were visible when they overlapped with other layers. Once the final layer was created, another attribute field, called SWAP Habitats was created. Then, the associated SWAP Habitat chapters for each polygon were calculated for this field based on the SWAP Habitat Chapters from the component layers. This ranked Union is conceptualized in Figure 3.4.




Figure 3.4. Conceptualization of the ranking of the individual component layers that make up the final habitat layer.

Habitat Sub-chapters

Habitat sub-chapters will follow for each of the 15 habitat types, with the following sections.

General description

Here we describe general characteristics of the habitat type and any major sub-types. We reference the Minnesota Ecological Systems and Native Plant Community Classes and any other sources used to compile the habitat type in the habitat layer.

Conservation Context

This section briefly summarizes information about past and present extent and condition of habitat type.

Species in Greatest Conservation Need

We give a summary of the number of SGCN species that are associated with this habitat type and descriptions of representative SGCN from various species groups.

Primary Stressors in this Habitat

The primary stressors that may negatively impact the habitat are described. Each stressor is depicted by an 'icon' to help the reader connect content from stressors to actions.

Priority Conservation Strategies

The five conservation strategies for the Habitat Goal are shared in each habitat sub-chapter.

Conservation Actions

Conservation actions are listed under the most pertinent strategy that they implement and tagged with icons to signify the stressor(s) they address. Frequently a conservation action can address more than one stressor. In those cases, the conservation action is identified by multiple stressor icons.

Case Studies, Key Partnerships and Resources

Throughout the sub-chapters we highlight case studies and key partnerships and provide links to important resources pertinent to the habitat. Case studies feature examples of conservation projects that effectively address one or more of the stressors. These give a brief summary of the project as well as links to read additional information. The case studies and key partnerships described are not an exhaustive list, but rather a set of examples to help readers engage in conservation actions.

References

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, e14434. <https://doi.org/10.1111/cobi.14434>

Prairie and Other Grasslands

Habitat Description

Prairie ecosystems are dominated by native graminoids (grasses and grasslike plants) with a species-rich component of forbs (herbaceous plants other than grasses or sedges). Prairie native plant communities are often found on Mollisols, which have a dark colored surface horizon that is high in organic matter. They are base-rich mineral soils that span the soil moisture spectrum from dry sand-gravel prairies on coarse droughty soils, to wet prairies on poorly drained sites with a variety of soil textures. Prairies are primarily distributed in the western and southern parts of the state. This includes the Prairie Parkland Province, and in the southernmost portions of the Anoka Sandplain Subsection in the Eastern Broadleaf Forest Province.

Ecological processes including fire and grazing have shaped prairie and other grasslands of Minnesota. Frequent fire helps to reduce invasion by trees and shrubs. In tallgrass prairie, average fire intervals were thought to be 10 years or less (Zouhar, 2021). Fire is critical for maintaining the dominance of herbaceous species by preventing trees and shrubs from becoming large enough to survive fire. The burning process also removes litter and stimulates grass production. Grazing by [bison \(*Bos bison*\)](#) and [elk \(*Cervus canadensis*\)](#) was a vital ecological process on pre-European settlement prairies. Grasses, which grow from lateral meristems at the base of the plant, are well adapted to grazing, which generally removes only the upper portion of the plant. Megafauna grazing facilitated forb diversity and abundance on native prairies by suppressing grasses from dominating the plant community.

Grassland types cover a wide spectrum, ranging from prairie remnants to many types of restored and working grasslands. The following are reference descriptions of Minnesota's prairies and grasslands (for more information visit DNR's [Minnesota Prairie](#) website).

Native (Remnant) Prairie

Grasslands dominated by native prairie vegetation, typically occurring where the sod has never been broken. These grass-dominated communities host a large diversity of forbs (wildflowers) and wildlife. Prairies are often further classified by soil moisture, species composition, and geography.

Dry Prairies

Usually dominated by grasses such as big bluestem (*Andropogon gerardii*), prairie dropseed (*Sporobolus heterolepis*), and little bluestem (*Schizachyrium scoparium*). The most common forbs are in the families Asteraceae and Fabaceae. Woody species are limited to dwarf shrubs such as leadplant (*Amorpha canescens*) and prairie rose (*Rosa arkansana*). Dry prairies can vary widely in species composition based on soils and topography. This plant community is likely to be found in areas featuring conditions that preclude plants needing mesic or wet conditions. These range from wind-blown sand dunes, to sandy riverine outwash, to excessively drained gravelly ridges or slopes. Most sites containing dry prairie experience frequent and severe growing-season moisture deficits due to soil type and/or topography.

Mesic Prairies

Frequently dominated by tallgrass species such as big bluestem and indiangrass (*Sorghastrum nutans*), with prairie dropseed often a codominant or subdominant component. Forb cover is variable by soil type, with plants in the family Asteraceae and Fabaceae frequently present. Prairie rose and leadplant are the most common woody species, with patches of wolfberry (*Symphoricarpos occidentalis*) occasionally present. Various other native shrubs are rarely present and typically are not abundant. Mesic prairies cover a wide gradient of soil types and topography, with much species overlap between dry prairies and wet prairies. Changes between mesic prairie and other prairie types are often subtle, with hydrological and topographical features differentiating the cover types.

Wet Prairies

Typically feature big bluestem and prairie cordgrass (*Spartina pectinata*) as major species and support a variety of sedges (*Carex spp.*). Wet prairies include numerous forbs, dwarf shrubs (such as prairie rose), and true shrubs like red-osier dogwood (*Cornus sericea*) and willows (*Salix spp.*). Wet prairies are typically found in shallow depressional areas within the larger prairie landscape, most often on poorly drained loam soils. Wet prairies can also occasionally be found on loamy sands in areas with a shallow water table.

Bluff Prairies (goat prairies)

Characteristic on dry, steep south and west facing slopes on the bluffs in the Driftless region of southeastern Minnesota. Species composition is similar to dry prairies, consisting of species tolerant of lower quality rocky soils. These include big bluestem, little bluestem, sideoats grama (*Bouteloua curtipendula*), hairy grama (*Bouteloua hirsuta*), leadplant, hoary puccoon (*Lithospermum canescens*), pasqueflower (*Pulsatilla patens*), and prairie smoke (*Geum triflorum*). Bluff (goat) prairies require regular disturbance, and the restriction of fire and grazing on these prairies has led to the encroachment of red cedar (*Juniperus virginiana*) and other native and non-native woody species including prickly ash (*Zanthoxylum americanum*) and multiflora rose (*Rosa multiflora*).

Brush Prairies

Characterized by an abundance of taller shrubs, oak “grubs” and sprouts, and quaking aspen (*Populus tremuloides*) suckers. In brush-prairies, herbaceous prairie plants are still a major component of the vegetation, but the woody components are more prevalent than in other prairie types. In the absence of fire, both savannas and brush-prairies rapidly succeed to woodland; brush-prairie moves to woodland faster than does savanna. Today, most brush-prairies occur in the Tallgrass Aspen Parklands Province in northwestern Minnesota.

Reconstructed and Restored Prairies

Grasslands resulting from efforts to re-build prairies by planting native prairie seeds or plants (plugs) back into areas previously converted to another land use, typically agricultural fields. Prairie reconstructions seek to support biologically diverse native plants and wildlife (including SGCN). This restores ecological function and increases resistance to invasion by non-native plants, while being cost-effective to manage. In recent years, much attention has been given to data collection and monitoring of reconstructed and restored prairies to compare restoration efforts against remnant prairies (see the [Prairie Reconstruction Initiative](#)).

Reconstructed prairies are often more species-rich than conservation plantings, though less so than remnant prairies. It is challenging to produce seed for many native species found in remnant prairies, often limiting the number of species available to practitioners for prairie reconstructions. Though modern conservation plantings generally feature native grassland species, they are often tailored to programmatic goals. In contrast, prairie reconstructions are usually focused on reconstructing overall ecosystem function. Consequently, prairie reconstructions will often include more grass and forb diversity than conservation plantings and are typically planted at a 1:1 grass-forb ratio.

Conservation Plantings

These Grasslands are generally planted through conservation programs, such as the Conservation Reserve Program (CRP) administered by Farm Service Agency through the United States Department of Agriculture, funded by the federal [Farm Bill](#), the [Reinvest in Minnesota \(RIM\)](#) program administered by the State Board of Water and Soil Resources, and numerous other programs administered by a variety of state/federal agencies and NGOs.

Grasslands planted through conservation programs vary widely in their composition. Each program typically has unique specifications for seed mix requirements that lead to variations in habitat quality. For instance, a seed mix meant to control soil erosion might consist of 90% grasses, while a mix meant to attract pollinators might contain 75% wildflowers (forbs). Though nearly all new conservation plantings use native species in the seed mix, many completed in the 1980s and 1990s contained nonnative species such as smooth brome (*Bromus inermis*), alfalfa (*Medicago sativa*), and bird's-foot trefoil (*Lotus corniculatus*). Unfortunately, these species are known invaders of native prairie. Despite being nonnative and problematic adjacent to native prairies, these older plantings featuring nonnative species do still provide marginal habitat for many grassland wildlife species and can contribute to habitat complexes alongside higher quality remnant and reconstructed prairies.

Other Grasslands

These include old agricultural fields, hayfields, pastures, and roadsides, along with open space and park grasslands in developed areas. They provide various benefits, but do not have the full biodiversity and function of native prairies. Termed "surrogate grasslands" in prior MN SWAPs, they were described in the 2005 Plan as "grasslands that have developed as a result of human activities since settlement". These sites are typically dominated by non-native, cool-season grasses. They occur in areas that once supported prairie or forest communities and are found across the state but are less common in the northeast. Dominant non-native grasses include smooth brome (*Bromus inermis*), quackgrass (*Agropyron repens*), redtop (*Agrostis gigantea*), timothy (*Phleum pratense*), and Kentucky bluegrass (*Poa pratensis*). Reed canary grass (*Phalaris arundinacea*), is an increasing species that can dominate and exclude other species on wetter sites. The forb component of surrogate grasslands is also dominated by invasive species, such as yellow sweet clover (*Melilotus officinalis*), white sweet clover (*M. alba*), alfalfa (*Medicago sativa*), dandelions (*Taraxacum spp.*), bird's-foot trefoil (*Lotus corniculatus*), and canada thistle (*Cirsium arvense*). Native forbs may also occur in these grasslands, especially goldenrods, milkweeds, and asters. Left unmanaged, these grasslands are often encroached on by native and invasive trees and shrubs. Encroachment by non-native species such as Siberian elm (*Ulmus pumila*) and Russian olive (*Elaeagnus angustifolia*), and by natives such as green ash (*Fraxinus pennsylvanica*), cottonwood (*Populus deltoides*), and sumacs (*Rhus spp.*), can convert grasslands to woodlands or shrublands, reducing or eliminating their habitat value for many grassland species.

Savanna

This type of habitat hosts few trees in a grassland matrix; in Minnesota these are typically dominated by oaks but sometimes by jack pine (*Pinus banksiana*). For additional information, see the *Savanna* Sub-chapter.

Habitat Map

To depict Prairie and Other Grassland habitat (see Figure 3.5), we compiled spatial data from several sources: DNR's Native Plant Communities and the Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Upland Prairie (UP)

[UPn12 Northern Dry Prairie \(PDF\)](#)

[UPn23 Northern Mesic Prairie \(PDF\)](#)

[UPs13 Southern Dry Prairie \(PDF\)](#)

[UPs23 Southern Mesic Prairie \(PDF\)](#)

Wetland Prairie (WP)

[WPn53 Northern Wet Prairie \(PDF\)](#)

[WPs54 Southern Wet Prairie \(PDF\)](#)

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Northern Tallgrass Prairie, Eastern North American Ruderal Meadow and Shrubland, and Northern Great Plains Ruderal Exotic Grassland and Shrubland.




Figure 3.5. A map depicting Prairie and Other Grassland habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological Systems of Upland Prairie and Wetland Prairie as well as three types from the Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

Temperate prairie is a globally rare habitat. Tallgrass Prairies are Critically Endangered (the highest level of imperilment short of collapse), according to the International Union for the Conservation of Nature (IUCN) Red List of Ecosystems. The list evaluates the extent, degradation, and biotic processes of ecosystems globally (Comer et al., 2022).

The prevalence of native, remnant prairies in Minnesota has dramatically declined from once covering over a third of the state (about 18 million acres; Marschner, 1974) to about 250,000 acres today, or about 1–2 percent of the historical area. This is largely due to land conversion (DNR, 2018). Conversion of remnant prairies continues, with over 200 acres lost per year on average, and 4,248 acres documented as converted since 1987 (MBS, 2017). Thus, remaining remnant prairie is of the highest priority for protection and restoration. Indeed, many of the largest remaining tracts of remnant prairie are managed for conservation purposes. These remaining tracts provide a space for public engagement with the prairie community including its SGCNs. They are also essential for growing our understanding of the prairie community through research and adaptive management.

The scarcity of remnant prairie underscores the importance of restored prairie and other grasslands in providing vital ecosystem functions including habitat for wildlife and plant SGCN. Restoring, enhancing, and managing other grasslands complements the function of the prairie at an ecosystem level by providing connectivity for animals to move, buffering sensitive communities like wetlands and streams, and enhancing watershed, soil health, and water quality. Focusing grassland habitat restoration around building larger complexes of both remnant prairie and other grasslands will be crucial for restoring and maintaining ecosystem health into the future.

As an example, even in protected, relatively high-quality remnant prairies, most grassland bird species' population trends mirror the negative statewide trends from Minnesota transects within the North American Breeding Bird Survey data, according to DNR's long-term Sustaining Prairie in a Changing Environment (SPICE) study (see Table 3.3). This suggests that mechanisms other than local-scale habitat protection are needed to address the declining trends in grassland bird populations. Protection or management focused on reserves, while important, is not enough. For grassland birds at least, conservation efforts should be expanded into the surrounding landscape. A similar conclusion can be inferred for other wildlife, such as insect pollinators, because of threats posed by insecticides (Goebel et al., 2022) and habitat fragmentation in the surrounding landscape.

Prairies and other grasslands have also been impacted by a change in disturbance regimes including a loss of fire and grazing. Ecosystem productivity and species diversity decline when litter and thatch are not periodically removed by fire, grazing, or haying (Anderson, 1990, Knapp et al., 2009). Minnesota remnant prairies have been shown to significantly decline in native species richness in the absence of fire (Larson et al., 2020). Grasses have a positive feedback relationship with fire, which creates strong barriers to tree and shrub establishment (Ratajczak et al., 2014). Before European settlement fires were frequent, estimated to occur every 1-5 years. It is widely recommended that prescribed fire be implemented every 3-5 years to increase native plant diversity and decrease encroachment by woody plant species.

Table 3.3. Bird declines detected in Minnesota, based on Minnesota transects from the North American Breeding Bird Survey and DNR’s Sustaining Prairies in a Changing Environment (SPICE) study from 2008 through 2024.

Bird Species	Breeding Bird Survey (MN)	SPICE Survey
bobolink	Decline of 4%	Probably Stable
clay-colored sparrow	Decline of 1.4%	Increase of 0.5%
common yellowthroat	Probably Stable	Probably Stable
dickcissel	Increase of 21%	Increase of 11%
grasshopper sparrow	Decline of 4%	Decline of 7%
Henslow's sparrow	Probably Stable*	Increase of 17%
LeConte’s sparrow	Decline of 7.5%	Decline of 10%
red-winged blackbird	Decline of 1.6%	Decline of 5%
ring-necked pheasant	Probably Stable	Decline of 2.5%
savannah sparrow	Decline of 6%	Decline of 7%
sedge wren	Decline of 3%	Decline of 6%
swamp sparrow	Probably Stable	Probably Stable
upland sandpiper	Decline of 1%	Decline of 12%
western meadowlark	Decline of 3%	Decline of 2.5%

Historically, native ungulates like bison, elk, and white-tailed deer (*Odocoileus virginianus*) were widely abundant and were a significant influence on prairie composition and diversity (Knapp et al., 1999). Following European settlement, all three species were effectively extirpated from the prairie ecosystem, to be replaced by cattle, sheep, and goats. Concurrently, fire suppression began in earnest, and many grasslands were intensively grazed and even overgrazed. The absence of fire coupled with intensive, sustained grazing represented a large shock to the ecological dynamics of the prairie ecosystem.

Subsequently, changes to policy and scale in the agricultural economy resulted in many small, family farms being consolidated into larger commodity operations from the 1960s–2000s. This led to widespread reductions in livestock grazing in many areas, many of which have not been grazed since. The sudden removal of all grazing pressure following decades of intensive grazing represents another shock to the ecology of the prairie ecosystems.

In locations where fire and grazing have both been removed, remaining prairies are at risk of conversion to woody plant communities. As such, many remnant prairies have been degraded to some degree resulting from the removal of any disturbance regime.

In the bluff prairies of the driftless region of Southeastern Minnesota, native grazers including bison, elk, and white-tailed deer (*Odocoileus virginianus*) were present before European settlement and likely contributed to disturbance to an unknown degree (Hartman and Steele, 2015). As elk, bison, and most white-tailed deer were extirpated, intermittent grazing was taken over by domesticated livestock including sheep, cattle, and goats (Hartman and Steele, 2015). Grazing by livestock increased from the early to mid-1900’s and likely became

very intensive during the droughts of the 1930's as farmers struggled to find enough forage (Hartman and Steele, 2015). These habitats likely became overgrazed and became subject to varying degrees of erosion, but varying and diverse grazing systems helped to maintain the overall structure of these habitats (Hartman and Steele, 2015). From the 1960's to present, farm practices changed including the reduction of diversity of livestock and grazing of bluff prairies and associated savannas altogether.

Today, Minnesota's prairie region is dominated by cropland, though hayfields, developed areas, riparian areas, streams, rivers, wetlands, and lakes are also present. Working lands, most of which are privately owned, such as ranches, farms, and lands used for energy production, are essential for both the state's economy and natural resource stewardship. These lands produce food, fiber, fuel, and other goods, while also offering the opportunity for animal and plant conservation when managed sustainably. Multi-faceted approaches including diverse financial incentives, technical guidance, and effective landowner engagement can lead to the application of more conservation practices to benefit SGCN and their key habitats. Private landowners can benefit economically and ecologically by engaging in sustainable practices that are compatible with their management objectives, and such practices also provide additional "stackable benefits" such as carbon sequestration, soil health, and water quality.

Climate Profile

A project through the Midwest Climate Adaptation Science Center and University of Wisconsin assessed the climatic factors of importance to grasslands and how changes in those factors under future climate projections may impact the health and composition of grasslands in Minnesota. The project created maps climate exposure, of the primary factors that influence grassland plant communities using climate projections from the [Minnesota CliMAT \(Climate Mapping and Analysis Tool\)](#). In addition, it included a systematic literature review to assess how climatic stressors affect grassland and wetland habitats within Minnesota, based on empirical statistical tests extracted from manuscripts in the upper Midwest and Great Plains (Barfknecht et al., in prep.).

Climate exposure

Plant community layers were based on the Midwest Terrestrial Habitat System (MWTSH) layer that uses attributes based on the National and International Vegetation Classification (NVC and IVC, respectively) designations for all natural landcover types, and [LANDFIRE](#) classifications for all non-natural landcover types. Central Tallgrass Prairie (CTP) communities are largely within the western, central, and southeastern portion of the state, but are mainly concentrated in the southern portions of the state in the Eastern Broadleaf Forest Province (Rollins, 2009; Tracey et al., 2024; Fig. 3.6). In comparison, Northern Tallgrass Prairie (NTP) communities largely reside within the entire western, central, and southwestern portions of the state, including the Tallgrass Aspen Parklands, Prairie Parkland Provinces, and are scattered in the southwestern portions of the Laurentian Mixed Forest Province. CTP communities are characterized by deeper and richer soils compared to NTP communities.

Climate projections within CTP communities show uniform increases in drought whereas projected changes in precipitation and temperature are more geographically complex (Figure 3.6; Liess et al., 2022 & 2023). Across the southeastern portion of the state, regions that support CTP communities are predicted to experience higher drought and growing season temperature and lower spring and growing season precipitation. The

greatest warming trends are expected to occur in the western regions of CTP. In comparison, NTP communities are expected to exhibit higher drought in the southwestern portion of the state with decreased spring and growing season precipitation and increased growing season temperature in the northwest.




Figure 3.6. The extent of NTP and CTP communities, illustrated by points to highlight geographic range, do not accurately portray the area of these communities. Projections illustrated are the differences in projected climate variables based on an ensemble of five climate models for the late century period (2060-2079) compared to historical simulations (1995-2014) under the SSP-585 (high emissions) scenario ([Minnesota CliMAT](#)). The factors depicted include growing season drought (maximum consecutive days with less than 0.01 inches of precipitation), growing season maximum temperatures (°F), and spring and growing season precipitation (inches).

Literature review

The systematic literature review identified 42 studies focused on grassland systems with 912 total statistical tests (Table 3.4). Studies were reviewed to assess statistical results for plant-climate relationships and summarized using a vote counting approach to determine directionality and consensus of each relationship. Across all studies, positive results were assigned a “+1”, insignificant results were assigned a “0”, and negative results were assigned a “-1”. Results for each plant-climate relationship were summarized across all studies and then averaged.

Table 3.4. Plant-climate relationship tests (912 tests) from 42 grassland research papers based on the vote counting approach. Columns represent the respective number of tests for each plant-climate relationship category (Tests), number of studies (Studies), and the number of significantly positive (POS), insignificant (NULL), and significant negative (NEG) tests. The overall directionality of these tests (SUM) and the average vote counting (VC Mean) are presented and represent the consensus response.

Plant-climate relationship	Tests	Studies	POS	NULL	NEG	SUM	VC Mean
Diversity and drought	116	6	3	86	26	-23	-0.207
Diversity and precipitation	189	18	66	107	16	50	0.265
Diversity and temperature	63	7	6	41	16	-10	-0.159
Functional trait and drought	48	5	5	31	12	-7	-0.146
Functional trait and precipitation	49	5	16	26	7	9	0.184
Functional trait and temperature	43	2	6	36	2	1	0.116
Population and drought	15	1	7	7	1	6	0.400
Population and precipitation	192	8	38	131	4	34	0.078
Population and temperature	4	2	2	2	0	2	0.500
Productivity and drought	90	7	11	56	23	-12	-0.133
Productivity and precipitation	67	10	35	27	5	30	0.448
Productivity and temperature	36	2	8	27	1	7	0.194

Based on these findings, drought appears to reduce overall grassland plant diversity, functional traits (such stem density and total number of stems), and plant productivity (Table 3.4), but with varying responses based on species and populations due to local adaptability (specifically based on big bluestem; Galliart et al., 2019). In experimental studies of precipitation, Koerner et al. (2014) and Koerner and Collins (2014) observed lower species richness, evenness, diversity, stem density, stem number, and aboveground net primary productivity in rainfall amendment (lower precipitation) treatments established in tallgrass prairie plots at Konza Prairie, KS compared to ambient control (higher precipitation) treatments, representing a large-scale pattern across multiple grasslands. Similarly, Serafini et al. (2019) observed lower plant species richness and diversity and both total and relative productivity under rainout (lower precipitation) treatments in grasslands in southeastern Ontario compared to control treatments, CA. At Cedar Creek Natural History Area, MN, USA, Tilman and Haddi (1992) and Tilman (1996) observed that severe drought and low precipitation conditions reduce plant diversity and productivity, often driven by the loss of foundation and rare species.

Although less pronounced, there was some evidence for warmer temperatures to be positively associated with productivity and functional traits, while negatively associated with plant diversity.

In summary, an increased incidence of drought would be expected to negatively affect many grassland plant species and hinder overall grassland plant diversity, while increased precipitation will promote plant diversity, functional traits, and productivity. Increased temperatures may induce a loss of plant diversity.

Species in Greatest Conservation Need

Prairies and Other Grasslands provide habitat for 222 animal and 119 plant SGCN as primary or secondary habitat (see Table 3.5.) Primary habitats are those that species rely on and use most consistently. Loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Table 3.5. Numbers of species of SGCN associated with Prairie and Other Grasslands habitat as either primary or secondary habitat*

Species Group	Primary Habitat	Secondary Habitat	Total
Amphibians	1	0	1
Birds	22	7	29
Mammals	12	1	13
Reptiles	11	0	11
Terrestrial Invertebrates			
Bees			83
Beetles (terrestrial)			4
Butterflies			15
Flies			2
Moths			56
Snails (terrestrial)		2	2
Spiders	4	2	6
Plants	119		119
Total	169	12	341

*Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with only one or a few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants).

Prairie and other grasslands support significantly more SGCN than any other habitat. This is likely due in part to their significant reduction over the past 150 years. The 54 vertebrate SGCN represents 41% of all vertebrate SGCN; the 119 plants represent 27% of all plant SGCN; and the 168 terrestrial invertebrates represent nearly 64% of all terrestrial invertebrates. Examples of selected SGCN are described below.

In addition to implications for individual SGCN animals, some species perform keystone roles for prairie ecosystem function. For example, the digging behavior of american badgers (*Taxidea taxus*), [northern pocket gophers \(*Thomomys talpoides*\)](#), and [richardson's ground squirrels \(*Spermophilus richardsonii*\)](#) create patches of bare soil and nesting sites for insects such as some wild bees and tiger beetles. Other mammals, birds, and reptiles may use their burrows for shelter and to hunt prey. Soil disturbance, burrowing, seed eating and

dispersal, and micro-grazing by rodents all create unique micro-habitats which help to support prairie diversity. Lastly, many SGCN are important parts of the prairie food web as predators, prey, or herbivores.

Amphibians

The [great plains toad \(*Anaxyrus cognatus*\)](#) is the only amphibian SGCN that is primarily associated with open grassland habitats that provide summer forage and overwintering habitat. Remnant tallgrass prairie, wet meadows, and even agricultural fields are all suitable habitat. Shallow wetlands provide breeding habitat; even small depressions in agricultural fields that accumulate small pools of water following heavy rain storms may be used for breeding. The toads spend much of their time in underground burrows throughout the rest of the year and may estivate during hot and dry periods.

Birds

Grassland birds are declining at greater rates than any other habitat guild, with an overall 53% decline since 1970 (Rosenberg et al., 2019). The great imperilment of grassland birds is likely indicative of the status of other wildlife and plants that use prairie. Birds are very detectable and common subjects of monitoring and are widely viewed as indicators of ecosystem health and function. There is growing work focused on other species, such as butterflies, that suggests threats and population declines extend beyond birds (Edwards et al., 2025).

The decline of grassland habitat, and subsequently grassland bird populations is not unique to Minnesota. Indeed, scientists have been sounding the alarm for decades regarding steep population-level declines in grassland bird species (Brennan and Kuvlesky, 2005; Rosenberg et al., 2019; Bernath-Plaisted et al., 2023). These declines are likely the result of landscape-scale conversion of grassland habitat to cropland as well as the disruption of historic fire and grazing regimes (Knopf, 1994; Samson et al., 2004; Augustine et al., 2021; Bernath-Plaisted et al., 2023). With ~80% of the remaining grasslands in the United States being privately owned and managed (Bernath-Plaisted et al., 2023), future trends for grassland bird populations may be largely dependent upon private land conservation efforts.

Ultimately, these continent-wide population declines in grassland bird species underscore the critical need to protect remaining intact grasslands, while subsequently investing heavily in prairie restoration and reconstruction efforts, both on public and privately owned land.

Three bird SGCN, [sprague's pipit \(*Anthus spragueii*\)](#), [chestnut-collared longspur \(*Calcarius ornatus*\)](#), and [baird's sparrow \(*Centronyx bairdii*\)](#), are native prairie specialists that were common in portions of western Minnesota prior to settlement by people of European descent, but are now extremely rare. Pipits and longspurs prefer dry prairie sites with short and mid-height grasses. Typically, natural disturbance such as fire or grazing are necessary to provide suitable grassland structure for these species. Baird's sparrows are largely restricted to native prairie with structural diversity maintained by natural disturbance, but occupy areas with taller grasses than the previous two species. However, they have been found to use non-native grasslands in other parts of their breeding range (Green et al., 2020).

All other grassland bird SGCN may be found in both native and non-native (surrogate) grasslands if suitable microhabitat features are present. Species found in mesic grasslands with taller grasses include sedge wrens (*Cistothorus stellaris*), leConte's (*Ammodramus leconteii*) and [henslow's sparrows \(*Centronyx henslowii*\)](#), bobolinks (*Dolichonyx oryzivorus*), and eastern meadowlarks (*Sturnella magna*). Grasslands with a short grass

component, often maintained by grazing or fire, are preferred by upland sandpipers (*Bartramia longicauda*), western kingbirds (*Tyrannus verticalis*), and grasshopper sparrows (*Ammodramus savavvarum*). Sharp-tailed grouse (*Tympanuchus phasianellus*), [lark sparrows \(*Chondestes grammacus*\)](#), and field sparrows (*Spizella pusilla*) occupy grasslands with scattered low shrubs or small trees.

For most grassland bird SGCN, larger expanses of prairie and other grasslands are much more important than isolated grassland fragments. Reconstructed prairies and conservation plantings may provide suitable habitat for SGCN birds if they have suitable structure. However, monotypic (single species) grasslands with very dense, tall vegetation typically do not provide suitable habitat for most grassland SGCN birds.

Mammals

Prairies and other grasslands are the primary habitat for the largest number of SGCN mammals (12). The extensive loss and degradation of this habitat across western and southern Minnesota is likely responsible for the declining populations and vulnerability of so many species. Historically, one of most visible, abundant and iconic inhabitants of Minnesota's grasslands were elk (*Cervus canadensis*). Reintroduced in the 1930s, the free roaming population now consists of only 200-250 animals in northwestern Minnesota. Most of the SGCNs are small to mid-sized mammals that are limited to grassland habitats. These prairie-specialist mammals include but are not limited to the white-tailed jack rabbit (*Lepus townsendii*), American badger, [prairie vole \(*Microtus ochrogaster*\)](#), [Northern grasshopper mouse \(*Onychomys leucogaster*\)](#), [plains pocket mouse \(*Perognathus flavescens*\)](#), and Richardson's ground squirrel; each can also be found in savanna habitats.

The important ecological roles that small and mid-sized mammals play in grassland systems were summarized by Ayodele (2025)(see the DNR [Prairie Wildlife](#) webpage). They include nutrient cycling, soil disturbance, grazing, dispersal of seeds and mycorrhizal fungi, and serving as both predators and prey for other animals. SGCN mammals can be vital to populations of SGCN from other groups. For example, Richardson's ground squirrels are widely cited as being strongly associated with the population health of key SGCN predators such as the [burrowing owl \(*Athene cunicularia*\)](#) and Swainson's hawk (*Buteo swainsoni*) (Houston and Bechard, 1984; Conway, 2018), while white-tailed jackrabbit's selective grazing activities play an important role in the distribution of grassland plants while their burrows can act as seed pits or refuges (Schlater et al., 2021). Both these SGCN species have experienced population declines. White-tailed jackrabbit populations, for example have significantly declined over the past century, primarily due to eradication efforts (e.g., hunting and poisoning) aimed at reducing crop damage in the early 1900s (Brown et al., 2018) and the loss of the species' preferred habitats, including grasslands, pastures, hayfields, and small grains. Today, the species' population is only a fraction of what it was throughout its range compared to the previous century (Brown et al., 2020), and it is rarely seen in Minnesota ([DNR ARS 2024](#)). Despite this decline in the species' population, little research has been conducted on it (Schlater et al., 2021).

Other grasslands such as pastures also provide habitat for several SGCN. Most of these species are adapted to prairie but some find appropriate habitat features in these grasslands, particularly those who are seeking sparser or shorter vegetation structure. For instance, plains pocket mice require sparse grassland vegetation. Prairie voles prefer relatively dry upland prairies and pastures with a high diversity of forbs. Richardson's ground squirrels are usually found in short grass prairie or pasture where they can see over the vegetation. Some SGCN will use surrogate grasslands that contain certain soil textures for burrowing, such as plains pocket mice using exposed, sandy soils.

Reptiles

Eleven of the 13 reptile species that have been designated SGCN use prairies and other grasslands as one of their primary habitats. Other related open habitats commonly used by these species that depend on the sun for thermoregulation include savannas and cliffs, talus, and rock outcrops. Included are seven snake species, two lizards (six-lined racerunner (*Aspidoscelis sexlineatus*) and [common five-lined skink \(*Plestiodon fasciatus*\)](#)), and two turtles ([Blanding's turtle \(*Emydoidea blandingii*\)](#) and [wood turtle \(*Glyptemys insculpta*\)](#)). Among the seven snake SGCN, the [plains hog-nosed snake \(*Heterodon nasicus*\)](#) is limited almost entirely to prairies, open grasslands, and savannas. This habitat specialist prefers well-drained, loose loamy or sandy soils, sparsely-vegetated habitats and frequently uses the burrows of pocket gophers. Gophersnakes (*Pituophis catenifer*) also are closely tied with all open habitats including prairies, savannas, and old fields; sites with sandy soils are especially important (Moriarty and Hall, 2014).

Insects

It has long been known that fragmentation and destruction of grassland reduces insect species diversity and abundance (Collinge, 2000). Several prairie specialists have already been extirpated in Minnesota. These include the [arogos skipper \(*Atrytone arogos*\)](#), [poweshiek skipperling \(*Oarisma poweshiek*\)](#), [ottoe skipper \(*Hesperia ottoe*\)](#), [assiniboia skipper \(*Hesperia assiniboia*\)](#), [garita skipperling \(*Oarisma garita*\)](#), and [Uhler's arctic \(*Oeneis uhleri varuna*\)](#). Many insect SGCN are prairie specialists. These insect SGCN often require habitat features provided by native or perhaps restored prairie that are not found in other grassland habitats, such as a specific host plant or microhabitat structure limited to prairie. Insects that require a specific host plant as larvae to complete their life cycle include the [regal fritillary \(*Speyeria idalia*\)](#) and prairie violet (*Viola pedatifida*), and bird's foot violet (*Viola pedata*); [red-tailed leafhopper \(*Aflexia rubranura*\)](#) and prairie dropseed; and the [phlox moth \(*Schinia indiana*\)](#) whose host plant is prairie phlox (*Phlox pilosa*) which is typically found in remnant native prairie ([Rare Species Guide](#)).

In addition, some butterflies may require a particular microhabitat structure. For example, several skippers, including the [Dakota skipper \(*Hesperia dacotae*\)](#), require bunchgrasses characteristic of the prairie habitat as opposed to sod-forming grasses, which often characterize other grasslands. All butterflies require flowering forbs as nectar sources on which adults feed.

Though actions that benefit prairie plant communities generally benefit prairie dwelling insects as well, it is important to note that any given management action will negatively impact some insect species and positively impact others depending on location, method, and time of year. Insects may winter above or below ground as eggs, larvae, pupae, or adults. Many species are capable of flight in adult stage for only a short time and are immobile or less mobile the rest of the year. Therefore, the seasons during which insects are vulnerable to above-ground (e.g., fire, mowing, haying, grazing) or below-ground (e.g., digging, disking) management practices differ by species. As emphasized by Leone et al. (2022), prairie management should be conceptualized as a mosaic of practices and conditions across prairie landscapes and through time, rather than a "one size fits all" approach.

Plants

Prairie systems support a number of endangered, threatened, and special concern plants with a wide range of ecological retirements. Plants like [sullivant's milkweed \(*Asclepias sullivantii*\)](#) or [small white lady slipper](#)

[Cyperpedium candidum](#) require the varying hydrology of wet prairies associated with intact wetlands. Species of inland dunes like [sea-side three-awn \(Aristida tuberculosa\)](#), [blue toadflax \(Nuttallanthus canadensis\)](#), and [goat's rue \(Tephrosia virginiana\)](#) colonize sparsely vegetated areas of wind-deposited fine sand. In the Southeast, prairies support many species on the edge of their ranges such as Indian plantains (*Arnoglossum* spp.) and [wild quinine \(Parthenium integrifolium\)](#). In the Western prairies, many species of the great plains have their most eastern populations in Minnesota's dry prairies. In the past, many short statured plants like [cross-leaved milkwort \(Polygala cruciata\)](#) or [prairie shooting star \(Dodecatheon meadia\)](#) were advantaged by fire and grazing in prairie systems which would have created openings for these plants. Other species, like [kinnickinnick dewberry \(Rubus multiflorus\)](#), or [kitten-tails \(Synthesis bullii\)](#) occur in the transitional communities between prairies and forests that were maintained in greater abundance by past fire and grazing.

Due to these microhabitat requirements, most rare SGCN plant species appear to be restricted to remnant prairies. Restoration plantings of the surrounding landscape may benefit remnant species by increasing connectivity for pollinators and buffering communities containing plant SGCN from invasive species and edge effects. These include: [earleaf false foxglove \(Aralia auriculata\)](#), [roundstem false foxglove \(Aralia gattereri\)](#), [winter bentgrass \(Agrostis hyemalis\)](#), [clasping milkweed \(Asclepias amplexicaulis\)](#), [prairie milkweed \(Asclepias hirtella\)](#), [narrow leaved milkweed \(Asclepias stenophylla\)](#), [sullivant's milkweed \(Asclepias sullivantii\)](#), [slender milkvetch \(Astragalus flexuosus var. Flexuosus\)](#), [missouri milkvetch \(Astragalus missouriensis var. Missouriensis\)](#), [spike oat \(Avenula hookeri\)](#), [plains wild indigo \(Baptisia leucophaea\)](#), [prairie moonwort \(Botrychium campestre\)](#), [frenchman's bluff moonwort \(Botrychium gallicomontanum\)](#), [plains reedgrass \(Calamagrostis montanensis\)](#), [hooker's sedge \(Carex hookerana\)](#), [blunt sedge \(Carex obtusata\)](#), [hill's thistle \(Cirsium pumilum var. Hillii\)](#), [white prairie clover \(Dalea candida var. Oligophylla\)](#), and [blanketflower \(Gaillardia aristata\)](#).

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, intensive agriculture, subsistence, livelihoods, and recreation. Habitat loss or alteration remains the primary threat to most, if not all SGCN. In this section we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. Species in Greatest Conservation Need.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Pressure from urban and exurban development encroaches on grasslands, with particular pressure on sand barrens and sand prairies near large urban areas. Development accounts for 25% of documented prairie acres lost between 1987 and 2015 (MBS, 2017). In addition, the conversion of prairie habitat to development often alters hydrology, impacting adjacent prairie patches.

According to the Minnesota State Demographic Center, our state’s population is projected to grow from 5.78 million in 2024 to 6.11 million in 2075. The addition of 330,000 people over the next 50 years will add to current development pressure and will likely threaten conversion of some remaining grasslands. Specifically, MNDOT estimates that 70% of population growth over the next 20 years will be focused on the seven-county metro area, putting substantial pressure on remaining dry prairie habitats in this region.



Crop Production

Conversion of native prairie to agricultural use has drastically reduced the prevalence of native prairie in the last 150 years, and in some economic markets remains an ongoing threat. Conversion to row crops remains the largest land use conversion of documented remnant prairies in Minnesota, accounting for 62% of acres lost between 1987 and 2015 (MBS, 2017).

Agricultural practices can alter hydrology in prairie systems, by increasing runoff from heavy rain events and introducing sediment and chemical residue into streams and wetlands. Draining of fields with pattern drain tiling alters the water table depth and may increase flash flooding during heavy rain events (see also Water Management and Use).

Abundant and widespread fertilizer use has long been a feature of Minnesota’s agricultural system. However, the overuse of fertilizer, particularly nitrogen, is likely leading to further degradation of Minnesota’s remaining

prairie communities. When overapplied, nitrogen can change the bacterial composition and nitrogen availability of the soil and is not restricted to the area of application (McLeod et al., 2016). This additional nitrogen can promote and facilitate the spread of invasive species (Morghan and Seastedt, 1999; McLeod et al., 2016), with reed canary grass (Rickey and Anderson, 2004) and smooth brome (Clark, 2008) being notable issues in Minnesota.

Pesticide use from agricultural practices have been identified as a leading factor in the declines of grassland dependent species (Mineau et al., 2005; Gibbs et al., 2009; Gibbons et al., 2015; Forister et al., 2016; Sánchez-Bayo and Wyckhuys, 2019). Here in Minnesota, even in protected prairie remnants, extirpation of prairie-dependent invertebrates coincides with increases in applications of agricultural insecticides. Neonicotinoids (systemic insecticides often applied as seed coats to corn, soybeans, and other crops) have become one of the most widely applied class of insecticides and have been widely posited as a primary driver of insect declines and presumably other wildlife (Mason et al., 2013; Goulson, 2013; Gibbons et al., 2015; Pecenka and Lundgren, 2015; Gilburn et al., 2015; Forister et al., 2016; Basley and Goulson, 2018; Olaya-Arenas and Kaplan, 2019; Li et al., 2020; Roy and Chen, 2023).

Conversely, broad-spectrum insecticides such as organophosphates (especially chlorpyrifos) and pyrethroids are contact insecticides that are often applied via aerial spraying. Often targeting soybean aphid and other crop pests, they are known to be easily transported into non-target prairie preserves from neighboring and even distant crop fields (Foreman et al., 2000; Mackay et al., 2014). Two of these, chlorpyrifos (organophosphate) and bifenthrin (pyrethroid), have been frequently documented far within prairies designated as Critical Habitat for listed butterflies including Dakota skipper, and whose applications are at least correlated with their declines (Runquist et al., 2024).

Mechanical harvest through haying can be a very effective method to enhance grassland stands by removing weeds, stimulating root development in grasses, and reducing the duff layer to allow native grass species to increase in density. However, haying can have negative impacts on ground nesting birds and other wildlife species through nest destruction and habitat loss. Haying should be planned during times that will minimize impacts to ground-nesting birds. The nesting season is generally considered to be between April 15th and August 1st. As a result, haying should be conducted in August or September. To avoid significant impacts to wildlife, it is recommended that no more than fifty percent of a field be hayed in any given year to maintain adequate cover for wildlife habitat. Flushing bars and mowing patterns that drive wildlife to the outer edges of the field should also be used.



Livestock Management

Livestock operation management can either benefit or degrade a grassland ecosystem and directly influences the plants and wildlife that utilize the associated habitat. Properly managed grazing can stimulate plant diversity and create diverse vegetative structure across the landscape. Overgrazing, if left unchecked, can reduce diversity in vegetative cover and structure, and have severe negative impacts on biodiversity, ecosystem function, soil health, and forage production.

Over the last 30 years, advances in range science, grazing management practices, technology, and increased education and outreach efforts within the agricultural community have allowed livestock producers to manage their herds to meet economic goals while preserving healthy grasslands for future generations. However,

overgrazing is still widespread in Minnesota and, coupled with increased cyclical drought and changing weather patterns, the negative ecological impacts from overgrazing can be significantly increased.

Overgrazing favors plants that are grazing increasers (species tolerant to intensive grazing), which eventually take over and are often undesirable, non-native, or even invasive. Under heavy continuous grazing, plant diversity can decrease by reducing or eliminating plant species intolerant to intensive grazing such as big bluestem (*Andropogon gerardii*), porcupine grass (*Hesperostipa spartea*), little bluestem (*Schizachyrium scoparium*), Indiangrass (*Sorghastrum nutans*), prairie dropseed (*Sporobolus heterolepis*), prairie onion (*Allium stellatum*), leadplant (*Amorpha canescens*), prairie clovers (*Dalea spp.*), narrow-leaved purple coneflower (*Echinacea angustifolia*), stiff sunflower (*Helianthus pauciflorus*), and blazing stars (*Liatris spp.*).

Low diversity grasslands also limit native species of pollinators that have evolved alongside native plants. In addition, overgrazing reduces grass and forb heights and can increase bare ground on many sites. Along with increasing the risk of erosion, these conditions are generally not as favorable for species adapted to the tallgrass prairie such as bobolinks and Henslow's sparrow, preferring mid-height to tall vegetation with 1-3 years of thatch.

Care must be taken when treating livestock for pests and parasites, as treatments often contain insecticides. Some insecticides persist in the environment weeks after treatment and can harm beneficial insects and contaminate soil and water via the animals' feces.



Mining and Quarrying

Sand and gravel mining in particular affects dry prairies, especially sand prairies, by converting habitat in order to extract the sub-surface aggregate material. Mining accounted for 12% of conversions of documented remnant prairie from 1987 to 2015 (MBS 2017), and conversions from prairie to mining appear to be increasing over time.

The construction of roads, trails, and railroads requires aggregate material. In areas with remnant dry prairie, the substrate beneath the dry prairie is often targeted for extraction to build these lines of infrastructure. It is impossible to repair these dry prairies once they have been mined for aggregate.

Dry prairies in closer proximity to highly developed areas and/or major roadways are at serious risk for sand and gravel mining. Developers and road authorities typically try to source aggregate material near the project site to reduce costs, putting prairies near emerging projects at high risk. As developmental pressures increase, the risk of these prairies being lost will increase substantially.



Wind and Solar Energy Infrastructure

According to [Our Minnesota Climate](#) "The amount of energy our state produces from renewable resources like solar power and wind has risen 60% over the past decade. Today, renewable energy accounts for 28% of Minnesota's electricity generation, with 52% of our energy coming from carbon-free sources like renewables, nuclear, and hydropower." These successes in renewable energy production can be complex in the context of wildlife and habitat impacts. Direct impacts to birds and bats are a particular concern, especially during migration, with wind turbine collisions documented as a high source of mortality for both groups. Loss and his colleagues (2013b) estimated between 140,000 and 328,000 birds are killed annually at wind turbines

across the contiguous United States; mortality increased with the height of the turbine hub. Foliage and tree-roosting bats, such as hoary bats and eastern red bats, also are frequently found dead below wind turbines (Arnett et al. 2008). Although variable in timing, most mortality occurs during late summer and fall, corresponding with migration and mating seasons. Estimates place the loss at approximately 888,000 bats per year (Smallwood 2013). To date, lighting and noise appear to not influence the level of mortality.

Similarly land-use impacts in and around solar arrays and wind turbines can negatively impact native plant communities (predominantly prairie) and rare features. Taking lessons from traditional power infrastructure and development, we can be mindful to wildlife considerations in the siting and design of emerging alternative energy resources.



Roads, Trails, and Railroads

Railroads, roads, and trails, while essential, fragment grassland habitats. They can also alter hydrology, increase sedimentation and runoff, and act as a vector for the introduction of invasive species. The associated impacts from initial construction as well as routine maintenance and infrastructure installation can have detrimental impacts to habitat within the right of way for the life of the road, trail or railroad.



Utility Corridors

Utility corridors transect prairie and other grasslands throughout the state. These corridors can contain a variety of utility infrastructure. This includes pipelines, telephone lines, and power lines. These lines can be aerial, above ground, or below ground. Regardless of type, utility corridors can fragment habitat based on their size, purpose, and the upkeep that is required to maintain them.

The initial construction of the associated utility infrastructure as well as the long-term maintenance of that infrastructure can have lasting impacts on native plant communities and the SGCN associated with them. Increased funding for rural broadband, conversion of aboveground powerlines to belowground, and development of new utilities (solar, wind, etc.) continue to threaten grassland habitats.

These corridors and the increased travel from vehicles and equipment that they require act as vectors for invasive plant species. The required vegetative management practices often include herbicide applications, removal of woody vegetation, and/or mowing. This increased disturbance increases the likelihood of establishing invasive plant species within the corridors and then spreading to adjacent lands.



Hunting and Collecting Animals

Small mammals can be impacted by deliberate trapping or shooting. For example, while Northern pocket gophers are rare, the Plains pocket gopher is very abundant and is unprotected, meaning that hunting and trapping is not limited. Several other SGCN, Richardson's ground squirrels and Franklin's ground squirrels and American badgers are still to this day considered pests. The burrowing nature of these species is concerning to some agricultural producers and may result in increased control measures.



Gathering Plants and Fungi

Harvesting and foraging rules in Minnesota vary depending on the type of land. Minnesota's state parks, state recreation areas, state waysides, state forest recreation areas, and state trails have restrictions on taking plants, mushrooms, and berries. Harvesting can cause numerous impacts, which can negatively impact natural resources and other visitors' experiences, such as: disturbing soils and trampling other plants, which can lead to invasive species taking hold; removing food sources otherwise available for wildlife; reducing the natural reproduction of some species; and preventing other people from having the same experience.

A large threat to Minnesota grasslands is the overcollection and, in some cases, illegal collection of native plants for sale in seed production. The rarest species, with the smallest amount of seed produced, and those with the highest market value are the most vulnerable from overcollection. One of the most popular plants for collection is narrow-leaved coneflower (*Echinacea angustifolia*).

Another type of plant harvesting is for the use of medicinal, herbal, or food purposes. Species that are most vulnerable to overharvesting include species with edible tubers such as prairie turnip (*Pedimelum esculentum*) and wild/prairie onion (*Allium stellatum*). Natural areas close to large population centers are most vulnerable to over harvesting and exacerbated by trends of foraging on social media.



Timber Harvest

Timber harvest in the Tallgrass aspen parkland region may alter patch dynamics between aspen and prairie stands by increasing root suckering and clonality of aspen.



Recreation

Minnesotans are known for their love of the outdoors. However, recreational overuse can threaten sensitive grassland plants and grassland communities. Though heavy foot traffic would likely be required to alter a plant community, popular hiking areas that see intense use may reach a threshold where certain rare plants are trampled, and the level of foot traffic is high enough to degrade the plant community. Foot traffic also introduces the risk of spreading invasive species. The greater recreational use in a given area, the higher the likelihood of invasive species introduction on the site.



Fire Management

Fire is a necessary and natural feature of Minnesota's native grasslands. Fire-adapted ecosystems, which includes all of Minnesota's grassland habitats, have evolved to thrive with fire disturbance. Sediment cores from the Northern Great Plains indicate that grasslands, drought, and fire have been a key feature of this landscape for much of the past 10,000 years (Nelson and Hu, 2008; Grimm et al., 2011). If litter or thatch are not periodically removed by fire, grazing, or haying, ecosystem productivity and species diversity decline (Anderson, 1990; Knapp et al., 2009).

In the modern era, landscape conversion and development has led to most remaining grasslands of all types being highly fragmented, scattered, and (in most cases) small in size. Though fire is necessary to maintain grassland habitat, the current nature of small, isolated grasslands means that, without careful planning, many

SGCN populations could be critically reduced and even extirpated by fire. (see Chapters SGCN Amphibians, SGCN Reptiles, SGCN Terrestrial Invertebrates).

Lack of fire is a stressor for many prairie plant communities, including dry and mesic tallgrass prairie, bluff prairies, brush-prairies, and savannas. Fire suppression leads to woody species encroachment, such as eastern red cedar (*Juniperus virginiana*) and/or Siberian elm (*Ulmus pumila*); for additional information, see Problematic Native Species and Invasive Species, respectively. If left unchecked, these woody species can result in wholesale ecosystem conversion (DeSantis et al., 2011).



Invasive Species (Problematic Non-native Species)

Invasive plant species can have negative impacts to native plant communities in prairie and grassland ecosystems. While both native and non-native plant species can possess invasive characteristics and be problematic in prairies and grasslands, non-native invasive plant species are of particular concern. Non-native plants did not evolve within the ecosystem that they now inhabit and were likely imported for their ability to establish easily and spread quickly. As a result, they are able to outcompete native plant species for available resources while decreasing biodiversity and degrading already limited habitat for SGCN.

A prevalent example of this is smooth brome (*Bromus inermis*). Smooth brome is a perennial cool season grass native to Europe and Asia. It has become quite dominant in prairies across much of the state. In remnant prairies, smooth brome has been shown to reduce native plant richness (Jones et al., 2023; Otfinowski et al., 2007), reduce native cover (Fink and Wilson, 2011), and alter soil dynamics (Vinton, 2006; Bell et al., 2023).

Like many non-native cool season grasses, smooth brome is difficult to manage once it is established. Work is ongoing to develop effective management strategies to reduce smooth brome in grassland habitats. Anecdotal evidence to date suggests that late fall (post-hard frost) herbicide applications and late spring burns can help to suppress smooth brome and allow native prairie species room to compete. Treatments will likely need to be repeated through time. Careful consideration should be exercised in using these techniques, particularly on remnant prairies.

While already impacting many remnant prairies, smooth brome is predicted to become more problematic in prairie ecosystems under increased nitrogen deposition (Clark, 2008), increased atmospheric carbon dioxide concentrations, and increased spring rainfall (Prevey, 2014; Printz and Hendrickson, 2015). Understanding effective management practices for reducing smooth brome in tallgrass remnant prairies is increasingly important. Likewise, Kentucky bluegrass (*Poa pratensis*) is another invasive cool-season grass that causes degradation similar to smooth brome (Gannon, 2024).

From 1960 to present, invasive woody species including common buckthorn (*Rhamnus cathartica*) and bush honeysuckle (*Lonicera spp.*) have proliferated and continued to invade natural areas including bluff prairies and adjacent savanna (Hartman and Steele, 2015). Invasive forb species such as sweetclover (*Melilotus officinalis*) are also common invasive species in these habitat types (Hartman and Steele, 2015). As buckthorn and honeysuckles continue to spread, all prairie habitat types are now being impacted by these woody invaders. These species are an impending challenge for prairie land managers, particularly in areas that are difficult to burn.



Problematic Native Species

Native plant species can possess characteristics associated with invasive species (non-native problematic species), such as encroaching into prairie habitat. These plant species are often referred to as pioneer or volunteer species. These characteristics can make them problematic under certain conditions. This is especially true in the absence of disturbance such as fire, thinning, or grazing. Historically disturbance was a limiting factor for many of these problematic native species. Frequent disturbance through fire limited the spread of these species by killing them before maturity and reproductive age.

An example of this is eastern red cedar (*Juniperus virginiana*). Its seeds are readily consumed and transported by birds and small mammals. Young red cedar trees will not tolerate fire and are killed. In the absence of fire this native tree species can encroach into prairie and grassland ecosystems and grow to maturity. Once established mature cedar trees develop an extensive fibrous root system that can extend laterally up to 20 ft from the root crown. Mature trees are able to take advantage of moisture and nutrients in the soil profile and eventually shade out many if not all of the native prairie vegetation at the ground level. This prevents grasses and forbs from growing thus eliminating the fine fuels needed to carry fire and kill the trees. This self-reinforcing feedback loop allows cedar to continue to spread once firmly established.

In prairie and grassland ecosystems these eastern red cedar groves eliminate native perennial grasses and forbs, fragment habitat, provide hunting perches for raptors that prey on ground nesting birds, and generally degrade the biodiversity in the area they inhabit.

Another increasing species is reed canary grass, previously classified as an invasive species but more recently thought to be native in at least some of Minnesota based upon research at the University of Minnesota. Reed canary grass is particularly damaging in wet prairies. The thick sod layer and dense growth can rapidly outcompete desirable native species, and it is common to see former wet prairies transition to a near monoculture of reed canary grass in areas where this species is present. To date, land managers have been unable to find an effective management strategy to reduce or eliminate reed canary grass.



Air-borne Pollution

Experiments under controlled greenhouse conditions have shown that increased levels of atmospheric carbon dioxide may favor the competitiveness of woody plants and cool season invasives by stimulating photosynthesis and growth. It may also reduce mild to moderate abiotic stress in woody plants if root growth and nutrients are not limited.

Nitrogen levels in the atmosphere are driven by natural processes as well as transportation and agricultural sources, primarily fertilizer (Burke et al., 2002; Wall and Pearson, 2013). A study indicates that increased atmospheric deposition of nitrogen reduced plant species numbers by 17% by increasing soil acidity and increasing above ground production of certain plants that reduces light availability to other plant species. This can lead to competitive exclusion. This suggests that chronic but low-level nitrogen deposition may have a greater impact on diversity than previously thought (Clark, Tilman, 2008).



Light and Noise Pollution

Nearly half of the United States land surface has light-polluted skies (Falchi et al. 2016). Sources of light pollution at night include interior and exterior lights on homes, buildings, billboards, parking lots, and streets as well as lighted sports fields, industrial plants, and airports (DarkSky International 2025a). Studies also point to the ecological impacts created during the daytime by polarized light that reflects off surfaces such as asphalt roads, glass panes, and solar panels (Chock et al. 2020; Horvath et al. 2009). Because most of the nation's large urban areas (with multiple artificial light sources every night), are embedded in the Temperate Deciduous Forest biome, biota occupying temperate forests are the most vulnerable to these impacts (Aubrecht et al. 2010).

Impacts of light pollution include disrupting the behavior of amphibians whose breeding calls are delivered at night, drawing nocturnal insects to artificial lights that increase their predation risk, and disorienting birds that use environmental cues to guide their nighttime migration ([DarkSky International 2025a, 2025b](#)). In addition, the phenology, growth rates and biomass of plants are impacted (Bucher et al. 2023). Bright day and dark night cycles synchronize the internal clocks of plants and animals, and alterations to these cycles can impact hormone production, reproductive behavior, neural activity, and metabolic functions (Bumgarner and Nelson 2021).

Noise pollution from aircraft, vehicles, boats, and other human causes is pervasive. Anthropogenic noise was detected in 36% of national parks surveyed (Buxton et al. 2019), and 12% of wilderness areas had anthropogenic noise levels 3 decibels higher than levels predicted to occur naturally (Buxton et al. 2017). Exposure to increasing noise levels can change the spatial distribution and movement patterns of wildlife, cause avoidance of feeding and nesting areas, and interrupt sleeping patterns and communications (Kok et al. 2021). For plants, noise can reduce seedling recruitment (Phillips et al. 2021).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). According to the Minnesota Pollution Control Agency and the Minnesota Department of Commerce these changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years. Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al., 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity,

compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; Runkle, J., et al (2022)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan (Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN), five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Examples of conservation actions are grouped below under these five strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Also note that some strategies, such as Strategy 3, collaborating with partners, could truly be applied to all actions to most broadly and effectively implement them.

Priority Conservation Actions for Prairie and Other Grasslands



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Protect remaining remnant prairie from conversion to other land uses.
	To protect prairie remnants from pressure for mining resources, increase funding for acquisitions and payments through programs such as Prairie Bank.
	Prevent further fragmentation of grassland habitat. Support diverse financial incentives to help landowners avoid conversion of grasslands into row crops where remnant prairie and/or SGCN occur. Implement landowner outreach and engagement activities. Prioritize access for landowners to a variety of programs for implementation of land protection options such as conservation easements. Support conservation programs that provide

	economic benefits for set aside and conservation programs and assist in public outreach. Improve access to USDA Farm Bill and other wildlife resources for landowners.
	Control or prevent the invasion of grasslands by trees and shrubs with mowing and cutting woody vegetation, prescribed fire, grazing, and careful use of herbicides with consideration for SGCN.
	Control spread of invasive species on native-dominated sites. Ensure equipment is cleaned prior to moving mowers, haying equipment, and seed harvesters among sites to prevent spreading invasive species.
	Livestock treated with systemic insecticides for pest or parasite control (such as ivermectin, doramectin, or eprinomectin) should not be released on conservation lands for at least 30 days following treatment. If parasite treatment is needed during the grazing period, the best practice is to remove affected animals for treatment and return them to the site after the recommended amount of time.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

Stressor	Action
	Promote and participate in prairie reconstruction and restoration to expand the amount of prairie and provide a diversity of native plants to sustain biodiversity and provide ecosystem benefits including healthy soil, carbon storage, improved hydrology, watershed health, and cleaner water. Projects that restore more natural hydrology, such as breaking up drain tile, can also benefit streams and downstream habitats. Planning projects to be resilient under anticipated changes in climate regimes will be the most successful (see also Climate section).
	To protect nesting SGCN birds in hayfields and mowed grasslands, consider adopting these bird-friendly measures: use flushing bars, lengthen cutting rotations, and avoid early-season mowing and haying
	Where people recreate, provide outreach and education on cleaning gear and equipment to reduce the introduction and spread of invasive species. DNR's Prevent the spread webpage includes specific actions for different land-based activities like biking, hiking, and off-highway vehicle riding. At trailheads, add invasive species prevention messages, such as those on the boot brush kiosks developed by the PlayCleanGo: Stop Invasive Species in Your Tracks program. Consider maintaining buffers around and minimizing disturbance to bluff prairies when designing and maintaining trails.

	Avoid soil compaction in areas occupied by mammal and reptile SGCN.
	Incentivize conservation grazing practices that benefit grassland and SGCN habitat. Promote the use of rotational grazing, adaptive grazing, and other regenerative practices to livestock producers. Promote grazing as a habitat management tool to build higher quality habitat, not just cattle production. Support soil health practices to improve soil health (structure, water retention, etc.), prevent erosion, and lower inputs (reseeding, fertilizer, etc.). Increase landowner knowledge of the methods and benefits of various regenerative grazing practices and provide diverse funding opportunities to transition to sustainable grazing and incentivize better habitat management on private lands. Specific practices will depend on the targeted SGCN, site conditions, landscape context, and other objectives.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

Stressor	Action
	Increase incentives and planning, and potentially ordinances, to include higher density housing, such as multi-dwelling housing, and create zones of no net growth in areas where grasslands are pressured by urban and exurban development. Awareness, education, economic incentives, community support, and working with local governments can all be part of this effort.
	Support more regular and well-planned fires in the prairie. Support the formation of Prescribed Burn Associations. Maintain and encourage the collaboration of agencies and nonprofits to meet prescribed fire objectives on a larger landscape scale. Ensure that burn plans incorporate best practices for SGCN including noting the timing of butterfly and moth larval emergence and use of burn units, rotations, and fire exclusion areas to avoid negatively impacting butterfly and moth populations when they are vulnerable to fire. Creating permanent non-fire refugia within fire managed sites has shown benefits to Lepidopteran populations setting aside areas for refugia from burning (Kwilosz and Knutson, 1999; Swengel and Swengel, 2007).
	Increase incentives to develop solar in and on developed areas (e.g, atop buildings and parking lots) rather than in undeveloped lands. Site wind and solar developments on low-use croplands and prohibit development in areas with prairie remnants. Consult with parties earlier in planning processes, including with counties. Explore options with wind energy companies to modify wind easements that are only intended to preclude

	development that would block wind to allow opportunities for acquisition and easements for conservation.
	Implement best practices for wildlife-friendly outdoor lighting, especially near grassland preserves. Consider noise pollution impacts on surrounding prairie and grassland habitats. Consider solutions advised by conservation organizations such as DarkSky . Best practices include using outdoor lighting only where and when it is actually needed, use of motion detectors and/or automatic timers so lights only turn on when they are most needed, placing light covers or shields so that light only shines down to the ground, and keeping landscape lighting low to the ground, and away from white or reflective surfaces. If lighting is necessary, swap out lights for less bright and warmer (yellow or red) bulbs, or cover existing bulbs with a filter that does the same. Consider closing your curtains at night to keep your light indoors (Xerces Society Firefly Friendly Lighting). Also consider solutions recommended by National Audubon's Lights Out Program .



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions

Stressor	Action
	Rigorous effectiveness monitoring/research to assess SGCN responses to management practices/regimes to help inform evidence-based adaptive management.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Stressor	Action
	Develop more evidence-based guidance for conservation grazing that is based on effectiveness monitoring and research, including not only vegetative targets but also responses by Species in Greatest Conservation Need.
	Research, better understand, and share information about the potential to use climate-smart regenerative agricultural practices to improve soil health, prevent soil erosion, build carbon storage, and reduce the carbon footprint of agricultural production to benefit SGCN.

	Encourage counties to eliminate bounties on ground squirrels and gophers where the Northern pocket gopher is found.
	Promote the coexistence of burrowing SGCN and livestock grazing. Emphasize the benefits of diverse grassland communities, discourage activities related to varmint hunting and pest control.

Case Study: Regenerative Grazing, Lonetree Ranch, and the Audubon Conservation Ranching Program

Audubon Conservation Ranching (ACR) is the flagship grassland habitat program of the National Audubon Society. With most remaining grasslands privately owned and operated as working lands, partnering with landowners is the most effective conservation collaboration to impact habitat and bird populations on a landscape scale. North America's grassland bird populations have experienced a steep decline of more than 53% over the last 50 years. By partnering with regenerative-minded farmers and ranchers through the ACR Program, Audubon is working to stabilize and grow those populations across the continent by building resilient lands that support wildlife, ecosystem services, people, and agriculture. Promoting regenerative grazing practices such as rotational grazing, reduced chemical use, cover crops, and native plant establishment, can lead to improved forage quality, drought resiliency, soil stability, carbon sequestration, animal health, and water quality. These key ecological benefits lend to long-term sustainability and improved bottom lines of each individual farm and ranch. To be certified in the ACR program, landowners must meet habitat management, environmental sustainability, and animal welfare standards. Once qualified by a third-party audit, producers then use the Audubon Certified Bird-Friendly seal for packaging and promotion of beef or bison products, which recognizes lands managed for birds and biodiversity. Consumers can then support conservation when they select products from these certified bird-friendly lands.

Lonetree Ranch is an excellent example of outstanding ACR certified bird-friendly land. This family-owned ranch, based in southwest Wyoming and Northeast Utah spans 9,000 acres and raises organic, grass-fed cattle. In the middle of sagebrush country, the Taylor family manages their sage dominated acres for forage and habitat for species such as the greater sage grouse, mule deer, pronghorn, elk, and moose. Their priority is to increase the biodiversity of their vegetative community to establish resiliency and ensure consistent forage is available for their cattle and wildlife. To support their biodiversity goals, Lonetree Ranch partnered with Audubon and the Wyoming Department of Conservation to rejuvenate sagebrush habitat. When sagebrush goes too long without disturbance, such as fire, drought, or grazing, it grows so densely that other plants can't survive in the understory. Project partners performed sagebrush mowing to open the canopy, increasing forb presence and diversity, while also encouraging younger sagebrush to establish. To further improve the bird-friendliness of Lonetree Ranch, Audubon assisted with a fencing project to replace barbed wire fences with wildlife friendly alternatives and improve rotational grazing. This was critical since the Lonetree Ranch is a migratory corridor for mule deer and pronghorn, but they can get entangled when jumping fences, causing serious injury or death.

To determine impacts on grassland bird populations, the ACR program incorporates breeding bird monitoring. Audubon utilizes the Bird-Friendliness Index (BFI), which combines avian point count data with remotely sensed environmental data to evaluate the capacity of a landscape to support an abundant, diverse, and resilient grassland bird community. Soil monitoring is also incorporated into the ACR program to support adaptive management practices and business decisions. Sampling includes collecting one-meter-long soil cores representing the major soils in the farm landscape to be described and analyzed for bulk density, soil carbon, and total nitrogen. These monitoring options build the foundation of the adaptive, regenerative grazing that ACR promotes to manage a mosaic of grassland habitats for birds.

In Minnesota, Audubon began certifying farms through the ACR program in 2024. With funding from the National Fish and Wildlife Foundation, USDA NRCS Conservation Innovation Grant, and the Minnesota Environment and Natural Resources Trust Fund Audubon is dedicated to providing technical and financial assistance to participating ACR producers in addition to monitoring the impacts of the applied regenerative practices on birds, vegetation, and soil health.

References for Prairies and Other Grasslands

- Anderson, R.C. 1990. The historic role of fire in the North American grassland. In: Collins, S.L.; Wallace, L.L., eds. *Fire in North American tallgrass prairies*. Norman, OK: University of Oklahoma Press: 8–18.
- Augustine, D., Davidson, A., Dickinson, K., & Van Pelt, B. (2021). Thinking Like a Grassland: Challenges and Opportunities for Biodiversity Conservation in the Great Plains of North America. *Rangeland Ecology & Management*, 78, 281–295. <https://doi.org/10.1016/j.rama.2019.09.001>
- Ayodele, Ayomiposi Joy. 2025. Small mammal communities as indicators of habitat health: A review. *International Journal of Science and Research Archive* 15 (02); 576-594.
[https://www.researchgate.net/publication/392198024_Small_mammal_communities_as_indicators_of_habitat_health_A_review#:~:text=Abstract,eDNA\)%20and%20automated%20monitoring%20systems.](https://www.researchgate.net/publication/392198024_Small_mammal_communities_as_indicators_of_habitat_health_A_review#:~:text=Abstract,eDNA)%20and%20automated%20monitoring%20systems.)
- Barfknecht, D, Cariveau, A, Jennelle, C, and B Zuckerberg. In prep. Future climate change vulnerability of native Minnesota grasslands and wetlands.
- Basley K, Goulson D. (2018). Effects of Field-relevant concentrations of Clothianidin on Larval Development of the Butterfly *Polyommatus Icarus* (Lepidoptera, Lycaenidae). *Environ Sci Technol* 52:3990–3996
- Bell, J.K., S.D. Siciliano, and E.G. Lamb. 2023. Seasonality and bacterial community assembly process dominate prairie ecosystem service disruption during invasion. *Soil biology and Biochemistry*. 184:109120.
- Bernath-Plaisted, J.S., M.D. Correll, S.G. Somershoe, A.M. Dwyer, A. Bankert, A. Beh, H. Berlanga, W.A. Boyle, J.L. Cruz-Romo, T.L. George, J. Herkert, N. Koper, A. Macias-Duarte, A.O. Panjabi, O.M. Ramirez-Flores, B. Robinson, I. Ruvalcaba-Ortega, J. Sibbing, E.H. Strasser, M. Titulaer, W.E. Van Pelt, and T. VerCauteren. 2023. Review of conservation challenges and possible solutions for grassland birds of the North American Great Plains. *Rangeland Ecology & Management*, 90(9), 165-185.
- Brennan, L. A., & Kuvlesky, W. P. (2005). North American grassland birds: An unfolding conservation crisis? *Journal of Wildlife Management*, 69(1), 1–13. [https://doi.org/10.2193/0022-541X\(2005\)069<0001:NAGBAU>2.0.CO;2](https://doi.org/10.2193/0022-541X(2005)069<0001:NAGBAU>2.0.CO;2)
- Brown DE, Beatty G, Brown JE, Smith AT. 2018. History, Status, and Population Trends of Cottontail Rabbits and Jackrabbits in the Western United States.
- Brown DE, Smith AT, Frey JK, and Schweiger BR. 2020. A review of the ongoing decline of the white-tailed jackrabbit. *Journal of Fish and Wildlife Management* 11:341-352.
- Bucher, S.F., L. Uhde, A. Weigelt, S. Cesarz, N. Eisenhauer, A. Geler, C. Kyba, C. Romermann, T. Shatwell and J. Hines. 2023. Artificial light at night decreases plant diversity and performance in experimental grassland communities. <https://doi.org/10.1098/rstb.2022.0358>.
- Bumgarner, J.R. and R. Nelson. 2021. Light at Night and Disrupted Circadian Rhythms Alter Physiology and Behavior. *Integrative and Comparative Biology* 61 (3); 1160-1169.
<https://academic.oup.com/icb/article/61/3/1160/6206372>.

- Burke, I. C., Lauenroth, W. K., Cunfer, G., Barrett, J. E., Mosier, A., & Lowe, P. (2002). Nitrogen in the Central Grasslands Region of the United States: Current anthropogenic additions of nitrogen to ecosystems of the US central grasslands far outweigh loss of nitrogen through crop removal, resulting in increased nitrogen fluxes with the potential to alter regional-scale biogeochemical cycling. *BioScience*, 52(9), 813–823. [https://doi.org/10.1641/0006-3568\(2002\)052\[0813:NITCGR\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2002)052[0813:NITCGR]2.0.CO;2)
- Buxton R.T., McKenna M.F., Mennitt D.J., *et al.* 2017a. Noise pollution is pervasive in US protected areas. *Science* **356**: 531–33.
- Buxton, R.T., M.F. McKenna, D.Mennitt, E. Brown, K. Kristrup, K.R. Crooks L.A. Angeloni, and G. Wittemyer. 2019. Anthropogenic noise in US national parks – sources and spatial extent. *Frontiers in Ecology and the Environment* (17) 559-564. <https://doi.org/10.1002/fee.2112>.
- Clark CM, Tilman D. Loss of plant species after chronic low-level nitrogen deposition to prairie grasslands. *Nature*. 2008 Feb 7;451(7179):712-5. doi: 10.1038/nature06503. PMID: 18256670.
- Collinge, S.K. (2000). Effects of grassland fragmentation on insect species loss, colonization, and movement patterns. *Ecology*, 81(8), 2211-2226.
- Conway, C.J., 2018. Spatial and temporal patterns in population trends and burrow usage of Burrowing Owls in North America. *Journal of Raptor Research*, 52:129-142. <https://meridian.allenpress.com/rapt/article-pdf/52/2/129/2514315/i0892-1016-52-2-129.pdf>
- Comer, P. J., Hak, J. C., & Seddon, E. (2022). Documenting at-risk status of terrestrial ecosystems in temperate and tropical North America. *Conservation Science and Practice*, 4(2), e603. <https://doi.org/10.1111/csp2.603>.
- DarkSky International. 2025a. [DarkSky International | Protecting the night skies for present and future generations](#). Accessed June 2025. <https://darksky.org>
- DarkSky International. 2025b. Artificial Light at Night: State of the Science 2025. DOI:10.5281/zeno.
- DeSantis et al. (2011). Drought and fire suppression lead to rapid forest composition change in a forest-prairie ecotone. *Forest Ecology and Management*. Volume 261, Issue 11.
- Edwards, C. B., E. F. Zipkin, E. H. Henry, N. M. Haddad, M. L. Forister, K. J. Burls, S. P. Campbell, E. E. Crone, J. Diffendorfer, M. R. Douglas, R. G. Drum, C. E. Fallon, J. Glassberg, E. M. Grames, and R. Hatfield. 2025. Rapid butterfly declines across the United States during the 21st century. 1094:1090–1094.
- Fabio Falchi, Pierantonio Cinzano, Dan Duriscoe, Christopher C. M. Kyba, Dhristopher D. Elvidge, Kimberly Daugh, Boris A. Portnov, Nataliya A. Rybnikova, and Riccard Furgoni. 2016. The new world atlas of artificial night sky brightness. *Science Advances*. 10 June 2016. Vol 2(6): DOI: [10.1126/sciadv.1600377](https://doi.org/10.1126/sciadv.1600377)
- Finch, D. M., Butler, J. L., Runyon, J. B., Fetting, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4

- Forister ML, Cousens B, Harrison JG et al. (2016). Increasing neonicotinoid use and the declining butterfly fauna of lowland California. *Biol Lett* 12:20160475
- Foreman et al. (2000). Pesticides in the atmosphere of the Mississippi River Valley, part II. *Science of the Total Environment* Vol 248, Issues 2-3.
- Galliat, M., Bello, N., Knapp, M., Poland, J., St Amand, P., Baer, S., & Johnson, L. (2019). Local adaptation, genetic divergence, and experimental selection in a foundation grass across the US Great Plains' climate gradient. *Global Change Biology*, 25(3), 850-868.
- Gibbons D, Morrissey C, Mineau P. (2015). A review of the direct and indirect effects of neonicotinoids and fipronil on vertebrate wildlife. *Environ Sci Pollut Res Int* 22:103–118
- Gibbs KE, Mackey RL, Currie DJ. (2009). Human land use, agriculture, pesticides and losses of imperiled species. *Divers Distrib* 15:242–253
- Gilburn AS, Bunnefeld N, Wilson JM et al. (2015). Are neonicotinoid insecticides driving declines of widespread butterflies? *PeerJ* 3:e1402
- Goebel, K. M., N. M. Davros, D. E. Andersen, and P. J. Rice. 2022. Tallgrass prairie wildlife exposure to spray drift from commonly used soybean insecticides in Midwestern USA. *Sci Total Environ* 818:151745.
- Goulson D (2013) Review: an overview of the environmental risks posed by neonicotinoid insecticides. *J Appl Ecol* 50:977–987.
- Green, M. T., P. E. Lowther, S. L. Jones, S. K. Davis, and B. C. Dale. 2020. Baird's Sparrow (*Centronyx bairdii*) in A. F. Poole, editor. *Birds of the World*. Version 1.0. Cornell Lab of Ornithology, Ithaca, New York.
<https://birdsoftheworld.org/bow/home>
- Grimm, E.C., Donovan, J.J., Brown, K.J. 2011. A high-resolution record of climate variability and landscape response from Kettle Lake, northern Great Plains, North America. *Quaternary Science Reviews*, 30(19-20), 2626-2650.
- Hartman, P., & Steele, K. (2015). *Ecological Site Description: Dolomite Colluvium Bluff Prairie*. United States Department of Agriculture.
https://efotg.sc.egov.usda.gov/references/Public/MN/105XY001_DolomiteColluviumBluffPrairie.pdf
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Houston C. S. and M. J. Bechard. 1984. Decline of the ferruginous hawk in Saskatchewan. *American Birds* 2: 166-170 <https://sora.unm.edu/sites/default/files/journals/nab/v038n02/p00166-p00170.pdf>
- Jill J. Gannon, Todd A. Grant, Sara C. Vacek, Cami S. Dixon and Clinton T. Moore *Ecological Restoration*, March 2024, 42 (1) 64-76; DOI: <https://doi.org/10.3368/er.42.1.64>

Jones, Seth A., Edward S. DeKeyser, Cami Dixon, and Breanna Kobiela. 2023. "Invasive Species Change Plant Community Composition of Preserved Prairie Pothole Wetlands" *Plants* 12, no. 6: 1281.

<https://doi.org/10.3390/plants12061281>

Knapp, A.K., J.M. Briggs, S.L. Collins, D.C. Hartnett, L.C. Johnson, and E.G. Towne. 1999. The keystone role of bison in North American tallgrass prairie. *BioScience* 49:39–50

Knapp, E.E., Estes, B. L., Skinner, C.N. 2009. Ecological effects of prescribed fire season: a literature review and synthesis for managers. JFSP Synthesis Reports 4.

Knopf, F. L. (1994). Avian Assemblages on Altered Grasslands. *Studies in Avian Biology*, 15(1), 247–257.

<https://pubs.usgs.gov/publication/70129554>

Koerner, S. E., & Collins, S. L. (2014). Interactive effects of grazing, drought, and fire on grassland plant communities in North America and South Africa. *Ecology*, 95(1), 98-109.

Koerner, S. E., Collins, S. L., Blair, J. M., Knapp, A. K., & Smith, M. D. (2014). Rainfall variability has minimal effects on grassland recovery from repeated grazing. *Journal of Vegetation Science*, 25(1), 36-44.

Kok A.C.M, Boris W. Berkhout, Nora V. Carlson, Neil P. Evans, Nicola Khan, Dominique A. Potvin,. Andrew N. Radford, Marion Sebire, Saeed Shafiei Sabet, Graeme Shannon, and Claudia A. F. Wascher. 2023. [How chronic anthropogenic noise can affect wildlife communities](#). *Frontiers in Ecology and Evolution* 04 (11)

<https://doi.org/10.3389/fevo.2023.1130075>.

Kwilosz, John R., and Randy L. Knutson. "Prescribed Fire Management of Karner Blue Butterfly Habitat at Indiana Dunes National Lakeshore." *Natural Areas Journal*, vol. 19, no. 2, 1999, pp. 98–108. JSTOR,

<http://www.jstor.org/stable/43911819>

Larson, D. L., D. L. Hernandez, J. L. Larson, J. B. Leone, and N. Pennarola. 2020. Management of remnant tallgrass prairie by grazing or fire: effects on plant communities and soil properties. *Ecosphere* 11(8):e03213. 10.1002/ecs2.3213

Leone, J. B., Pennarola, N. P., Larson, J. L., Oberhauser, K., & Larson, D. L. 2022. [Divergent responses of butterflies and bees to burning and grazing management in tallgrass prairies](#). *Ecology and Evolution*, 12, e9532.

<https://doi.org/10.1002/ece3.9532> + [corrections](#)

Li Y., Ruiqing M., and M. Khanna. 2020. Neonicotinoids and decline in bird biodiversity in the United States. *Nature Sustainability* 3 (12):1027-1035.

Liess, S., Roop, H. A., Twine, T. E., Fernandez, A., Dolma, D., Gorman, J., & Neff, P. D. (2023). Fine-scale Climate Projections over Minnesota for the 21st Century. In *AGU Fall Meeting Abstracts* (Vol. 2023, No. 2594, pp. A23Q-2594).

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>

Mackay et al. (2014). Fate in the Environment and Long-Range Atmospheric Transport of the Organophosphorus Insecticide, Chlorpyrifos and Its Oxon. Reviews of Environmental Contamination and Toxicology, Vol 231.

Marschner, F.J. (1974). The original vegetation of Minnesota, compiled from U.S. General Land Office Survey notes by Francis J. Marschner [map] Redrafted from the original by P.J. Burwell and S.J. Haas under the direction of M.L. Heinselman. North Central Forest Experiment Station, USDA. St. Paul, MN.

Mason R, Tennekes H, Sanchez-Bayo F, Uhd Jepsen P. (2013). Immune suppression by neonicotinoid insecticides at the root of global wildlife declines. J Environ Immunol Toxicol 1:3

McLeod, M. L., Cleveland, C. C., Lekberg, Y., Maron, J. L., Philpott, L., Bru, D., & Callaway, R. M. (2016). Exotic invasive plants increase productivity, abundance of ammonia-oxidizing bacteria and nitrogen availability in intermountain grasslands. Journal of Ecology, 104(4), 994-1002.

Mineau P, Downes CM, Kirk DA et al. (2005) Patterns of bird species abundance in relation to granular insecticide use in the Canadian prairies. Ecoscience 12:267–278

Minnesota Biological Survey. (2017). The conversion of documented prairie native plant communities in Minnesota until 2015: an analysis of the DNR Native Prairie Communities layer [compiled by Dustin Graham]. Department of Natural Resources, St. Paul, MN

Minnesota Department of Natural Resources (DNR). 2025. DNR Pollinator Best Management Practices and Habitat Restoration Guidelines. V2

Minnesota Department of Natural Resources (DNR). 2018. [Minnesota Prairie Conservation Plan](https://www.dnr.state.mn.us/prairieplan/index.html). <https://www.dnr.state.mn.us/prairieplan/index.html>; accessed 02/10/2025.

Minnesota Department of Natural Resources. (2025). Ecological Classification System. <https://www.dnr.state.mn.us/ecs/index.html>

Morghan, K. R., & Seastedt, T. R. (1999). Effects of soil nitrogen reduction on nonnative plants in restored grasslands. Restoration Ecology, 7(1), 51-55.

Moriarty, John J. and Carol D. Hall. 2014. Amphibians and Reptiles in Minnesota. University of Minnesota Press.

Nelson, D.M., Hu, F.S. 2008. Patterns and drivers of Holocene vegetational change near the prairie–forest ecotone in Minnesota: revisiting McAndrews’ transect. New Phytologist, 179, 449-459.

Olaya-Arenas P, Kaplan I. (2019). Quantifying Pesticide exposure risk for Monarch caterpillars on milkweeds Bordering Agricultural Land. Front Ecol Evol 7:223

Otfinowski R, Kenkel NC, Catling PM. 2007. The biology of Canadian weeds. 134. *Bromus inermis* Leyss. Canadian Journal of Plant Science 87:183–198.

Pecenka JR, Lundgren JG. (2015). Non-target effects of clothianidin on monarch butterflies. *Naturwissenschaften* 102:19.

Phillips, J.N., S. E. Termondt, and C. D, Francis. 2021. Long-term noise pollution affects seedling recruitment and community composition, with negative effects persisting after removal. *Proceeding of the Royal Society*. <https://doi.org/10.1098/rspb.2020.2906>.

Ratajczak, Z., Nippert, J. B., Briggs, J. M., & Blair, J. M. 2014. Fire dynamics distinguish grasslands, shrublands and woodlands as alternative attractors in the Central Great Plains of North America. *Journal of Ecology*, 102(6), 1374-1385.

Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. 2025. Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwcas.umn.edu/node/821>

Rickey, M. A., & Anderson, R. C. (2004). Effects of nitrogen addition on the invasive grass *Phragmites australis* and a native competitor *Spartina pectinata*. *Journal of Applied Ecology*, 41(5), 888-896.

Rollins, M. G. (2009). LANDFIRE: a nationally consistent vegetation, wildland fire, and fuel assessment. *International Journal of Wildland Fire*, 18(3), 235-249.

Rosenberg, K. V., Dokter, A. M., Blancher, P. J., Sauer, J. R., Smith, A. C., Smith, P. A., Stanton, J. C., Panjabi, A., Helt, L., Parr, M., & Marra, P. P. (2019). Decline of the North American avifauna. *Science*, 366(6461), 120–124. <https://doi.org/10.1126/science.aaw1313>

Roy C.L, and D. Chen. 2023. High population prevalence of neonicotinoids in sharp-tailed grouse and greater prairie-chickens across an agricultural gradient during spring and fall. *Science of the Total Environment* 856:159120.

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>

Runquist, E., Nordmeyer, C. & Stapleton, S. (2024). Widespread annual occurrence of pesticides within designated critical habitats for endangered prairie butterflies. *J Insect Conserv* **28**, 539–552. <https://doi.org/10.1007/s10841-024-00572-5>

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, e14434. <https://doi.org/10.1111/cobi.14434>

Samson, F. B., Knopf, F. L., & Ostlie, W. R. (2004). Great Plains ecosystems: past, present, and future. *Wildlife Society Bulletin*, 32(1), 6–15. [https://doi.org/10.2193/0091-7648\(2004\)32\[6:GPEPPA\]2.0.CO;2](https://doi.org/10.2193/0091-7648(2004)32[6:GPEPPA]2.0.CO;2)

Sánchez-Bayo F, Wyckhuys KAG. (2019). Worldwide decline of the entomofauna: a review of its drivers. *Biol Conserv* 232:8–27

Schlater SM, Ringenberg JM, Bickford N, Ranglack DH. 2021. White-tailed Jackrabbits: A review and call for research. *The Southwestern Naturalist*, 65(2):161-172.

Serafini, J., Grogan, P., & Aarssen, L. (2019). Summer precipitation limits plant species richness but not overall productivity in a temperate mesic old-field meadow. *Journal of Vegetation Science*, 30(5), 832-844.

Swengel, A.B., Swengel, S.R. Benefit of permanent non-fire refugia for Lepidoptera conservation in fire-managed sites. *J Insect Conserv* 11, 263–279 (2007). <https://doi.org/10.1007/s10841-006-9042-9>

Swengel, A.B., Swengel, S.R. Benefit of permanent non-fire refugia for Lepidoptera conservation in fire-managed sites. *J Insect Conserv* 11, 263–279 (2007). <https://doi.org/10.1007/s10841-006-9042-9>

Tilman, D. (1996). Biodiversity: population versus ecosystem stability. *Ecology*, 77(2), 350-363.

Tilman, D., & El Haddi, A. (1992). Drought and biodiversity in grasslands. *Oecologia*, 89, 257-264.

Tracey, C., Faber-Langendoen, F., Thomas Bonnot, T. (2024), Midwest Terrestrial Habitat System. <https://www.sciencebase.gov/catalog/item/665e271bd34e19fd55a9733d>

Wall, D., and T. Pearson, 2013. “D3. Atmospheric Deposition of Nitrogen in Minnesota Watersheds.” In *Nitrogen in Minnesota Surface Waters: Conditions, trends, sources, and*. Minnesota Pollution Control Agency.

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Zouhar, Kristin. 2021. Fire regimes of plains grassland and prairie ecosystems. In: *Fire Effects Information System*, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Missoula Fire Sciences Laboratory (Producer). Available: www.fs.usda.gov/database/feis/fire_regimes/PlainsGrass_Prairie/all.html [2025, May 8].

Savanna

Habitat Description

Savannas typically occur where fire frequency or intensity is somewhat lower than in prairie landscapes, yet higher than in forested areas. At such sites, more fire-tolerant shrubs and trees can persist, forming brush-prairie and savanna communities. Savannas typically have scattered trees, sometimes clumps of trees, above a layer of prairie grasses and forbs. Bur oak (*Quercus macrocarpa*) is the most common and widespread tree, but Northern pin oak (*Quercus ellipsoidalis*) and, in the extreme southeastern part of the state, black oak (*Quercus velutina*) are also typical. Small, open-grown, often gnarled bur oaks are the most distinctive savanna tree species. Savannas where jack pine (*Pinus banksiana*) is the predominant tree species occur on deep sand substrates in the northern half of the state.

Habitat Map

To depict savanna habitat (see Figure 3.7), we compiled spatial data from several sources: DNR's Native Plant Communities and the Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Upland Prairie (UP)

[UPn13 Northern Dry Savanna \(PDF\)](#)

[UPn24 Northern Mesic Savanna \(PDF\)](#)

[UPs14 Southern Dry Savanna \(PDF\)](#)

[UPs24 Southern Mesic Savanna \(PDF\)](#)

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Midwest Woodland and Savanna.




Figure 3.7. A map depicting Savanna habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological Systems of Upland Prairie as well as one type from the Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

Oak savanna is one of Minnesota's rarest wildlife habitats. Prior to European settlement, oak savanna was one of the most common ecosystems in Minnesota, covering roughly 10 percent of the state. Fire suppression, agriculture, and urban growth have reduced the extent of oak savanna coverage to approximately 0.02% of its former range in the Midwestern United States (DNR). Statewide, it is estimated that approximately 9,800 to 11,100 acres of the State's original extent of Oak Savanna (all subtypes) remains. Dry Barrens Oak Savanna in Minnesota is even rarer, with acreage totaling less than 5400 acres. Owing to its' rarity, Dry barrens oak savanna is considered Critically Imperiled (S1) or Critically Imperiled to Imperiled (S1S2) at the State level and Imperiled at a global level (G2). The savanna is a transitional ecosystem between the tallgrass prairie and woodland environments, so it is an important habitat for both woodland and prairie wildlife (see also the [Prairie and Other Grasslands Chapter](#)).

Before European settlement, native grazers, including bison (*Bison bison*), [elk \(*Cervus canadensis*\)](#), and white-tailed deer (*Odocoileus virginianus*), likely contributed to the disturbance and maintenance of oak savanna habitats (Hartman and Steele, 2015). As elk and bison were extirpated and white-tailed deer populations significantly reduced, intermittent grazing was taken over by domesticated livestock including sheep, cattle, and goats (Hartman and Steele, 2015).

Species in Greatest Conservation Need

Savannas provide habitat for 100 animal and 8 plant SGCN as primary or secondary habitat (see Table 3.6.) Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Table 3.6. Numbers of species of SGCN associated with Savanna as either primary or secondary habitat*.

Species Groups	Primary Habitat	Secondary Habitat	Total
Birds	3	4	7
Mammals	5	13	18
Reptiles	12	0	12
Terrestrial Invertebrates			
Bees			4
Beetles (terrestrial)			6
Butterflies			8
Moths			40
Spiders			5
Plants	8		8
Total	28	17	108

* Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants). Examples of selected SGCN are described below.

Birds

Native savanna that meets native plant community criteria (native prairie with scattered native trees or shrubs) is now rare in Minnesota because of conversion to other land uses or through succession to oak woodland due to fire suppression. Most SGCN birds that historically occurred in savanna also make use of surrogate habitats with savanna-like structure. Native savanna often has a range of woody cover, and bird SGCN will vary along this gradient. Open areas dominated by grassland, with scattered shrub or small trees, may support typical grassland birds, while larger patches of trees can provide suitable habitat for woodland or forest species.

Red-headed woodpeckers (*Melanerpes erythrocephalus*) are dependent on large dead trees and snags without bark. Typically, suitable dead trees are adjacent to open areas. Eastern whip-poor-wills (*Antrostomus vociferus*) nest in savanna with a semi-open tree canopy, sand or gravel soils, and an open ground layer. [Loggerhead](#)

[shrikes \(*Lanius ludovicianus*\)](#) require areas of shorter grass adjacent to scattered trees for perches and nesting. Shrikes are now found more often in savanna-like habitat, often along roadsides or fencelines.

Savannas that are predominantly grassland, with only widely-scattered small trees or shrubs, are preferred by field sparrows (*Spizella pusilla*), [lark sparrows \(*Chondestes grammacus*\)](#), and Eastern meadowlarks (*Sturnella magna*), and may be suitable for other grassland bird SGCN (see Prairie/Grassland chapter). [Bell's vireos \(*Vireo bellii*\)](#) breed in dense, shrubby thickets, primarily in southeastern Minnesota. Suitable shrub thickets may occur within upland savanna, prairie, shrub swamps, or along forest or wetland edges. However, Bell's vireos are restricted to shrub patches within or at the edges of open landscapes, rather than under a forest canopy.

Mammals

The open landscape of savannas provides primary habitat for five SGCN mammals. Considered a sentinel or keystone species of grasslands and savannas, the American badger has a significant impact on the surrounding landscape. Residing in underground burrows, and digging through loose, loamy or sandy soils for prey, its excavations provide numerous benefits including aerating the soil, creating shelters for other animals, and improving nutrient cycling. Three other mammals are closely associated with savanna habitats, including the [least weasel \(*Mustela nivalis*\)](#), the [plains pocket mouse \(*Perognathus flavescens*\)](#), and the [plains spotted skunk \(*Spilogale interrupta*\)](#); prairies and other grasslands are primary habitats for each of these species as well. Despite intensive surveys for the spotted skunk in the 1990s, only 4 animals were found resulting in the species listing as a state threatened.

Reptiles

Savannas are especially important habitats for reptiles; 12 of the 13 reptile SGCN are present in these open woodlands of oaks and jack pine. Prairies and other grasslands are nearly as important with 11 of the 13 SGCN, followed close behind by Cliffs, Talus and Rock Outcrops. Lacking canopies, all three of these open habitats provide the necessary thermal requirements for these cold-blooded animals.

[Common five-lined skinks \(*Plestiodon fasciatus*\)](#), six-lined racerunners (*Aspidoscelis sexlineatus*), Eastern hog-nosed snakes (*Heterodon platirhinos*), milk snakes (*Lampropeltis Triangulum*), and [lined snakes \(*Tropidoclonion lineatum*\)](#) are most common in and around woodland edge habitats and savannas. Five-lined skinks use rock fissures and cracks in bedrock outcrops as hibernacula, but the suitability of this habitat is threatened by the encroachment of Eastern red cedar (*Juniperus virginiana*). Several of these reptiles (six-lined racerunners, [north american racers \(*Coluber constrictor*\)](#), and lined snakes) prefer more open areas. Other species (common five-lined skinks, eastern hog-nosed snakes, and milk snakes) prefer edges, using openings as basking areas. Key habitat features for these species include down woody debris (for cover, nesting sites, and basking sites) and burrows or crevices as overwintering sites. For many, the most iconic species of these open habitats are gophersnakes (*Pituophis catenifer*) which are particularly fond of the loose, sandy soils of savannas which also support a rich abundance of burrowing small mammals for food (Moriarty and Hall, 2014).

Plants

Savanna communities with their herbaceous prairie ground layer support many of the same endangered, threatened, and special concern plant species found in respective open prairie habitats (see *Prairie and Other Grasslands* sub-chapter), such as [spike oat \(*Avenula hookeri*\)](#), [plains reedgrass \(*Calamagrostis montanensis*\)](#),

[blunt sedge \(*Carex obtusata*\)](#), and others. Dry barrens communities, with sand soils and arid conditions also host a number of annual and perennial species adapted to, and dependent on these conditions. Annuals such as [seaside three-awn \(*Aristida tuberculosa*\)](#), [annual skeletonweed \(*Shinnersoseris rostrata*\)](#), and [purple sandgrass \(*Triplasis purpurea*\)](#) require shifting sands and this habitat can be threatened by encroaching vegetation (often native) that stabilizes the sands. Some rare perennial species are also primarily found in barrens and dune communities in Minnesota, including [Indian ricegrass \(*Achnatherum hymenoides*\)](#), [winter bentgrass \(*Agrostis hyemalis*\)](#), [slender hairgrass \(*Deschampsia flexuosa*\)](#), [beach heather \(*Hudsonia tomentosa*\)](#), [narrow-leaved pinweed \(*Lechea tenuifolia*\)](#), and [drummond's campion \(*Silene drummondii*\)](#). Some of the rare species found in savanna barrens are at the edge of their ranges in Minnesota, and in the state only savanna barrens satisfy their habitat requirements. Indian ricegrass, for example, is a western species of dry plains that only finds suitable habitat at a few barrens locations in Minnesota.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildlife following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

A large extent of savanna and upland shrublands have been impacted by development for housing and agricultural use, particularly as these habitats ranged across parts of the state with the highest density of human development.



Crop Production

Conversion of savanna habitat to agriculture in conjunction with other forms of human development is a historical and potential future concern.



Livestock Management

As elk and bison were extirpated and white-tailed deer populations significantly reduced, grazing by livestock including sheep, cattle, and goats increased from the early to mid 1900's and likely became very

intensive during the droughts of the 1930's as farmers struggled to find enough forage (Hartman and Steele, 2015). Some savanna habitats likely became overgrazed and became subject to varying degrees of erosion.



Tree Plantations

Pine plantations are abundant in some areas formerly occupied by savannah habitat. Campaigns for increased pressure for tree planting both on former cropland but also pasture for carbon sequestration pose potential for additional plantations in sensitive habitat.



Fire Management

Historically the open environment of oak savanna was maintained largely by fire. Savannas occurred in physical proximity to prairies, but where features such as streams, lakes, and steep topography impeded the spread of those fires. Savannas form where fire occurs frequently enough to prevent trees and shrubs from dominating and shading out sun-loving herbaceous plants, but where frequency and severity are low enough to allow fire-tolerant trees to become established and sometimes reach maturity. All savannas are highly sensitive to fire suppression, quickly succeeding to woodland and eventually to forest in the absence of fire. In the absence of fire, woodland species and invasive shrubs gradually fill the savanna, closing the canopy and shading the prairie grasses and forbs.

Fire frequencies in this habitat have changed from XX to YY over ZZZ for such and such reasons.



Invasive Species (Problematic Non-native Species)

From 1960 to present, invasive species proliferated and continued to invade natural areas including bluff prairies and adjacent savanna (Hartman and Steele 2015). Invasive woody species include common buckthorn (*Rhamnus cathartica*), bush honeysuckle (*Lonicera* sp.), and multiflora rose (*Rosa multiflora*). Forbs such as sweetclover (*Melilotus officinalis*) and grasses such as kentucky bluegrass (*Poa pratensis*) and smooth brome (*Bromus inermis*) are also common invasive species in this habitat type.



Diseases and Pathogens

Bur oak blight is a native fungal disease that affects some bur oaks causing leaves to brown and die. Over time, bur oak blight may weaken the tree, promoting infestation by twolined chestnut borer or Armillaria root disease. Wetter springs may increase the prevalence and severity of this disease as fungal spores spread from the diseased leaves by rainwater, exacerbated by earlier spring rainstorms brought on by Minnesota's changing climate. Oaks in open upland areas are more at risk, and the disease is most problematic for stressed bur oak; healthy bur oaks can tolerate bur oak blight for a decade or more. (consider cross-listing with Changes in Precipitation and Hydrology related to Climate Change)



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of

snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years ([MPCA 2025](#)).

Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al. 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan (Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN), five strategies were identified:

-  **Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.**
-  **Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.**
-  **Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.**
-  **Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.**
-  **Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.**

Examples of conservation actions are grouped below under these five strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Also note that some strategies, such as Strategy 3, collaborating with partners, could truly be applied to all actions to most broadly and effectively implement them.

Priority Conservation Actions for Savannas

 Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.	
Stressor	Action
	Due to their rarity on the landscape, priority should be given to protecting savannas from development.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

Stressor	Action
	To address fire suppression, seek to restore fire to overgrown savannas and brush-prairies, by increasing fire frequency on public lands and conducting prescribed fires. Also monitor the effects of fire suppression and prescribed fire. Barriers are limited agency budgets, lack of capacity and information, permitting, and lack of trained personnel. Solutions lie in supporting inter-agency groups working to restore fire, indigenous people and Tribes that are working to restore fire, cost-shared 'strike-teams' that send fire crews across lands held by multiple land management entities.
	To effectively restore sites degraded by invasive species, such as Kentucky bluegrass, smooth brome, buckthorn, honeysuckle, and multiflora rose, a combination of several conservation actions will most likely be needed (Hartman and Steele, 2015). Monitor the understory and edges of your savanna for these invasive species—and take fast action to control them through prescribed fire, hand or mechanical removal, or selective/spot herbicide treatment, all of which can help the system to be as healthy and resilient as possible. The goal for conservation actions should be to reintroduce disturbance regimes that would have naturally kept woody species in check. This may include prescribed fire or grazing. However, persistent or particularly dense stands of non-native woody species may require a combination of mechanical and chemical treatments to effectively re-establish these natural disturbance regimes. Even as natural disturbance regimes are re-established, ongoing monitoring may be needed to spot treat non-desirable species.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

Stressor	Action
  	Encourage the application of habitat conservation funds towards the protection, restoration, and enhancement of savanna and woodlands, which are highly imperiled but because they are less extensive might not have the emphasis that prairies and forests do in some funding programs.

Stressor	Action
	Reintroduction of elk and bison to these natural systems may be difficult due to societal reasons, but the reintroduction of domestic livestock may be more manageable and when paired with conservation grazing practices, could help to mimic the natural grazing patterns of their native counterparts.
	Encourage managers of savannas, including in urban and suburban parks, to maintain dead trees and trees with dead branches to provide habitat for insects and cavity nesting birds



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions

Stressor	Action
	Also monitor the effects of fire suppression and prescribed fire. Barriers are limited agency budgets, lack of capacity and information, permitting, and lack of trained personnel. Solutions lie in supporting inter-agency groups working to restore fire, indigenous people and Tribes that are working to restore fire, cost-shared ‘strike-teams’ that send fire crews across lands held by multiple land management entities.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Stressor	Action
	Education to natural resource professionals on how to identify a savannah remnant and appropriate management scenarios will be important to mitigate the threat of inappropriate tree planting within savannah habitat.
	Increased spring and early summer rain can increase the prevalence of burr oak blight. As stressed trees are more susceptible to burr oak blight, sometimes only some trees in an area will become affected and can be removed or treated (see this handout for more information). However, if an area has a lot of burr oak blight, possibly consider other strains

	or types of trees may be better adapted to the changing climatic conditions (additional resources on forests and climate change can be found here .)
	Distribute and encourage the use of the following technical resources for managing high quality oak savannas: 1) the High Value Conservation Forests for Dry Barrens Oak Savanna , part of Forest Certification , which outlines, based on the condition of the site, best practices for prescribed fire, forestry, grazing, and restoration; and 2) the Woodlands of Minnesota Landowner Handbook – Oak Savanna which is more directed at private landowners.

References for Savanna

- Finch, D. M., Butler, J. L., Runyon, J. B., Fetting, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniat, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing.
https://doi.org/10.1007/978-3-030-45367-1_4
- Hartman, P., & Steele, K. (2015, January 7). *Ecological Site Description: Dolomite Colluvium Bluff Prairie*. United States Department of Agriculture.
https://efotg.sc.egov.usda.gov/references/Public/MN/105XY001_DolomiteColluviumBluffPrairie.pdf
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Minnesota Department of Natural Resources. 2021. Woodlands of Minnesota Landowner Handbook-Oak Savanna. <https://www.lrl.mn.gov/docs/2022/other/220745.pdf>
- Moriarty, John J. and Carol D. Hall. 2014. *Amphibians and Reptiles in Minnesota*. University of Minnesota Press.
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. P., & LeDee, O. E. (2025). *Effects of climate change on midwestern ecosystems: Temperate flooded and swamp forest* (p. 110pp). Midwest Climate Adaptation Science Center. Available: <https://mwccasc.umn.edu/node/821>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.
<https://statesummaries.ncics.org/chapter/mn/>
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, e14434.
<https://doi.org/10.1111/cobi.14434>
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report].
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Upland Conifer Forest and Woodlands (Fire-dependent)

Habitat Description

Upland coniferous forests occur primarily on mesic to dry sites with coarse sandy or gravelly soils or with thin soils over bedrock in the Laurentian Mixed Forest Province, also as small patches in the Driftless Region in the Eastern Broadleaf Forest Province in the southeastern part of the state. Dominated by pines (*Pinus spp.*), spruce (*Picea spp.*), balsam fir (*Abies balsamea*), or Eastern white cedar (*Thuja occidentalis*), these fire-dependent forests also can contain hardwoods such as quaking aspen (*Populus tremuloides*), paper birch (*Betula papyrifera*), oaks (*Quercus spp.*), and red maple (*Acer rubrum*) (see the Upland Deciduous Forest and Woodland Chapter). Historically, fire was the major source of tree mortality; in a modern context, timber harvests are influential in determining forest ages and patch sizes.

Fire has shaped this forest type, and the dominant trees of this forest, like most plants characteristic of fire-dependent communities, are adapted to survive repeated surface fires or to regenerate successfully following stand-replacing fire. Many plants have seeds or vegetative structures designed to survive fire or are opportunists that can take advantage of short periods when nutrients are relatively abundant and light levels are high. Fires vary greatly in intensity from crown fires, which kill most of the canopy trees, to moderate surface fires, which kill few canopy trees. Fire exerts a strong influence on the pattern of plant reproduction by exposing mineral soil seedbeds, triggering seed dispersal, and increasing the amount of light reaching the forest floor; it also played a significant role in nutrient cycling and nutrient availability.

The frequency of wildfires in this habitat historically varied from an average of every 20 years to every 100 years across forest types in the fire-dependent system, with the northern forest generally experiencing longer intervals than those of the central floristic region ([Laurentian Mixed Forest Fire-Dependent Forest/Woodland System Summary](#)). Some of the northern forests are estimated to have had stand-replacing fires every 170 to 220 years with surface fires about 210 to 260 years with a combined rotation period for all fires of 100 to 115 years. In contrast, central region forests are estimated to have experienced stand-replacing fires every 110 to 130 years and surface fires about every 30 years, with a combined frequency of fire every 22 to 25 years.

Wind was and continues to be another important natural disturbance, as evidenced by the 1999 windstorm that caused extensive damage over nearly 500,000 acres in the Superior National Forest in northern Minnesota, as well as countless acres in Ontario and elsewhere across northern Minnesota (Parke & Larson, 1999). Winds averaged 60 mph and were clocked as high as 80 to 100 mph; and it was estimated that anywhere from approximately 25 million trees on the Superior National Forest and 48 million region-wide were blown down. In addition to the forest, many homes and businesses were impacted.

Plants in this forest type must also survive frequent drought and potentially long periods between disturbance events when nutrients are tied up in plant material and light levels decrease beneath increasingly dense tree

canopies. Evergreen or over-wintering leaves are particularly characteristic; these adaptations enable plants to conserve scarce nutrients in response to low nutrient levels and the thickened outer membranes help to reduce water loss or even herbivory. In general, the species in fire dependent communities have lower nutrient and water requirements and higher light requirements.

Habitat Map

To depict Upland Conifer Forest and Woodland habitat (see Figure 3.8), we compiled spatial data from several sources: DNR's Native Plant Communities and the Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological System

In the Fire-dependent Forest system, some Native Plant Community Classes split up at the level of Types, with some fitting into this Conifer Forest and Woodland Chapter with others in the Upland Deciduous Forest and Woodland Chapter. Types are indicated with a letter following the number in the alphanumeric code and are indented in the list.

Fire-dependent Forest

[FDc12-Central Poor Dry Pine Woodland](#)

[FDc23-Central Dry Pine Woodland](#)

[FDc24-Central Rich Dry Pine Woodland](#)

[FDc25-Central Dry Oak-Aspen \(Pine\) Woodland](#)

FDc25a Jack Pine - Oak Woodland

[FDc34-Central Dry-Mesic Pine-Hardwood Forest](#)

FDc34a Red Pine - White Pine Forest

[FDn12-Northern Dry-Sand Pine Woodland](#)

[FDn22-Northern Dry-Bedrock Pine \(Oak\) Woodland](#)

FDn22a-Jack Pine Woodland (bedrock);

FDn22b-Red Pine - White Pine Woodland (Northeastern Bedrock);

FDn22d-Red Pine - White Pine Woodland (Eastcentral Bedrock))

[FDn32-Northern Poor Dry-Mesic Mixed Woodland](#)

[FDn33-Northern Dry-Mesic Mixed Woodland](#)

FDn33a Red Pine - White Pine Woodland

FDn33c Black Spruce Woodland

[FDn43-Northern Mesic Mixed Forest](#)

FDn43a-White Pine - Red Pine Forest

FDn43c-Upland White Cedar Forest

[FDs27-Southern Dry-Mesic Pine-Oak Woodland](#)

FDs27b White Pine - Oak Woodland (Sand)

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Laurentian Pine – Oak Forest and Woodland, Laurentian Pine Barrens, Laurentian Subboreal Dry Pine – Black Spruce Woodland, and Laurentian Subboreal Mesic Balsam Fir – Spruce – Hardwood Forest.




Figure 3.8. A map depicting Upland Conifer Forest and Woodland habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological Systems of Fire-dependent Forest/Woodland as well as four types from the Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

Today, fire has been essentially replaced with timber harvest, resulting in a different distribution of forest patch sizes, ages, and composition. Moreover, conversion of upland conifer forests to other uses (such as agriculture, development and transportation corridors) has fragmented this habitat across the Laurentian Mixed Forest Province. The remaining forests in this region often lack the ecological complexity of pre-European settlement forests because of many factors (such as, grazing, invasive plants and animals, edge

effects, changes in native animal populations, and consumptive uses), whose relative importance varies in different regions of the state. Even in the largely forested areas of northern Minnesota, rural sprawl has greatly reduced the extent of large, contiguous forest areas; most forest areas are within 25 kilometers (15.5 miles) of small housing settlements (Radeloff et al., 2005).

Historically, wildfire played an important role at both the site and landscape level in these forests, but due to land use changes and fire suppression, wildfire is not common today. Following a crown fire, a variety of biological legacies or features of the predisturbance ecosystem that persist after disturbance (Perry & Amaranthus, 1997) remain in the young regenerating forest. These biological legacies, which include standing dead trees, large trees that escaped the fire, down logs, and small patches of unburned vegetation, provide important habitat features for SGCN. At the landscape level, wildfires create a shifting mosaic of native plant communities with a variety of ages and structural characteristics that provide habitats for SGCN.

Throughout the Laurentian Mixed Forest Province, the extent of upland conifer forest has been reduced by two-thirds since settlement by people of European descent and most have been replaced by aspen-dominated forests (Friedman & Reich, 2005). Fire has been replaced by clear-cut logging as the major stand-replacing disturbance. Most of the remaining upland conifer forests (pine) have been thinned or originated following clear-cutting and typically lack the structural diversity of stands originating following fire. Forest management has resulted in a decrease in the amount of old forests, a decline that will continue for decades. Management to reduce spruce budworm outbreaks (harvesting balsam fir at young ages) may reduce populations of the warblers that eat spruce budworms. Even in the Boundary Waters Canoe Area Wilderness, the huge blowdown of 1999 is resulting in the conversion of pine forests to forests dominated by other tree species.

Climate changes can indirectly affect all of Minnesota's forest communities by interacting with invasive species, insect pests and diseases, fires, and deer herbivory (Galatowitsch et al., 2009; Handler et al., 2014). Droughts will lead to an increased risk for wildfires. Minnesota's conifer forests are particularly vulnerable to the impacts of climate change, with broadleaf temperate trees able to outcompete coniferous boreal tree species, in addition to the other negative impacts warming can have on these forests (Frelich et al., 2021).

Species in Greatest Conservation Need

Upland conifer forests and woodlands provide habitat for 64 animal and 35 plant SGCN as primary or secondary habitat (see Table 3.7.) Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Table 3.7. Numbers of species of SGCN associated with the Upland Conifer Forest and Woodlands as either primary or secondary habitat¹.

Species Group	Primary Habitat	Secondary Habitat	Total
Amphibians	0	3	3
Birds	7	9	16
Mammals	1	18	19
Reptiles	2	0	2
Terrestrial Invertebrates			
Bees			5
Beetles (terrestrial)			2
Butterflies			5
Moths			7
Snails (terrestrial)	1	4	5
Plants	35		35
Total	46	34	99

1. Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Animals with more general habitat requirements are associated with multiple habitats, while specialists are associated with fewer. Plants were only associated with their most primary habitat. Detailed tables associating each of the SGCN with each of the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C for animals and Appendix D for plants. Examples of some of the SGCN and their significance are described below.

Amphibians

Three salamander SGCN – [spotted \(*Ambystoma maculatum*\)](#), [four-toed \(*Hemidactylium scutatum*\)](#), and Eastern newt (*Notophthalmus viridescens*) – are strongly associated with mature forest habitats. They primarily occupy deciduous forests but will also used mixed deciduous-conifer forest types. Access to wetlands are critical for these species, which use them in the spring to breed. [Eggs are laid in or near the water, and larvae remain in the pools until metamorphosis occurs.](#) Four-toed salamanders specifically lay their eggs in moss hummocks next to wetlands. Four-toed salamanders occur most frequently in mature upland forests on glacial moraine landscapes with frequent isolated wetlands that include an alder margin and moss hummocks adjacent to [pockets of open water](#). Two salamanders that are not SGCN are also prevalent in this habitat (Eastern red-

backed salamander (*Plethodon cinereus*) and blue-spotted salamander (*Ambystoma laterale*)), further emphasizing the importance of Upland Conifer Forests to these amphibians for all activities outside of the breeding season.

Birds

Many SGCN birds of upland forests are adapted to the wide range of habitat conditions and landscape patterns created by natural disturbance regimes such as fire. Black-backed (*Picoides arcticus*) and American three-toed woodpeckers (*Picoides dorsalis*) are most frequently found in recently burned upland coniferous forests, where they feed on wood-boring beetles. Prior to fire suppression, new patches of burned forest occurred near older burn areas within a decade or less, providing a steady supply of dead trees for woodpeckers. Black-backed and three-toed woodpeckers will also utilize unburned forest where suitable dead trees are available. Wildfires also created a mosaic of new patches of dense young forests of jack pine and spruce, ideal habitat for spruce grouse (*Canachites canadensis*), which feed primarily on needles of these species. The fire regimes in upland coniferous forests allowed the development of extensive old-growth forests with habitat features such as large snags, down trees with root wads, stumps, very large trees, and small openings in the canopy, which provide optimal habitat for winter wrens. Older stands of white spruce and balsam fir provide optimal habitat for bay-breasted warblers (*Setophaga castanea*) and Cape May warblers (*Setophaga tigrina*), which feed on spruce budworms and increase during periodic outbreaks of budworm in older spruce and fir. Black-throated blue warblers (*Setophaga caerulescens*) require mature closed-canopy forests, often on steep slopes, with a well-developed understory and dense shrub layer. In fire-dependent upland forests, black-throated blue warblers require forest communities on the mesic side of the spectrum (almost exclusively found in northern mesic mixed forest – FDn43). Occupied forest stands typically have a mix of coniferous and deciduous trees but may occur in conifer-dominated forests if other structural characteristics (mature trees, dense shrub layer, etc.) are present. [American goshawks \(*Accipiter gentilis*\)](#) prefer forests with a relatively open flight path between the canopy and subcanopy. Because they have home ranges that extend over several native plant communities, the landscape mosaic is particularly important. The Eastern whip-poor-will (*Antrostomus vociferus*) may reach its highest population density in the state in pine-dominated (particularly jack pine or young red pine (*Pinus resinosa*)) woodlands and forests in north-central Minnesota. Key habitat components are dry sand or gravel soils and an open ground layer lacking dense saplings or shrubs.

Mammals

Among the 19 mammal SGCN that utilize Upland Conifer Forests and Woodlands, only one is a habitat specialist. The rock vole (*Microtus chrotorrhinus*) may be found in a variety of habitats but in Minnesota it is largely restricted to northern conifer and hardwood forests where the understory includes boulder fields, moss-covered rocks, talus, and downed logs and a nearby water source. In addition to Upland Conifer Forests, Upland Deciduous Forests and Cliffs, Talus and Rock Outcrops have all been identified as primary habitats. Included among the other 18 SGCN that can be found in Upland Conifer Forests are all 7 bat SGCN and top predators such as the [Canada lynx \(*Lynx canadensis*\)](#) and [gray wolf \(*Canis lupus*\)](#).

Reptiles

Upland Conifer Forests and Woodlands are designated primary habitat for only two reptile species. The [wood turtle \(*Glyptemys insculpta*\)](#) is largely an aquatic species, found in small or medium-sized rivers running through both conifer and deciduous uplands. They are found in the adjacent riparian habitat basking, foraging ,

and laying their eggs but rarely wander more than about 1,300 feet from the water. The Eastern hog-nosed snake (*Heterodon platirhinos*) utilizes a variety of upland habitats but demonstrates a preference for open fields, field/forest edges, and sandy pine stands. Such conditions can be found in many habitats including Upland Conifer Forests and Woodlands.

Plants

Among the 35 plant SGCN that are found in upland conifer forests, Slender Hair Grass, ([Deschampsia flexuosa](#)) is a particularly enigmatic plant often found near the shoreline of Lake Superior in association with pine forests. Little is known about this species biology, ecology or management needs. It emphasizes the importance of information gathering, such as management effectiveness monitoring, for the long-term conservation of plant SGCN. Of the total number of vascular plant SGCN identified for the 2025-2035 SWAP (447); the SGCN found in the upland conifer forest represent approximately just 8% of all SGCN plants. [Eastern hemlock \(*Tsuga canadensis*\)](#) is a slow-growing, long-lived, and late successional conifer with a range occurring from New England to the Great Lakes and extending down the Appalachian Mountains (Godman & Lancaster, 1990). While capable of living to over 500 years old, few seedlings reach maturity, with fewer than half of these seeds being fertile (Godman & Lancaster, 1990), and germination is dependent on specific substrate requirements; mineral soils or rotting logs can provide the moisture requirements for seedlings (Harmon et al., 1986; Olson et al., 1959; Rooney & Waller, 1998), hence mature hemlock trees facilitate seedling regeneration (Holmes, 2009; Tyrrell & Crow, 1994). Historic reports show hemlock in north central and northeastern Minnesota, but logging and fire (Lawson, 1942) and the early animal hide tanning industry (Stearns, 1997) have shrunk Minnesota's population down to an estimated 40-50 wild trees in small stands around Duluth ([DNR Rare Species Guide](#)).

Minnesota's largest population of hemlock grows on steep slopes that are prone to landslides (Mooers and Watrus, 2005). Groundwater seepage has likely contributed to this population's persistence, but recent large rain events have caused slope failures, washing away mature trees and seedlings (Perry, 2024). Minnesota's stands are failing to regenerate naturally, given the challenges of reduced seed source and facilitation of mature trees to establish seedlings. Enacted protections around endangered species support a hands-off approach, making research into artificial regeneration methods more difficult (Ellingson, 2017). Increasing populations of hemlock in Minnesota will likely require reintroduction or assisted migration (Pollack, 2024).

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key 'stressors' that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the "Stressors" section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as 'stressors' can also be used to advance conservation goals. Broad terms such as 'fire management' reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense

wildlife following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Residential, commercial, and industrial development can all directly alter forest ecosystems by removing, degrading, or fragmenting forest cover and habitat. Fragmentation of forests is a growing concern, and occurs when a large, contiguous forest land mass is divided into smaller tracts through sale and subdivision, road construction, or clearing of forests for agriculture, housing, or other commercial developments. This often creates smaller forest pockets, which are interspersed with non-forest land activities. Fragmentation can inhibit the natural migration of many plant and animal species, increase the risk of wildfires at the wildland urban interface, restrict public recreational access, reduce the habitat value and ecological complexity of forest lands, and contribute to the spread of invasive species. Because of their natural beauty, diversity, recreation opportunities, and abundant natural resources, forested landscapes are appealing places to live; even in the largely forested areas of northern Minnesota, rural development has greatly reduced the extent of large, contiguous forest areas; most forest areas are within 25 kilometers (15.5 miles) of small housing settlements (Radeloff et al., 2005). Forested lands adjacent to lakes and rivers are especially attractive sites for development.



Crop Production

Forests and woodlands, especially in the southern portions of the Laurentian Mixed Forest Province, where Upland Coniferous forests are most common, are susceptible to conversion for agricultural use. Although large-scale row crops are not common, small farmsteads with cattle and hayfields are scattered throughout the landscape.



Livestock Management

Minnesota currently has over 650,000 acres of mesic, upland deciduous, and riparian forests on farms that are grazed by livestock (Zamora et al., 2017; Garrett et al., 2004). Concerns around soil and water quality can arise when forests are grazed. Historically, grazing by livestock in forested landscapes likely became very intensive during the droughts of the 1930's as farmers struggled to find enough forage (Hartman and Steele, 2015). These habitats likely became overgrazed and became subject to varying degrees of erosion, while the forest vegetation is subject to trampling, grazing, fraying, and bark stripping all of which can impact regeneration. (Hartman and Steele, 2015).



Mining and Quarrying

Forests, when they co-occur with mineral or sand and gravel deposits, are at risk of degradation due to the direct and indirect impacts of mining activities such as soil compaction, introduction of invasive species, human disturbance, waste discharge and pollution, forest fragmentation, and in some cases, permanent forest loss (Sonter et al., 2018). Mining projects can trigger additional land use changes with the establishment of associated infrastructure such as improved road access. Renewed interest in mining copper,

nickel, cobalt and precious metals in northern Minnesota has heightened concern about the potential impacts in the northern forest region.



Roads, Trails, and Railroads

Roads, trails, and railroads can create fragmentation and edge effects in large tracts of forest. Increase in the number of roads and trails, particularly for motorized vehicles, increases human disturbance and introduction of invasive species further into the forest. Repeated use and soil compaction associated with roads and trails also can modify the vegetation within the area of use. Temporary winter logging trails may become rutted while materials used to stabilize the trails for heavy vehicles may remain in place, leaving behind wood planks and wood chips. Trails that provide winter access to upland deciduous stands can disturb forest wildlife, especially species that begin nesting in late winter, such as many forest raptors.



Utility Corridors

Utility corridors can cause fragmentation and habitat degradation, such as by introducing invasive plants into forests and woodlands, as well as direct disturbance and oftentimes permanent deforestation and habitat loss incurred during their installation.



Tree Plantations

Red pine plantations have been planted on sites where red pine and jack pine forests and woodlands occurred naturally. These plantations are designed to maximize stand-level growth of red pine, and do not typically resemble the more biodiverse, naturally occurring woodlands where red pine often occurred with white and jack pine and several other less abundant species. In general, monoculture plantations have lower value for wildlife and biodiversity than natural forests (Hua et al., 2022). Conversions to plantations happened historically, based on planting records, and are ongoing with current standard practices that include clear-cutting natural stands and planting at plantation-level stocking densities.



Timber Harvest

Even-aged harvests followed by natural or artificial regeneration are typical in upland conifer forests, which produce even-age class distributions. Timber management may to an extent mimic stand-replacing fires and natural insect outbreaks that historically occurred in these forests. However, the rotation ages may be shorter than prehistoric disturbance intervals and lack the representation of mature and older trees in those areas. Also, strip thinning or individual tree selection are commonly applied, which also reduce stand diversity; variable density thinning, which promotes structural diversity in a stand (Palik & Kastendick, 2023), is uncommon.

Timber harvest is used to address forest health concerns, such as dwarf mistletoe in black spruce, and *Diplodia* (*Diplodia sapinea*), a fungus that infects red pine, especially young red pines (see also the Diseases and Pathogens stressor). Due to concerns with *Diplodia*, clear-cuts are typically applied without the retention of red pine reserve trees and planted with rows of young red pine seedlings. The planted stock may be infected or become infected from the fungus present in older trees nearby.

Timber harvest in this fire-dependent conifer forest can increase the amount of young forest on the landscape, change age-class distributions, reduce species and structural diversity in stands, convert forest types (i.e., jack pine to red pine), reduce patch sizes, and cause habitat fragmentation. Clear-cuts creating even age-class distribution up to rotation age can reduce the amount of mature and older forest.



Recreation

In addition to the impacts created by recreational trails (discussed above), recreational activities themselves can proliferate invasive species. Seeds of invasive plants such as common tansy (*Tanacetum vulgare*) can become embedded in the wheels of vehicles or the shoes of individuals and eventually become established along compacted trails where conditions for their growth may be more favorable. Earthworm egg cases can also be spread in mud, which are problematic for moonwort (*Botrychium*) species. Increased use of roads and trails also increases human disturbance in the forest, reducing the amount of inaccessible area.



Fire Management

A history of altered fire intervals, fire-suppression, and exclusion of cultural burning practices has resulted in an accumulation of fuels. In some areas the historical understory vegetation was more open to allow for ease of hunting and to encourage berry producing shrubs such as blueberries. Fire suppression for extended periods of time favors shade-tolerant species such as red maple and oaks. Over time this leads to mesophication of the plant community and results in declining diversity of plant and animal species (Hanberry et al., 2012; Nowacki & Abrams, 2008). Prescribed fire is infrequently applied.



Invasive Species (Problematic Non-native Species)

Invasive species are an ongoing stressor in Minnesota's forest and woodland systems. Invasive plant species, such as buckthorn (*Rhamnus cathartica*) and Tartarian honeysuckle (*Lonicera tartarica*) often outcompete native plant species for light, space, and water and repress understory plant species leading to modified habitats unsuitable for SGCN. Dense canopies formed by invasive shrubs also inhibit the regeneration of native tree species due to the increase in shade.

Invasive animals, primarily invertebrates, cause profound ecological changes in forests (Frelich et al. 2019). Eurasian earthworms upend the traditional soil profile (Hale et al. 2005; Fahey et al. 2013), remove the accumulated layers of organic leaf litter (Holdsworth et al. 2008, 2012), and alter the community of soil mycorrhizal fungi (Paudel et al. 2016). These changes cascade through the ecosystem, including ground layer vegetation (Hale et al. 2006; Holdsworth et al. 2007), leaf litter fauna (McCay and Scull 2019), salamanders (Maerz et al. 2009; Ziemba et al. 2016), and likely other vertebrates.

The spongy moth (*Lymantria dispar*) is an invasive forest pest that feeds on more than 300 species of deciduous trees and shrubs, including aspen, oak, and birch. Repeated removal of leaves, or defoliation, stresses trees and can leave them vulnerable to disease or other pest infestations that can kill trees. Once it becomes established in a location, the spongy moth has cycles of large population outbreaks every eight to 12 years, leading to widespread defoliation and nuisance from caterpillars.

European pine sawfly (*Neodiprion sertifer*) primarily feeds on two and three-needle pines but can also infest five-needle pines when populations are high or when they are growing adjacent to their preferred host type.



Problematic Native Species

Overbrowsing by deer can severely limit natural tree regeneration (Matonis et al., 2011; De Jager et al., 2013), especially for preferred species, such as white pine and oak. Elevated deer populations also reduce populations of palatable ground layer plants, including orchids (Knapp and Wiegand, 2014), members of the lily family (Balgooyen and Waller, 1995; Augustine and Frelich 1998; Fletcher et al., 2001), and rare species (McGraw and Furedi, 2005). Decades of cumulative impacts have resulted in profound changes to the composition of native plant communities (Wiegmann and Waller, 2006; Frerker et al., 2014; Nuttle et al., 2014; Flagel et al., 2016; Anderson et al. 2017; Sabo et al. 2017). The effects of deer browsing often interact with those of invasive earthworms (Fischelli et al., 2013; Davalos et al., 2014).

[Eastern](#) spruce budworm is a native caterpillar and is the most destructive pest of spruce-fir forests in eastern North America. Outbreaks may be related to factors such as annual weather variation and climate change. Spruce budworm outbreaks attract a variety of predator specialists including the bay-breasted, Cape May, and Tennessee warblers.



Diseases and Pathogens

Diseases and pathogens are ongoing and future stressors in forest and woodland systems. [Diplodia](#) (*Diplodia sapinea*) is a fungus that causes dead branches and branch tips (shoot blight), decreased growth, top-kill, and sometimes death in jack pine and red pine. It is not known if the pathogen is native to Minnesota, but it has been present since the 1970's. It is most detrimental to seedlings and saplings growing in the shade. Another similar fungal pathogen is [Sirococcus conigenus](#). In one study, sapling pines developing near previously infected older pines were at greater risk for becoming infected and dying from these two fungal pathogens (Haugen and Ostry, 2013); thus standard practice is to remove all mature pines from most harvests where pines are the target replacement trees. It is assumed present even if it is not there.



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years ([Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025](#)). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, heightened disease and pathogen risks, winter burn, and deer herbivory (Galatowitsch et al., 2009; Handler et al., 2014).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation,

interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). Minnesota's conifer forests are particularly vulnerable to the impacts of climate change, with broadleaf temperate trees able to outcompete coniferous boreal tree species, in addition to the other negative impacts warming can have on these forests (Frelich et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). More frequent droughts will lead to an increased risk for wildfires. Restoration efforts can be disrupted by droughts, which are becoming more frequent with changing climatic conditions, causing seedling mortality and shifts in community composition if there are multiple seasons with insufficient rain. Prolonged droughts also can stress trees, increasing susceptibility to disease, pests, or pathogens. For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan (Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN), five strategies were identified:

-  **Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.**
-  **Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.**
-  **Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.**
-  **Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.**
-  **Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.**

Examples of conservation actions are grouped below under these five strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Also note that some strategies, such as Strategy 3, collaborating with partners, could truly be applied to all actions to most broadly and effectively implement them.

Potential Conservation Actions for Upland Conifer Forests and Woodlands

 **Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.**

Stressor	Action
     	Identify priority areas for protected area designations to facilitate long-term conservation outcomes and protection against conversion of forest to different land uses. Use the Conservation Action Network or other sources of information to highlight high biodiversity areas or those providing connectivity between other protected areas that are likely to provide disproportionately high conservation value.

	
   	Consider projected effects on high value conservation forests, old-growth forests, and Important Bird Areas when making decisions and policy that are projected to impact upland coniferous forests (e.g., mining and quarrying or oil and gas permitting, planning for new development of energy sources and any associated new road or trail construction). High conservation value forests can be vulnerable to degradation by mining activities..



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

Stressor	Action
	Encourage the use of existing corridors for new trails, roads, and utility corridor development to minimize the construction of new roads and trails that would increase forest fragmentation.
 	Limit the use and operation of heavy equipment when road or site conditions are vulnerable to damage. For instance, depending on the site characteristics, conducting timber harvests during frozen ground conditions can reduce soil compaction and rutting impacts. Increases in winter temperatures are reducing the duration of the winter season.
	Support forest management approaches that are informed by ecological processes, increase structural and compositional complexity and diversity, and assist in resilience towards future stressors (e.g., climate change, disease outbreaks, invasive species) in forests (refer to Native Plant Classification silviculture strategies for forest management). Promote forest management for a diversity of forest growth stages across the landscape to promote vegetation characteristics that maintain habitat for a wide range of plant and animal species, promote resiliency to insect and disease outbreaks, and mitigate the adverse effects of climate change. Ensure older forest and old forest qualities are retained on the landscape in amounts and areas that provide critical habitat to SGCN animals and plants; encourage management that sustains mature and old-growth forests in forest planning efforts. Maintain or enhance habitat connectivity between forest patches to allow wildlife movement. Consider flexibility in stocking goals in natural stands (i.e., longer time frames to meet goals, non-crop trees can be a significant portion of goals) and protecting important biodiversity areas. Explore a range

	of silvicultural methods to regenerate jack pine in central Minnesota where the nutrient poor soils initially slow the growth of young seedlings.
	Apply knowledge of SGCN habitat use into forest planning efforts to help balance habitat for SGCN and other values according to management requirements and objectives. Consider natural area designations and other ways to enhance and preserve key mature stands. Consider the ecological values of various forest ages and how they relate to SGCN habitat when determining desired future conditions and planning sustainable management to advance those conditions.
	Retain old and dead standing trees (snags) in forests when practical as habitat for wildlife, including cavity-nesting birds.
	To adapt to a warming climate, consider strategies for maintaining conifers in the Laurentian Mixed Forest Province, focusing on species, such as white pine and jack pine, that are predicted to be well-adapted to a warming climate. Collaborate with various partners to select sites in northern Minnesota ('conifer strongholds') where replanting would be most successful.
	Where people recreate, provide outreach and education on cleaning gear and equipment to reduce the introduction and spread of invasive species. DNR's Prevent the spread webpage includes specific actions for different land based activities like biking, hiking and off-highway vehicle riding. At trailheads, add invasive species prevention messages, such as those on the boot brush kiosks developed by the PlayCleanGo: Stop Invasive Species in Your Tracks program.
	Increase fire frequency on public lands, exploring opportunities to implement forest management practices that use fire, letting some wildfires burn and application of prescribed fire, such as in the 1999 blowdown in the Boundary Waters Canoe Area Wilderness, would improve forest health. Support forest centered burn crews. Integrate with forest management as a means to prepare forests for prescribed burning and to prevent catastrophic fires.
	Enact measures to ensure that conifer seedlings prone to overbrowsing by deer are adequately protected when deemed necessary. Bud capping, fencing, and slash walls are ways to mitigate potential impacts from deer browse.
	Implement control strategies for invasive species, such as buckthorn, and ensure sufficient timeframe for follow-up management that extends for many years. Support funding for these activities and consider expanding restoration resources to make sure they can be applied to woodland systems, which are sometimes in between funding opportunities aimed at prairies or forests.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

Stressor	Action
	Support local governments in measures that minimize fragmentation and encroachment into high quality forests and woodlands such as protecting high quality areas, zoning that guides housing developments and lot sizes that support local forests, and land use planning and development guidance.
	Encourage sustainable forest management through planning when and where timber harvest can be used as an effective tool to address a range of values and priorities, including forest health and productivity, critical ecological functions, and habitat restoration from historical disturbance events. Incorporate considerations of forest compositional and structural diversity at meaningful scales, historic size and return intervals of disturbance regimes, biodiversity, rare features, wildlife needs, and diversity of native plant community growth stages. Ensure that a balance of forest habitat conditions (e.g. younger and older forest, small and large patch sizes) are available for plant and animal species across the landscape over time.
	Collaborate across forest ownerships and managers to implement sustainable forestry at the site and landscape level to create habitat conditions important to SGCN, such as unfragmented older mesic hardwood forests to create larger forest patches and reduce forest fragmentation. Support forest management approaches that are informed by ecological processes, increase structural and compositional complexity and diversity, and assist in resilience towards future stressors (e.g., climate change, disease outbreaks, invasive species). Promote forest management strategies that mimic landscape disturbance patterns (for example, retaining large blocks of mature habitat and creating more small forest openings as gaps or strips), maintain or enhance connectivity among forest patches to enable wildlife movement, and retain biological legacies (at site level) such as large trees with cavities. Consult Native Plant Community Classification silvicultural strategies for forest stand prescriptions .
	Incorporate climate adaptation into forestry planning, including climate-adapted silviculture. Extending the duration of grants and contracts for restoration projects could help to allow conservation practitioners to plant when conditions are more favorable, in light of droughts being likely to become more frequent.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions

Stressor	Action
	Monitor the responses of SGCN to various types of forest management at the site-level and landscape-scale. Conduct research and model anticipated changes with different forest management scenarios for impacts on wildlife habitat. Develop models of habitat configurations for various SGCN and consider how current harvest forecasts and conversion rates (i.e., one forest type to another; natural origin habitat to plantation) relate to those models in terms of sustainability for maintaining habitat for SGCNs. Support the collection of accurate and reliable forest stand inventory data for planning.
	Communicate and research the benefits of alternative management techniques (i.e., variable density thinning, natural seeding), reference to natural disturbance intervals, natural regeneration methods, and to increase habitat quality for SGCN. Promote consideration of flexibility in stocking goals in natural stands (i.e., longer time frames to meet goals, non-crop trees can be a significant portion of goals) and the ability to protect important biodiversity areas from harvest.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Stressor	Action
	Promote and consider further development of tax incentives that enable landowners and farmers to gain value for maintaining conservation values on their properties. Provide increased levels of incentive payments for woodland owners to help protect woodlands and forests from further reduction due to development pressure. Support and promote education and technical assistance to maintain or restore forest habitat.
	Managing for an increased diversity of dominant species in a site can assist in resilience toward future disease outbreaks, susceptibility to some pathogens, and invasive species. To assist in the management of woodlands and forests that are resilient to pathogens and invasive species may require additional funding. Also consider pairing management of non-native invasives with other incentivized management activities and prioritizing comingled activities to create efficiencies.

References for Upland Coniferous Forest

- Anderson, R.C., Anderson, M.R. and Corbett, E.A., (2017). White-tailed deer (*Odocoileus virginianus*) and fire effects on flowering diversity of tallgrass prairie forbs. *The Journal of the Torrey Botanical Society*, 144(3), pp.243-253
- Augustine, D. J., & Frelich, L. E. (1998). Effects of White-Tailed Deer on Populations of an Understory Forb in Fragmented Deciduous Forests. *Conservation Biology*, 12(5), 995–1004. <https://doi.org/10.1046/j.1523-1739.1998.97248.x>
- Balgooyen, C. P. and D. M. Waller. (1995). The use of *Clintonia borealis* and other indicators to gauge impacts of white-tailed deer on plant communities in northern Wisconsin, USA. *Natural Areas Journal*, 15(4), 308-318. <https://www.cabidigitallibrary.org/doi/full/10.5555/19960610148>
- Dávalos, A., Nuzzo, V. and Blossey, B., (2014). Demographic responses of rare forest plants to multiple stressors: the role of deer, invasive species and nutrients. *Journal of Ecology*, 102(5), pp.1222-1233.
- De Jager, N.R., Cogger, B.J. and Thomsen, M.A., (2013). Interactive effects of flooding and deer (*Odocoileus virginianus*) browsing on floodplain forest recruitment. *Forest Ecology and Management*, 303, pp.11-19.
- Ellingson, E. K. (2017). *The Genetic Diversity and Conservation Potential of Eastern Hemlock (Tsuga Canadensis (L.) Carrière) in Minnesota* (1968601810) [M.S., University of Minnesota]. <http://library.dnr.state.mn.us/login?url=https://www.proquest.com/dissertations-theses/genetic-diversity-conservation-potential-eastern/docview/1968601810/se-2?accountid=28003>
- Fahey, T.J., Yavitt, J.B., Sherman, R.E., Maerz, J.C., Groffman, P.M., Fisk, M.C. and Bohlen, P.J., (2013). Earthworm effects on the incorporation of litter C and N into soil organic matter in a sugar maple forest. *Ecological Applications*, 23(5), pp.1185-1201.
- Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weyand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Fisichelli, N.A., Frelich, L.E., Reich, P.B. and Eisenhauer, N., (2013). Linking direct and indirect pathways mediating earthworms, deer, and understory composition in Great Lakes forests. *Biological Invasions*, 15, pp.1057-1066
- Flagel, D.G., Belovsky, G.E. and Beyer, D.E., (2016). Natural and experimental tests of trophic cascades: gray wolves and white-tailed deer in a Great Lakes forest. *Oecologia*, 180, pp.1183-1194
- Fletcher, D.J., Shipley, L. A., McShea, W. J., & Shumway, D. L. (2001). Wildlife herbivory and rare plants: The effects of white-tailed deer, rodents, and insects on growth and survival of Turk's cap lily. *Biological Conservation*, 101(2), 229–238. [https://doi.org/10.1016/S0006-3207\(01\)00070-2](https://doi.org/10.1016/S0006-3207(01)00070-2)

- Frelich, L. E., Montgomery, R. A., & Reich, P. B. (2021). Seven Ways a Warming Climate Can Kill the Southern Boreal Forest. *Forests*, 12(5), 560. <https://doi.org/10.3390/f12050560>
- Frelich, L.E., Blossey, B., Cameron, E.K., Dávalos, A., Eisenhauer, N., Fahey, T., Ferlian, O., Groffman, P.M., Larson, E., Loss, S.R. and Maerz, J.C., (2019). Side-swiped: ecological cascades emanating from earthworm invasions. *Frontiers in Ecology and the Environment*, 17(9), pp.502-510.
- Frerker, K., Sabo, A. and Waller, D., (2014). Long-term regional shifts in plant community composition are largely explained by local deer impact experiments. *PloS one*, 9(12), p.e115843.
- Friedman, S. K., & Reich, P. B. (2005). Regional Legacies of Logging: Departure from Presettlement Forest Conditions in Northern Minnesota. *Ecological Applications*, 15(2), 726–744. <https://doi.org/10.1890/04-0748>
- Galatowitsch, S., Frelich, L., & Phillips-Mao, L. (2009). Regional climate change adaptation strategies for biodiversity conservation in a midcontinental region of North America. *Biological Conservation*, 142(10), 2012–2022. <https://doi.org/10.1016/j.biocon.2009.03.030>
- Garrett, H.E. *et al.* (2004). Hardwood silvopasture management in North America. In: Nair, P.K.R., Rao, M.R., Buck, L.E. (eds) *New Vistas in Agroforestry*. Advances in Agroforestry, vol 1. Springer, Dordrecht. https://doi.org/10.1007/978-94-017-2424-1_2.
- Godman, R. M., & Lancaster, K. (1990). *Tsuga canadensis* (L.) Carr. Eastern hemlock. In *Silvics of North America* (pp. 604–612).
- Hale, C.M., Frelich, L.E. and Reich, P.B., (2006). Changes in hardwood forest understory plant communities in response to European earthworm invasions. *Ecology*, 87(7), pp.1637-1649.
- Hale, C.M., Frelich, L.E., Reich, P.B. and Pastor, J., (2005). Effects of European earthworm invasion on soil characteristics in northern hardwood forests of Minnesota, USA. *Ecosystems*, 8, pp.911-927.
- Hanberry, B. B., Palik, B. J., & He, H. S. (2012). Comparison of historical and current forest surveys for detection of homogenization and mesophication of Minnesota forests. *Landscape Ecology*, 27(10), 1495–1512. <https://doi.org/10.1007/s10980-012-9805-5>
- Handler, S., Duveneck, M. J., Iverson, L., Peters, E., Scheller, R. M., Wythers, K. R., Brandt, L., Butler, P., Janowiak, M., Shannon, P. D., Swanston, C., Barrett, K., Kolka, R., McQuiston, C., Palik, B., Reich, P. B., Turner, C., White, M., Adams, C., ... Ziel, R. (2014). Minnesota forest ecosystem vulnerability assessment and synthesis: a report from the Northwoods Climate Change Response Framework project. *Gen. Tech. Rep. NRS-133*. Newtown Square, PA; U.S. Department of Agriculture, Forest Service, Northern Research Station. 228 p., 133, 1–228. <https://doi.org/10.2737/NRS-GTR-133>
- Harmon, M. E., Franklin, J. F., Swanson, F. J., Sollins, P., Gregory, S. V., Lattin, J. D., Anderson, N. H., Cline, S. P., Aumen, N. G., Sedell, J. R., Lienkaemper, G. W., Cromack, K., & Cummins, K. W. (1986). Ecology of Coarse Woody Debris in Temperate Ecosystems. In *Advances in Ecological Research* (Vol. 15, pp. 133–302). Elsevier. [https://doi.org/10.1016/S0065-2504\(08\)60121-X](https://doi.org/10.1016/S0065-2504(08)60121-X)

Hartman, P., & Steele, K. (2015). *Ecological Site Description: Dolomite Colluvium Bluff Prairie*. United States Department of Agriculture.

https://efotg.sc.egov.usda.gov/references/Public/MN/105XY001_DolomiteColluviumBluffPrairie.pdf

Haugen, L. M., & Ostry, M. E. (2013). Long-term impact of shoot blight disease on red pine saplings. *Northern Journal of Applied Forestry*, 30(4): 170-174., 30, 170–174. <https://doi.org/10.5849/njaf.13-024>

He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>

Holdsworth, A.R., Frelich, L.E. and Reich, P.B., (2007). Effects of earthworm invasion on plant species richness in northern hardwood forests. *Conservation Biology*, 21(4), pp.997-1008.

Holdsworth, A.R., Frelich, L.E. and Reich, P.B., (2008). Litter decomposition in earthworm-invaded northern hardwood forests: role of invasion degree and litter chemistry. *Ecoscience*, 15(4), pp.536-544.

Holdsworth, A.R., Frelich, L.E. and Reich, P.B., (2012). Leaf litter disappearance in earthworm-invaded northern hardwood forests: role of tree species and the chemistry and diversity of litter. *Ecosystems*, 15, pp.913-926.

Holmes, S. A. (2009). *The influence of concurrent disturbances on plant community dynamics in northern hemlock-hardwood forests* (304950493) [Ph.D., Michigan Technological University].

<http://library.dnr.state.mn.us/login?url=https://www.proquest.com/dissertations-theses/influence-concurrent-disturbances-on-plant/docview/304950493/se-2?accountid=28003>

Hua, F., Bruijnzeel, L. A., Meli, P., Martin, P. A., Zhang, J., Nakagawa, S., Miao, X., Wang, W., McEvoy, C., Peña-Arancibia, J. L., Brancalion, P. H. S., Smith, P., Edwards, D. P., & Balmford, A. (2022). The biodiversity and ecosystem service contributions and trade-offs of forest restoration approaches. *Science*, 376(6595), 839–844. <https://doi.org/10.1126/science.abl4649>

Knapp, W. M., & Wiegand, R. (2014). Orchid (Orchidaceae) decline in the Catoctin Mountains, Frederick County, Maryland as documented by a long-term dataset. *Biodiversity and Conservation*, 23(8), 1965–1976. <https://doi.org/10.1007/s10531-014-0698-2>

Lawson, E. L. (1942, January). What happened to the hemlock? *Minnesota Conservation Volunteer*, 3(16), 64–66.

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>

Maerz, J.C., Nuzzo, V.A. and Blossey, B., (2009). Declines in woodland salamander abundance associated with non- native earthworm and plant invasions. *Conservation Biology*, 23(4), pp.975-981.

- Matonis, M.S., Walters, M.B. and Millington, J.D., (2011). Gap-, stand-, and landscape-scale factors contribute to poor sugar maple regeneration after timber harvest. *Forest Ecology and Management*, 262(2), pp.286-298. DNR 2024a
- McCay, T.S. and Scull, P., (2019). Invasive lumbricid earthworms in northeastern North American forests and consequences for leaf-litter fauna. *Biological Invasions*, 21, pp.2081-2093.
- McGraw, J. B., & Furedi, M. A. (2005). Deer Browsing and Population Viability of a Forest Understory Plant. *Science*, 307(5711), 920–922. <https://doi.org/10.1126/science.1107036>
- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). (2025). Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state’s greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/Iraq-3sy25.pdf>
- Mooers, H. D., & Watrus, N. J. (2017). *Results of Deer Creek, Carlton County, Minnesota, Groundwater Seepage Investigation*. <https://hdl.handle.net/11299/183617>
- Nowacki, G. J., & Abrams, M. D. (2008). The Demise of Fire and “Mesophication” of Forests in the Eastern United States. *BioScience*, 58(2), 123–138. <https://doi.org/10.1641/B580207>
- Nuttle, T., Ristau, T.E. and Royo, A.A., (2014). Long-term biological legacies of herbivore density in a landscape-scale experiment: forest understoreys reflect past deer density treatments for at least 20 years. *Journal of Ecology*, 102(1), pp.221-228
- Olson, J., Stearns, F., & Nienstaedt, H. (1959). Eastern Hemlock: Growth Cycle and Early Years. <https://Portal.Ct.Gov//Media/CAES/DOCUMENTS/Publications/Circulars/C205pdf.Pdf?La=en>.
- Palik, B. J., & Kastendick, D. N. (2023). Restoration of woody plant communities in red pine plantations on abandoned agricultural land in northern Minnesota, USA. *Forest Ecology and Management*, 541, 121095. <https://doi.org/10.1016/j.foreco.2023.121095>
- Parke, P. S., & Larson, N. J. (1999). The boundary waters windstorm of July 4, 1999. *NWS, Duluth, MN*. [Available online at www.crh.noaa.gov/dlh/science/].
- Paudel, S., Longcore, T., MacDonald, B., McCormick, M.K., Szlavecz, K., Wilson, G.W. and Loss, S.R., (2016). Belowground interactions with aboveground consequences: Invasive earthworms and arbuscular mycorrhizal fungi. *Ecology*, 97(3), pp.605-614.
- Perry, D., & Amaranthus, M. (1997). Disturbance, recovery, and stability. In K. A. Kohm & J. F. Franklin (Eds.), *Creating a forestry for the 21st century: The science of ecosystem management* (pp. 31–56). Island Press.

- Perry, E. (2024). *Eastern hemlock (Tsuga canadensis) monitoring results at Hemlock Ravine* (p. 24). Minnesota Department of Natural Resources.
- Pollack, K. D. (2024). *Hunting for Hemlock: Locating, restoring, and expanding eastern hemlock (Tsuga canadensis (L.) Carr.) presence in Minnesota*. University of Minnesota.
- Radeloff, V. C., Hammer, R. B., Stewart, S. I., Fried, J. S., Holcomb, S. S., & McKeefry, J. F. (2005). The Wildland–Urban Interface in the United States. *Ecological Applications*, 15(3), 799–805. <https://doi.org/10.1890/04-1413>
- Rooney, T. P., & Waller, D. M. (1998). Local and regional variation in hemlock seedling establishment in forests of the upper Great Lakes region, USA. *Forest Ecology and Management*, 111(2–3), 211–224. [https://doi.org/10.1016/S0378-1127\(98\)00333-8](https://doi.org/10.1016/S0378-1127(98)00333-8)
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Sabo, A.E., Frerker, K.L., Waller, D.M. and Kruger, E.L., (2017). Deer-mediated changes in environment compound the direct impacts of herbivory on understorey plant communities. *Journal of Ecology*, 105(5), pp.1386-1398
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2025). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>
- Sonter, L. J., Ali, S. H., & Watson, J. E. M. (2018). Mining and biodiversity: Key issues and research needs in conservation science. *Proceedings of the Royal Society B: Biological Sciences*, 285(1892), 20181926. <https://doi.org/10.1098/rspb.2018.1926>
- Stearns, F. W. (1997). *History of the Lake States Forests: Natural and human impacts*. (General Technical Report Nos. 8–29). United States Department of Agriculture Forest Service.
- Tyrrell, L. E., & Crow, T. R. (1994). Dynamics of dead wood in old-growth hemlock–hardwood forests of northern Wisconsin and northern Michigan. *Canadian Journal of Forest Research*, 24(8), 1672–1683. <https://doi.org/10.1139/x94-216>
- Wiegmann, S. M., & Waller, D. M. (2006). Fifty years of change in northern upland forest understories: Identity and traits of “winner” and “loser” plant species. *Biological Conservation*, 129(1), 109–123. <https://doi.org/10.1016/j.biocon.2005.10.027>
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S.

Geological Survey Scientific Investigations Report 2023–5064, p. 55).

<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Zamora, D., Wyatt, G., Buttler, M., Ford, M., Magner, J., Reichenbach, M., Burkett, E., Current, D., Walter, D. (2017). Silvopasture: Establishment and Management Principles for Minnesota. Retrieved from the University Digital Conservancy, <https://hdl.handle.net/11299/214976>.

Ziemba, J.L., Hickerson, C.A.M. and Anthony, C.D., (2016). Invasive Asian earthworms negatively impact keystone terrestrial salamanders. *PloS one*, 11(5), p.e0151591.

Upland Deciduous Forests and Woodlands (Fire-dependent)

Habitat Description

Upland deciduous forests, in the fire-dependent ecological system, are found throughout the Laurentian Mixed Forest and Eastern Broadleaf Forest provinces, and much less commonly in the Prairie Parkland and Tallgrass Aspen Parklands provinces. Fire-dependent forests can include aspens (*Populus spp.*), oaks (*Quercus spp.*), birches (*Betula spp.*), pines (*Pinus spp.*), spruce (*Picea spp.*), and balsam fir (*Abies balsamea*). Please see the Upland Conifer Forest sub-chapter for information on conifer-dominated forests. Soils, landforms, and disturbances, primarily fire, drive the distribution, patch size, ages, and composition of upland forests.

Because the forest canopy blocks most of the sunlight before it reaches the understory, most understory plants have evolved some degree of shade tolerance. The amount of light reaching the understory also has a strong effect on the structural complexity of the forest. As these forests age, they become structurally more complex as canopy trees die, producing gaps in the canopy that allow growth of less shade-tolerant shrubs and trees. Standing large dead trees become more frequent, eventually becoming large down logs, which add to the structural diversity of older forests. Fire-dependent forests tend to have a more open canopy than Mesic Hardwood forests and a denser layer of shrubs such as hazelnuts (*Corylus spp.*), dogwoods (*Cornus spp.*), and mountain maple (*Acer spicatum*).

Fire-dependent upland deciduous forests include forests with a blend of dominance by tree species including oaks, aspen, and maples. Although there is a continuum, we can break the narrative to focus on oak-dominant and aspen-dominant.

Oak-dominant forests and woodlands:

Upland hardwood oak forest habitats occur on xeric (dry) to relatively mesic (moist) forest sites throughout the Eastern Broadleaf Forest Province. This habitat is found in portions of the western half of the Laurentian Mixed Forest Province but is uncommon in the northeast. On drier sites, northern pin oak (*Quercus ellipsoidalis*), bur oak (*Q. macrocarpa*), white oak (*Q. alba*), and, in the southeast, black oak (*Q. velutina*) are important canopy species. Associated canopy trees include black cherry (*Prunus serotina*), paper birch (*Betula papyrifera*), aspens, and shagbark hickory (*Carya ovata*). These forests occur on nutrient-poor, well-drained sandy soils on outwash plains, river terraces, and beach ridges, and in the past were strongly influenced by fire. The canopy of modern dry oak forests is relatively open, allowing for a dense shrub layer, typically dominated by american hazel (*Corylus americana*). Many of the drier oak forests contain open-grown trees indicative of a more open woodland or savanna prior to fire suppression.

Aspen-dominated Forest and Woodlands (see also Mesic Hardwoods for Aspen Forests)

Fire-dependent upland hardwood forest (aspen) is characterized by a canopy dominated by quaking aspen (*Populus tremuloides*), big-toothed aspen (*P. grandidentata*), paper birch, or a mixture of these species. These shade-intolerant tree species are the dominant trees in the early stages of a wide variety of native plant communities in fire-dependent and mesic hardwood forest systems. Thus, aspen forest is a cover type that may eventually develop into many other native plant communities. In the absence of disturbance, such as fire or timber harvest, aspen acres may decline across the landscape as they succeed to longer-lived species.

Aspen forests typically have a nearly complete canopy of aspen or birch, but the canopy is not as dense as that of sugar maple. As a result of higher light levels penetrating the canopy, these forests usually have a well-developed shrub layer dominated by hazelnuts (*Corylus* spp.) or dogwoods (*Cornus* spp.). The coverage and diversity of the herbaceous plant layer are highly variable depending on site conditions and stand history.

As aspen forests age, they typically increase in structural diversity. Historically, most aspen stands in northern Minnesota had a conifer component, which increased as the stand aged. Today, most aspen stands have little or no conifer understory, due to past management and slash fires. Still, many older aspen stands are relatively structurally diverse, with large trees, snags, down logs, and an understory containing more shade-tolerant hardwoods or conifers that will become the canopy dominants in the absence of a stand-replacing disturbance.

Habitat Map

To depict Upland Deciduous Forest and Woodland habitat (see Figure 3.9), we compiled spatial data from several sources: DNR's Native Plant Communities and the Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological System

In the Fire-dependent Forest system, some Native Plant Community Classes split up at the level of Types, with some fitting into this Upland Deciduous Forest and Woodland Chapter with others in the Conifer Forest and Woodland Chapter. For the classes that split, we specify in parentheses the Types that we would include here.

Fire-dependent Forest

[FDc25-Central Dry Oak-Aspen \(Pine\) Woodland](#)

FDc25b Oak – Aspen Woodland

[FDc34-Central Dry-Mesic Pine-Hardwood Forest](#)

FDc24b Oak – Aspen Forest

[FDn22-Northern Dry-Bedrock Pine \(Oak\) Woodland](#)

FDn22c Pin Oak Woodland (Bedrock)

[FDn33-Northern Dry-Mesic Mixed Woodland](#)

FDn33b Aspen – Birch Woodland

[FDn43-Northern Mesic Mixed Forest](#)

FDn43b Aspen – Birch Forest

[FDs27-Southern Dry-Mesic Pine-Oak Woodland](#)

FDs27c Black Oak – White Oak Woodland (Sand)

[FDs36-Southern Dry-Mesic Oak-Aspen Forest](#)

[FDs37-Southern Dry-Mesic Oak \(Maple\) Woodland](#)

[FDs38-Southern Dry-Mesic Oak-Hickory Woodland](#)

[FDw24-Northwestern Dry-Mesic Oak Woodland](#)

[FDw34-Northwestern Mesic Aspen-Oak Woodland](#)

[FDw44-Northwestern Wet-Mesic Aspen Woodland](#)

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Midwest Oak – Hickory Forest, Eastern North American Native Ruderal Forest, and Northern Tallgrass Oak – Aspen Woodland.




Figure 3.9. A map depicting Upland Deciduous Forest and Woodland habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological Systems of Fire-dependent Forest/Woodland as well as three types from the Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

Historically, fire was a particularly important disturbance agent in upland forests. With the decline of fire frequency post Euro-American settlement, most upland deciduous forests remain on sandy, gravelly, or otherwise droughty sites where succession to closed-canopy mesic hardwood forest communities has been slowed by harsh growing conditions. In the past, fires in the deciduous woodlands could kill stands of trees and other aboveground vegetation under the right climate, fuel, and topographic settings. However, even intense fires in these deciduous woodlands did not generate the kinds of conflagrations possible in closed-canopy coniferous forests. By comparison, fires in upland deciduous woodlands were more regenerative than destructive.

Historically, wildfire played an important role at both the site and landscape level in these forests, but due to land use changes and fire suppression, wildfire is less common today. Following a crown fire, a variety of biological legacies, or features of the pre-disturbance ecosystem that persist after disturbance, remain in the young regenerating forest (Perry and Amaranthus 1997). These biological legacies, which include standing dead trees, large trees that escaped the fire, down logs, small patches of unburned vegetation, and surviving propagules and organisms (such as buried seeds, surviving roots, mycorrhizal fungi and other soil microbes and invertebrates) are important ecologically and can provide habitat features for SGCN. At the landscape level, wildfires create a shifting mosaic of native plant communities with a variety of ages and structural characteristics that provide habitats for SGCN.

Plants that occur in this community have seeds or vegetative structures that can survive fire and are good at colonizing burned sites. Many plants are opportunists that can take advantage of the short periods following fire when nutrients are relatively abundant and light levels are high. Such plants must also survive frequent drought and potentially long periods between fires when light levels decrease beneath increasingly dense shrub and tree canopies. The most evident characteristic of plants in this community is their ability to sprout prolifically. The trees, shrubs, and many of the herbs can store considerable amounts of carbohydrates below ground in roots, rhizomes, or other specialized organs and then sprout vigorously after aerial stems are destroyed by fire. These plants seem to be particularly plastic in allocating resources to underground or aboveground tissues, depending on the impact of fire on their overall vigor.

With fire less frequent in these forest stands, forest management activities including timber harvests are now the most prevalent form of disturbance. Sustainable forest management includes planning when and where timber harvest can be used as an effective tool to address a range of values and priorities, including forest health and productivity, critical ecological functions, and habitat restoration from historical disturbance events. During timber harvest planning, it's important to consider forest compositional and structural diversity at meaningful scales and pursue efforts to replicate natural disturbance regimes that define these plant communities. Planning efforts must strive to ensure that a balance of forest habitat conditions (e.g. younger and older forest, small and large patch sizes) are available for plant and animal species across the landscape over time.

Conversion to other uses such as agriculture, urban development, and transportation corridors, has fragmented forests across the state; this is most evident in southeastern and central Minnesota where few large patches of upland forest remain. The remaining forests in these regions typically lack the ecological complexity of pre-European settlement forests due to several factors such as, grazing, invasive plants and

animals, edge effects, changes in native animal populations, and consumptive uses. Even in the largely forested areas of northern Minnesota, rural sprawl has greatly reduced the extent of large, contiguous forest areas; most forest areas are within 25 kilometers (15.5 miles) of small housing settlements (Radeloff et al., 2005).

Today, aspen forest is the most abundant forest habitat in Minnesota and much more widespread than it was prior to settlement by people of European descent. An analysis of General Land Office bearing tree records from the late 1800s and Forest Inventory and Analysis plots from the 1990s shows that in the Laurentian Mixed Forest Province, aspen forest communities have increased nearly tenfold (Friedman and Reich 2005). However, it is thought that aspen forests are structurally less diverse than they were historically, and on average younger than that of pre-European settlement aspen forests. Currently, about 60% of the aspen age class distribution is 1-40 years old, and about 18.5% of the cover type is over 60 years old statewide (US Department of Agriculture, Forest Service, FIA Program, 2025). The location of older aspen stands shift across the landscape over time as old stands are harvested or succeed to other tree species, and young-mature aspen stands that aren't harvested age. Older, more structurally diverse aspen forests have high value to some SGCN, such as American goshawk (*Astur atricapillus*). The configuration of that habitat on the landscape (e.g. tract size, proximity to other forest types) also relates to its suitability and significance as habitat for SGCN.

Historically, fires in the oak habitat were more regenerative than destructive. The typical cycle involved top-killing of plants and vegetative recovery by resprouting. Fires enhanced plant reproduction by exposing mineral soil, triggering seed dispersal, breaking seed dormancy, and increasing light and heat conditions on the ground. In the absence of fire, relatively mesic or fire-sensitive species such as bitternut hickory (*Carya cordiformis*), basswood (*Tilia americana*), and red maple (*Acer rubrum*) are increasing in abundance in this habitat. Without fire, there is little natural oak regeneration in most dry oak stands.

Some oak forest stands, particularly if disturbed by grazing or selective logging, have been colonized by invasive species such as buckthorn (*Rhamnus cathartica*) and tartarian honeysuckle (*Lonicera tartarica*). Like other forest habitats, most oak forest habitats in the Eastern Broadleaf Forest Province and southern and western portions of the Laurentian Mixed Forest Province have been fragmented by agriculture and development.

Species in Greatest Conservation Need

Upland deciduous forests and woodlands provide primary or secondary habitat for 68 animal and 33 plant SGCN (see Table 3.8). Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Table 3.8. Numbers of Species in Greatest Conservation Need associated with the Upland Deciduous Forest and Woodlands as either primary or secondary habitat¹

Species Group	Primary Habitat	Secondary Habitat	Total
Amphibians	3	0	3
Birds	3	9	12
Mammals	2	20	22
Reptiles	9	0	9
Terrestrial Invertebrates			
Bees			9
Beetles (terrestrial)			1
Moths			4
Snails (terrestrial)	3	3	6
Spiders	1	1	2
Plants	33		33
Total	54	33	101

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants).

Of the 131 terrestrial vertebrate SGCN identified in the 2025-2035 SWAP, 35% (46) occur in upland deciduous forests and woodlands. A large proportion (69%) of the 32 mammal SGCN occur in this habitat. Like other fire-dependent habitats, plants found in the upland deciduous forests only represent 7% of the 447 that are designated as SGCN. Examples of selected SGCN are described below.

Amphibians

Three salamander SGCN – [spotted \(*Ambystoma maculatum*\)](#), [four-toed \(*Hemidactylium scutatum*\)](#), and Eastern newt (*Notophthalmus viridescens*) – are strongly associated with mature forest habitats. They primarily occupy deciduous forests but will also use mixed deciduous-conifer forest types. Access to wetlands, such as vernal pools, are critical for these species, which use them in the spring to breed. Eggs are laid in or near the water, and larvae remain in the pools until metamorphosis occurs. Four-toed salamanders specifically lay their eggs in moss hummocks next to wetlands. Eggs are laid in or near the water, and larvae remain in the pools until

metamorphosis occurs. Four-toed salamanders specifically lay their eggs in moss hummocks next to wetlands. Four-toed salamanders occur most frequently in mature upland forests on glacial moraine landscapes with frequent isolated wetlands that include an alder margin and moss hummocks adjacent to pockets of open water.

Birds

SGCN birds more typical of Mesic Hardwood Forest can also occupy dry-mesic fire-dependent forests in certain landscape settings (e.g., when part of a larger forested site where forest types co-occur, depending on slope, aspect, and soils). For example, dry-mesic oak forests often occur on south- or west-facing slopes, forming an important component of forested stream valleys important to [acadian flycatchers \(*Empidonax virescens*\)](#), [cerulean warblers \(*Setophaga cerulea*\)](#), and [louisiana waterthrushes \(*Parkesia motacilla*\)](#). Such forests must have the same structural components important to these species as Mesic Upland Forest (i.e., mature trees, closed-canopy, structurally diverse understory/subcanopy, etc.).

Similarly, SGCN found in savanna (e.g., eastern whip-poor-will (*Antrostomus vociferus*), red-headed woodpecker (*Melanerpes erythrocephalus*) or dry conifer woodlands also occur in semi-open deciduous woodlands that have the necessary habitat components. Dry sand or gravel soils and an open ground layer are important to whip-poor-wills, while red-headed woodpeckers require dead trees and snags without bark (see Savanna chapter).

Young (early successional) upland deciduous forest, including edges, is an important component of golden-winged warbler (*Vermivora chrysoptera*) habitat. However, a mosaic of seral stages is important, including mature forest. Suitable early successional habitat may occur in several forest types – not just Upland Deciduous. Aspen and birch are frequent dominants, although other tree species can be suitable for golden-winged warblers. Early successional habitat may occur in monocultures following disturbance (e.g., fire, windthrow, silvicultural activities), in naturally fragmented mosaics (e.g., upland forest interspersed with wetland openings, shrub swamps, lowland conifers, lake or river shores) as well as along road or utility corridors through forested landscapes.

Mammals

Upland Deciduous Forests and Woodlands support a very similar mammalian SGCN fauna as Upland Conifer Forests. Once again, all 7 bat SGCN are reported in Upland Deciduous Forests but it is considered primary habitat for just one, the Tri-colored bat (*Perimyotis subflavus*). This cave-hibernating species commonly roosts during the spring, summer and fall among the leaves of live or recently dead hardwood trees. As noted by the [USFWS](#), tri-colored bats have been devastated by the impacts of white-nose syndrome, causing an estimated decline of more than 90% in the species' hibernating colonies across North America. Two other species that are more likely to be seen in Upland Deciduous Forests include the [North American least shrew \(*Cryptotis parvus*\)](#) and plains [spotted skunk \(*Spilogale interrupta*\)](#). Both species are also found in a range of other habitats, especially open or brushy edge habitats.

Reptiles

Eastern hog-nosed snakes (*Heterodon platirhinos*) are most common in and around woodland edge habitats. Key habitat features for these species include down woody debris (for cover, nesting sites, and basking sites)

and burrows or crevices as overwintering sites. Eight other reptile SGCN's also utilize Upland Deciduous Forests and Woodlands as primary habitat including six snakes: [North American racer \(*Coluber constrictor*\)](#), [timber rattlesnake \(*Crotalus horridus*\)](#), Eastern hog-nosed snake (*Heterodon platirhinos*), smooth greensnake (*Opheodrys vernalis*), [Western ratsnake \(*Pantherophis obsoletus*\)](#), and gophersnake (*Pituophis catenifer*). All of these species can be found in a wide variety of habitats. The North American racer, [five-lined skink \(*Plestiodon fasciatus*\)](#) and timber rattlesnake use deciduous forests and woodlands as foraging habitat, if they are close to rock outcrops and cliffs that are important for basking, overwintering and denning.

Plants

Fire dependent upland forests and woodlands are habitat to a number of unique and rare plant species that help make up the complexity of these native plant communities (see Appendix D). For example, [black huckleberry, \(*Gaylussacia baccata*\)](#), is a rare, fire adapted species with a limited range in Minnesota. Much of its habitat is fragmented and lacks the fire needed to maintain its habitat. Without fire, fire-sensitive and more shade tolerant species can invade black huckleberry's habitat. Prescribed fires are needed to approximate natural wildfires and allow black huckleberry to persist.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the Stressors section in Chapter 1. Species in Greatest Conservation Need.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Residential, commercial, and industrial development can all directly alter forest ecosystems by removing, degrading, or fragmenting forest cover and habitat. Fragmentation of forests is a growing concern, and occurs when a large, contiguous forest land mass is divided into smaller tracts through sale and subdivision, road construction, or clearing of forests for agriculture, housing, or other commercial developments. This often creates smaller forest pockets, which are interspersed with non-forest land activities. Fragmentation can inhibit the natural migration of many plant and animal species, increase the risk of wildfires at the wildland urban interface, restrict public recreational access, reduce the habitat value and ecological complexity of forest lands, and contribute to the spread of invasive species. Because of their natural beauty, diversity, recreation opportunities, and abundant natural resources, forested landscapes are appealing places to live; even in the largely forested areas of northern Minnesota, rural development has greatly reduced the extent of large, contiguous forest areas; most forest areas are within 25 kilometers (15.5 miles) of small housing settlements (Radeloff et al., 2005). Forested lands adjacent to lakes and rivers are especially attractive sites for development.

Development can also indirectly impact forests, through modification of ecosystem hydrology, introduction of nonnative species, altered disturbance regimes, reduced connectivity to other habitats, increased abundances of competitors and predators, altered microhabitat conditions, increased direct human disturbance, and avoidance by wildlife that evolved to use large chunks of habitat embedded in a landscape dominated by native ecosystems.



Crop Production

Historically, many of the upland deciduous and mesic hardwood forests were converted to cropland, leaving a patchwork of widely scattered wood lots, averaging 40 to 80 acres in size, across central and southeastern Minnesota. The conversion of forests to agricultural use is ongoing in some locations.



Livestock Management

Minnesota currently has over 650,000 acres of mesic, upland deciduous, and riparian forests on farms that are grazed by livestock (Zamora et al., 2017, Garrett et al., 2004). Properly managed grazing can stimulate plant diversity and create diverse vegetative structure across the landscape. Overgrazing, if left unchecked, can reduce diversity in vegetative cover and structure, and have severe negative impacts on biodiversity, ecosystem function, soil health, and forage production.

Historically, grazing by livestock in forested landscapes likely became very intensive during the droughts of the 1930's as farmers struggled to find enough forage (Hartman & Steele, 2015). These habitats likely became overgrazed and became subject to varying degrees of erosion, while the forest vegetation was subject to trampling, grazing, fraying, and bark stripping all of which can impact regeneration. (Hartman and Steele, 2015). From the 1960's to present, grazing practices have changed. This includes reductions in the types of livestock raised by agricultural producers to primarily cattle, as well as the elimination of grazing on many bluff prairies and associated savannas altogether. This reduction in grazing has led to an increase of woody vegetative cover and the reduction of bluff prairie and savanna habitats. Best management practices associated with silvopasture, or the practice of intentionally combining management of trees, forage, and grazing as one integrated practice, has the potential to contribute to the restoration of Oak-dominant forests and woodland habitats.



Mining and Quarrying

Forests, when they co-occur with mineral, oil, or gas deposits, are at risk of degradation due to the direct and indirect impacts of mining activities such as permanent forest loss, soil compaction, introduction of invasive species, human disturbance, waste discharge and pollution, and forest fragmentation (Sonter et al., 2018). Mining projects can trigger additional, cascading land use change with the establishment of associated infrastructure such as improved road access.



Roads, Trails, and Railroads

Roads, trails, and railroads can create fragmentation and edge effects in large tracts of forest. Increase in the number of roads and trails, particularly for motorized vehicles, increases human disturbance and introduction of invasive species further into the forest. Repeated use and soil compaction associated with roads and trails also can modify the vegetation within the area of use. Temporary winter logging trails may become rutted while materials used to stabilize the trails for heavy vehicles may remain in place, leaving behind wood planks and wood chips. Trails that provide winter access to upland deciduous stands can disturb forest wildlife, especially species that begin nesting in late winter, such as many forest raptors.



Utility Corridors

Utility corridors transect forested habitats throughout the state. These corridors can contain a variety of utility infrastructure. This includes pipelines, telephone lines, and power lines. These lines can be aerial, above ground, or below ground. Regardless of type, utility corridors can fragment habitat based on their size, purpose, and the upkeep that is required to maintain them.

The initial construction of the associated utility infrastructure as well as the long-term maintenance of that infrastructure can have lasting impacts on native plant communities and the SGCN associated with them. Increased funding for rural broadband, conversion of aboveground powerlines to belowground, and development of new utilities (solar, wind, etc.) continue to threaten forested habitats.

These corridors and the increased travel from vehicles and equipment that they require act as vectors for invasive plant species. The required vegetative management practices often include herbicide applications, removal of woody vegetation, and/or mowing. This increased disturbance increases the likelihood of establishing invasive plant species within the corridors and then spreading to adjacent lands.



Timber Harvest

Even-aged harvests (e.g. clear or seed-tree cuts) are mostly used in fire-dependent forests and are similar to stand replacing fires of historical disturbance regimes. Methods to enhance biological legacies at local and landscape scales in even-aged harvests need more study and application.

There are needs for providing sufficient late-successional upland deciduous forest stands to balance management for production and forest health that removes trees before they are old. This is particularly notable for oaks, which become more susceptible to disease with age and can spread disease from old oaks to seedlings. Oak forests are commonly managed with full even-aged harvests early enough to provide for stump-sprouting, which creates a set of ecological characteristics that differ from the forests with more natural disturbance and regeneration patterns and might not provide optimal habitat conditions for SGCN. Many wildlife species benefit from managing for within-stand species and structural diversity in large habitat patch sizes to provide habitat for area-sensitive species and to minimize habitat fragmentation effects.



Recreation

In addition to the impacts created by recreational trails (discussed above), recreational activities themselves can proliferate invasive species. Seeds of invasive plants such as common tansy (*Tanacetum vulgare*) and garlic mustard (*Alliaria petiolata*) can become embedded in the wheels of vehicles or the shoes of individuals and eventually become established along compacted trails where conditions for their growth may be more favorable. Earthworm egg cases can also be spread in mud, which are problematic for moonwort (*Botrychium*) species. Increased use of roads and trails also increases human disturbance in the forest, reducing the amount of inaccessible area.



Fire Management

A history of fire suppression, insufficient fire frequencies, and the exclusion of cultural burning practices contribute to the ongoing homogenization of composition and structure in upland, fire-dependent forest habitats (Hanberry et al., 2012). In the absence of fire, open forests can transition into closed forest ecosystems, leading to decreased structural diversity and a decline in the diversity of associated plant and animal species. Woody densification also creates a positive feedback loop of mesophication, where cool, damp, and shaded conditions increase ecosystem moisture and further reduce flammability (Nowacki and Abrams, 2008). Oaks, in particular, have declined dramatically and have been outcompeted by fire-sensitive and shade-tolerant canopy trees (e.g., elms, boxelder, maples, American basswood, and ashes). Today, in the absence of fire, even-aged timber harvests (e.g., clear or seed-tree cuts) are frequently used in fire-dependent

forests. Oak regeneration following clearcuts can be challenging. Poor oak seedling recruitment (especially if a poor acorn crop preceded the harvest), competition with other tree species, and deer browsing can all reduce successful regeneration (Jacobson et al. 2019).



Invasive Species (Problematic Non-native Species)

Invasive plant species, such as buckthorn (*Rhamnus cathartica*) and Tartarian honeysuckle (*Lonicera tartarica*) often outcompete native plant species for light, space, and water and repress understory plant species leading to modified habitats unsuitable for SGCN. Dense canopies formed by invasive shrubs also inhibit the regeneration of native tree species due to the increase in shade. While a concern statewide, upland deciduous forests in central and southern Minnesota are currently most impacted by these invasive shrubs.

Invasive animals, primarily invertebrates, cause profound ecological changes in forests (Frelich et al., 2019). First, Eurasian earthworms upend the traditional soil profile (Hale et al., 2005; Fahey et al., 2013), remove the accumulated layers of organic leaf litter (Holdsworth et al., 2008, 2012), and alter the community of soil mycorrhizal fungi (Paudel et al., 2016). These changes cascade through the ecosystem, including ground layer vegetation (Hale et al., 2006, Holdsworth et al. 2007), leaf litter fauna (McCay and Scull, 2019), salamanders (Maerz et al., 2009, Ziemba et al., 2016), and likely other vertebrates.

Second, the spongy moth (*Lymantria dispar*) is an invasive forest pest that feeds on more than 300 species of deciduous trees and shrubs, including aspen, oak, and birch. Repeated removal of leaves, or defoliation, stresses trees and can leave them vulnerable to disease or other pest infestations that can kill trees. Once it becomes established in a location, the spongy moth has cycles of large population outbreaks every eight to 12 years, leading to widespread defoliation and nuisance from caterpillars.

Other invasive insects defoliate trees but typically do not cause significant mortality. Elm zigzag sawfly (*Aproceros leucopoda*) is an invasive defoliator of elm trees that was reported in Minnesota in July 2024 for the first time.



Problematic Native Species

Overbrowsing by deer can severely limit natural tree regeneration (Matonis et al., 2011; De Jager et al, 2013), especially for preferred species, such as oak. Elevated deer populations also reduce populations of palatable ground layer plants, including orchids (Knapp and Wiegand, 2014), members of the lily family (Balgooyen and Waller 1995; Augustine and Frelich, 1998; Fletcher et al., 2001), and rare species (McGraw and Furedi, 2005). Decades of cumulative impacts have resulted in profound changes to the composition of native plant communities (Wiegmann and Waller, 2006; Frerker et al., 2014; Nuttle et al., 2014; Flagel et al., 2016; Anderson et al., 2017; Sabo et al., 2017). The effects of deer browsing often interact with those of invasive earthworms (Fisichelli et al., 2013; Davalos et al. 2014).



Diseases and Pathogens

Oaks are susceptible to several types of disease and pathogens, such as oak wilt (caused by the invasive fungus *Bretziella fagacearum*), sudden oak death (caused by the water mold *Phytophthora ramorum*), and bur oak blight (caused by a native fungal pathogen, *Tubakia iowensis*). Individual trees tend to be more

susceptible when under stress, which may interact then with other threats, such as if future climatic conditions bring more frequent late summer droughts. Older oak trees are more susceptible to interacting stressors including drought, heavy seasonal precipitation, insects, and diseases. Wetter springs may increase the prevalence and severity of fungal diseases as spores spread from the diseased leaves by rainwater, exacerbated by earlier spring rainstorms brought on by Minnesota's changing climate.



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years ([Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025](#)). Furthermore, these shifts in thermal regimes are ecologically significant. Warming can indirectly affect all of Minnesota's forest communities by altering the impact of invasive species, insect pests and diseases, winter burn, drought, and deer herbivory (Galatowitsch et al. 2009; Handler et al. 2014).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Restoration activities could also be disrupted by causing seedling mortality and shifts in community composition if there are multiple seasons with insufficient rain. Short funding windows for restoration efforts exacerbate the problem by inducing restoration plantings even in dry periods. Prolonged drought has stressed trees in the south east making some particularly susceptible to native pests and pathogens. Changes in precipitation interact with the age class distribution of tree species such as oak, in which old trees are more susceptible to stress from drought, heavy seasonal precipitation, in turn making trees vulnerable to insects and diseases.

Heavy rain events have been increasing in frequency and severity in Minnesota in recent decades. Forests on steep slopes, common in the state’s southeastern counties, are vulnerable to landslides during these events.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan (Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN), five strategies were identified:

- 

Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.
- 

Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.
- 

Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.
- 

Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.
- 

Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.


Examples of conservation actions are grouped below under these five strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Also note that some strategies, such as Strategy 3, collaborating with partners, could truly be applied to all actions to most broadly and effectively implement them.

Potential Conservation Actions for Upland Deciduous Forests and Woodlands



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action

	Consider projected effects on areas of biological significance such as high value conservation forests, old-growth forests, and Important Bird Areas when making decisions and policy that are projected to impact upland deciduous forests (e.g., mining and quarrying or oil and gas permitting, planning for new development of energy sources and any associated new road or trail construction).

Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

Stressor	Action
	Explore the potential use of silviopasture management regimes (the practice of intentionally combining management of trees, forage, and grazing as one integrated practice) to help restore some upland deciduous forests that do not spring ephemerals or other sensitive SGCN species (UMN Extension: Silvopasture: Establishment and Management Principles for Minnesota 2013)
	Attempt to minimize forest fragmentation when planning the location of new roads and trails.
	Encourage the use of existing corridors for new trails, roads, and utility corridor development to minimize the construction of new roads and trails that would increase forest fragmentation.
	Apply knowledge of SGCN habitat use into landscape forest planning efforts, to help balance habitat for SGCN and other values according to management requirements and objectives. Consider natural area designations and other ways to enhance and preserve key mature stands. Conduct research and model anticipated changes with different forest management scenarios for impacts on wildlife habitat. Consider the ecological values of various forest ages and how they relate to SGCN habitat when determining desired future conditions and planning sustainable management to advance those conditions.
	Limit the use and operation of heavy equipment when road or site conditions are vulnerable to damage. For instance, depending on the site characteristics, conducting timber harvests during frozen ground conditions can reduce soil compaction and rutting impacts.

	Provide information on important habitat types, landscape configurations, and conditions most beneficial for SGCN. Support forest management that reflects ecological disturbance goals and SGCN habitat needs such as stable amounts of older forest. Communicate and document the benefits of alternative management techniques (i.e., variable density thinning, natural seeding), reference to natural disturbance intervals, natural regeneration methods, and to increase habitat quality for SGCN. Promote consideration of more flexibility in stocking goals in natural stands (i.e., longer time frames to meet goals, non-crop trees can be a significant portion of goals) and the ability to protect really important biodiversity areas from harvest.
	Retain old and dead standing trees (snags) in forests when practical as habitat for wildlife, including cavity-nesting birds.
	Where people recreate, provide outreach and education on cleaning gear and equipment to reduce the introduction and spread of invasive species. DNR's Prevent the spread webpage includes specific actions for different land based activities like biking, hiking and off-highway vehicle riding. At trailheads, add invasive species prevention messages, such as those on the boot brush kiosks developed by the PlayCleanGo: Stop Invasive Species in Your Tracks program.
	Implement control strategies for invasive species, such as buckthorn, and ensure sufficient timeframe for follow-up management that extends for many years. Support funding for these activities, and consider expanding restoration resources to make sure they can be applied to woodland systems, which are sometimes in between funding opportunities aimed at prairies or forests.

	Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

Stressor	Action
	Support local governments in measures that minimize fragmentation and encroachment into high quality forests and woodlands such as protecting high quality areas, zoning that guides housing developments and lot sizes that support local forests, and land use planning and development guidance.
	Collaborate across forest ownerships and managers to implement sustainable forestry at the site and landscape level to create habitat conditions important to SGCN, such as unfragmented older mesic hardwood forests to create larger forest patches and reduce forest fragmentation. Support forest management approaches that are informed by ecological processes, increase structural and compositional complexity and diversity, and assist in resilience towards future stressors (e.g., climate change, disease outbreaks,

	invasive species). Promote forest management strategies that mimic landscape disturbance patterns (for example, retaining large blocks of mature habitat and creating more small forest openings as gaps or strips), maintain or enhance connectivity among forest patches to enable wildlife movement, and retain biological legacies (at site level) such as large trees with cavities. Consult Native Plant Community Classification silvicultural strategies for forest stand prescriptions .
	Engage in forest planning efforts and support the collection of accurate and reliable forest stand inventory data for planning. On state lands, participate in Section Forest Resource Management Plans (SFRMPs) and designation of Management Opportunity Areas (MOAs) address values such as biodiversity, rare features, diversity of native plant community growth stages, and wildlife needs that can't be addressed through individual site-level management (DNR 2025). In addition, High Conservation Value Forests identified through the Forest Stewardship Council certification (FSC 2025) highlight “globally, regionally or nationally significant concentrations of biodiversity values, including rare, threatened, and endangered species and their habitats,” or “globally, regionally or nationally significant large landscape level forests...” or “rare, threatened or endangered ecosystems” (DNR 2015).
	Manage for a diversity of forest growth stages across the landscape to promote vegetation characteristics that maintain habitat for a wide range of plant and animal species, promote resiliency to insect and disease outbreaks, and mitigate the adverse effects of climate change. Ensure mature, older forest and old forest qualities are retained on the landscape in amounts and areas that provide critical habitat to SGCN animals and plants; encourage management that sustains mature and old-growth forests in forest planning efforts. Apply management practices to create young, early successional forest on the landscape to provide vegetation characteristics and edge habitat between young and mature forest that support plant and animal species.
	Droughts are becoming more frequent and can challenge the success of restoration plantings; extending the duration of grants and contracts for restoration projects could help to allow conservation practitioners to plant when conditions are more favorable.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions

Stressor	Action
	Monitor the responses of SGCN to various types of forest management at the site-level and landscape-scale. Collaborate with MFRC site-level guideline monitoring and NPC silvicultural case studies to incorporate SGCN monitoring (Silviculture Case Studies). Develop models of habitat configurations for various SGCN and consider how current

	harvest forecasts and conversion rates (i.e., one forest type to another; natural origin habitat to plantation) relate to those models in terms of sustainability for maintaining habitat for SGCNs.
	Collaboratively explore techniques to promote forest regeneration for oak, with an emphasis on forests of high conservation value.
	Explore opportunities to burn upland deciduous forest stands with a significant oak component to fill o investigate and then monitor the results for oaks as well as other vegetative components of the forest stands. Include other considerations to improve outcomes such as a good acorn crop year prior to the burn and management to stem the growth of competing species.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Stressor	Action
	Promote and consider further development of tax incentives that enable landowners, homeowners, and farmers to gain value for maintaining conservation values on their properties. Provide increased levels of incentive payments for woodland owners to help protect woodlands and forests from further reduction due to development pressure.
	Managing for an increased diversity of dominant species, and age classes when appropriate, in a site may assist in resilience toward future disease outbreaks, susceptibility to some pathogens, and invasive species. For instance, oak forests with a higher diversity of additional species can be more resilient to oak wilt's overall impact. For more information, see DNR Oak Wilt Management and DNR Oak Decline .
	Seek additional funds to assist with the management of woodlands and forests to increase their resilience to pathogens and invasive species. Also consider pairing invasives management with other incentivized management activities to create efficiencies.

References for Upland Deciduous Forest and Woodlands

Anderson, R.C., Anderson, M.R. and Corbett, E.A., 2017. White-tailed deer (*Odocoileus virginianus*) and fire effects on flowering diversity of tallgrass prairie forbs. *The Journal of the Torrey Botanical Society*, 144(3), pp.243-253

Augustine, D. J., & Frelich, L. E. (1998). Effects of White-Tailed Deer on Populations of an Understory Forb in Fragmented Deciduous Forests. *Conservation Biology*, 12(5), 995–1004. <https://doi.org/10.1046/j.1523-1739.1998.97248.x>

- Balgooyen, C. P. and D. M. Waller. (1995). The use of *Clintonia borealis* and other indicators to gauge impacts of white-tailed deer on plant communities in northern Wisconsin, USA. *Natural Areas Journal*, 15(4), 308-318. <https://www.cabidigitallibrary.org/doi/full/10.5555/19960610148>
- Dávalos, A., Nuzzo, V. and Blossey, B., 2014. Demographic responses of rare forest plants to multiple stressors: the role of deer, invasive species and nutrients. *Journal of Ecology*, 102(5), pp.1222-1233.
- De Jager, N.R., Cogger, B.J. and Thomsen, M.A., 2013. Interactive effects of flooding and deer (*Odocoileus virginianus*) browsing on floodplain forest recruitment. *Forest Ecology and Management*, 303, pp.11-19.
- Fahey, T.J., Yavitt, J.B., Sherman, R.E., Maerz, J.C., Groffman, P.M., Fisk, M.C. and Bohlen, P.J., 2013. Earthworm effects on the incorporation of litter C and N into soil organic matter in a sugar maple forest. *Ecological Applications*, 23(5), pp.1185-1201.
- Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weyand, D. M. Finch, C. F. Miniat, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Fisichelli, N.A., Frelich, L.E., Reich, P.B. and Eisenhauer, N., 2013. Linking direct and indirect pathways mediating earthworms, deer, and understory composition in Great Lakes forests. *Biological Invasions*, 15, pp.1057-1066
- Flagel, D.G., Belovsky, G.E. and Beyer, D.E., 2016. Natural and experimental tests of trophic cascades: gray wolves and white-tailed deer in a Great Lakes forest. *Oecologia*, 180, pp.1183-1194
- Fletcher, D. J., Shipley, L. A., McShea, W. J., & Shumway, D. L. (2001). Wildlife herbivory and rare plants: The effects of white-tailed deer, rodents, and insects on growth and survival of Turk's cap lily. *Biological Conservation*, 101(2), 229–238. [https://doi.org/10.1016/S0006-3207\(01\)00070-2](https://doi.org/10.1016/S0006-3207(01)00070-2)
- Friedman, S. K., & Reich, P. B. (2005). Regional Legacies of Logging: Departure from Presettlement Forest Conditions in Northern Minnesota. *Ecological Applications*, 15(2), 726–744. <https://doi.org/10.1890/04-0748>
- Frerker, K., Sabo, A. and Waller, D., 2014. Long-term regional shifts in plant community composition are largely explained by local deer impact experiments. *PloS one*, 9(12), p.e115843.
- Frelich, L.E., Blossey, B., Cameron, E.K., Dávalos, A., Eisenhauer, N., Fahey, T., Ferlian, O., Groffman, P.M., Larson, E., Loss, S.R. and Maerz, J.C., 2019. Side-swiped: ecological cascades emanating from earthworm invasions. *Frontiers in Ecology and the Environment*, 17(9), pp.502-510.
- Galatowitsch, S., Frelich, L., & Phillips-Mao, L. (2009). Regional climate change adaptation strategies for biodiversity conservation in a midcontinental region of North America. *Biological Conservation*, 142(10), 2012–2022. <https://doi.org/10.1016/j.biocon.2009.03.030>
- Garrett, H.E. et al. (2004). Hardwood silvopasture management in North America. In: Nair, P.K.R., Rao, M.R., Buck, L.E. (eds) *New Vistas in Agroforestry. Advances in Agroforestry*, vol 1. Springer, Dordrecht. https://doi.org/10.1007/978-94-017-2424-1_2.

- Hale, C.M., Frelich, L.E., Reich, P.B. and Pastor, J., 2005. Effects of European earthworm invasion on soil characteristics in northern hardwood forests of Minnesota, USA. *Ecosystems*, 8, pp.911-927.
- Hale, C.M., Frelich, L.E. and Reich, P.B., 2006. Changes in hardwood forest understory plant communities in response to European earthworm invasions. *Ecology*, 87(7), pp.1637-1649.
- Handler, S., Duveneck, M. J., Iverson, L., Peters, E., Scheller, R. M., Wythers, K. R., Brandt, L., Butler, P., Janowiak, M., Shannon, P. D., Swanston, C., Barrett, K., Kolka, R., McQuiston, C., Palik, B., Reich, P. B., Turner, C., White, M., Adams, C., ... Ziel, R. (2014). Minnesota forest ecosystem vulnerability assessment and synthesis: a report from the Northwoods Climate Change Response Framework project. *Gen. Tech. Rep. NRS-133*. Newtown Square, PA; U.S. Department of Agriculture, Forest Service, Northern Research Station. 228 p., 133, 1–228. <https://doi.org/10.2737/NRS-GTR-133>
- Hanberry, B. B., Palik, B. J., & He, H. S. (2012). Comparison of historical and current forest surveys for detection of homogenization and mesophication of Minnesota forests. *Landscape Ecology*, 27(10), 1495–1512. <https://doi.org/10.1007/s10980-012-9805-5>.
- Hartman, P., & Steele, K. (2015). *Ecological Site Description: Dolomite Colluvium Bluff Prairie*. United States Department of Agriculture. https://efotg.sc.egov.usda.gov/references/Public/MN/105XY001_DolomiteColluviumBluffPrairie.pdf
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Holdsworth, A.R., Frelich, L.E. and Reich, P.B., 2007. Effects of earthworm invasion on plant species richness in northern hardwood forests. *Conservation Biology*, 21(4), pp.997-1008.
- Holdsworth, A.R., Frelich, L.E. and Reich, P.B., 2008. Litter decomposition in earthworm-invaded northern hardwood forests: role of invasion degree and litter chemistry. *Ecoscience*, 15(4), pp.536-544.
- Holdsworth, A.R., Frelich, L.E. and Reich, P.B., 2012. Leaf litter disappearance in earthworm-invaded northern hardwood forests: role of tree species and the chemistry and diversity of litter. *Ecosystems*, 15, pp.913-926.
- Jacobson, Keith, Michelle Martin, and Mike Wachholz. 2019. Oak Regeneration Success but Recruitment Failure after Clearcut with Reserves on a Southern Dry-Mesic Oak Forest (DNR). Great Lakes Silviculture Library.
- Knapp, W. M., & Wiegand, R. (2014). Orchid (Orchidaceae) decline in the Catoctin Mountains, Frederick County, Maryland as documented by a long-term dataset. *Biodiversity and Conservation*, 23(8), 1965–1976. <https://doi.org/10.1007/s10531-014-0698-2>
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Maerz, J.C., Nuzzo, V.A. and Blossey, B., 2009. Declines in woodland salamander abundance associated with non- active earthworm and plant invasions. *Conservation Biology*, 23(4), pp.975-981.

- Matonis, M.S., Walters, M.B. and Millington, J.D., 2011. Gap-, stand-, and landscape-scale factors contribute to poor sugar maple regeneration after timber harvest. *Forest Ecology and Management*, 262(2), pp.286-298.
- McCay, T.S. and Scull, P., 2019. Invasive lumbricid earthworms in northeastern North American forests and consequences for leaf-litter fauna. *Biological Invasions*, 21, pp.2081-2093.
- McGraw, J. B., & Furedi, M. A. (2005). Deer Browsing and Population Viability of a Forest Understory Plant. *Science*, 307(5711), 920–922. <https://doi.org/10.1126/science.1107036>
- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). 2025. Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state’s greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>
- Nowacki, G. J., & Abrams, M. D. (2008). The Demise of Fire and “Mesophication” of Forests in the Eastern United States. *BioScience*, 58(2), 123–138. <https://doi.org/10.1641/B580207>
- Nuttle, T., Ristau, T.E. and Royo, A.A., 2014. Long-term biological legacies of herbivore density in a landscape-scale experiment: forest understoreys reflect past deer density treatments for at least 20 years. *Journal of Ecology*, 102(1), pp.221-228
- Paudel, S., Longcore, T., MacDonald, B., McCormick, M.K., Szlavecz, K., Wilson, G.W. and Loss, S.R., 2016. Belowground interactions with aboveground consequences: Invasive earthworms and arbuscular mycorrhizal fungi. *Ecology*, 97(3), pp.605-614.
- Perry, D.A and M.P. Amaranthus. 1997. Disturbance, recovery, and stability. Pages 31-56 in K.A. Kohm and J.F. Franklin, editors. *Creating a Forestry for the 21st Century*. Washington DC: Island Press.
- Radeloff, V. C., Hammer, R. B., Stewart, S. I., Fried, J. S., Holcomb, S. S., & McKeefry, J. F. (2005). The Wildland–Urban Interface in the United States. *Ecological Applications*, 15(3), 799–805. <https://doi.org/10.1890/04-1413>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Sabo, A.E., Frerker, K.L., Waller, D.M. and Kruger, E.L., 2017. Deer-mediated changes in environment compound the direct impacts of herbivory on understorey plant communities. *Journal of Ecology*, 105(5), pp.1386-1398
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, e14434. <https://doi.org/10.1111/cobi.14434>
- U.S. Department of Agriculture, Forest Service, Forest Inventory and Analysis (FIA) Program. (2022).

Wiegmann, S. M., & Waller, D. M. (2006). Fifty years of change in northern upland forest understories: Identity and traits of “winner” and “loser” plant species. *Biological Conservation*, 129(1), 109–123.
<https://doi.org/10.1016/j.biocon.2005.10.027>

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55).
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Zamora, D., Wyatt, G., Buttler, M., Ford, M., Magner, J., Reichenbach, M., Burkett, E., Current, D., Walter, D. (2017). Silvopasture: Establishment and Management Principles for Minnesota. Retrieved from the University Digital Conservancy, <https://hdl.handle.net/11299/214976>.

Ziemba, J.L., Hickerson, C.A.M. and Anthony, C.D., 2016. Invasive Asian earthworms negatively impact keystone terrestrial salamanders. *PloS one*, 11(5), p.e0151591.

Mesic Hardwood Forests

Habitat Description

Mesic (moist) hardwood forest habitat can be found across the state. It is present throughout portions of the Laurentian Mixed Forest Province where it occurs on level to rolling landscapes with fine-textured soils that retain water, and in the Eastern Broadleaf Forest Province where rugged terrain, water bodies such as lakes and rivers, and moist soil provide protection from wildfires. It can be found to a much lesser extent in the Prairie Parkland, where it is primarily limited to well-drained soils found in the rolling till plains, and in the Tallgrass Aspen Parklands (DNR, 2005). Regardless of their location, the common features are uplands with soils that retain water and settings where wildfires are infrequent. A continuous, often dense, canopy of deciduous trees, typically including sugar maple (*Acer saccharum*), American basswood (*Tilia americana*), Northern red oak (*Quercus rubra*), quaking aspen (*Populus tremuloides*) and bigtooth aspen (*Populus grandidentata*), is also characteristic of this habitat and varies across native plant community classes. Other canopy trees include American elm (*Ulmus americana*), red elm (*Ulmus rubra*), black ash (*Fraxinus nigra*), green ash (*Fraxinus pennsylvanica*), bitternut hickory (*Carya cordiformis*), and hackberry (*Celtis occidentalis*).

Because the forest canopy blocks most of the sunlight before it reaches herbaceous plants and seedlings on the forest floor, most understory plants have evolved some degree of shade tolerance. Some species, known as spring ephemerals, have adapted to the low light conditions by developing and flowering before the canopy has fully leafed out. Mature mesic forest stands commonly have several nearly closed layers of woody plants, including a well-defined forest canopy, subcanopy, and shrub layer. These layers combine to produce largely continuous cover. As these forests age, they become structurally more complex as canopy trees die, producing gaps in the canopy that allow the growth of less shade-tolerant shrubs and trees. Standing large dead trees become more frequent, eventually becoming large down logs, which add to the structural diversity of older forests.

Natural disturbance in this habitat is characterized by the death of individual trees, which occurs at a rather constant rate in older forests. Stand-regenerating disturbances such as wildfires and catastrophic windthrow were rare historically in this mesic habitat, having average frequencies of once every 360 to more than 1,000 years. Disturbances that resulted in the partial loss of canopy trees, such as light surface fires and moderate windthrow, were far more frequent. Historically, surface fire was more important in the north, and wind was more important in central and southern Minnesota.

Typical sites are buffered from seasonal drought by fine-textured soils with impermeable soil horizons capable of retaining rainfall or snowmelt below the surface. Usually, these soils are well drained and are waterlogged or saturated only after spring snowmelt or heavy, prolonged rains. Essential nutrients, especially nitrogen, are mineralized from decaying organic matter at relatively high rates and quickly become available again for uptake by plants during the spring and early summer months. As a result, nutrients and organic matter accumulate at the soil surface in leaf litter and humus.

Mesic Hardwood Forest in some locations contain vernal pools, small, ephemeral wetlands that occur within Minnesota's forests and provide habitat to an array of SGCN; see the Vernal Pool sub-chapter.

Mesic hardwood forests have diverse characteristics that vary considerably depending on site conditions and location. In addition to the sugar maple-basswood-northern red oak forests of the Big Woods region, the blend of dominant trees can vary considerably in abundance, ranging from drier, more fire-dependent aspen-dominated habitats to more mesic sites dominated by oaks. Aspen dominated forests and oak forests are briefly described below.

Aspen-dominant Forest

Mesic aspen forest is characterized by a canopy dominated by quaking aspen, big-toothed aspen, paper birch (*Betula papyrifera*), or a mixture of these species. These shade-intolerant tree species are the dominant trees in the early stages of a wide variety of native plant communities in fire-dependent and mesic hardwood forest systems across a range of dry to mesic soils. Thus, aspen forest is a cover type that may eventually develop into many other native plant communities. In the absence of disturbance, such as fire or timber harvest, aspen acres may decline across the landscape as they succeed to longer-lived species.

Aspen forests typically have a nearly complete canopy of aspen or birch, but the canopy is not as dense as that of sugar maple. As a result of higher light levels penetrating the canopy, these forests usually have a well-developed shrub layer dominated by hazelnuts (*Corylus* spp.) or dogwoods (*Cornus* spp.). The coverage and diversity of the herbaceous plant layer are highly variable depending on site conditions and stand history.

See also the *Upland Deciduous Forests and Woodlands (Fire-dependent) Chapter*.

Oak-dominant Forest

Oak forest habitat on more mesic sites is dominated by Northern red oak, white oak (*Quercus alba*), and bur oak (*Quercus macrocarpa*). Tall, straight, single-stemmed trees and a less dense shrub layer characterize the oak habitat on these sites.

Habitat Map

To depict Mesic Hardwood Forest habitat (see Figure 3.10), we compiled spatial data from several sources: DNR's Native Plant Communities and the Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Mesic Hardwood Forest (MH)

[MHc26 Central Dry-Mesic Oak-Aspen Forest \(PDF\)](#)

[MHc38 Central Mesic Cold-Slope Hardwood-Conifer Forest \(PDF\)](#)

[MHc36 Central Mesic Hardwood Forest \(Eastern\) \(PDF\)](#)

[MHc37 Central Mesic Hardwood Forest \(Western\) \(PDF\)](#)

[MHc47 Central Wet-Mesic Hardwood Forest \(PDF\)](#)

[MHn45 Northern Mesic Hardwood \(Cedar\) Forest \(PDF\)](#)

[MHn35 Northern Mesic Hardwood Forest \(PDF\)](#)

[MHn47 Northern Rich Mesic Hardwood Forest \(PDF\)](#)

[MHn44 Northern Wet-Mesic Boreal Hardwood-Conifer Forest \(PDF\)](#)

[MHn46 Northern Wet-Mesic Hardwood Forest \(PDF\)](#)

[MHs37 Southern Dry-Mesic Oak Forest \(PDF\)](#)

[MHs38 Southern Mesic Oak-Basswood Forest \(PDF\)](#)

[MHs39 Southern Mesic Maple-Basswood Forest \(PDF\)](#)

[MHs49 Southern Wet-Mesic Hardwood Forest \(PDF\)](#)

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Laurentian Hardwood Forest, North – Central Beech – Maple – Basswood Forest, and Laurentian Hemlock – White Pine – Hardwood Forest.




Figure 3.10. A map depicting Mesic Hardwood Forest habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological Systems of Mesic Hardwood Forest as well as three types from the Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

The extent of upland deciduous hardwood forests has been greatly reduced in southern and west-central Minnesota since settlement by people of European descent. The extensive mesic hardwood forests of the Big Woods Subsection have been reduced by a factor of more than 100. However, in the northern parts of the Laurentian Mixed Forest Province, the extent of maple-basswood forests has increased because of fire suppression. Like other forest habitats, most maple-basswood habitats in the Eastern Broadleaf Forest Province and southern and western portions of the Laurentian Mixed Forest Province have been fragmented by agriculture and development. In many locations, the remaining forests typically lack the ecological complexity of pre-European settlement forests because of several factors (for example, grazing, invasive plants and animals, edge effects, changes in native animal populations, and consumptive uses).

Forest planning is critical to ensure that a balance of forest habitat conditions (e.g., younger and older forest, small and large patch sizes) are available for plant and animal species across the landscape and can support conservation goals. For instance, aspen and birch are short-lived tree species and naturally transition towards other forest types (e.g., maple, basswood, oak) in the absence of significant disturbance. Given that some species require younger forest habitat or mature habitat of shorter-lived tree species, it can be important to maintain those conditions through the application of management tools like timber harvest and prescribed fire. Conversely, other plant and animal species require habitats characterized by a non-stand replacing or smaller-scale disturbance, including larger blocks of older forest and mature forests of later successional tree species. For these forest species, large-scale, whole canopy removal may negatively affect local habitat conditions in the short-term.

Aspen-dominated Forest

As aspen forests age, they typically increase in structural diversity. Historically, most aspen stands in northern Minnesota had a conifer component, which increased as the stand aged. Today, most aspen stands have little or no conifer understory, due to past management and slash fires. Still, many older aspen stands are relatively structurally diverse, with large trees, snags, down logs, and an understory containing more shade-tolerant hardwoods or conifers that will become the canopy dominants in the absence of a stand-replacing disturbance.

Today, aspen forest is the most abundant forest habitat in Minnesota and much more widespread than it was prior to settlement by people of European descent. An analysis of General Land Office bearing tree records from the late 1800s and Forest Inventory and Analysis plots from the 1990s shows that in the Laurentian Mixed Forest Province, aspen forest communities have increased nearly tenfold (Friedman and Reich, 2005). However, it is thought that aspen forests are structurally less diverse than they were historically, and on average younger than that of pre-European settlement aspen forests. Currently, about 60% of the aspen age class distribution is 1-40 years old, and about 18.5% of the cover type is over 60 years old statewide (US Department of Agriculture, Forest Service, FIA Program, 2025). The location of older aspen stands shift across the landscape over time as old stands are harvested or succeed to other tree species, and young-mature aspen stands that aren't harvested age.

Older, more structurally diverse aspen forests have high value to some SGCN, such as American Goshawk, and the configuration of that habitat on the landscape (e.g. tract size, proximity to other forest types) also relates to its suitability and significance as habitat for SGCN.

Oak-dominated forest

Mesic Hardwood oak forests had fewer severe fires prior to settlement by people of European descent than did dry oak forests, and more mesic, fire-sensitive trees such as basswood, green ash, bitternut hickory, and big-toothed aspen are usually present with the oaks. Nevertheless, those fires that did occur were responsible for the establishment and maintenance of oaks on these sites. Without fire or appropriate forest management to replace fire, most mesic oak forests will succeed to maple-basswood forests.

Some oak forest stands, particularly if disturbed by grazing or selective logging, have been colonized by invasive species such as buckthorn (*Rhamnus cathartica*) and Tartarian honeysuckle (*Lonicera tartarica*). Like other forest habitats, most oak forest habitats in the Eastern Broadleaf Forest Province and southern and western portions of the Laurentian Mixed Forest Province have been fragmented by agriculture and development.

Climate Profile for Mesic Hardwood Forest

The anticipated impacts of climate change on Mesic Forests were profiled in the report “Effects of Climate Change on Midwestern Ecosystems: Appalachian-Interior-Northeast Mesic Forest” published by the Midwest Climate Adaptation Science Center (Ratcliffe et al., 2025). The most impactful changes in climate for this system include more frequent droughts and milder winters. Mesic forests have generally been experiencing moister conditions (mesophication) as well as fire suppression over the past century, favoring shade-loving moisture-dependent species. While precipitation is expected to increase overall, particularly during the springtime, the increase in late summer drought may shift conditions to be more favorable for more drought-tolerant and fire-adapted species.

Dryer and hotter summers mean droughts are expected to increase in both frequency and severity, primarily impacting soil moisture and increasing tree mortality, overall leading to changes in species composition. Wildfire risk will also be heightened in these dryer conditions. In the Driftless Region, drought conditions are expected to increase by over 800%.

Warmer winters will also have impacts as reduced snowpack leads to changes in habitat availability and growing season. Warmer winters will promote the migration of invasive species better suited for warmer climates. Pests and pathogens may also do more damage under future climate scenarios as winter kills will be less impactful.

Key Climate Change Impacts:

- **Habitat Structure:** Climate-induced erosion from flooding and drier soils will negatively impact the slopes and hillsides where hardwood mesic forests are commonly found. Drought and warmer winters will also impact other soil metrics such as pH and nutrient availability, further impacting habitat structure.

- **Community Composition:** This habitat type is adapted primarily wet conditions, which makes it particularly susceptible to the warmer, dryer conditions climate change may bring. Regions like the Driftless Area may see these impacts sooner and more concentrated compared to Northern Regions like the Erie Drift Plains, which is aided by the lake-effect. Throughout the state, drought resistant species like *Quercus* will become better adapted and more common, while species adapted to wetter conditions may decline. Wildfire will also become more common during the warm and dry summers, supporting fire-adapted species. Lastly, warmer winters may mean less snowpack and increased freeze-thaw cycles, leading to plant mortality and changes in species composition.
- **Invasive Species:** Climate change conditions will favor invasive species, especially fire-dependent and drought-adapted species. Wildfire will present opportunity for more shade-intolerant and fire-dependent invasive species to colonize this habitat. Warmer winters will allow invasive species from warmer regions to begin to colonize colder regions.
- **Pests and Pathogens:** Pest populations are expected to increase with climate change. Drought will make hosts more susceptible to pests, while warmer winters are expected to kill less pests than they had in the past and allow for their northward expansion. Pest mortality may increase in some cases, for example during more frequent and intense heatwaves.
- **Herbivory:** Mild winters will increase white-tailed deer (*Odocoileus virginianus*) populations as food becomes more available and winter mortality decreases. This may lead to a transition to more browse-resistant species, which in some cases suppresses climate-resilient species.

Please note that this entire Climate Profile is derived from Ratcliffe et al., (2025).

Climate Focal Wildlife Species: Cerulean Warbler

This profile is excerpted from “Effects of Climate Change on Midwestern Ecosystems: Appalachian-Interior-Northeast Mesic Forest” published by the Midwest Climate Adaptation Science Center (Ratcliffe et al. 2025).

The cerulean warbler (*Setophaga cerulea*) is a small, Nearctic-Neotropical migratory songbird considered to be a Bird of Conservation Concern (Buehler and others, 2020; U.S. Fish and Wildlife Service, 2021). Like many long-distance migratory birds, cerulean warblers demonstrate a limited capacity to advance migratory arrival in response to advancing spring conditions and may be vulnerable to phenological asynchrony (i.e., peak nesting and insect availability) (Connare and Islam, 2022). This is of special concern given cerulean warblers’ heavy reliance on Lepidoptera (Auer and others, 2016) and the vulnerability of insects to climate change (Edwards and others, 2025). The species is associated with mature, deciduous forests with canopy gaps (Buehler and others, 2020). In the breeding range, they demonstrate preference for white oak (*Quercus alba*) and bitternut hickory (*Carya cordiformis*), avoiding northern red oak (*Quercus rubra*) and red maple (*Acer rubrum*) (Buehler and others, 2020), potentially reflecting higher lepidopteran abundance in forests dominated by these tree types (Connare and Islam, 2022). Novel interspecific interactions are also of concern. For example, as the northern parula (*Setophaga americana*) breeding range expands northward, hybridization with cerulean warbler may become more common (Trimbath and others, 2019).

In the overwintering range, cerulean warblers may be found in old-growth forests, shade coffee plantations, and low-mid elevation forests of the Andes (Buehler and others, 2020). The U.S. Gulf Coast and Mississippi Valley provide important stopover habitat to meet migratory demands of trans-Gulf crossings (Raybuck and others, 2022). Climate change threatens suitable habitats across the nonbreeding range. In the Andes, studies predict a moderate contraction (19–43%)

in suitable range, driven by climate change, and an upslope elevational shift (Brodie et al. 2024). Both climate-driven flood hazard and human alterations to the Mississippi River system threaten the viability of bottomland hardwood forests (Munoz and others, 2018). Targeted forest management and restoration may be viable strategies to offset declines in the species, provided it increases stand structural diversity, creates small canopy openings, and promotes preferential tree species (e.g., white oak) (Kellner and others, 2016; Nareff and others, 2019).

Species in Greatest Conservation Need

Mesic Hardwoods provide primary or secondary habitat for 49 animal SGCN and 52 plant SGCN (see Table 3.9). Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants).

Table 3.9. Numbers of Species in Greatest Conservation Need associated with Mesic Hardwood Forest as either primary or secondary habitat¹.

Species Group	Primary Habitat	Secondary Habitat	Total
Amphibians	2	0	2
Birds	5	2	7
Mammals	2	14	16
Reptiles	5	0	5
Terrestrial Invertebrates			
Bees			8
Beetles (terrestrial)			1
Snails (terrestrial)	4	4	8
Spiders	2	0	2
Plants	52		52
Total	72	20	101

1. Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Of the 131 terrestrial vertebrate SGCN identified in the 2025-2035 SWAP, 23% (30) occur in mesic hardwoods. Half of the mammal 32 SGCN species are associated with this habitat. The 52 plant SGCN found in mesic hardwoods represent 12% of the 447 SGCN. Examples of selected SGCN are described below.

Amphibians

Two salamander SGCN – [spotted salamander \(*Ambystoma maculatum*\)](#) and [four-toed salamander \(*Hemidactylium scutatum*\)](#) – are strongly associated with vernal pools (see Vernal Pool sub-chapter) and the mature forest habitats surrounding them. Temporary forested wetlands are critical for these species, which need them in the spring to breed. Eggs are laid in or near the water, and larvae remain in the pools until metamorphosis occurs. Four-toed salamanders specifically lay their eggs in moss hummocks next to wetlands. Four-toed salamanders occur most frequently in mature upland forests on glacial moraine landscapes with frequent isolated wetlands that include an alder margin and moss hummocks adjacent to open water.

Birds

[Acadian flycatchers \(*Empidonax virescens*\)](#), black-throated blue warblers (*Setophaga caerulescens*), [cerulean warblers \(*Setophaga cerulea*\)](#), [hooded warblers \(*Setophaga citrina*\)](#), [Louisiana waterthrushes \(*Parkesia motacilla*\)](#), and [red-shouldered hawks \(*Buteo lineatus*\)](#) generally require large areas of contiguous mature to old-growth hardwood forest. Acadian flycatchers favor relatively undisturbed forests and experience high rates of brood parasitism and nest depredation in fragmented landscapes. They have habitat requirements very similar to Louisiana waterthrushes but are less dependent on swiftly flowing streams. Where permanent streams are not present, acadian flycatchers are still usually associated with steep-sided ravines. Cerulean warblers need large, tall trees with horizontal heterogeneity in the canopy, and hooded warblers need mature forests with significant treefall gaps that provide shrubby undergrowth for nesting.

Louisiana waterthrushes are dependent on small swiftly flowing streams bordered by mature closed-canopy deciduous forests, typically in steep-sided valleys. Mesic hardwoods are the most important forest type for Louisiana waterthrush through most of their Minnesota range (e.g., Blufflands ECS subsection), although floodplain forests gain importance farther north (e.g., Mille Lacs Uplands ECS subsection). A key characteristic of suitable Louisiana waterthrush habitat is high quality forest in both the valley bottom and on both adjoining slopes. Several mesic hardwood native plant community classes provide suitable habitat, although Southern Wet-mesic Hardwood Forest on the valley bottom is particularly important (unpublished MBS data). Although the minimum width of forest corridors necessary for Louisiana waterthrushes is unknown, occupied forested stream valleys in Minnesota were a minimum of 300 meters wide.

Largely restricted to northeastern Minnesota, black-throated blue warblers require mature closed-canopy forests, often on steep slopes, with a well-developed understory and dense shrub layer. In Mesic Hardwood forests, black-throated blue warblers are often found in mixed deciduous-coniferous stands. However, suitable forest structure is more important than the exact species composition of canopy trees.

Mammals

The mammalian SGCN fauna is quite similar across all upland forest habitats, including Upland Conifer Forests (22 mammal SGCN), Upland Deciduous Forests (19 mammal SGCN) and Mesic Hardwoods (16 mammal SGCN). Missing in the Mesic Hardwoods are several species more closely associated with boreal habitats including moose (*Alces alces*), snowshoe hare (*Lepus americanus*), [Canada lynx \(*Lynx canadensis*\)](#), rock vole (*Microtus chrotorrhinus*), and [Eastern heather vole \(*Phenacomys ungava*\)](#). For some species with home ranges that extend over several native plant communities, such as [gray wolves \(*Canis lupus*\)](#), the landscape mosaic can be as or more important than the individual habitats.

[Woodland voles \(*Microtus pinetorum*\)](#) are the one species not found in fire-dependent upland forests. They require moist, light soil or humus in forests to construct burrows and sub-surface runways. Grazing by cattle, which compacts the soil, and the presence of invasive non-native earthworms, which destroy the humus, may make forests within its limited range in southeastern Minnesota unsuitable for this species. They are also found in Deciduous Wet Forest and Riparian and Floodplain Forests. Currently their distribution is limited to extreme southeastern Minnesota. Mesic Hardwoods also provide primary habitat to the cave-hibernating Tri-colored bats (*Perimyotis subflavus*). Live or recently dead leaves of hardwood trees provide their roosting habitat during the spring, summer, and fall months. As noted by the [USFWS](#), the species population has

declined more than 90% in recent years due to the impacts of white-nose syndrome. As a result, the species has been proposed for federal listing as a Threatened species

Reptiles

The reptilian SGCN fauna found in Mesic Hardwoods is similar to that found in Upland Deciduous Forest and Woodlands but includes only five species compared to the 9 found in Upland Deciduous Forests; none are unique to the Mesic Hardwoods. Included are one lizard ([Common Five-lined Skink, *Plestiodon faxciatus*](#)), one turtle ([Wood Turtle, *Glyptemys insculpta*](#)), and three snakes ([North American racer \(*Coluber constrictor*\)](#); [Timber rattlesnake \(*Crotalus horridus*\)](#); and [Western ratsnake \(*Pantherophis obsoletus*\)](#)). With the exception of the wood turtle, the remaining SGCN depend on unique microhabitat features that are found in the Mesic Hardwoods of southeastern Minnesota. These include rocky outcrops, forested rocky bluffs, bluff prairies, and rock crevices that are common in the Mississippi River blufflands, as well as granite outcrops, that are common along portions of the Minnesota River valley. These features provide multiplate basking sites for thermoregulation and dens for overwintering, two of the SGCN most essential habitat requirements. Limited to the southeastern corner of the state, the Western ratsnake spends considerable time in trees (6 to 12 meters above ground) and may take cover in tree cavities (Moriarty and Hall, 2014).

Plants

The relative lack of stand-replacing disturbance events and rich growing conditions of mesic hardwood forests supports a diverse assemblage of plant species. Among them are some of three smallest plants in Minnesota- a group of tiny ferns known as moonworts and grapeferns (*Botrychium* and *Sceptridium*). Some of the rarest of these <4" tall ferns, including goblin fern (*Botrychium mormo*), [narrow triangle moonwort \(*Botrychium angustisegmentum*\)](#), and [blunt-lobed grapefern \(*Sceptridium oneidense*\)](#), inhabit the shaded groundlayer of mature maple-basswood forests. Changes to local habitat conditions, either from canopy opening/removal or leaf litter consumption by non-native earthworms, threaten the ability of these unique and often cryptic plants to persist in an area. Given their small stature, moonwort spore dispersal distances are quite limited, making local conservation efforts critical.

[American ginseng \(*Panax quinquefolium*\)](#) is a perennial herb native to hardwood forests in eastern North America. Wild ginseng became quite rare in Minnesota due to the widespread clearing of mesic hardwood forests for agriculture and development, and unsustainable ginseng harvest levels. The roots are commercially harvested, with interstate and international trade regulated by the [Convention on International Trade in Endangered Species of Wild Fauna and Flora \(CITES\)](#), and harvest further regulated in Minnesota under [Minnesota Rules, Chapter 6282](#). Under the CITES treaty, the export of wild ginseng root can only be allowed if a state can demonstrate that continued export will not cause ginseng populations within that state to decline any further. Currently, Minnesota is considered in compliance. However, populations are currently so depleted and fragmented that adequate recovery is unlikely without further restrictions on harvest. The DNR hosts a [webpage](#) to help collectors assure the future of wild ginseng in Minnesota.

[Butternut \(*Juglans cinerea*\)](#) was once common throughout the Eastern United States, including Minnesota. A lethal fungal disease, [butternut canker \(*Ophiognomonia clavignenti-juglandacearum*\)](#), reached Minnesota in the 1970's, and as of today, nearly all butternut trees in the state are dead or dying. Active research into treatment options and the potential for augmentations using naturally occurring resistant genotypes is ongoing.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures that result from human inhabitation, sustenance, livelihoods, and recreation. The list below reflects a list of threats in a way that is organized and referenced globally with modifications for this Plan (see Threats section; Salafsky et al., 2024). These are the underlying pressures that have led or continue to lead to habitat loss or degradation. Notes about how these issues may specifically impact this habitat are included below. Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “*Stressors*” section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildlife following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Residential, commercial, and industrial development can all directly alter forest ecosystems by removing, degrading, or fragmenting forest cover and habitat. Fragmentation of forests is a growing concern, and occurs when a large, contiguous forest land mass is divided into smaller tracts through sale and subdivision, road construction, or clearing of forests for agriculture, housing, or other commercial developments. This often creates smaller forest pockets, which are interspersed with non-forest land uses. Fragmentation can inhibit the natural migration of many plant and animal species, increase the risk of wildfires at the wildland urban interface, restrict public recreational access, reduce the habitat value and ecological complexity of forest lands, and contribute to the spread of invasive species. Because of their natural beauty, diversity, recreation opportunities, and abundant natural resources, forested landscapes are appealing places to live; even in the largely forested areas of northern Minnesota, rural development has greatly reduced the extent of large, contiguous forest areas; most forest areas are within 25 kilometers (15.5 miles) of small housing settlements (Radeloff et al., 2005). Forested lands adjacent to lakes and rivers are especially attractive sites for development.



Crop Production

Historically, many of the mesic hardwood forests were converted to cropland, leaving a patchwork of widely scattered wood lots, averaging 40 to 80 acres in size, across central and southeastern Minnesota. The conversion of forests to agricultural use is ongoing in some locations.



Livestock Management

Minnesota currently has over 650,000 acres of mesic, upland deciduous, and riparian forests on farms that are grazed by livestock (Zamora et al., 2017; Garrett et al., 2004); heavy grazing could impair soil health and water quality.



Mining and Quarrying

Forests, when they co-occur with mineral or sand and gravel deposits, are at risk of degradation due to the direct and indirect impacts of mining activities such as permanent forest loss, soil compaction, introduction of invasive species, human disturbance, waste discharge and pollution, and forest fragmentation (Sonter et al., 2018). Mining projects can trigger additional, cascading land use change with the establishment of associated infrastructure such as improved road access.



Roads, Trails, and Railroads

Roads, trails, and railroads can create fragmentation and edge effects in large tracts of forest. Increase in the number of roads and trails, particularly for motorized vehicles, increases human disturbance and introduction of invasive species further into the forest. Repeated use and soil compaction associated with roads and trails also can modify the vegetation within the area of use (Zenner et al., 2008). Temporary winter logging trails may become rutted while materials used to stabilize the trails for heavy vehicles may remain in place, leaving behind wood planks and wood chips. Trails that provide winter access to upland deciduous stands can disturb forest wildlife, especially species that begin nesting in late winter, such as many forest raptors.



Utility Corridors

Utility corridors transect forests and woodlands throughout the state. These corridors can contain a variety of utility infrastructure. This includes pipelines, telephone lines, and power lines. These lines can be aerial, above ground, or below ground. Regardless of type, utility corridors can fragment habitat based on their size, purpose, and the upkeep that is required to maintain them.

The initial construction of the associated utility infrastructure as well as the long-term maintenance of that infrastructure can have lasting impacts on native plant communities and the SGCN associated with them as well as the direct disturbance and oftentimes permanent deforestation and habitat loss incurred during their installation.

These corridors and the increased travel from vehicles and equipment that they require act as vectors for invasive plant species. The required vegetative management practices often include herbicide applications, removal of woody vegetation, and/or mowing. This increased disturbance increases the likelihood of establishing invasive plant species within the corridors and then spreading to adjacent lands.



Timber Harvest

As forest composition and structure vary across mesic hardwood native plant communities and local conditions, so do the silvicultural practices typically used during the management of these sites. More moist and rich forest conditions support longer-lived, shade tolerant tree species that are often

harvested with uneven-aged approaches (e.g., gap or patch cuts, thinning) in comparison to drier and more nutrient poor sites with shorter-lived, shade intolerant tree species where even-aged harvests (e.g. clearcut or shelterwood harvest) are more prevalent. While each of these silvicultural strategies can incorporate elements of natural disturbance regimes (e.g., gap harvest= small wind throw event, clearcut= major blowdown), when they are applied with a focus on maximizing economic utilization they may result in degraded or simplified habitat conditions. For instance, lower overstory retention leads to a loss of structural complexity in post-harvest stands. Additionally, the scale and rate of harvest activities in mesic hardwood forests shapes landscape-scale age class distributions and the amount of forest habitats in suitable condition and configuration for a range of plant and animal species.

Typical commercial timber harvest intervals may be shorter than historic disturbance intervals, creating needs for providing sufficient late-successional forest stands to balance management for timber production and forest health that removes trees before they are old. This is particularly notable for oaks, which become more susceptible to disease with age and can spread disease from old oaks to seedlings, so oak-dominated forests and woodlands are commonly managed with complete, even-aged harvests. Such harvests can create a set of ecological characteristics that differ from mesic forests with more natural disturbance and regeneration patterns and might not provide optimal habitat conditions for SGCN. Many wildlife species benefit from managing for within-stand species and structural diversity in large habitat patches to provide habitat for area-sensitive species and to minimize habitat fragmentation effects.

Harvesting mesic sites in non-frozen conditions can cause soil compaction and rutting impacts.



Recreation

In addition to the impacts created by recreational trails (discussed above), recreational activities themselves can proliferate invasive species. Seeds of invasive plants such as common tansy (*Tanacetum vulgare*) and garlic mustard (*Alliaria petiolata*) can become embedded in the wheels of vehicles or the shoes of individuals and eventually become established along compacted trails where conditions for their growth may be more favorable. Earthworm egg cases can also be spread in mud, which are problematic for moonwort (*Botrychium*) species. Increased use of roads and trails also increases human disturbance in the forest, reducing the amount of inaccessible area.



Fire Management

Some mesic forests have originated from forests originally dominated by oaks. However, there is an inherent difference in soils and topographic features that promote the establishment of Mesic Hardwood Forest ecosystems. On the landscape they are buffered from seasonal drought by fine-textured soils capable of holding or perching rainfall. Over time, as climates became cooler and wetter, conditions were more favorable for maples, basswood, and other fire-sensitive hardwoods. These conditions were enhanced by the suppression of fire. Now, as temperatures continue to rise and droughts become more frequent, oaks are more likely to thrive in many areas.



Invasive Species (Problematic Non-native Species)

The arrival of Eurasian earthworms is causing profound ecological changes to previously worm-free mesic hardwood forests (Frelich et al., 2019). They upend the traditional soil profile (Hale et al., 2005;

Fahey et al, 2013), remove the accumulated layers of organic leaf litter (Holdsworth et al., 2008, 2012), and alter the community of soil mycorrhizal fungi (Paudel et al., 2016). These changes cascade through the ecosystem, including ground layer vegetation (Hale et al., 2006; Holdsworth et al., 2007), leaf litter fauna (McCay and Scull 2019), salamanders (Maerz et al., 2009; Ziemba et al., 2016), and likely other vertebrates. One rare plant in particular, the goblin fern, is threatened with widespread extirpation due to earthworms (Gundale, 2002; Zlonis and Henderson, 2018). On the other hand, invasive plant species generally benefit from earthworms (Nuzzo et al., 2009; Whitfeld et al., 2014). They can also stress sugar maples, leading to canopy dieback (Bal et al., 2018), and they may slow the climate-induced replacement of boreal tree species by temperate species (Frelich et al., 2012). In addition to the earthworm species such as nightcrawlers (*Lumbricus* species) that have been well-studied in northern Minnesota forests, there is the concern that jumping worm species (*Amyntas* and *Metaphire* species) could arrive in these forests and cause additional impacts. Invasive plant species, such as buckthorn and Tartarian honeysuckle often outcompete native plant species and repress understory plant species leading to modified habitats unsuitable for SGCN. While a concern statewide, forests in central and southern Minnesota are currently most impacted these invasive shrubs. Where they co-occur with SGCN plants might be a good place to prioritize control efforts.

The spongy moth (*Lymantria dispar*) is an invasive forest pest that feeds on more than 300 species of deciduous trees and shrubs, including aspen, oak, and birch. Repeated removal of leaves, or defoliation, stresses trees and can leave them vulnerable to disease or other pest infestations that can kill trees. Once it becomes established in a location, the spongy moth has cycles of large population outbreaks every eight to 12 years, leading to widespread defoliation and nuisance from caterpillars.

Other non-native insects defoliate trees but typically do not cause significant mortality. Elm zigzag sawfly (*Aproceros leucopoda*) is a non-native defoliator of elm trees that was reported in Minnesota in July 2024 for the first time.



Problematic Native Species

Over browsing by deer can severely limit natural tree regeneration (Matonis et al., 2011; De Jager et al., 2013). Elevated deer populations also reduce populations of palatable ground layer plants, including orchids (Knapp and Wiegand, 2014), members of the lily family (Balgooyen and Waller, 1995; Augustine and Frelich, 1998; Fletcher et al., 2001), and rare species (McGraw and Furedi, 2005). Decades of cumulative impacts have resulted in profound changes to the composition of native plant communities (Wiegmann and Waller, 2006; Flagel et al., 2016; Frerker et al., 2014; Nuttle et al., 2014; Anderson et al., 2017; Sabo et al., 2017).



Diseases and Pathogens

Oaks are susceptible to several types of disease and pathogens, such as oak wilt (caused by the non-native fungus *Bretziella fagacearum*), sudden oak death (caused by the water mold *Phytophthora ramorum*), and bur oak blight (caused by a native fungal pathogen, *Tubakia iowensis*). Individual trees tend to be more susceptible when under stress, which may interact then with other threats, such as if future climatic conditions bring more frequent late summer droughts. Older oak trees are more susceptible to interacting stressors including drought, heavy seasonal precipitation, insects, and diseases. Wetter springs may increase the prevalence and severity of fungal diseases as spores spread from the diseased leaves by rainwater, exacerbated by earlier spring rainstorms brought on by Minnesota's changing climate.



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al., 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

Projections include increased temperatures for the growing season as well as increased winter temperatures (Ratcliffe et al., 2025). Increased wintertime temperatures may bring additional freeze-thaw cycling, put spring buds at risk of being impacted by frost, and increase the probability of phenological mismatch between plants and animals reliant on them (such as insects). Climate change will affect different types of trees and forests; for instance, monotype aspen stands are less resilient than mixed species forests, with moderate-high vulnerability to climate change (Handler et al., 2017).

Changes in Precipitation and Hydrology related to Climate



From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter,

drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

If late summer droughts become more frequent under future climate conditions, this could pose a challenge to oak heath, exacerbating the prevalence of disease and pathogens. Restoration activities could also be disrupted if there are multiple seasons with insufficient rain for a successful seeding or seedling planting. The ability for different seeds to emerge and thrive under altered climatic patterns may also influence community composition post-restoration. As noted above, as temperatures rise and drought events increase, oak is likely to become a more prevalent component of upland deciduous forests. In addition to precipitation and hydrology, extreme weather events (e.g., large wind events, ice storms, heavy snowfall) are expected to increase as temperatures increase, as well as the risk of wildfire (Ratcliffe et al., 2025).

Heavy rain events have been increasing in frequency and severity in Minnesota in recent decades. Forests on steep slopes, especially in southeastern Minnesota, are vulnerable to landslides during these events. An increase in rain events has already impacted the mesic hardwood forest at Nerstrand Big Woods State Park in Rice County. Much wetter than it was 100 years ago, the park has experienced tree mortality in approximately 12%, or 200 acres of its property. The trees impacted include maples, basswood, and oaks.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan (Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN), five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Examples of conservation actions are grouped below under these five strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Also note that some strategies, such as Strategy 3, collaborating with partners, could truly be applied to all actions to most broadly and effectively implement them.

Potential Conservation Actions for Mesic Hardwood Forest



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Identify high priority sites for designation as protected areas to facilitate long-term conservation outcomes and protection against conversion of forest to different land uses. High biodiversity areas or those providing connectivity between other protected areas may provide disproportionately high conservation value.
	Apply knowledge of SGCN habitat use into forest planning efforts to help balance habitat for SGCN and other values according to management requirements and objectives. Consider natural area designations and other ways to enhance and preserve key mature stands.
	Encourage sustainable forest management through planning when and where timber harvest can be used as an effective tool to address a range of values and priorities, including forest health and productivity, critical ecological functions, and habitat restoration from

Stressor	Action
	historical disturbance events. Incorporate considerations of forest compositional and structural diversity at meaningful scales, historic size and return intervals of disturbance regimes, biodiversity, rare features, wildlife needs, and diversity of native plant community growth stages. Ensure that a balance of forest habitat conditions (e.g. younger and older forest, small and large patch sizes) are available for plant and animal species across the landscape over time.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.

Stressor	Action
	Explore the potential use of silvopasture management regimes (the practice of intentionally combining management of trees, forage, and grazing as one integrated practice) to help restore some upland deciduous forests that do not have spring ephemerals or other sensitive SGCN species (UMN Extension: Silvopasture: Establishment and Management Principles for Minnesota 2013).
	Consider projected effects on high value conservation forests, old-growth forests, and Important Bird Areas when making decisions and policy that are projected to impact upland deciduous forests (e.g., mining and quarrying or oil and gas permitting, planning for new development of energy sources and any associated new road or trail construction).
	Encourage the use of existing corridors for new trails, roads, and utility corridor development to minimize the construction of new roads and trails that would increase forest fragmentation.
	Limit the use and operation of heavy equipment when road or site conditions are vulnerable to damage. For instance, depending on the site characteristics, conducting timber harvests during frozen ground conditions can reduce soil compaction and rutting impacts. Avoid rutting and damage to peat soils when harvesting spruce tops.
	Consider the ecological values of various forest ages and how they relate to SGCN habitat when determining desired future conditions and planning sustainable management to advance those conditions.
	Manage for a diversity of forest growth stages across the landscape to promote vegetation characteristics that maintain habitat for a wide range of plant and animal species, promote resiliency to insect and disease outbreaks, and mitigate the adverse effects of climate change. Ensure older forest and old forest qualities are retained on the landscape in amounts and areas that provide critical habitat to SGCN animals and plants; encourage management that sustains mature and old-growth forests in forest planning efforts. Use

Stressor	Action
	management practices to create young, early successional forests on the landscape to provide vegetation characteristics and edge habitat between young and mature forests that support plant and animal species across at least part of their life cycle.
	Retain old and dead standing trees (snags) in forests when practical as habitat for wildlife, including cavity-nesting birds. Retain old and dead standing trees (snags) in forests when practical as habitat for wildlife, including cavity-nesting birds, such as detailed in the Minnesota Forest Resource Council's Site-Level Forest Management Guidelines .
	Well-managed and contained fires will need to become a more important forest management tool in the near future to improve the chances of oak regeneration as mesic forests become less resilient and oak becomes more dominant in this changing climate scenario (Frelich et al., 2019).
	Managing for an increased diversity of dominant species in a site can assist in resilience toward future disease outbreaks, susceptibility to some pathogens, and invasions by non-native species. Assisting in the management of woodlands and forests that are resilient to pathogens and invasive species may require additional dedicated and certain funding. Also consider pairing invasives management with other incentivized management activities and prioritizing comingled activities to create efficiencies.
	Implement control strategies for non-native invasive species and ensure sufficient timeframe for follow-up management that extends for many years. Invasive shrubs require control strategies, prioritization, and funding to fight encroachment of additional invasive shrubs, such as buckthorn. Strategies need to be long-term, spanning many years to be most effective. Support funding for these activities and consider expanding restoration resources to make sure they can be applied to woodland systems, which are sometimes in between funding opportunities aimed at prairies or forests.
	Incorporate climate change into forestry planning, including climate-adapted silviculture. Implement strategies and approaches for climate-informed forest management. Reference resources from the Northern Institute of Applied Science , such as their adaptation workbook , or the Tree Species Projections for Ecological Sections in Minnesota from the Climate Change Response Network.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.

Stressor	Action
	Support local governments in measures that minimize fragmentation and encroachment into high quality forests and woodlands such as protecting high quality areas, zoning that guides housing developments and lot sizes that support local forests, and land use planning and development guidance.
	Support and participate in forest planning efforts by providing information on important habitat types, landscape configurations, and conditions most beneficial for SGCN (https://mn.gov/frc/landscape/). Pursue opportunities to collaborate across forest ownerships and encourage the implementation of sustainable forestry at the site and landscape levels to create and maintain habitat conditions important to SGCN (e.g., unfragmented older mesic hardwood forests). Values such as regional biodiversity, rare species protection, native plant community growth stage diversity, and habitat connectivity between forest patches to allow wildlife movement can't be met through individual site-level management and need to be considered at larger landscape scales (DNR 2025).



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.

Stressor	Action
	Monitor the responses of SGCN to various types of forest management at the site-level and landscape-scale. Collaborate with MFRC site-level guideline monitoring and NPC silvicultural case studies to incorporate SGCN monitoring (https://silvlib.cfans.umn.edu/all-case-studies). Develop models of habitat configurations for various SGCN and consider how current harvest forecasts and conversion rates (i.e., one forest type to another; natural origin habitat to plantation) relate to those models in terms of sustainability for maintaining habitat for SGCNs.
	Support research on ways to remove or reduce prevalence and/or impacts of earthworms in invaded forests. Examples include the Minnesota Invasive Terrestrial Plants and Pests Center work with invasive jumping worms (https://mitppc.umn.edu/research/research-projects).
	Conduct research and model anticipated changes with different forest management scenarios for impacts on wildlife habitat.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Stressor	Action
	Promote and consider further development of tax incentives that enable landowners and farmers to gain value for maintaining conservation values on their properties. Provide increased levels of incentive payments for woodland owners to help protect woodlands and forests from further reduction due to development pressure.
	Support and promote education and technical assistance to maintain or restore forest habitat.
	Where people recreate, provide outreach and education on cleaning gear and equipment to reduce the introduction and spread of invasive species. DNR's Prevent the spread webpage includes specific actions for different land based activities like biking, hiking and off-highway vehicle riding. At trailheads, add invasive species prevention messages, such as those on the boot brush kiosks developed by the PlayCleanGo: Stop Invasive Species in Your Tracks program. Prioritize high recreation use areas where invasive species have not yet established.
	Provide information on important habitat types, landscape configurations, and conditions most beneficial for SGCN. Support forest management that reflects ecological disturbance goals and SGCN habitat needs such as stable amounts of older forest and maintenance of within-stand tree diversity. Communicate and document the benefits of alternative management techniques (i.e., variable density thinning, natural seeding), reference to natural disturbance intervals, natural regeneration methods, and to increase habitat quality for SGCN. Promote consideration of more flexibility in stocking goals in natural stands (i.e., longer time frames to meet goals, non-crop trees can be a significant portion of goals) and the ability to protect really important biodiversity areas from harvest. Support forest management approaches that are informed by ecological processes, increase structural and compositional complexity and diversity, and assist in resilience towards future stressors (e.g., climate change, disease outbreaks, invasion by non-native species) in forests. Refer to the Native Plant Community silvicultural strategies for forest stand prescriptions as well as Forestry for Minnesota Birds ForestryForMNBirdsGuidebook.pdf
	Research and communicate the benefits of alternative management techniques (i.e., variable density thinning, natural seeding), reference to natural disturbance intervals, natural regeneration methods, and to increase habitat quality for SGCN. Promote consideration of more flexibility in stocking goals in natural stands (i.e., longer time frames to meet goals, non-crop trees can be a significant portion of goals) and the ability to protect important biodiversity areas from harvest.

Stressor	Action
	Engage with stakeholders, including Minnesota DNR Division of Fish and Wildlife, to inform goal- setting processes to establish ecologically and socially desirable deer population levels across the state. Encourage bud capping, fencing, and slash walls to mitigate potential impacts from deer browse.

References for Mesic Hardwood Forests

Anderson, R.C., Anderson, M.R. and Corbett, E.A., (2017). White-tailed deer (*Odocoileus virginianus*) and fire effects on flowering diversity of tallgrass prairie forbs. *The Journal of the Torrey Botanical Society*, 144(3), pp.243-253

Augustine, D. J., & Frelich, L. E. (1998). Effects of White-Tailed Deer on Populations of an Understory Forb in Fragmented Deciduous Forests. *Conservation Biology*, 12(5), 995–1004. <https://doi.org/10.1046/j.1523-1739.1998.97248.x>

Bal, T.L., Storer, A.J. and Jurgensen, M.F., (2018). Evidence of damage from exotic invasive earthworm activity was highly correlated to sugar maple dieback in the Upper Great Lakes region. *Biological invasions*, 20, pp.151-164.

Balگوoyen, C. P. and D. M. Waller. (1995). The use of *Clintonia borealis* and other indicators to gauge impacts of white-tailed deer on plant communities in northern Wisconsin, USA. *Natural Areas Journal*, 15(4), 308-318. <https://www.cabidigitallibrary.org/doi/full/10.5555/19960610148>

De Jager, N.R., Cogger, B.J. and Thomsen, M.A., (2013). Interactive effects of flooding and deer (*Odocoileus virginianus*) browsing on floodplain forest recruitment. *Forest Ecology and Management*, 303, pp.11-19.

Fahey, T.J., Yavitt, J.B., Sherman, R.E., Maerz, J.C., Groffman, P.M., Fisk, M.C. and Bohlen, P.J., (2013). Earthworm effects on the incorporation of litter C and N into soil organic matter in a sugar maple forest. *Ecological Applications*, 23(5), pp.1185-1201.

Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniat, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4

Flagel, D.G., Belovsky, G.E. and Beyer, D.E., (2016). Natural and experimental tests of trophic cascades: gray wolves and white-tailed deer in a Great Lakes forest. *Oecologia*, 180, pp.1183-1194

Fletcher, D.J., Shipley, L. A., McShea, W. J., & Shumway, D. L. (2001). Wildlife herbivory and rare plants: The effects of white-tailed deer, rodents, and insects on growth and survival of Turk's cap lily. *Biological Conservation*, 101(2), 229–238. [https://doi.org/10.1016/S0006-3207\(01\)00070-2](https://doi.org/10.1016/S0006-3207(01)00070-2)

Frelich, L.E., Blossey, B., Cameron, E.K., Dávalos, A., Eisenhauer, N., Fahey, T., Ferlian, O., Groffman, P.M., Larson, E., Loss, S.R. and Maerz, J.C., (2019). Side-swiped: ecological cascades emanating from earthworm invasions. *Frontiers in Ecology and the Environment*, 17(9), pp.502-510.

Frelich, L.E., Peterson, R.O., Dovčiak, M., Reich, P.B., Vucetich, J.A. and Eisenhauer, N., (2012). Trophic cascades, invasive species and body-size hierarchies interactively modulate climate change responses of ecotonal temperate–boreal forest. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1605), pp.2955-2961.

Frerker, K., Sabo, A. and Waller, D., (2014). Long-term regional shifts in plant community composition are largely explained by local deer impact experiments. *PloS one*, 9(12), p.e115843.

Garrett, H.E. *et al.* (2004). Hardwood silvopasture management in North America. In: Nair, P.K.R., Rao, M.R., Buck, L.E. (eds) *New Vistas in Agroforestry. Advances in Agroforestry*, vol 1. Springer, Dordrecht.
https://doi.org/10.1007/978-94-017-2424-1_2.

Gundale, M.J., (2002). Influence of exotic earthworms on the soil organic horizon and the rare fern *Botrychium mormo*. *Conservation Biology*, 16(6), pp.1555-1561.

Hale, C.M., Frelich, L.E. and Reich, P.B., (2006). Changes in hardwood forest understory plant communities in response to European earthworm invasions. *Ecology*, 87(7), pp.1637-1649.

Hale, C.M., Frelich, L.E., Reich, P.B. and Pastor, J., (2005). Effects of European earthworm invasion on soil characteristics in northern hardwood forests of Minnesota, USA. *Ecosystems*, 8, pp.911-927.

Handler et al. (2017). Climate change field guide for Northern Minnesota Forests: site-level considerations and adaptation. DOI : <https://doi.org/10.32747/2017.6949547.ch>

He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>

Holdsworth, A.R., Frelich, L.E. and Reich, P.B., (2007). Effects of earthworm invasion on plant species richness in northern hardwood forests. *Conservation Biology*, 21(4), pp.997-1008.

Holdsworth, A.R., Frelich, L.E. and Reich, P.B., (2008). Litter decomposition in earthworm-invaded northern hardwood forests: role of invasion degree and litter chemistry. *Ecoscience*, 15(4), pp.536-544.

Holdsworth, A.R., Frelich, L.E. and Reich, P.B., (2012). Leaf litter disappearance in earthworm-invaded northern hardwood forests: role of tree species and the chemistry and diversity of litter. *Ecosystems*, 15, pp.913-926.

Knapp, W. M., & Wiegand, R. (2014). Orchid (Orchidaceae) decline in the Catoctin Mountains, Frederick County, Maryland as documented by a long-term dataset. *Biodiversity and Conservation*, 23(8), 1965–1976.
<https://doi.org/10.1007/s10531-014-0698-2>

- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Maerz, J.C., Nuzzo, V.A. and Blossey, B., (2009). Declines in woodland salamander abundance associated with non-native earthworm and plant invasions. *Conservation Biology*, 23(4), pp.975-981.
- Matonis, M.S., Walters, M.B. and Millington, J.D., (2011). Gap-, stand-, and landscape-scale factors contribute to poor sugar maple regeneration after timber harvest. *Forest Ecology and Management*, 262(2), pp.286-298.
- McCay, T.S. and Scull, P., (2019). Invasive lumbricid earthworms in northeastern North American forests and consequences for leaf-litter fauna. *Biological Invasions*, 21, pp.2081-2093.
- McGraw, J. B., & Furedi, M. A. (2005). Deer Browsing and Population Viability of a Forest Understory Plant. *Science*, 307(5711), 920–922. <https://doi.org/10.1126/science.1107036>
- Minnesota Department of Natural Resources. (2005). Field Guide to the Native Plant Communities of Minnesota: The Eastern Broadleaf Forest Province. Ecological Land Classification Program, Minnesota County Biological Survey, and Natural Heritage and Nongame Research Program. DNR St. Paul, MN.
- Moriarty, J.J., Hall, C.D. (2014). Amphibians and Reptiles in Minnesota. University of Minnesota Press.
- Nuttle, T., Ristau, T.E. and Royo, A.A., (2014). Long-term biological legacies of herbivore density in a landscape-scale experiment: forest understoreys reflect past deer density treatments for at least 20 years. *Journal of Ecology*, 102(1), pp.221-228
- Nuzzo, V.A., Maerz, J.C. and Blossey, B., (2009). Earthworm invasion as the driving force behind plant invasion and community change in northeastern North American forests. *Conservation Biology*, 23(4), pp.966-974.
- Paudel, S., Longcore, T., MacDonald, B., McCormick, M.K., Szlavecz, K., Wilson, G.W. and Loss, S.R., (2016). Belowground interactions with aboveground consequences: Invasive earthworms and arbuscular mycorrhizal fungi. *Ecology*, 97(3), pp.605-614.
- Radeloff, V. C., Hammer, R. B., Stewart, S. I., Fried, J. S., Holcomb, S. S., & McKeefry, J. F. (2005). The Wildland–Urban Interface in the United States. *Ecological Applications*, 15(3), 799–805. <https://doi.org/10.1890/04-1413>
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Appalachian-Interior-Northeast Mesic Forest. Midwest Climate Adaptation Science Center. (97 pages) Available: <https://mwcasc.umn.edu/node/821>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Sabo, A.E., Frerker, K.L., Waller, D.M. and Kruger, E.L., (2017). Deer-mediated changes in environment compound the direct impacts of herbivory on understorey plant communities. *Journal of Ecology*, 105(5), pp.1386-1398

- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, e14434. <https://doi.org/10.1111/cobi.14434>
- Sonter, L. J., Ali, S. H., & Watson, J. E. M. (2018). Mining and biodiversity: Key issues and research needs in conservation science. *Proceedings of the Royal Society B: Biological Sciences*, 285(1892), 20181926. <https://doi.org/10.1098/rspb.2018.1926>
- Whitfeld, T.J., Roth, A.M., Lodge, A.G., Eisenhauer, N., Frelich, L.E. and Reich, P.B., (2014). Resident plant diversity and introduced earthworms have contrasting effects on the success of invasive plants. *Biological invasions*, 16, pp.2181-2193.
- Wiegmann, S. M., & Waller, D. M. (2006). Fifty years of change in northern upland forest understories: Identity and traits of “winner” and “loser” plant species. *Biological Conservation*, 129(1), 109–123. <https://doi.org/10.1016/j.biocon.2005.10.027>
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55). <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>
- Zamora, D., Wyatt, G., Buttler, M., Ford, M., Magner, J., Reichenbach, M., Burkett, E., Current, D., Walter, D. (2017). Silvopasture: Establishment and Management Principles for Minnesota. Retrieved from the University Digital Conservancy, <https://hdl.handle.net/11299/214976>.
- Zenner, E. K., & Berger, A. L. (2008). Influence of skidder traffic and canopy removal intensities on the ground flora in a clearcut-with-reserves northern hardwood stand in Minnesota, USA. *Forest Ecology and Management*, 256(10), 1785–1794. <https://doi.org/10.1016/j.foreco.2008.05.030>
- Ziemba, J.L., Hickerson, C.A.M. and Anthony, C.D., (2016). Invasive Asian earthworms negatively impact keystone terrestrial salamanders. *PloS one*, 11(5), p.e0151591.
- Zlonis, K.J. and Henderson, B.W., (2018). Invasive earthworm damage predicts occupancy of a threatened forest fern: implications for conservation and management. *Forest Ecology and Management*, 430, pp.291-298.

Deciduous Wet Forest

Habitat Description

Black ash (*Fraxinus nigra*) is the dominant canopy tree in swamp forests statewide; yellow birch (*Betula alleghaniensis*), red maple (*Acer rubrum*), aspens (*Populus spp.*), and balsam poplar (*Populus balsamifera*) are common associates in the Laurentian Mixed Forest Province; and basswood (*Tilia americana*), elms (*Ulmus spp.*), paper birch (*Betula papyrifera*), and yellow birch in the Eastern Broadleaf Forest Province. Conifer species, especially white cedar (*Thuja occidentalis*) and balsam fir (*Abies balsamea*) are sometimes present in the canopy and understory (see also the Coniferous Wetland Forest chapter).

In southeastern Minnesota, this habitat occurs as small patches in areas of groundwater seepage, usually at the base of steep slopes on level river terraces. In such settings, the water table is almost always within reach of plant roots but does not remain above the mineral soil surface for long periods during the growing season.

In northern Minnesota, larger basins connected to shallow aquifers create more expansive ash swamps with occasional springs and moving groundwater.

Understories are characterized by patches of shrubs, including speckled alder (*Alnus incana*), mountain maple (*Acer spicatum*), dogwoods (*Cornus spp.*), gooseberries or currants (*Ribes spp.*), and winterberry (*Ilex verticillata*). Mosses and upland forest herbs occur on raised hummocks, down logs, and tip-up mounds, and sedges and wetland forbs occur in wet or mucky hollows.

Soils in these deciduous wet forests are peaty or mucky mineral soils.

Habitat Map

To depict Deciduous Wet Forest habitat (see Figure 3.11), we compiled spatial data from several sources: DNR's Native Plant Communities, the MN National Wetland Inventory, and the Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Wet Forest (WF)

[WFn55 Northern Wet Ash Swamp \(PDF\)](#)

[WFn64 Northern Very Wet Ash Swamp \(PDF\)](#)

[WFW54 Northwestern Wet Aspen Forest \(PDF\)](#)

[WFS57 Southern Wet Ash Swamp \(PDF\)](#)

[WFS55 Southern Wet Aspen Forest \(PDF\)](#)

National Wetlands Inventory

From the [National Wetland Inventory for Minnesota \(NWI\) Layer](#) we included the Hardwood Wetland Simplified Plant Community Class type.

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Eastern North American Ruderal Flooded and Swamp Forest.




Figure 3.11. A map depicting Deciduous Wet Forest habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological Systems of Wet Forest, one type from the National Wetland Inventory, and one type from the Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

Deciduous wet forests are found in shallow depressions or in narrow zones along the margins of lakes, wetlands, and peatlands throughout Minnesota. These swamp forests have high plant species diversity. Due to their seasonally saturated conditions, most deciduous wet forest ecosystems have not been largely developed or degraded and are still generally intact. However, threats to these habitats include altered hydrology from a variety of sources (e.g., ditching, utility right-of-ways) and invasive species impacts, especially including emerald ash borer (*Agrilus planipennis*) expansion. The emerald ash borer is an invasive insect from Asia (UMN 2024) that has killed millions of ash trees throughout the eastern half of the U.S. and southeastern Canada.

The expanding presence of emerald ash borer in Minnesota, and responsive potential increase in [forest management activities](#) in these wet forests ([DNR 2020](#)) may eventually lead to a significant reduction in black ash as a predominant tree in wet deciduous forests. The loss of ash that is not replaced by other trees can lead to a transition of a wet forest to a wetland dominated more by shrubs and herbaceous vegetation and result in declines in forest-dependent wildlife species (Grinde et al., 2021).

Climate Profile for Deciduous Wet Forest

Climate impacts to Deciduous Wet Forest are described in the section on the Laurentian-Acadian Alkaline Swamp in the report “Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest” published by the Midwest Climate Adaptation Science Center (Ratcliffe et al., 2025). The most impactful changes in climate for this system include increases in spring rainfall of 25% over historical by late century, with more frequent and larger flood events to increase the duration of and level of flooding in these forests. Additionally, higher temperatures combined with less summer precipitation creates vapor pressure deficits (VPD; driven by higher evapotranspiration) projected to be 72% higher by end of century over historical (1971-2000). The most impactful changes in climate for this system include more frequent and larger flood events, also with more frequent late summer drought periods.

A major concern for Deciduous Wet Forests is more frequent and more intense flooding events. More precipitation will mean more sediment deposition as well as habitat changes. This projected increase in flooding will favor pioneer and early successional species, while limiting the growth of more stable species. Further habitat change will occur during more frequent summer droughts, which will impact plant growth and favor drought dependent species. These climate-induced changes may promote the spread of invasive species. Lastly, pests and pathogen outbreaks may have a greater impact on Deciduous Wet Forests as pest ranges expand, populations increase, and hosts become more susceptible to infection. Large increase in flood frequency and intensity is projected to occur in the spring, as both winter and spring precipitation is projected to increase. Coupled with warmer temperatures leading to an earlier snowmelt, large floods are predicted to continue to greatly impact Deciduous Wet Forests.

This habitat will also be impacted by warmer and dryer summers, which will cause soil drying and an overall reduction of habitat in these water-dependent ecosystems.

Key Climate Change Impacts:

- **Habitat Structure:** More frequent flood and droughts will alter species composition in this habitat type, especially among wetland adapted species. Windthrow events may become more frequent due to more flooding, less frozen soils, and more pest pressure.
- **Community Composition:** Plant communities will be greatly impacted by drier summer and more frequent flooding. Regeneration will become increasingly difficult under these conditions, and better adapted species will begin to dominate these habitats.
- **Invasive Species:** Flooding events create opportunities for invasive species to establish their populations, and as these flooding events become more common, so do the invasive species.
- **Pests and Pathogens:** Climate change will both increase the number of pests and pathogens and will make hosts more susceptible to the pests and pathogens. In some cases, like heatwaves, pests and pathogens may also see declines in population.
- **Herbivory:** Deer populations are expected to grow as climate change brings warmer winters and greater food availability. This increase in deer population will favor browse-tolerant species as well as disrupt tree regeneration.

Species in Greatest Conservation Need

Deciduous wet forests provide primary or secondary habitat for 27 animal and 7 plant SGCN (see Table 3.10). Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associate each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants).

Table 3.10. Numbers of Species in Greatest Conservation Need associated with the Deciduous Wet Forest as either primary or secondary habitat¹.

Species Group	Primary Habitat	Secondary Habitat	Total
Amphibians	0	3	3
Birds	0	4	4
Mammals	1	11	12
Reptiles	3	0	3
Aquatic Invertebrates			
Dragonflies & damselflies			1
Terrestrial Invertebrates			
Moths			1
Snails (terrestrial)	0	1	1
Spiders	0	2	2
Plants	7		7
Total	11	21	34

1. Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Of the 131 terrestrial vertebrate SGCN identified in the 2025-2035 SWAP, 17% (22) occur in deciduous wet forests. 38% of the 32 mammal SGCN occur in this habitat. The 7 plant SGCN found in this wet forest habitat represent less than 2% of all plant SGCN. Examples of selected SGCN are described below.

Amphibians

Two salamander SGCN – [spotted salamander \(*Ambystoma maculatum*\)](#) and [four-toed salamander \(*Hemidactylium scutatum*\)](#) – are associated with vernal pools (see Vernal Pool sub-chapter) and the mature Deciduous Wet Forests. The forests were identified as secondary habitat for the salamanders who breed in these areas each spring. Eggs are laid in or near the water, and larvae remain in the pools until metamorphosis occurs. Four-toed salamanders specifically lay their eggs in moss hummocks next to wetlands. Four-toed

salamanders occur most frequently in mature upland forests on glacial moraine landscapes with frequent isolated wetlands that include an alder margin and moss hummocks adjacent to pockets of open water.

Birds

Most SGCN birds found in other deciduous forest types may also utilize Wet Forests, but none exclusively. [Cerulean warblers \(*Setophaga cerulea*\)](#), [Louisiana waterthrushes \(*Parkesia motacilla*\)](#), and [red-shouldered hawks \(*Buteo lineatus*\)](#) may be present in Wet Forests, but primarily when part of a mosaic of large tracts of mature Floodplain or Mesic Hardwood forests. In such cases, key habitat features such as flowing streams or seepage channels for Louisiana waterthrushes, or large-diameter trees near wetland openings for red-shouldered hawks, must be present. Red-headed woodpeckers (*Melanerpes erythrocephalus*) may frequent black ash Deciduous Wet Forests when suitable dead trees are present.

Mammals

Eleven of the 12 mammal SGCN that occur in Deciduous Wet Forest were found in multiple other habitats ranging from upland fire-dependent communities and riparian forests to prairies and other grasslands, savannas, and even urban and other developed lands. These habitat generalists include five bat species: the [big brown bat \(*Eptesicus fuscus*\)](#), silver-haired bat (*Lasionycteris noctivagans*), Eastern red bat (*Lasiurus borealis*), Hoary bat (*Lasiurus cinereus*) and [little brown myotis \(*Myotis lucifugus*\)](#). A cave-hibernating species, the little brown bat has been heavily impacted by white-nose syndrome.

The [woodland vole \(*Microtus pinetorum*\)](#) is the only species for which Deciduous Wet Forest has been designated its primary habitat. Other primary habitats for the species include Mesic Hardwoods and Riparian and Floodplain Forests. They prefer loose, well-drained soils and humus layers to construct burrows and sub-surface runways. Grazing by cattle, which compacts the soil, and the presence of invasive non-native earthworms, which destroy the humus, may make forests within its' limited range in southeastern Minnesota unsuitable for this species.

Reptiles

Deciduous Wet Forest provides primary habitat for three reptile SGCN: [wood turtle \(*Glyptemys insculpta*\)](#), Eastern hog-nosed snake (*Heterodon platirhinos*) and [Western ratsnake \(*Pantherophis obsoletus*\)](#). Deciduous Wet Forest is just one of 5 forested habitats that have been designated primary habitat for the wood turtle. Each is suitable as long as fast-moving rivers and streams are embedded within deciduous or coniferous forests. Nevertheless, it is not entirely restricted to forests. It has been found in open habitats including prairies, grasslands, and savannas which, if located near flowing waters, can provide critical foraging and nesting habitat.

Even though the Eastern hog-nosed snake feeds almost exclusively on amphibians it also is not restricted to wet habitats but occupies forest edges and sites with sandy soils where the ground is conducive to burrowing. A total of 5 different habitats were identified as primary habitat for the snake. Four habitats have been designated as providing primary habitat for the Western Ratsnake: Upland Deciduous Forest, Mesic Hardwoods, Deciduous Wet Forest, and Savanna. All four can be found in Fillmore and Houston County where the species resides.

Plants

The relatively nutrient rich and seasonally saturated soils of deciduous wet forests present a challenging and competitive habitat for plant species. These sites can host some unique and rare plants that occupy specific niches in the ground layer of swamps and seepages. Bog bluegrass (*Poa paludigena*) occurs only in groundwater-connected forested wetland habitats, where it grows on mossy logs and hummocks above deeper mucky pools under shady black ash canopies. The annual, spring ephemeral forb, [false mermaid](#) (*Floerkea proserpinacoides*) grows in localized seepage habitats beneath mature black ash, basswood, and American elm overstories. Both species and their respective wet forest habitats are sensitive to changes in local hydrology as well as impacts from invasive exotic species.

Lichens

Deciduous wet forests are also home to unique lichen species. Lichen ‘species’ are a complex that consists of two or more, typically obligate, symbionts. The dominant partner is a fungus, which creates the main body of the organism. The endosymbiont is either a green alga or a cyanobacteria that lives within the fungal tissue. [Caloplaca parvula](#) is a Minnesota State Endangered species endemic to the Great Lakes region. It has been documented in two counties in Northeast Minnesota and is found almost exclusively on black ash trees. This species is at risk of being lost from the state as emerald ash borer continues to spread throughout Minnesota.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Roads, Trails, and Railroads

Roads, trails, and railroads can create fragmentation and edge effects. Increase in the number of roads and trails, particularly for motorized vehicles, increases human disturbance and introduction of invasive species into forests. Repeated use and soil compaction associated with roads and trails also can modify the vegetation and hydrology in the area.



Utility Corridors

Utility corridors transect deciduous wet forests throughout the state. These corridors can contain a variety of utility infrastructure. This includes pipelines, telephone lines, and power lines. These lines can be above ground, or below ground. Regardless of type, utility corridors can cause fragmentation and habitat degradation based on their size, purpose, and the upkeep that is required to maintain them.

The initial construction of the associated utility infrastructure as well as the long-term maintenance of that infrastructure can have lasting impacts on native plant communities and the SGCN associated with them. Pipelines have the potential to change the hydrology of deciduous wet forests through impoundment of water and changing of surface and groundwater flow.

These corridors and the increased travel from vehicles and equipment that they require act as vectors for invasive plant species. The required vegetative management practices often include herbicide applications, removal of woody vegetation, and/or mowing. This increased disturbance increases the likelihood of establishing invasive plant species within the corridors and then spreading to adjacent lands.



Timber Harvest

Timber harvest activity in deciduous wet forest habitats has generally been limited in scope and intensity due to the constraints of operating on saturated soils that don’t reliably freeze in winter. When timber harvesting does occur, uneven aged silvicultural practices are generally applied (e.g., strip or gap

harvest, hardwood thinning) and retain greater amounts of canopy trees to maintain evapotranspiration rates and local hydrologic conditions. The expanding presence of emerald ash borer in Minnesota, and a potential increase in [forest management activities](#) in these wet forests ([DNR 2020](#)) may lead, eventually, to a significant reduction in black ash as a predominant tree in wet deciduous forests. Pre-salvage timber harvest of ash has increased with the spread of EAB (Holt et al., 2021) and is used as a strategy to retain some timber value before EAB-driven disturbance. Supporting forest regeneration, including ash regeneration, after timber harvest remains critical (Higgins et al., 2025; Kolka et al., 2018).



Recreation

Recreation can spread invasive species, inadvertently being moved on people's shoes or gear. Moving wood for campfires can inadvertently move pests such as emerald ash borer in the firewood, so firewood is not to be moved across the state.



Invasive Species (Problematic Non-native Species)

Invasive species are ongoing stressor in forest and woodland systems. Specifically pertinent to black ash forests, emerald ash borer is an invasive insect from Asia (UMN 2024) that has killed millions of ash trees throughout the eastern half of the U.S. and southeastern Canada. Minnesota's deciduous wet forests can contain nearly pure stands of black ash trees, where they help maintain the water table. Black ash trees often form dense stands and have a relatively high rate of evapotranspiration (releasing water vapor from their leaves into the atmosphere) and this draws down the water table. Loss of these trees, such as by emerald ash borer mortality, the wet forest habitat may become more inundated and change over to grass, cattails, and shrubs, threatening the plants and animals that rely on black ash forest habitats.



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al., 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Higher growing season temperatures combined with less summer precipitation creates vapor pressure deficits (VPD; driven by higher evapotranspiration) projected to be 72% higher by the end of the century compared to historical levels (1971-2000; Ratcliffe et al., 2025). [Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts](#) (He et

al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota’s average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months (as much as 25%), which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

See Climate Profile inset for more information.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan (Protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN), five strategies were identified:

- 
Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.
- 
Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.
- 
Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.
- 
Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.
- 
Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Examples of conservation actions are grouped below under these five strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Also note that some strategies, such as Strategy 3, collaborating with partners, could truly be applied to all actions to most broadly and effectively implement them.

Potential Conservation Actions for Deciduous Wet Forest


Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.

Stressor	Action
	Explore a range of silvicultural treatments in an effort to utilize standing timber commercially ahead of anticipated ash mortality due to emerald ash borer and to establish non-ash species. When harvesting mature ash trees and promoting other tree species in wet forests, avoid clearcutting, as the maintenance of some larger trees will maintain site hydrology. Removing groups of ash (0.1 to 0.5 acre in size) is best, and strip clearcuts and strip shelterwoods are methods successful at promoting other species if those other species are present (e.g., red maple and balsam fir) (DNR 2025). Retain some ash, since they will provide habitat for wildlife when they die. Field studies on the Chippewa National Forest concluded that group selection, individual tree selection, and

Stressor	Action
	shelterwood harvest are good options. See also this handbook for private landowners managing ash woodlands (UMN 2019).
	For forests depleted by emerald ash borer, encourage more diverse tree species regeneration in an effort to maintain ecosystem functionality into the future. Consider the use of more southern species when establishing new species in black ash stands impacted by emerald ash borer, such as swamp white oak and hackberry (Palik et al., 2021).
	Buy local and burn local - be careful not to move firewood across the state (DNR 2025b), because that can inadvertently move invasive pest species such as emerald ash borer that can be in the firewood. Follow best practices and buy firewood adjacent to where it will be burned.
	Managing for forests that are resilient to pathogens and invasive species may require additional dedicated and stable funding. Implementing control strategies for invasive species, such as buckthorn, requires a sufficient timeframe for follow-up management that extends for many years. Support funding for these activities and consider expanding restoration resources to make sure they can be applied to woodland systems, which are sometimes in between funding opportunities aimed at prairies or forests.
	Manage for an increased diversity of dominant species to assist in establishing resiliency toward future disease outbreaks, susceptibility to some pathogens, and invasions by invasive species. In response to emerald ash borer dieback, promote a diversity of trees featuring non-ash species. Increasing forest diversity while maintaining water regulation and groundwater conditions suitable for forest growth are key strategies.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

SPACE

Stressor	Action
	In places where people recreate, do outreach and education on cleaning gear and equipment to reduce the introduction and spread of invasive species. At trailheads for motorized and non-motorized recreation, add invasive species prevention messages, such as those on the boot brush kiosks developed by the PlayCleanGo: Stop Invasive Species in Your Tracks program. DNR's Prevent the

Stressor	Action
	spread webpage includes specific actions for different land based activities like biking, hiking and off-highway vehicle riding.

References for Deciduous Wet Forest

Finch, D.M. et al. (2021). Effects of Climate Change on Invasive Species. In: Poland, T.M., Patel-Weyand, T., Finch, D.M., Miniati, C.F., Hayes, D.C., Lopez, V.M. (eds) *Invasive Species in Forests and Rangelands of the United States*. Springer, Cham. https://doi.org/10.1007/978-3-030-45367-1_4

Grinde, Alexis, R., Melissa B. Youngquist, Robert A. Slesak, Stephen R. Kolbe, Josh D. Bednar, Brian J. Palik, and Anthony W. D’Amato (2022). “Potential Impacts of Emerald Ash Borer and Adaptation Strategies on Wildlife Communities in Black Ash Wetlands.” *Ecological Applications* 32(4): e2567. <https://doi.org/10.1002/eap.2567>

He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>

Higgins, H., D’Amato, A. W., & Siegert, N. W. (2025). Effects of harvest treatments anticipating emerald ash borer invasion on northern hardwood forests in New England, USA. *Forest Ecology and Management*, 588, 122748. <https://doi.org/10.1016/j.foreco.2025.122748>

Holt, J. R., Smetzer, J. R., Borsuk, M. E., Laflower, D., Orwig, D. A., & Thompson, J. R. (2022). Emerald ash borer intensifies harvest regimes on private land. *Ecological Applications*, 32(2), e2508. <https://doi.org/10.1002/eap.2508>

Kastendick, Doug, Brian Palik, Owen Tegtmeier, and Mary Ulrich. 2024. Evaluating the ecological impacts of EAB and climate change on black ash forests (Chippewa NF). Great Lakes Silviculture Library. University of Minnesota (<https://silvlib.cfans.umn.edu/evaluating-ecological-impacts-eab-and-climate-change-black-ash-forests-chippewa-nf>).

Kolka, R., D’Amato, A., Wagenbrenner, J., Slesak, R., Pypker, T., Youngquist, M., Grinde, A., & Palik, B. (2018). Review of Ecosystem Level Impacts of Emerald Ash Borer on Black Ash Wetlands: What Does the Future Hold? *Forests*, 9(4), 179. <https://doi.org/10.3390/f9040179>

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>

Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). (2025). Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state’s greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>

DNR 2025. Emerald Ash Borer Management.

<https://www.dnr.state.mn.us/invasives/terrestrialanimals/eab/management.html>. Accessed 5/28/2025

DNR 2025b. Firewood Information. <https://www.dnr.state.mn.us/firewood/index.html>. Accessed 5/28/2025

DNR. Sustainable Timber Harvest Development of the DNR 10-yr Stand Exam List. (2020). chrome-extension://efaidnbnmnnibpcajpcgclclefindmkaj/https://files.dnr.state.mn.us/forestry/subsection/harvest-analysis/sth-10-year-stand-exam-list-report.pdf

Palik, B. J., D'Amato, A. W., Slesak, R. A., Kastendick, D., Looney, C., & Kragthorpe, J. (2021). Eighth-year survival and growth of planted replacement tree species in black ash (*Fraxinus nigra*) wetlands threatened by emerald ash borer in Minnesota, USA. *Forest Ecology and Management*, 484, 118958.

<https://doi.org/10.1016/j.foreco.2021.118958>

Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. (2025). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwcasc.umn.edu/node/821>

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.

<https://statesummaries.ncics.org/chapter/mn/>

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, e14434.

<https://doi.org/10.1111/cobi.14434>

University of Minnesota. (2019). Managing Ash Woodlands: Recommendations for Minnesota Woodland Owners. <https://hdl.handle.net/11299/205052>. Accessed 5/28/2025

University of Minnesota. (2024). Emerald Ash Borer in Minnesota. <https://extension.umn.edu/tree-and-shrub-insects/emerald-ash-borer> Accessed 5/28/2025

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55).

<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Coniferous Forest Wetland

Habitat Description

The Coniferous Forest Wetland habitat type includes wet cedar forests; forested rich peatlands (relatively nutrient rich often due to groundwater inputs) and conifer swamps and forested bogs, which are more acidic, associated with nutrient-poor substrates and a lack of connection to groundwater flow. Coniferous forest wetlands are found throughout the Laurentian Mixed Forest Province, with concentrations of large peatlands in the Agassiz Lowlands and Tamarack Lowlands ecological subsections. These forests are also present to a lesser degree in the Eastern Broadleaf Forest and Tallgrass Aspen Parklands Provinces. The soils are peat or mucky mineral soil that is usually saturated with water deficient in oxygen and low in nutrients. Coniferous forest wetlands occur as part of large peatland complexes ([Patterned Peatlands](#)) in association with open bogs (see also Non-forested Wetlands Sub-chapter) as well as in shallow basins along lakes, wetlands, or streams.

Coniferous forest wetlands are dominated by black spruce (*Picea mariana*), tamarack (*Larix laricina*), or white cedar (*Thuja occidentalis*). Tree height and density vary from nearly closed canopies with moderate heights (>30') on richer sites to scattered, stunted coniferous trees (<30') in the most nutrient-poor black spruce bogs. The understory of this habitat is characterized by a mossy ground layer with an abundance of forbs, sedges, and broad-leaved evergreen shrubs. Brown mosses predominate in the richer environments, whereas the more acid-loving species of Sphagnum dominate the bogs. Typical shrubs include labrador tea (*Rhododendron groenlandicum*), leatherleaf (*Chamaedaphne calyculata*), and bog rosemary (*Andromeda glaucophylla*). Richer examples of this habitat (that is, swamps more so than bogs) also support species characteristic of surrounding upland forests, but these species are limited to tree bases and moss hummocks elevated above the water table.

Plant adaptations to the harsh growing conditions in lowland conifer forests include evergreen leaves (conifers and ericaceous shrubs), reliance on ectomycorrhizal fungi to facilitate nutrient uptake, capture of insects to provide additional nutrients (pitcher plants [*Sarracenia purpurea*] and sundews [*Drosera* spp.]), and secondary compounds in leaves to reduce herbivory.

Habitat Map

To depict Coniferous Forest Wetland habitat (see Figure 3.12), we compiled spatial data from several sources: DNR's Native Plant Communities, the MN National Wetland Inventory, and the Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods* in Chapter 3. *Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Acid Peatland (AP)

[APn81 Northern Poor Conifer Swamp \(PDF\)](#)

[APn80 Northern Spruce Bog \(PDF\)](#)

Forested Rich Peatland (FP)

[FPn63 Northern Cedar Swamp \(PDF\)](#)

[FPn62 Northern Rich Spruce Swamp \(Basin\) \(PDF\)](#)
[FPn71 Northern Rich Spruce Swamp \(Water Track\) \(PDF\)](#)
[FPn72 Northern Rich Tamarack Swamp \(Eastern Basin\) \(PDF\)](#)
[FPn81 Northern Rich Tamarack Swamp \(Water track\) \(PDF\)](#)
[FPn82 Northern Rich Tamarack Swamp \(Western Basin\) \(PDF\)](#)
[FPw63 Northwestern Rich Conifer Swamp \(PDF\)](#)
[FPs63 Southern Rich Conifer Swamp \(PDF\)](#)
[FPn73 Northern Rich Alder Swamp \(PDF\)](#)

Wet Forest (WF)

[WFn53 Northern Wet Cedar Forest \(PDF\)](#)

National Wetlands Inventory

From the [National Wetland Inventory for Minnesota \(NWI\) Layer](#) we included the Coniferous Bog and Shrub Wetland Simplified Plant Community Class types.

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Eastern North American Boreal-Subboreal Bog and Acidic Fen, Laurentian-Acadian Alkaline Swamp, and Laurentian Acadian Wet Meadow and Shrub Swamp.




Figure 3.12. A map depicting Coniferous Forest Wetland habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological Systems of Wet Forest, Forested Rich Peatland, and Acid Peatland, two types from the National Wetland Inventory, and three types from the Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

Despite numerous attempts at drainage in the early 20th century, lowland conifer forests still cover vast areas, primarily in large peatlands complexes in the northern part of the Laurentian Mixed Forest Province. Minnesota's coniferous wetlands, specifically peatlands, are not only unique because of their expanse, remoteness, and unique landscape forms (e.g., patterned water tracks and raised bogs), they also are a significant component of the global carbon cycle. Like all vegetation, peatland plants absorb carbon dioxide for photosynthesis. Unlike vegetation on drier or mesic soils, the water-logged, oxygen-deficient peatland soils reduce the rate of plant decomposition, thereby slowing the release of carbon dioxide, a major contributor to greenhouse gas emissions. Although peatlands only cover 3% of the world's land area, they collectively store nearly 30% of the world's carbon, thus reducing the amount of carbon in the atmosphere and helping to mitigate the impact of warming temperatures due to the release of greenhouse gas emissions. In Minnesota alone, peatlands cover at least 10% of the land area and store over 4 billion metric tons of carbon which accounts for at least 37% of the state's stored terrestrial carbon (Walker, 2011).

According to the DNR Native Plant Community (NPC) Field Guides (listed under Associated Native Plant Community Classes above), prior to European settlement catastrophic disturbances within the Coniferous Forest Wetland habitat type were relatively uncommon. These events occurred approximately every 365 to over 1,000 years. Smaller disturbances resulting in partial mortality of the forest canopy were somewhat common and usually consisted of patchy blowdowns and surface fires. The frequency of these events varied greatly across the habitat type from 40 to 90 years on many sites.

Much of the coniferous forest wetland habitat in Minnesota occurs on publicly managed lands. Sites with merchantable densities or volumes of trees are identified for timber harvesting using even-aged silvicultural practices (generally clearcutting) to produce pulpwood or firewood, address forest health concerns, and incidentally create young forest habitat. Black spruce, and to a lesser extent tamarack, are the main commercially important tree species in this habitat, as white cedar is often difficult to regenerate and has historically had a lower harvest rate. Although the total amount of timber harvested each year in Minnesota has trended downward since 1998, the black spruce harvest has slowly increased; although very small, there also is a slight increase in the harvest and use of white cedar (Llorente, 2020). Together black spruce and tamarack account for approximately 9% of the total harvest.

Climate Profile for Coniferous Forest Wetland

Coniferous forest wetlands, as they are a group of several habitats, were profiled in two reports published by the United States Geological Survey Midwest Climate Adaptation Science Center in support of State Wildlife Action Plan revisions: "Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest" (Ratcliffe et al., 2025a), and "Effects of Climate Change on Midwestern Ecosystems: North American Bog and Fen" (Ratcliffe et al., 2025b). We draw content from both reports in this section, and all content in this section is derived from those reports.

For swamp forests, the most impactful changes in climate include more frequent and larger flood events and more frequent late summer droughts. For boreal-subboreal bogs and acidic fens, warming temperatures with declines in summer rainfall will increase evapotranspiration (vapor pressure deficits; VPD). Northern bogs and fens, already at the southern extent of their range, may be particularly stressed by these shifts in hydrologic conditions and warming trends (Ratcliffe et al., 2025b).

In the Northern Forest ecoregion that includes northeastern Minnesota, projected changes in annual temperature by end of century (2070-2099) are 6 – 11 degrees higher than historic (1970-2000), depending on the carbon emissions level and using the mean of 20 climate models (see Table 9 in Ratcliffe et al., 2025b). Rainfall projections include greater expected winter and spring rain and more summer drought.

Wet Forests

More flooding will mean more erosion and increase the frequency of succession cycles. Pioneer species will be favored under these conditions. Drought-tolerant species will also be favored as summers become drier and hotter. These changes may lead to less species diversity, especially in fragile wetland ecosystems. Pests, pathogens, and invasive species will all benefit from these changes while more sensitive native plants will be hurt the most. Lastly, herbivores will see population increases and do more damage to selected forage plants.

Key Climate Change Impacts:

- **Habitat Structure:** More frequent flooding will benefit flood-tolerant and early successional species. Early successional species will also benefit during the summer as wildfires become more common. Trees may become more vulnerable to windthrow events as soil conditions changes, and warmer temperatures increase pest and pathogen viability.
- **Community Composition:** Increases in flooding and droughts will impact the plant community. Warmer temperatures may force some species to move north, and extreme heat during summer will increase the risk of seedling mortality. Again, early successional species will benefit from these changes.
- **Invasive Species:** More frequent disturbance will provide more opportunities for invasive species to establish themselves.
- **Pests and Pathogens:** Pest and pathogen populations are expected to increase as warmer winters prevent large die-offs. Hosts are also expected to become more susceptible to pests and pathogens during large droughts and other climate-related events.
- **Herbivory:** Deer are expected to increase in population as winters become warmer, and food becomes more available. This will lead to overpopulation, negatively impacting understory plant populations, as well as natural regeneration. Lastly, deer-browse resistant species, like black spruce, will gain an advantage over other more selected species.

Eastern North American Boreal-Subboreal Bog and Acidic Fen

This habitat type will be hit particularly hard by dryer and hotter summers.

Key Climate Change Impacts:

- **Habitat Structure:** Drier conditions will mean lower water tables, which will hurt important species such as sphagnum moss (*Sphagnum spp.*). These drier conditions will mean reduced water retention and more frequent wildfire.
- **Community Composition:** Increase in temperatures are predicted to hurt species like black spruce and tamarack, while benefiting species such as highbush blueberry (*Vaccinium corybosum*).
- **Invasive Species:** Longer growing seasons and more available nutrients will benefit invasives such as reed canary grass (*Phalaris arundinacea*) and cattail (*Typha spp.*).

- **Pests and Pathogens:** Climate change will make pests and pathogens more viable and will make hosts more susceptible to their effects.

Species in Greatest Conservation Need

Coniferous forest wetlands provide primary or secondary habitats for 37 animal and 21 plant SGCN (see Table 3.11). Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Table 3.11. Numbers of Species in Greatest Conservation Need associated with the Coniferous Forest Wetlands as either primary or secondary habitat¹.

Species Groups	Primary Habitat	Secondary Habitat	Total
Birds	9	3	12
Mammals	1	6	7
Aquatic Invertebrates			
Dragonflies and damselflies			1
Terrestrial Invertebrates			
Butterflies			3
Moths			12
Snails (terrestrial)	1	1	2
Plants	21		21
Total	32	10	58

1. Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants).

Conifer forest wetlands provide primary or secondary habitat for 19 vertebrate SGCN, 15% of the total number of terrestrial vertebrate SGCN. Birds represent 12 of the total SGCN. Coniferous wetlands are also important to terrestrial invertebrate fauna with a total of 17 SGCN, including 12 moths. This habitat is the primary habitat for 21 plant SGCN (5% of the 447 that are designated as SGCN). Examples of selected SGCN are described below.

Birds

Connecticut warblers (*Oporornis agilis*) nest in tamarack and spruce swamps with varying amounts of shrubby understory. Boreal chickadees (*Poecile hudsonicus*) prefer mature black spruce, where they require cavities for nesting. Olive-sided flycatchers (*Contopus cooperi*) breed in lowland conifers, generally requiring a fairly open canopy with tall prominent trees and snags. Black-backed and American three-toed woodpeckers feed by

flaking bark off of dead trees, particularly tamarack and black spruce. Great gray owls require trees large enough to support their nests.

Mammals

[Northern bog lemmings \(*Synaptomys borealis*\)](#) are limited to lowland conifer forests and open peatlands in extreme northern Minnesota; they have been shown to disappear from peatlands altered by human activities. Little is known about the species. Current populations are believed to be isolated relics of a larger population that occurred much farther south when glaciers advanced during the Pleistocene epoch. Fewer than 12 records have been documented in Minnesota since the species was first found in Lake County in 1932. Those records have been limited to open bogs, shrub carrs, and black spruce swamps.

Since the [Eastern heather vole \(*Phenacomys ungava*\)](#) was first documented in Lake County in 1940, fewer than 20 specimens have been collected in Cook, Lake, and St. Louis Counties. Like the northern bog lemming it also prefers lowland conifer forests and open peatlands but may be found in upland fire-dependent habitats as well. Proximity to water and the presence of boulders, coarse woody debris, and ericaceous plants appear to be important habitat components. Larger species typical of conifer wetlands include the SGCN snowshoe hare, Canada lynx.

Invertebrates

Two butterflies, the disa alpine (*Erebia disa*) and the bog copper (*Lycaena epixanthe*), require lowland conifers with cranberries (*Vaccinium macrocarpon* or *V. oxycoccos*) and lowland black spruce forests, respectively.

Plant

Many unique and rare plant species rely on the range of habitats provided by coniferous forest wetlands. Plants adapted to survive in the relatively harsh growing conditions of these coniferous forest wetlands are especially sensitive to changes in local hydrology, which maintain a range of soil pH and moisture availability that supports local plant communities.

White cedar swamps and forests often host diverse assemblages of native orchid species, sometimes with more than a dozen orchid species present on a site- including ram's-head lady's-slippers (*Cypripedium arietinum*), hooker's orchid (*Platanthera hookeri*) and fairy slipper orchids (*Calypso bulbosa*). These long-lived and charismatic plant species generally occupy a relatively sparse ground layer, maintained by partial to full tree canopy cover. More acidic spruce bog habitats harbor other rare orchid species including [bog adder's-mouth \(*Malaxis paludosa*\)](#) and tall white bog orchid (*Platanthera dilatata*), as well as plants with unique peatland adaptations, like the carnivorous English and linear-leaved sundews ([Drosera anglica](#), [D. linearis](#)). These [orchid species](#) are typically long-lived perennial plants that rely on associations with soil mycorrhizal fungi for access to minerals, water, and carbohydrates and usually require insect pollination and reproduce through tiny seeds that rely on fungal partnerships to germinate and grow. While orchids are short in stature, there is evidence that long-distance wind dispersal can happen. Vegetative offshoots from parent plants have also been documented.

Lichens

Coniferous wet forests are also home to unique lichen species. Lichen ‘species’ are a complex that consists of two or more, typically obligate, symbionts. The dominant partner is a fungus, which creates the main body of the organism. The endosymbiont is either a green alga or a cyanobacteria that lives within the fungal tissue. [Orange-tinted fringe lichen \(*Heterodermia obscurata*\)](#) is a Minnesota State Species of Concern documented in only one county in Minnesota. It occurs in forested peatland systems. While this species seems secure in other parts of its global range, we know very little about it or why it is so rare in Minnesota. Please see also Appendix B. Lichen list of SGCN.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. Species in Greatest Conservation Need.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Mining and Quarrying

Peat mining, where layers of peat are removed and used horticulturally as a soil amendment and potting medium, is a unique stress to coniferous forested wetland communities. Minnesota is one of the top three states in the nation for active peat mining operations; currently there are 10 active leases covering over 8,000 peatland acres. Mining not only results in significant impacts to the local hydrology and habitat, it also reduces the site’s ability to store carbon. Although mining leases require restoration and cleanup once operations cease, peat mining is nearly irreversible. It takes hundreds to thousands of years to form the layers of organic matter known as peat.



Roads, Trails, and Railroads

Roads, trails, and railroads can fragment large tracts of forest habitat, create edge effects, and alter hydrology. An increase in the number of roads and trails increases the potential for human disturbance and can introduce invasive species further into the forest. Coniferous forest wetlands are particularly susceptible to changes in local hydrology when roads and trails impede or channel water movement. Depending on the position of conifer swamps relative to the road and direction of groundwater movement, either localized draining or flooding can occur. When culverts are placed, beaver activity and blockages can further alter hydrology and expand the affected area to entire local basins.

Repeated use and soil compaction associated with roads and trails can modify the vegetation within the area of use. Trails, for example, are often very swampy with cattails and sedges compared to the adjacent forest understory which is largely composed of sedges, mosses, and ericaceous shrubs.

In northern Minnesota an extensive network of snowmobile trails through conifer wetland forests provides winter access to sites that are remote and largely inaccessible throughout the remainder of the year. Activity along these trails can disturb forest wildlife, especially species that begin nesting in late winter, such as many forest raptors.



Utility Corridors

Utility corridors transect Coniferous Forest Wetland habitat throughout the state. These corridors can contain a variety of utility infrastructure. This includes pipelines, telephone lines, and power lines. These lines can be aerial, above ground, or below ground. Regardless of type, utility corridors can fragment habitat based on their size, purpose, and the upkeep that is required to maintain them.

The initial construction of the associated utility infrastructure as well as the long-term maintenance of that infrastructure can have lasting impacts on native plant communities and the SGCN associated with them. Increased funding for rural broadband, conversion of aboveground powerlines to belowground, and development of new utilities (solar, wind, etc.) continue to threaten Coniferous Forest Wetland habitats.

These corridors and the increased travel from vehicles and equipment that they require act as vectors for invasive plant species. The required vegetative management practices often include herbicide applications, removal of woody vegetation, and/or mowing. This increased disturbance increases the likelihood of establishing invasive plant species within the corridors and then spreading to adjacent lands.

Although large utility corridor projects may be permitted to remove state listed plant species, the number and scale of these activities makes those efforts challenging, despite environmental review and permitting processes intended to minimize the negative consequences of these infrastructure projects.



Gathering Plants and Fungi

Coniferous forest wetlands are an important habitat for many of Minnesota's 48 [orchid species](#). Due to their showy flowers and unique forms, orchids are often a target for illegal collectors. While it is difficult to quantify the amount of illegal orchid harvesting that occurs, it is a significant enough threat that Minnesota has a [statute](#) (M.S. 17.23) pertaining to the conservation of certain wildflower species (including orchids) that requires landowner permission prior to collection, transplanting, or sale. Orchids are long-lived perennial plants that form close relationships with soil fungal partners and typically do not survive transplanting efforts- it is much better to take photographs and enjoy them where they are found naturally.

Spruce tree tops are considered a non-timber or decorative forest product (used in seasonal decorations) and interest in this [harvest activity](#) has been increasing regionally. Depending on local site conditions, gathering spruce tops for seasonal decorations from lowland conifer habitats has the potential to damage residual trees, ground layer plants, and the underlying peatland soils. These activities typically occur from late summer through late fall, and equipment operations in wetland sites outside of frozen conditions carry an increased risk of damage. Logging operations also have the potential for rutting if done prior to sufficient frozen ground conditions.



Timber Harvest

The typical natural disturbance regime in lowland conifer forests consisted of patchy blowdowns and surface fires small-scale blowdowns resulting in partial mortality of the forest canopy. These disturbances occurred every 40 to 90 years. Today, due to economies of scale and difficult access, some lowland conifer harvest blocks can be hundreds of acres in size and occasionally impact local hydrology because of temporary reductions in evapotranspiration and resulting water table rise (also known as swamping). This combination of scale and intensity of existing forest management practices in lowland conifers does not align well with documented natural disturbance regimes for these plant communities, which rarely would have experienced stand-replacing events.

Additionally, silvicultural treatments to minimize the impact of eastern dwarf mistletoe (*Arceuthobium pusillum*) on the growth and vigor of lowland conifers result in simplified forest structures (e.g. cutting all black spruce >5' tall). Mistletoe, a native parasitic plant, provides benefits to many forest wildlife species as a food source and a nesting or roosting structure (see also Problematic Native Species).

Although the total amount of timber harvested each year in Minnesota has trended downward since 1998, the black spruce harvest has slowly increased; although small, there also is a slight increase in the harvest and use of white cedar (Llorente, 2020).

Timber harvest can induce shifts in lowland conifer forest age-class distributions towards younger forests on the landscape, diminished habitat patch sizes, and resulting fragmentation. There is also the potential for inadvertent harvest of primary old growth forests due to slow growth rates and narrow tree rings that make it difficult to estimate tree ages in some lowland conifer sites. The result is that stands may be older than forest inventory data suggests. Regeneration of the harvested conifers may be problematic in areas with extremely low pH levels, poor nutrients, and saturated conditions. White cedar regeneration is especially difficult.



Recreation

In addition to the impacts created by recreational trails (discussed above), recreational activities themselves can proliferate invasive species. Seeds of invasive plants such as common tansy (*Tanacetum vulgare*) can become embedded in the wheels of vehicles or the shoes of individuals and eventually become established along compacted trails where conditions for their growth may be more favorable. Earthworm egg cases can also be spread in mud, which are problematic for moonwort (*Botrychium*) species. Increased use of roads and trails also increases human disturbance in the forest, reducing the amount of inaccessible area.



Fire Management

Due to the wetland nature of lowland conifer sites, wildfire has not been a significant natural disturbance agent of these habitats. However, with changing precipitation patterns and increasing human activity, fires can start and spread into these habitats and often lead to dramatic changes. Under extreme drought conditions, the organic peat substrates of coniferous forest wetlands itself can ignite and be consumed, resulting in changes to the depth to local water tables and altering the wetland plant communities altogether. Due to remote access and extreme conditions that accompany peat fires, the suppression tactics used to extinguish them can be detrimental themselves. When surface water is limited or unavailable to

wildland firefighters, heavy equipment is sometimes used to turn and mix mineral and organic soils in an effort to smother peat fires (i.e. “mashed potatoes” approach). While these tactics are used as a last resort, the outcomes can severely impact a site and warrant substantial restoration efforts.



Dams and Water Management

One of the most extensive disturbances to conifer lowlands is the attempt by early settlers to convert these wetlands into productive agricultural and forestry lands by constructing ditches. Constructed without any consideration of natural drainage patterns, the efforts were largely unsuccessful in the large expansive peatlands of northern Minnesota (Tester et al., 2020). Nevertheless, thousands of kilometers of ditches (Krause et al., 2021) remain intact despite not being actively maintained for decades. Unfortunately, they have left some lasting impacts.

Ditches alter peatland soils and hydrology, an impact that can extend as far as 100-150 meters away from the ditch itself. Soils for example, become compacted, drier, and often subside while mosses that require cool, wet environs are replaced by vascular plants that prefer drier conditions. These disturbed soils also are less capable of storing carbon, Krause and his colleagues (2012) have estimated that the annual loss of carbon from the ditches crisscrossing the northern Minnesota peatlands averages 38,000 metric tons per year.



Invasive Species (Problematic Non-native Species)

Sites where lowland conifer soils have been disturbed, such as along ditches and on and along trails, can become pathways for the introduction of wetland invasive plants such as purple loosestrife (*Lythrum salicaria*). Hydrologic disturbance may enable species like glossy buckthorn (*Frangula alnus*), reed canary grass (*Phalaris arundinacea*), common reed (*Phragmites australis*), and cattails (*Typha* spp.) to establish, displacing native species and altering microhabitat structure (Faber-Langendoen et al., 2020; Kost et al., 2007 in Ratcliffe et al., 2025b).



Problematic Native Species

The larvae of Eastern larch beetle (*Dendroctonus simplex*), a native Minnesota insect, feeds on the inner bark of tamarack trees, which can result in tree mortality. Historically this beetle produced only small, localized outbreaks that lasted several years, but as climate change lengthens its growing and breeding season, beetle populations and activity have increased substantially. Since approximately 2000, an [ongoing Eastern larch beetle outbreak](#) has resulted in more than one million acres of tamarack mortality across northern Minnesota. While some SGCN bird species (e.g., black-backed woodpeckers) may benefit from heightened insect activity and resulting coarse woody debris habitat, the scale of these events and associated salvage harvesting operations continue to pose landscape and site level habitat concerns.

Eastern dwarf mistletoe (*Arceuthobium pusillum*) is a tiny unique plant of northern Minnesota peatlands that parasitizes black spruce trees by rooting directly into their branches. Mistletoe growth prompts the host tree to develop dense clusters of branches and foliage (known as witch’s brooms) and eventually leads to the host tree’s death. Mistletoe brooms and associated mortality pockets are used as habitat by [numerous wildlife species](#), including boreal chickadees and great gray owls, and often increase structural and compositional diversity in the local ecosystem. As dense mistletoe populations can result in areas of dead or dying spruce trees and reduced productivity, forest managers are encouraged to manage the impacts during timber harvest

planning and operations. Eastern dwarf mistletoe plants disperse their sticky seeds by launching them with internal hydrostatic pressure into adjacent areas, so forest managers cut all spruce trees over five feet tall to limit seed dispersal distances away from other suitable spruce tree hosts. Because sticky seeds also disperse via movement on wildlife species that use witch's broom habitats, the utility of those practices warrants further investigation due to its potential ineffectiveness and the tradeoff of creating simplified habitat structure.

[Eastern](#) spruce budworm is a native caterpillar and is the most destructive pest of spruce-fir forests in eastern North America. Periodic outbreaks occur in the same sites approximately every 25 to 40 years. They may be induced by factors such as annual weather variation and climate change. Management favors commercially thinning mid-age stands (35-45 years old; [UMN Extension](#)). Although spruce budworm can negatively impact a stand's marketability, populations of several birds favor forest patches that are infested, including the bay-breasted, cape may, and Tennessee warblers.

Finally, overbrowsing by deer can severely limit natural tree regeneration (Matonis et al., 2011; De Jager et al., 2013), especially for preferred species, such as white cedar (Cornett et al., 2000). Elevated deer populations also reduce populations of palatable ground layer plants, including orchids (Knapp and Wiegand, 2014), members of the lily family (Balgooyen and Waller, 1995; Augustine and Frelich, 1998; Fletcher et al., 2001), and rare species (McGraw and Furedi, 2005). Decades of cumulative impacts have resulted in profound changes to the composition of native plant communities (Wiegmann and Waller, 2006; Frerker et al., 2014; Nuttle et al., 2014; Flagel et al., 2016; Sabo et al., 2017). The effects of deer browsing often interact with those of invasive earthworms (Fisichelli et al., 2013; Davalos et al., 2014).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025b).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

With respect to the conifer forest wetlands in Minnesota, in the Northern Forest ecoregion that includes northeastern Minnesota, projected changes in annual temperature by end of century (2070-2099) are 6 – 11 degrees higher than historic (1970-2000), depending on the carbon emissions level and using the mean of 20

climate models (see Table 9 in Ratcliffe et al., 2025b). Warming temperatures could convert Minnesota's peatlands from a carbon sink to an active source of carbon emissions ([Peatlands Resilience Initiative](#)). A warmer climate also directly impacts the community's unique vegetative composition and phenology (Antala et al., 2022). Many key species may lose suitable habitat, as they are already near the southern extent of their range and have heightened vulnerability and limited adaptive capacity, making them especially at risk from climate change (e.g., *Larix laricina*, *Picea mariana*; Ratcliffe et al., 2025b). Among the changes projected are an increase in shrubs and grasses at the expense of peatland mosses such as *Sphagnum* sp., while the flowering season for many vascular plants will begin earlier. See Climate Profile section.

Changes in Precipitation and Hydrology related to Climate



Minnesota's climate is becoming wetter and more hydrologically variable. Between 1895 and 2020, the state's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). In addition to this overall rise, the frequency and intensity of heavy precipitation events have also grown. Since 2000, very heavy rainfall events—defined as 6 inches or more in a single day—have occurred two to three times more often than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information, 2022](#)). These extreme rainfall events have increased flooding, which disrupts ecosystems, damages infrastructure, and impairs water quality (Williams-Sether & Sanocki, 2025).

Looking ahead, climate models project continued increases in annual precipitation, particularly during winter and spring months. These wetter seasons are expected to elevate flood risks, while late summer is likely to become hotter and drier, intensifying drought conditions ([Climate Change in Minnesota](#)). By mid-century (2040–2059), average annual precipitation in Minnesota is projected to increase by up to 1.2 inches, depending on greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly contradictory pattern—wetter winters and springs punctuated by drier late summers—has far-reaching implications for water availability, wetland integrity, soil stability, and species that rely on seasonal hydrologic cycles (Runkle et al., 2022). For guidance on climate-adapted management, see the Climate subchapter in Chapter 6. Implementation.

These changes are especially consequential for Minnesota's rich peatland ecosystems. Acidic peatlands, which rely exclusively on precipitation for water inputs, are highly sensitive to shifts in rainfall timing and volume. Greater spring rainfall may raise water tables and favor more open peatland systems, while prolonged summer droughts could dry peat layers, encouraging invasion by forest species and accelerating peat decomposition. Ecosystem models predict significant declines in dominant conifer species such as black spruce and tamarack under these conditions (Ratcliffe et al., 2025b). Many of these peatlands also harbor rare and endemic plant species that are presumed to be highly vulnerable to altered hydrology and substrate changes.

Compounding these risks, hotter and drier summers are expected to elevate vapor pressure deficit (VPD)—a key driver of plant and soil water loss—by 32–66% by the end of the century (2070–2099), relative to historical conditions (1970–2000), depending on emissions levels (Ratcliffe et al., 2025b, Table 11). Increased VPD accelerates evaporation, lowers water tables, and exposes peat to oxygen, promoting aerobic decomposition and the release of stored nutrients and carbon. Sustained heat and drought also threaten the productivity and dominance of *Sphagnum* mosses, which are vital to peat accumulation and long-term carbon storage (Bu et al., 2011; Norby et al., 2019 in Ratcliffe et al., 2025b).

An increase in late summer droughts also raises the risk of peatland fires. These fires are especially difficult to manage due to the remoteness of large peatland areas and the tendency for deep organic soils to smolder underground for extended periods. For example, during the severe drought of 2021, a lightning strike near Greenwood Lake in Lake County ignited a wildfire that burned over 28,000 acres, including extensive peatland areas. In some locations, more than one meter of peat was consumed, transforming long-standing carbon sinks into active carbon sources ([USFS, 2024](#)).

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, to protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN, five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Examples of conservation actions are presented below grouped by the primary strategy that they implement and tagged with icons for the stressor(s) that they address. Some of the strategies, such as Strategy 3, collaborating with partners, could truly be applied to most broadly and effectively implement all actions. In this case, under Strategy 3 we highlight actions for which collaboration is particularly vital. If an action does not relate to a stressor, we mark this with a ‘n/a’ in the stressor column.

Potential Conservation Actions for Coniferous Forest Wetlands



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Identify priority areas for protected area designations to facilitate long-term conservation outcomes and protection against conversion of forest to different land uses. Examples could include high biodiversity areas, old growth forests, or those providing connectivity between other protected areas that may provide disproportionately high conservation value.

Stressor	Action
	On state lands, engage in forest planning efforts, such as Section Forest Resource Management Plans (SFRMPs), Sustainable Timber Harvest (STH) plans, designation of Management Opportunity Areas (MOAs), and designation of High Conservation Value Forests to assist in addressing values such as biodiversity, rare features, diversity of native plant community growth stages, and wildlife needs that can't be addressed through individual site-level management (see also DNR Forest Resource Planning).
	Encourage sustainable forest management through planning when and where timber harvest can be used to address a range of values and priorities, including forest health and productivity, critical ecological functions, and habitat restoration from historical disturbance events. Incorporate considerations of forest compositional and structural diversity at meaningful scales, historic size and return intervals of disturbance regimes, biodiversity, rare features, wildlife needs, and diversity of native plant community growth stages. Ensure that a balance of forest habitat conditions (e.g. younger and older forest, small and large patch sizes) are available for plant and animal species across the landscape over time.
	Support forest management approaches that are informed by ecological processes, increase structural and compositional complexity and diversity, and assist in resilience towards future stressors (e.g., climate change, disease outbreaks, invasion by non-native species). Promote forest management strategies that mimic landscape disturbance patterns (for example, retaining large blocks of mature habitat and creating more small forest openings as gaps or strips), maintain or enhance connectivity among forest patches to enable wildlife movement, and retain biological legacies (at site level) such as large trees with cavities. Consult Native Plant Community Classification silvicultural strategies for forest stand prescriptions .
	Apply knowledge of SGCN habitat use into landscape forest planning efforts, to help balance habitat for SGCN and other values according to management requirements and objectives. Consider natural area designations and other ways to enhance and preserve key mature stands



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

Stressor	Action
	Consider projected effects on high value conservation forests, old-growth forests, and Important Bird Areas when making decisions and policy that are projected to impact

Stressor	Action
	coniferous wetland forests (e.g., mining and quarrying or oil and gas permitting, planning for new development of energy sources and any associated new road or trail construction).
	Encourage the use of existing corridors for new trails, roads, and utility corridor development to minimize the construction of new roads and trails that would increase forest fragmentation.
	Limit the use and operation of heavy equipment when road or site conditions are vulnerable to damage. For instance, depending on the site characteristics, conducting timber harvests during frozen ground conditions can reduce soil compaction and rutting impacts. Avoid rutting and damage to peat soils when harvesting spruce tops.
	Carefully consider hydrological, habitat, and landscape level disruption to conifer wetland forests when planning timber harvest operations in these delicate systems. Consider forest productivity and the ability of lowland conifers to regenerate prior to timber harvest. Conduct logging operations in wetland forests only after the ground is sufficiently frozen to prevent rutting and hydrological impacts from road constructions and use.
	To prevent soil compaction and changes to site hydrology, minimize use of heavy equipment to suppress fires in conifer lowlands when possible.
	Implement best management practices for Eastern larch beetle management in addressing outbreaks. Encourage tree diversity by leaving other tree species such as black spruce and cedar on the site, or by planting or aerial seeding with a mixture of site-appropriate trees, including tamarack.
	Balance management of spruce budworm outbreaks with consideration of their benefits to warblers and other birds that serve an important role as budworm predators.
	Implement protective measures to reduce deer browsing on white cedar seedlings on sites that are suitable for cedar growth and establishment.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

Stressor	Action
	Increase outreach and capacity regarding enforcement to protect vulnerable orchid species (see Plant SGCN chapter).

Stressor	Action
	Collaborate across forest ownerships and managers to implement sustainable forestry at the site and landscape level to create habitat conditions important to SGCN, such as unfragmented older mesic hardwood forests to create larger forest patches and reduce forest fragmentation.
	Partner with other conservation organizations and natural resource agencies to protect intact peatlands from further degradation and to restore partially drained peatlands.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions

Stressor	Action
	Support the collection of accurate and reliable forest stand inventory data for planning.
	Monitor the responses of SGCN to various types of forest management at the site-level and landscape-scale. Collaborate with MFRC site-level guideline monitoring and NPC silvicultural case studies to incorporate SGCN monitoring. Develop models of habitat configurations for various SGCN and consider how current harvest forecasts and conversion rates (i.e., one forest type to another; natural origin habitat to plantation) relate to those models in terms of sustainability for maintaining habitat for SGCNs.
	Monitor harvest of nontimber forest products such as spruce tops to avoid rutting and damage to sensitive peat substrates. Consider limiting spruce top harvesting in areas of biodiversity significance or sensitive hydrology. While pressure and impacts have historically been dispersed and localized, monitoring can help alert where future management actions are needed to protect broader ecosystem functionality.
	Monitor eastern larch beetle and effects; predict susceptible areas.
	Investigate the efficacy of current management efforts to reduce the spread of Eastern dwarf mistletoe in addition to the benefits/impacts to wildlife species.
	Monitor, research, and manage impacts of emerging stressors like increased temperatures, interactions with pest species, and altered hydrology anticipated with changes in climate.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Stressor	Action
	Develop and share studies of alternative harvest methods for black spruce as an alternative to clear cuts. Support research and case studies utilizing shelterwood, gap and strip harvest techniques that retain greater habitat diversity (Anderson et al., 2020).
	Research techniques to promote forest regeneration for challenging species (i.e. white cedar) or in certain circumstances (i.e. where altered hydrology is a concern).
	Where people recreate, provide outreach and education on cleaning gear and equipment to reduce the introduction and spread of invasive species. DNR's Prevent the spread webpage includes specific actions for different land based activities like biking, hiking and off-highway vehicle riding. At trailheads, add invasive species prevention messages, such as those on the boot brush kiosks developed by the PlayCleanGo: Stop Invasive Species in Your Tracks program.
	Implement strategies and approaches for climate-informed forest management, including increasing forest biodiversity for increased resilience towards future changes, including disease or pathogen outbreaks, extreme weather events, and invasions by non-native species. Apply the Climate Change Field Guide for Northern Minnesota Forests (Handler et al. 2017) for forest management guidance regarding harvest strategies and subsequent regeneration options. Learn about how climate may impact our forests in the future with this resource: Minnesota Forest Ecosystem Vulnerability Assessment and Synthesis .

Case Study: Peatland Forest Management for Bird SGCN

In 2018 With funding from the Legislative-Citizen Commission on Minnesota Resources (LCCMR) the Natural Resources Research Institute, Minnesota Forest Resources Council, University of Minnesota-Twin Cities, and the Minnesota Department of Natural Resources teamed up to initiate [a project](#) examining the feasibility of several alternative silvicultural treatments in lowland conifer forests to maintain suitable habitat conditions for two SGCN bird species, Connecticut warbler and boreal chickadee. Study sites dominated by black spruce and tamarack were chosen to receive a combination of alternative management techniques. These included ½ acre gap harvests, 25 foot and 100-foot-wide strip harvests, and clearcutting. This led to the development of [alternative harvest methods](#) that will help inform sustainable forest management and enhance understanding of these complex habitats and valuable ecosystems that maintain biodiversity. Ongoing monitoring is planned to assess both bird and plant community responses.

References for Coniferous Forest Wetland

- Anderson, B.D., Windmuller-Campione, M.A., Russell, M.B., Palik, B.J., & Kastendick, D.N. (2020). Short- and long-term results of alternative silviculture in peatland black spruce in Minnesota, USA. *Forest Science*, 66(2), 256-265. <https://doi.org/10.1093/forsci/fxz078>
- Antala, M., Jaszczak, R., van der Tol, C. & Rastogi, A. (2022). Impact of climate change-induced alterations in peatland vegetation phenology and composition on carbon balance. *Science of the Total Environment*. Volume 827: June 2022, 154294. <https://www.sciencedirect.com/science/article/pii/S0048969722013869>.
- Augustine, D. J., & Frelich, L. E. (1998). Effects of White-Tailed Deer on Populations of an Understory Forb in Fragmented Deciduous Forests. *Conservation Biology*, 12(5), 995–1004. <https://doi.org/10.1046/j.1523-1739.1998.97248.x>
- Balgooyen, C. P. and D. M. Waller. (1995). The use of *Clintonia borealis* and other indicators to gauge impacts of white-tailed deer on plant communities in northern Wisconsin, USA. *Natural Areas Journal*, 15(4), 308-318. <https://www.cabidigitallibrary.org/doi/full/10.5555/19960610148>.
- Cornett, M. W., Frelich, L. E., Puettmann, K. J., & Reich, P. B. (2000). Conservation implications of browsing by *Odocoileus virginianus* in remnant upland *Thuja occidentalis* forests. *Biological Conservation*, 93(3), 359–369. [https://doi.org/10.1016/S0006-3207\(99\)00129-9](https://doi.org/10.1016/S0006-3207(99)00129-9)
- Dávalos, A., Nuzzo, V., & Blossey, B. (2015). Single and interactive effects of deer and earthworms on non-native plants. *Forest Ecology and Management*, 351, 28–35. <https://doi.org/10.1016/j.foreco.2015.04.026>
- De Jager, N. R., Cogger, B. J., & Thomsen, M. A. (2013). Interactive effects of flooding and deer (*Odocoileus virginianus*) browsing on floodplain forest recruitment. *Forest Ecology and Management*, 303, 11–19. <https://doi.org/10.1016/j.foreco.2013.02.028>
- Finch, D.M. et al. (2021). Effects of Climate Change on Invasive Species. In: Poland, T.M., Patel-Weynand, T., Finch, D.M., Miniati, C.F., Hayes, D.C., Lopez, V.M. (eds) *Invasive Species in Forests and Rangelands of the United States*. Springer, Cham. https://doi.org/10.1007/978-3-030-45367-1_4
- Fisichelli, N. A., Frelich, L. E., Reich, P. B., & Eisenhauer, N. (2013). Linking direct and indirect pathways mediating earthworms, deer, and understory composition in Great Lakes forests. *Biological Invasions*, 15(5), 1057–1066. <https://doi.org/10.1007/s10530-012-0350-6>
- Flagel, D. G., Belovsky, G. E., & Beyer, D. E. (2016). Natural and experimental tests of trophic cascades: Gray wolves and white-tailed deer in a Great Lakes forest. *Oecologia*, 180(4), 1183–1194. <https://doi.org/10.1007/s00442-015-3515-z>
- Fletcher, D. J., Shipley, L. A., McShea, W. J., & Shumway, D. L. (2001). Wildlife herbivory and rare plants: The effects of white-tailed deer, rodents, and insects on growth and survival of Turk's cap lily. *Biological Conservation*, 101(2), 229–238. [https://doi.org/10.1016/S0006-3207\(01\)00070-2](https://doi.org/10.1016/S0006-3207(01)00070-2)

- Frerker, K., Sabo, A., & Waller, D. (2014). Long-Term Regional Shifts in Plant Community Composition Are Largely Explained by Local Deer Impact Experiments. *PLOS ONE*, 9(12), e115843. <https://doi.org/10.1371/journal.pone.0115843>
- Handler, S., K. Marcinkowski, M. Janowiak, and C. Swanston. (2017). Climate change field guide for northern Minnesota forests: Site-level considerations and adaptation. USDA Northern Forests Climate Hub Technical Report #2. University of Minnesota College of Food, Agricultural, and Natural Resource Sciences, St. Paul, MN. 88p. Available at www.forestadaptation.org/MN_field_guide.
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Knapp, W. M., & Wiegand, R. (2014). Orchid (Orchidaceae) decline in the Catocin Mountains, Frederick County, Maryland as documented by a long-term dataset. *Biodiversity and Conservation*, 23(8), 1965–1976. <https://doi.org/10.1007/s10531-014-0698-2>
- Krause, Lima, Kevin J. McCullough, Evan S. Kane, Randall K. Kolka, Rodney A. Chimner and Erik A. Lilleskov. (2021). The impacts of historical ditching on peat volume and carbon in northern Minnesota USA peatlands. 2021. *Journal of Environmental Management* 296 (2021) 113090.
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Llorente, I. D. P. (2020). Addressing Climate Change Uncertainty in Forest Planning: A Multi-Stage Optimization Model with Multiple Future and Price Sensitive Demands (Doctoral Dissertation, University of Minnesota).
- Matonis, M. S., Walters, M. B., & Millington, J. D. A. (2011). Gap-, stand-, and landscape-scale factors contribute to poor sugar maple regeneration after timber harvest. *Forest Ecology and Management*, 262(2), 286–298. <https://doi.org/10.1016/j.foreco.2011.03.034>
- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). 2025. Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state's greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>
- McGraw, J. B., & Furedi, M. A. (2005). Deer Browsing and Population Viability of a Forest Understory Plant. *Science*, 307(5711), 920–922. <https://doi.org/10.1126/science.1107036>
- Nuttle, T., Ristau, T. E., & Royo, A. A. (2014). Long-term biological legacies of herbivore density in a landscape-scale experiment: Forest understoreys reflect past deer density treatments for at least 20 years. *Journal of Ecology*, 102(1), 221–228. <https://doi.org/10.1111/1365-2745.12175>

- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. (2025a). Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwccasc.umn.edu/node/821>
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. (2025b). Effects of Climate Change on Midwestern Ecosystems: North American Bog and Fen. Midwest Climate Adaptation Science Center. (67 pages) Available: <https://mwccasc.umn.edu/node/821>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Sabo, A. E., Frerker, K. L., Waller, D. M., & Kruger, E. L. (2017). Deer-mediated changes in environment compound the direct impacts of herbivory on understorey plant communities. *Journal of Ecology*, 105(5), 1386–1398. <https://doi.org/10.1111/1365-2745.12748>
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>
- Tester, John R., Susan M. Galatowitsch, Rebecca A. Montgomery, and John J. Moriarty. (2020). Minnesota's Natural Heritage. Second Edition.
- Walker, M., (2011). Restoration Strategies for Ditched Peatland and Scientific and Natural Areas. <https://www.legacy.mn.gov/projects/restoration-strategies-ditched-peatland-and-scientific-and-natural-areas>
- Weiskopf, Gretel. (2024). Measuring the effects of fire in northern Minnesota peatlands. USFS. <https://www.fs.usda.gov/inside-fs/delivering-mission/sustain/measuring-effects-fire-northern-minnesota-peatlands>
- Wiegmann, S. M., & Waller, D. M. (2006). Fifty years of change in northern upland forest understories: Identity and traits of “winner” and “loser” plant species. *Biological Conservation*, 129(1), 109–123. <https://doi.org/10.1016/j.biocon.2005.10.027>
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Riparian and Floodplain Forest

Habitat Description

Riparian and floodplain forest habitats occur primarily on floodplains and associated terraces along major rivers and their tributaries (see also Rivers and Streams Chapter). Floodplain and terrace forests are seasonally wet forests that flood following spring snowmelt as well as unusually heavy rains. These forests are found on sandy or silty alluvium (soil deposited by flowing water) associated with streams and rivers throughout the Eastern Broadleaf Forest Province and are extensive along the Mississippi, Minnesota, and St. Croix rivers. In the Laurentian Mixed Forest Province, these forests also occur along major rivers but are not as extensive as in the Eastern Broadleaf Forest Province. These forests also rarely occur in the Tallgrass Aspen Parklands and Prairie Parkland provinces.

The canopy of these forests is dominated by deciduous trees that are tolerant of saturated soils, prolonged inundation, frequent erosion, and sediment deposition. Species less tolerant of these conditions occur on terraces, which flood only in very wet years. In southern Minnesota, silver maple (*Acer saccharinum*) (which often occurs as nearly pure stands), black willow (*Salix nigra*), and eastern cottonwood (*Populus deltoides*) are common canopy dominants. Less common species include river birch (*Betula nigra*), elms (*Ulmus spp.*), green ash (*Fraxinus pennsylvanica*), and [swamp white oak \(*Quercus bicolor*\)](#). In the north, black ash and silver maple are important canopy trees with lesser amounts of green ash, American elm (*Ulmus americana*), bur oak (*Quercus macrocarpa*), and basswood (*Tilia americana*). Canopy coverage in these forests is highly variable; areas of continuous canopy are punctuated with large gaps, which may be vegetated with ephemeral herbaceous plants or may remain largely unvegetated if flood disturbance is repeated and severe. The understory is typically open, with few shrubs or saplings. Pools or mucky depressions in old river channels are often present on actively flooded sites.

Habitat Map

To depict Riparian and Floodplain Forest habitat (see Figure 3.13 we compiled spatial data from several sources: DNR's Native Plant Communities and the Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Floodplain Forest (FF)

[FFn57 Northern Terrace Forest \(PDF\)](#)

[FFn67 Northern Floodplain Forest \(PDF\)](#)

[FFs59 Southern Terrace Forest \(PDF\)](#)

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Laurentian-Acadian Floodplain Forest and Midwest Floodplain Forest.




Figure 3.13. A map depicting Riparian and Floodplain Forest habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological System of Floodplain Forest, and two types from the Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

Riparian and floodplain forest habitats occur along major rivers, rivers, and streams statewide (see also Rivers and Streams Chapter). Riparian and floodplain forests in southern Minnesota are among the most disturbed native habitats in the state. Even prior to European settlement, large swaths of forests were logged to provide fuel for riverboats. Since settlement by people of European descent, large areas of floodplain forests in southern Minnesota have been lost due to urbanization and conversion to agriculture. Now, these forests, which once formed continuous bands of habitat along the major rivers in southeastern Minnesota, persist as a broken chain of forest patches. In addition, the damming of major rivers has greatly reduced the annual pulse of flooding that maintained the ecological integrity of the floodplain forests. Other factors that have reduced the value of the habitat for wildlife include the loss of most canopy American elms from Dutch elm disease and the spread of reed canary grass (*Phalaris arundinacea*), which impedes establishment of seedlings of native plants. The recent loss of ash trees to emerald ash borer (*Agrilus planipennis*) and elm (*Ophiostoma spp.*) to Dutch elm disease, shifted the composition of the Mississippi River floodplain forest towards a composition of largely Silver Maple and Oak, particularly along pools 1 – 10. The loss of ash trees is also pronounced along the Minnesota River.

Protecting riparian forest along river corridors is important to facilitate movement of wildlife and the movement of nutrients and energy between the river and the surrounding landscape (see also the Rivers and Streams Chapter). In addition to the habitat they provide for many resident animals and nesting birds, millions of migrating birds depend on the Mississippi River and its associated floodplains each spring and fall as one of the most important migratory flyways in North America.

Climate Profile for Floodplain Forests

Riparian and Floodplain Forest was profiled in the report “Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest” published by the Midwest Climate Adaptation Science Center (Ratcliffe et al., 2025). The most impactful changes in climate for this system include more frequent and larger flood events (see Figure 3.14), also with more frequent late summer drought periods.

A major concern for Floodplain Forests is more frequent and more intense flooding events. More precipitation will mean more sediment deposition as well as habitat changes. This projected increase in flooding will favor pioneer and early successional species, while limiting the growth of more stable species. Large increase in flood frequency and intensity is projected to occur in the spring, as both winter and spring precipitation is projected to increase. Coupled with warmer temperatures leading to an earlier snowmelt, large floods are predicted to continue to greatly impact this system.

Floodplain Forests will also be impacted by warmer and dryer summers, which will cause soil drying and an overall reduction of habitat in these water-dependent ecosystems. Further habitat change will occur due to more frequent summer droughts, which will impact plant growth and favor drought dependent species. These climate-induced changes may promote the spread of invasive species.

Key Climate Change Impacts:

- **Habitat Structure:** Flooding events will cause erosion and remove valuable habitat components from Floodplain Forests such as streambanks and large woody debris, especially in flooding prone regions, such as agriculture dominated landscapes. Flooding also has a negative impact on tree mortality. Flooding may also increase habitat viability by bringing in new sediment, introducing water into usually dry areas, and carrying nutrients (as well as pollutants).
- **Community Composition:** Floodplain Forests will transition away from mast-producing oaks (*Quercus* spp.) and hickories (*Carya* spp.) to more flood resistant species, such as ashes (*Fraxinus* spp.), willows (*Salix* spp.) and maples (*Acer* spp.). Drought resistant species will be favored during warmer and dryer summers, while more moisture dependent species, like tamarack (*Larix laricina*) and black Spruce (*Picea mariana*).
- **Invasive Species:** As the climate warms, species from warmer regions may begin to move into the Midwest in search of cooler conditions, pushing out native species, like willows (*Salix* spp.) and cottonwoods (*Populus* spp.). Climate-driven disturbance such as flooding events, will increase the threat to native species, as opportunities for out competition by non-natives increases.
- **Pests and Pathogens:** Increasing temperatures will also likely cause ranges of some pest species expand, populations increase, and hosts become more susceptible to infection with rising temperatures. Plant communities will be hit harder by pests as warmer winters protect pest populations and enable additional annual broods. Pests like the spongy Moth (*Lymantria dispar*), emerald ash borer, and hickory bark beetle (*Scolytus quadrispinosus*) are predicted to benefit from warmer winters and drying summers, leading to increased tree mortality.
- **Herbivory:** Warmer winters will benefit white-tail deer (*Odocoileus virginianus*) populations because of boosted survival rates and better access to forage, furthering the currently observed trend of white-tailed deer population growth. (All of this Climate Profile content is from Ratcliffe et al., 2025.)




Figure 3.14. (from Ratcliffe et al. 2025). Projected changes in mean streamflow in the Midwest. Projections use gridded simulations of daily runoff based on CMIP5 climate models under the high-emissions (RCP 8.5) scenario. Values show the percent change in mean daily streamflow (cubic feet per second) for each season (winter, December–February; spring, March–May; summer, June–August; fall, September–November) at the end of the century (2070–2099) compared to historical data (1977–2006). Data and metric definitions are provided by the U.S. Stream Flow Metric Dataset (Authors (in review); U.S. Department of the Interior, Bureau of Reclamation, 2014; USDA Forest Service Office of Sustainability and Climate, 2022; Wenger and others, 2010).

Species in Greatest Conservation Need

Riparian and Floodplain Forests provide primary or secondary habitat for 25 animal and 22 plant SGCN (see Table 3.12). Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Table 3.12. Numbers of Species in Greatest Conservation Need associated with Riparian and Floodplain Forests as either primary or secondary habitat¹.

Species Groups	Primary Habitat	Secondary Habitat	Total
Amphibians	2	0	2
Birds	3	3	6
Mammals	1	9	10
Reptiles	2	0	2
Terrestrial Invertebrates			
Beetles (terrestrial)			2
Moths			1
Spiders	0	2	2
Plants	22		22
Total	30	14	47

1. Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants). Examples of selected SGCN are described below.

Twenty-five (19%) of the 131 terrestrial vertebrate SGCN are associated with Riparian and Floodplain Forests, including ten of the mammals (31% of the 32 mammal SGCN). 22 plant SGCN associate with Riparian and Floodplain Habitats as their primary habitat (5% of plant SGCN).

Amphibians

[Blanchard's cricket frog \(*Acris blanchardi*\)](#) are found in riparian areas right along the water's edge; open areas with muddy shorelines and dense emergent vegetation are preferred. They reside near water throughout the summer but, during dry conditions, they may seek out wetter conditions elsewhere. Crayfish burrows or small openings along the bank of a wetland or river provide sites for hibernation. Less than 2 inches in size, their breeding call resembles that of some birds, making visual identification essential. A state endangered species in Minnesota, currently populations are limited to sites in Hennepin and Winona counties.

Pickerel frogs (*Lithobates palustris*) are also found in Riparian Forests and Floodplains, preferring the edges of ponds, small rivers, and streams that flow through wooded areas or open habitats (Moriarty & Hall, 2014). They rarely venture far from the water’s edge or from nearby wet meadows. Winter months are spent hibernating in their water habitats. Pickerel frogs are restricted to the far southeastern corner of the state; county records are available from Goodhue, Fillmore, Houston, Mower, Olmsted, and Winona counties (Moriarty & Hall, 2014).

Birds

Floodplain forests in riparian zones provide key habitat to many SGCN. The only SGCN bird restricted to floodplain forest is the prothonotary warbler (*Protonotaria citrea*). They are most often found along medium- to large rivers such as the Mississippi, Minnesota, and St. Croix. A cavity-nesting species, it requires large trees or snags, often situated right over the water (Pfannmuller et al., 2017).

Climate Profile: Prothonotary Warbler

This profile is excerpted from “Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest” published by the Midwest Climate Adaptation Science Center (Ratcliffe et al., 2025).

“The prothonotary warbler (*Protonotaria citrea*) is a small cavity-nesting songbird that breeds in riparian and floodplain forests in the central Midwest and winters in mangrove forests of Central and South America. Climate-driven hydrological changes and sea-level rise threaten both its breeding and non-breeding habitats. The species has been documented expanding its breeding range northward, including into the Mississippi River Valley, and its population remains relatively stable (NatureServe, 2008).

Prothonotary warblers nest in tree cavities over standing water, making them highly sensitive to hydrological extremes (Cooper and others, 2009). High magnitude flooding with high water depths can directly impact reproductive success (Flaspohler, 1996; Graber and others, 1983). Prolonged floodplain inundation can also alter predator foraging behavior by limiting non-arboreal options, increasing warbler nest depredation by predators like raccoons (Cooper and others, 2009). However, partial flooding appears to provide optimal protection against predation (Cooper and others, 2009; Hoover, 2006; Hoover, 2009). Conversely, drier conditions expose dense ground cover, removing the protective influence of standing water and increasing rates of nest depredation (Cooper and others, 2009; Kirsch, 2009). Despite these direct influences of hydrology, altered hydrological regimes appear unlikely, except in the most severe instances of flooding, to reduce the abundance of most preferred nesting tree species, which include red maple (*Acer rubrum*), sugarberry (*Celtis laevigata*), and silver maple (*Acer saccharinum*) (Table 20) (Gabbe and others, 2002; Kirsch and Wellik, 2017; Wuebbles and others, 2021).

Brood parasitism by brown-headed cowbirds (*Molothrus ater*) is another major threat to the prothonotary warbler, affecting up to 50% of nests— reducing clutch size, hatching success, and nesting survival (Hoover, 2003b). Although brown-headed cowbird abundance has been in decline (Cox and others, 2012; Forrester and others, 2023), parasitism rates remain high, and climate-driven increases in precipitation could boost cowbird numbers, further pressuring warbler populations (Forrester and others, 2023; Rosamond and others, 2020). Moreover, as native bird populations decline due to higher precipitation, parasitism pressure may increase on small host populations (Rosamond and others, 2020). Prothonotary warblers show limited ability to adapt to brood parasitism, making them particularly vulnerable (Hoover, 2003a).

As forested wetlands face increasing climate stress, climate-informed restoration may be a key adaptation strategy for this species. For example, in the Cache River watershed, habitat restoration increased prothonotary warbler density and nesting success (Hoover, 2009). Additionally, artificial nesting structures offer a promising conservation strategy, as warblers readily use nest boxes, which have shown no recorded depredation (Mueller and others, 2019).”

As forested wetlands face increasing climate stress, climate-informed restoration may be a key adaptation strategy for this species. For example, in the Cache River watershed, habitat restoration increased protonotary warbler density and nesting success (Hoover, 2009). Additionally, artificial nesting structures offer a promising conservation strategy, as warblers readily use nest boxes. Other deciduous forest SGCN birds will occupy both upland and floodplain forests, with preferences often varying by region. Cerulean warblers require large tracts of mature deciduous forest. Although it is found in mesic hardwood stands in much of its Minnesota range this canopy-dwelling species occupies floodplain forests along the Minnesota, Mississippi, and St. Croix rivers. Similarly, Red-shouldered hawks are most frequent in floodplain forests along larger rivers in southeastern and east-central Minnesota. Oxbow lakes and other floodplain forest openings are an important feature in broad floodplains. Red-shoulders prefer nesting sites in large-diameter trees in high, thick canopies. Louisiana waterthrushes (*Parkesia motacilla*) are most closely associated with mesic hardwoods bordering small, clear water streams, but will also occupy mature floodplain forests, particularly along larger rivers in east-central Minnesota such as the Kettle and St. Croix. In these broad floodplain forests, smaller side channels seem to be a key feature. The Acadian flycatcher (*Empidonax virescens*) is much more frequent in mesic hardwoods but is occasional in true floodplain forests. The bird most tightly associated with mature riparian floodplain forests in Minnesota is the Prothonotary warbler. Although it is uncommonly found in floodplains of small streams and rivers, nearly all the records in Minnesota are along the Mississippi and Minnesota rivers. Although often thought of as species restricted to savannas or upland deciduous woodlands, red-headed woodpeckers are also found in floodplain forests in areas where excessive flooding or other disturbances have created openings with dead trees and snags lacking bark.

Many other birds also use river corridors, beyond those formally associated with it for a primary or secondary habitat, for breeding, migration, and wintering habitat. In particular, large, old trees provide important habitat for nesting herons, egrets, barred owls, wood ducks, and bald eagles. Riparian corridors provide vital migratory stopover habitat, and mortality during migration exceeds that during breeding or wintering periods for many adult migratory songbirds (Silllett and Holmes, 2002). Quality and availability of stopover habitat may be important factors affecting their populations. Some SGCN that do not specialize in riparian forest for breeding, but may rely on it for stopover habitat include Connecticut warbler (*Oporornis agilis*), black-throated blue warbler, olive-sided flycatcher, bay-breasted warbler, Swainson's thrush, and Cape May warbler. Because riparian forests are often linear and relatively contiguous, they are well suited to migration and may be especially important as stopover habitat compared to other habitats.

Climate Profile: Red-shouldered Hawk

This profile is excerpted from “Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest” published by the Midwest Climate Adaptation Science Center (Ratcliffe et al., 2025).

“The [red-shouldered hawk \(*Buteo lineatus*\)](#) is a medium-sized raptor found throughout the eastern and central U.S. In the Midwest, these hawks typically occupy mature forests (Henneman and Andersen, 2009; Woodford and others, 2008) with favorable proximity to wetlands to forage for amphibians and reptiles (Balcerzak and Wood, 2003).

During the breeding season, while cooler temperatures combined with heavy precipitation may reduce nestling survival, warmer temperatures can increase prey activity, foraging efficiency, and reproductive rates, suggesting a positive effect of climate change on the species (Bednarz and Dinsmore, 1982; Craighead and others, 1969; Dykstra and others, 2021; Henneman, 2006; Jacobs and Jacobs, 2002). Climate change-driven loss of wetlands and reductions in forests, especially coniferous forests, adversely affect their reproductive success and survival (Dykstra and others, 2021). Despite these challenges, red-shouldered hawks demonstrate adaptive capacity through their flexible diet and habitat use, and even successfully occupying suburban habitats (Dykstra and others, 2019).

Climate change may reduce suitable habitat for red-shouldered hawks in the Midwest Floodplain Forest by decreasing contiguous forest cover and increasing the frequency and size of canopy gaps. These gaps are associated with increased rates of nest predation and increased competition from red-tailed hawks (*Buteo jamaicensis*) (Bednarz and Dinsmore, 1982; Craighead and others, 1969; Henneman, 2006; Jacobs and Jacobs, 2002). More frequent flooding may have mixed effects on prey populations, potentially reducing some small mammal populations while creating canopy gaps that benefit others (Chamberlain and Leopold, 2003; Scharine and others, 2009). Moreover, key nesting trees such as american beech (*Fagus grandifolia*), oaks (*Quercus* spp.), and sugar maple (*Acer saccharum*) are likely to be negatively impacted by increased flooding and higher pest pressure from forest tent caterpillars (*Malacosoma disstria*) and spongy moths (*Lymantria dispar*) (Cuthrell and others, 2003; Jacobs and Jacobs, 2002).

Management implications. The goals and methods of forest management in response to climate change may offer the greatest opportunity for climate adaptation for this species. Although red-shouldered hawks tolerate small patches of open habitat, intensive and even standard harvest practices that reduce forest density produce unsuitable habitat for this species (Henneman and Andersen, 2009; Naylor and others, 2004). Active management that maintains relatively dense forests may include pest control, the establishment of fuel breaks to reduce fire risk, and the introduction of tree species suited to future conditions (Swanston and others, 2016). In addition, increasing tree species richness and diversity, increasing the quantity of down woody debris, and protecting forested wetlands can all foster the retention of red-shouldered hawks on the landscape (King and others, 2011).”

Mammals

Mammal SGCN that utilize Riparian and Floodplain Forests are nearly identical to those identified occupying Deciduous Wet Forests. The [woodland vole \(*Microtus pinetorum*\)](#) is the only species for which Riparian and Floodplains Forests has been designated its primary habitat. Other primary habitats for the species include Mesic Hardwoods and Deciduous Wet Forests. The voles prefer loose, well-drained soils and humus layers to

construct burrows and sub-surface runways. Grazing by cattle, which compacts the soil, and the presence of invasive non-native earthworms, which destroy the humus, may make forests within its' limited range in southeastern Minnesota unsuitable for this species. Loose, well-drained soils and humus layers are important for constructing burrows and sub-surface runways. Grazing by cattle, which compacts the soil, and the presence of invasive non-native earthworms, which destroy the humus, may make forests within its' limited range in southeastern Minnesota unsuitable for this species. A state Special Concern species, the vole is restricted to two southeastern counties: Winona and Houston.

Other SGCN mammals that utilize riparian and floodplain forests include six of Minnesota's seven bat SGCNs. The water's and riparian edges are ideal sites for drinking and foraging insects. The tri-colored bat (*Perimyotis subflavus*) is the only species not reported in floodplains.

Reptiles

Many other species in greatest conservation need rely on a combination of upland and stream habitats to complete their life cycles. Both wetlands and sandy uplands are necessary for the [Blanding's turtle \(*Emydoidea blandingii*\)](#) to complete its life cycle. Fluvial outwash plains, such as those in Weaver Bottoms along the Mississippi River, provide nesting habitat for blanding's turtles, [gopher snakes \(*Pituophis catenifer*\)](#), eastern hog-nosed snakes (*Heterodon platirhinos*), and map turtles (*Graptemys spp.*). The [wood turtle \(*Glyptemys insculpta*\)](#) hibernates in rivers, nests on undisturbed sandy banks, but spends much of its time in nearby upland forests. Floodplains and riparian are used as primary habitat when vegetative conditions are optimal and there is no flooding; this is especially true for males who tend to remain closer to water during the summer months. Fox snakes (*Elaphe vulpina*) live in forested riparian habitats See also the Rivers and Streams Chapter for more information on aquatic species such as mussels and fish.

Plants

[Dwarf trout lily \(*Erythronium propullans*\)](#) is a federally endangered species endemic to Floodplain Forests in southeastern Minnesota. The plants are typically found on the lower to mid-portions of the slopes descending toward the floodplain. Habitat loss due to development is a main threat to this species. Also, large-scale precipitation events, made more likely by climate change, have recently increased erosion in this species' floodplain habitats, threatening the persistence of some populations. Erosion control or shoreline stabilization measures could minimize the impacts of flash flood events.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Urban development disproportionately expands to riparian corridors and adjacent landscapes. Because of their natural beauty, diversity, recreation opportunities, and abundant natural resources, riparian corridors and adjacent landscapes are appealing places to live. Urban development can directly remove habitat, but it can also have indirect effects on protected habitat. Development in landscapes adjacent to riparian forest can affect habitat quality, particularly for many bird species. Urban development in the adjacent landscape can reduce corridor width, reduce connectivity to other habitats, increase abundances of competitors and predators, alter microhabitat conditions, increase density of nonnative plants, simplify habitat structure, increase direct human disturbance, and prompt avoidance by wildlife that evolved to use large chunks of habitat embedded in a landscape dominated by native ecosystems.



Timber Harvest

Timber harvest can result in failed regeneration and loss of a riparian forest if management activities do not take into consideration many of the unique aspects of these stands. These include the seasonal fluctuation and importance of changing water levels, invasive species, forest pathogens, and changing flood patterns due to climate change, dams, and watershed- wide land use changes. If a harvest is not well enough planned, a riparian forest may convert to a grassland (particularly reed canary grass) or a shrubland (particularly alder, or buckthorn (*Rhamnus cathartica*) in more upland areas). Many riparian and floodplain forests lack young trees to regenerate and succeed the older trees after they are harvested. Selective harvest practices that initially focus on logging the largest and healthiest trees, followed by a second harvest years later, can also result in a stand dominated by reed canary grass, and shrubs.



Recreation

Recreational use can contribute to altered hydrology through soil compaction on trails as well as the introduction of non-native invasive species by inadvertently bringing in seeds on footwear or gear,

including boats.



Dams and Water Management

Dams alter the hydrology and ecology of adjacent floodplain forest. For example, seasonal flood cycles are disrupted, the deposition of sediments and nutrients are reduced, water quality declines, the amount of forest cover decreases, and the natural regeneration of trees is disrupted and altered. Additional run-off from precipitation changes observed with climate change and land use changes are also problematic. This can be seen on the Mississippi River corridor south of the Minneapolis/St. Paul area. The resulting habitat may lack species and structural diversity and ability to regenerate following disturbances including flooding, disease or windstorms.



Invasive Species (Problematic Non-native Species)

Riparian areas and floodplain forests are particularly vulnerable to colonization by invasive species due to the constant influx of seed sources via waterways and the high frequency of flooding events that scour substrates and make open substrates for invasive species to colonize. Garlic mustard (*Alliaria petiolata*), a biennial herbaceous plant, does well in shaded, moist, and disturbed environment and is a common invasive in Minnesota's floodplain forests. In addition to its rapid establishment, studies have documented its ability to disrupt the symbiotic relationship of the roots of many native species with soil fungi, a mechanism that can reduce diversity in the herbaceous ground layer (Rodgers et al., 2022).

Emerald ash borer is an invasive insect that has killed millions of ash trees throughout the eastern half of the U.S. and southeastern Canada. In floodplain forests in northern Minnesota, black ash is often a dominant canopy tree. Its loss is predicted to have major impacts on the structure, composition, and nutrients in ash-dominated habitats. Black ash is also a culturally important tree to some Native American Tribes in the state.

Eurasian earthworms are another invasive disruptor of natural communities, including floodplains. They upend the traditional soil profile (Hale et al., 2005; Fahey et al., 2013), remove the accumulated layers of organic leaf litter (Holdsworth et al., 2008, 2012), and alter the community of soil mycorrhizal fungi (Paudel et al., 2016). These changes cascade through the ecosystem, including ground layer vegetation (Hale et al., 2006; Holdsworth et al., 2007), leaf litter fauna (McCay and Scull, 2019), and salamanders (Maerz et al., 2009; Ziemba et al., 2016). In addition, the presence of earthworms may also benefit invasive plant species (Nuzzo et al., 2009, Whitfeld et al., 2014). Jumping worms (*Amyntas* and *Metaphire* species) are newer species to Minnesota that are predicted to have negative impacts in Minnesota forests as they spread.



Problematic Native Species

Reed canary grass (*Phalaris arundinacea*) is a hardy, aggressive, cool-season grass that can form dense, monotypic stands in wetlands. Despite recent research suggesting that reed canary grass found in Minnesota is predominantly native, it can outcompete other native wetland species, posing challenges for wetland restoration and mitigation efforts. Source: University of Minnesota Extension ([Reed canary grass](#)).



Diseases and Pathogens

Dutch elm disease was first found in Minnesota during the summer of 1961 (French, 1993). Peak losses to all three native elm species that occur in the state (American elm, *Ulmus americana*; red or slippery elm, *U. rubra*; and rock elm, *U. thomasii*) occurred in the late 1970s and 1980s. American elm, in particular, was a dominant species in floodplains across the northeastern United States; today large elm trees are uncommon in floodplain forests and have largely been replaced by silver maple. When present, elms rarely reach heights over 10 meters; this is the height when they become susceptible to Dutch elm disease ([UMN extension](#)).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al., 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

Higher summer temperatures with decreased summer precipitation are projected to increase evaporation, decrease soil moisture, and increased water vapor deficits (Ratcliffe et al., 2025). See "Climate Profile" section above for more detailed information. Warming temperatures are not likely to directly adversely affect most of the tree species that comprise riparian forest. Indeed, many of the species that comprise riparian forest in Minnesota are southerly-distributed and may respond positively to warming temperatures. However, many of the invasive plants that compete with these trees, especially as seedlings, also are likely benefitting from warming temperatures; an example is garlic mustard.

The ranges of some birds are shifting in response to warming temperatures ([NAS 2019](#)). Some of the more southerly-distributed birds that use riparian forest in Minnesota and are predicted to move north as the climate warms include Acadian flycatcher, Louisiana waterthrush, and prothonotary warbler ([NAS 2019](#)).



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota’s average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. Flood timing has shifted so they occur both earlier and later in the season and with longer durations. Though the plants in these riparian forests evolved to survive in floods and even exploit flooding, these changing patterns can result in unsuitable conditions. They may not kill mature trees, but they impact the vegetation and structure of the vegetation on the forest floor, reduce regeneration of young trees, and favor invasives. In particular, these conditions make it difficult for many tree seedlings to establish and survive. Today we are witnessing reduced regeneration of riparian forest because of these changing flood patterns. One example of the detrimental impact of increased large-scale precipitation events can be seen in declines in the quality of habitat and subsequent population declines among populations of the federally endangered dwarf trout lily (*Erythronium propullans*). These high-intensity events have led to the erosion and scouring of finer, silty soils preferred by the species.

The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). Also see “Climate Profile” section above. By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al. 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, to protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN, five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Examples of conservation actions are presented below grouped by the primary strategy that they implement and tagged with icons for the stressor(s) that they address. Some of the strategies, such as Strategy 3, collaborating with partners, could truly be applied to most broadly and effectively implement all actions. In this case, under Strategy 3 we highlight actions for which collaboration is particularly vital.

Potential Conservation Actions for Riparian and Floodplain Forests



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Landscape and urban planning can ameliorate development pressure on floodplain forests and riparian areas by guiding where and how this development occurs and also considering where increased habitat protection and management would be most beneficial. Support local governments in measures that minimize fragmentation and encroachment into high quality forests such as protecting high quality areas, zoning that guides housing developments and lot sizes that support local forests, and land use planning and development guidance.

Stressor	Action
	Since many animals require both upland and stream habitats for their life cycles, it is essential to manage for connectivity of both habitat types. Planning at watershed scales should maintain the corridor characteristics of the floodplain and strong connectivity from upland to aquatic habitat. Protect habitats along river corridors to allow movement of wildlife and facilitate movement of nutrients and energy between the stream and surrounding landscape. Management options include modified lock or dam construction and operation, such as rock arches or weirs, and removal of obsolete dam structures, as well as larger scale watershed actions such as erosion control measures.
	Prioritize protection and management of habitat that provides the elements required by floodplain forest wildlife, but that are less prone to flooding. In particular, maintain and restore floodplain forest along the primary tributaries to the Mississippi River including the Root, Vermillion, Whitewater, and Zumbro rivers in southeastern Minnesota.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.

Stressor	Action
	Planting tree saplings prior to harvest may help prevent riparian forest from converting to problematic species after harvest (i.e. to grassland, (particularly reed canary grass) or shrubland (alder or buckthorn)). Planting a diversity of species provides an opportunity to maintain a riparian forest habitat and increase its resilience in the face of changing climatic conditions and greater flooding intensities and frequencies.
	Timber harvest activities in floodplain forests require careful planning for imperiled species that rely on mature riparian forest, such as cerulean warbler, acadian flycatcher, and red-shouldered hawk. Adequate acreage of suitable mature riparian forest should be available in the landscape before a harvest is implemented. Manage stands to retain biological legacies (at site level) such as large trees with cavities.
	Consider restocking floodplain forests with larger tree saplings rather than seedlings. Because there is minimal natural regeneration of forests along the Mississippi River floodplain planting tree seedlings is often a requirement to get sufficient regeneration to ensure lowland forest is maintained and persists at a site. Audubon has found success planting 3-gallon container trees along the Root River floodplain. The larger container stock has a better chance of survival than bare roots; they can survive long-term floods where the bare root seedlings would likely drown with 10-20 days of

Stressor	Action
	flooding. The larger trees are 4-6 in height and can be protected from deer browse with tree tubes. Once the leaves are above the level of reed canary grass and the deer browse, the trees have a very good chance of surviving to maturity. Fall plantings are encouraged because spring floods are uncertain and can impact access to sites where restoration is planned. The planting could occur prior to harvest, prior to anticipated disturbance (such as an area prone to flooding), or as the forest ages and trees are near dying. If the saplings survive and grow, they can reduce growth of reed canary and other problematic species that respond to an opening canopy. Challenges for this strategy include availability of the saplings, resources such as funding and time, and survival of the saplings if an area is prone to flooding, or because of wildlife browse (deer and small mammals).
	Explore countering increased flood intensities by raising the elevation of the habitat so it is less prone to flooding but still maintained as riparian forest. This might be most applicable and feasible during a planned habitat restoration effort. An example of this is thin layer placement on constructed islands along the main channel of the Mississippi River by the US Army Corps of Engineers.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.

Stressor	Action
	Maintaining the contiguity, resilience, and diversity of native habitats will depend on local people conserving privately owned lands because they have incentives to maintain ecological integrity on their land and in the surrounding landscape.
	Participate and collaborate in interdisciplinary and interagency forest planning, providing information on important habitat types, landscape configurations, and conditions most beneficial for SGCM. Pursue opportunities to collaborate across forest ownerships and encourage the implementation of sustainable forestry at the site and landscape levels to create and maintain habitat conditions important to SGCM. Values such as regional biodiversity, rare species protection, native plant community growth stage diversity, and certain wildlife habitat needs are best considered at larger landscape scales (DNR 2025).

(no activities were identified under Strategy 4)



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Stressor	Action
	Promote and consider further development of tax incentives that enable landowners to gain value for maintaining conservation values on their properties. Provide increased levels of incentive payments for woodland owners to help protect woodlands and forests from further reduction due to development pressure. Support and promote education and technical assistance to maintain or restore forest habitat.
	Encourage climate-smart forestry practices (see Case Studies).
	Where people recreate, provide outreach and education on cleaning gear and equipment to reduce the introduction and spread of invasive species. DNR's Prevent the spread webpage includes specific actions for different land based activities like biking, hiking and off-highway vehicle riding. At trailheads, add invasive species prevention messages, such as those on the boot brush kiosks developed by the PlayCleanGo: Stop Invasive Species in Your Tracks program.
	Research and develop management practices for minimizing the impacts of reed canary grass on open substrate, plant diversity, and new tree growth.

References for Riparian and Floodplain Forest

Fahey, T. J., Yavitt, J. B., Sherman, R. E., Maerz, J. C., Groffman, P. M., Fisk, M. C., & Bohlen, P. J. (2013). Earthworm effects on the incorporation of litter C and N into soil organic matter in a sugar maple forest. *Ecological Applications*, 23(5), 1185–1201. <https://doi.org/10.1890/12-1760.1>

Finch, D.M. et al. (2021). Effects of Climate Change on Invasive Species. In: Poland, T.M., Patel-Weynand, T., Finch, D.M., Miniati, C.F., Hayes, D.C., Lopez, V.M. (eds) *Invasive Species in Forests and Rangelands of the United States*. Springer, Cham. https://doi.org/10.1007/978-3-030-45367-1_4

French, David W. 1993. History of Dutch Elm Disease in Minnesota. Minnesota Agricultural Experiment Station, University of Minnesota. [MR229.pdf \(2.25 MB\)](#)

Hale, C. M., Frelich, L. E., & Reich, P. B. (2006). Changes in Hardwood Forest Understory Plant Communities in Response to European Earthworm Invasions. *Ecology*, 87(7), 1637–1649. [https://doi.org/10.1890/0012-9658\(2006\)87\[1637:CIHFUP\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1637:CIHFUP]2.0.CO;2)

Hale, C.M., Frelich, L.E., Reich, P.B. and Pastor, J., (2005). Effects of European earthworm invasion on soil characteristics in northern hardwood forests of Minnesota, USA. *Ecosystems*, 8, pp.911-927.

He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>

- Holdsworth, A. R., Frelich, L. E., & Reich, P. B. (2007). Effects of Earthworm Invasion on Plant Species Richness in Northern Hardwood Forests. *Conservation Biology*, 21(4), 997–1008. <https://doi.org/10.1111/j.1523-1739.2007.00740.x>
- Holdsworth, A. R., Frelich, L. E., & Reich, P. B. (2008). Litter decomposition in earthworm-invaded northern hardwood forests: Role of invasion degree and litter chemistry. *Écoscience*, 15(4), 536–544. <https://doi.org/10.2980/15-4-3151>
- Holdsworth, A. R., Frelich, L. E., & Reich, P. B. (2012). Leaf Litter Disappearance in Earthworm-Invaded Northern Hardwood Forests: Role of Tree Species and the Chemistry and Diversity of Litter. *Ecosystems*, 15(6), 913–926. <https://doi.org/10.1007/s10021-012-9554-y>
- Hoover, J. P. (2009). Prothonotary Warblers as indicators of hydrological conditions in bottomland forests. In *Tundra to Tropics: Proceedings of the Fourth International Partners in Flight Conference. Partners in Flight, McAllen, Tex* (pp. 128-137).
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Maerz, J. C., Nuzzo, V. A., & Blossey, B. (2009). Declines in Woodland Salamander Abundance Associated with Non-Native Earthworm and Plant Invasions. *Conservation Biology*, 23(4), 975–981. <https://doi.org/10.1111/j.1523-1739.2009.01167.x>
- McCay, T. S., & Scull, P. (2019). Invasive lumbricid earthworms in northeastern North American forests and consequences for leaf-litter fauna. *Biological Invasions*, 21(6), 2081–2093. <https://doi.org/10.1007/s10530-019-01959-1>
- Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). 2025. Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state's greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>
- Moriarty, John J. and Carol D. Hall. 2014. Amphibians and Reptiles in Minnesota. 2014. University of Minnesota Press.
- National Audubon Society (NAS). 2019. Survival by Degrees: 389 Bird Species on the Brink. <https://www.audubon.org/climate/survivalbydegrees>.
- Nuzzo, V. A., Maerz, J. C., & Blossey, B. (2009). Earthworm Invasion as the Driving Force Behind Plant Invasion and Community Change in Northeastern North American Forests. *Conservation Biology*, 23(4), 966–974. <https://doi.org/10.1111/j.1523-1739.2009.01168.x>
- Paudel, S., Longcore, T., MacDonald, B., McCormick, M. K., Szlavecz, K., Wilson, G. W. T., & Loss, S. R. (2016). Belowground interactions with aboveground consequences: Invasive earthworms and arbuscular mycorrhizal fungi. *Ecology*, 97(3), 605–614. <https://doi.org/10.1890/15-1085>

Pfannmuller, L., G. Niemi, J. Green, B. Sample, N. Walton, E. Zlonis, T. Brown, A. Bracey, G. Host, J. Reed, K. Rewinkel, and N. Will. (2017). The First Minnesota Breeding Bird Atlas (2009-2013).

Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. 2025. Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwccasc.umn.edu/node/821>

[Rodgers](#), Vikki L. , [Sara E. Scanga](#) , [Mary Beth Kolozsvary](#) , [Danielle E. Garneau](#) , [Jason S. Kilgore](#) , [Laurel J. Anderson](#) , [Kristine N. Hopfensperger](#) , [Anna G. Aguilera](#) , [Rebecca A. Urban](#) , [Kevyn J. Juneau](#). *BioScience*, Volume 72, Issue 6, June 2022, Pages 521–537, <https://doi.org/10.1093/biosci/biac012>

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>

Sillett, T. S., & Holmes, R. T. (2002). Variation in survivorship of a migratory songbird throughout its annual cycle. *Journal of Animal Ecology*, 71(2), 296–308. <https://doi.org/10.1046/j.1365-2656.2002.00599.x>

Whitfeld, T. J. S., Roth, A. M., Lodge, A. G., Eisenhauer, N., Frelich, L. E., & Reich, P. B. (2014). Resident plant diversity and introduced earthworms have contrasting effects on the success of invasive plants. *Biological Invasions*, 16(10), 2181–2193. <https://doi.org/10.1007/s10530-014-0657-6>

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Ziemba, J. L., Hickerson, C.-A. M., & Anthony, C. D. (2016). Invasive Asian Earthworms Negatively Impact Keystone Terrestrial Salamanders. *PLOS ONE*, 11(5), e0151591. <https://doi.org/10.1371/journal.pone.0151591>

Non-forested Wetlands

Habitat Description

This chapter describes a set of wetland plant communities dominated by herbaceous plants or low-growing woody plant species (shrubs). Non-forested wetlands are found throughout Minnesota and consist of several major types: marshes, wet meadows, fens, and bogs. Like all wetlands, non-forested wetlands occur on sites that are flooded or saturated at least seasonally, if not much of the year. Because of periodic or permanent water inundation, soils in these habitats often become anaerobic. Many dominant plants in wetland communities are tolerant of persistent water levels and have stems, leaves, and roots that contain intercellular air spaces (aerenchyma) that store oxygen and diffuse it from above-water structures to roots during waterlogged conditions.

Marshes occur on frequently inundated sites. These communities are typically inundated by nutrient-rich water. They include emergent marshes and open marshes. Emergent marshes are dominated by vascular plants, such as cattails (*Typha* spp.), that can survive indefinitely with their roots and lower stem submerged and their aerial shoots above water. In addition to cattails, emergent marshes are characterized by perennial emergent plants, such as bulrushes (*Scirpus* spp.) and arrowheads (*Sagittaria* spp.), mixed with annual forbs during low-water periods when substrates are exposed, and with floating-leaved and submergent aquatic plants in settings with persistent standing water. Emergent plants provide important habitat for a variety of wetland bird species. Plants with floating leaves, such as water lilies, dominate open marshes, which are sometimes classified as aquatic communities. Variation in species composition over time occurs in response to changes in hydrological conditions.

Wet meadows are graminoid-dominated wetlands that are subjected to periodic relatively brief inundation and drawdowns during the growing season. The dominant graminoids are broad-leaved species such as lake sedge (*Carex lacustris*), tussock sedge (*C. stricta*), and bluejoint (*Calamagrostis canadensis*). Peak water levels are persistent enough to prevent trees from becoming established. However, there may be little or no standing water present during much of the growing season. As a result, the substrate surface alternates between aerobic and anaerobic conditions. Any organic matter that accumulates over time is usually oxidized during periodic drawdowns and may even burn during severe droughts. Soils range from mineral soils to muck and peat. Because surface water is derived from runoff, stream flow, or groundwater, it is circumneutral (pH 6.0–8.0) and has high mineral and nutrient content. Wet meadows are present statewide on the fringes of wetland basins, along streams and drainageways, in drained beaver ponds, at higher elevations of shallow bays, and as semi-floating mats along sheltered lake shorelines. Wet meadows grade into lowland shrub communities where water levels are lower and less persistent.

Peatlands cover more than six million acres (or approximately 10% of the state), primarily in Northern Minnesota, and are a defining feature of its natural landscape (see [Minnesota Scientific and Natural Areas Patterned Peatlands](#)). These wetlands formed over thousands of years through the slow accumulation of partially decomposed plant material in saturated, low-oxygen conditions. Peatlands may be dominated by mosses such as sphagnum moss (*Sphagnum* spp.) or other plants such as sedges (*Carex* spp.). Some peatlands are forested, primarily by black spruce (*Picea mariana*) and tamarack (*Larix laricina*; also see

Conifer Forest Wetland sub-chapter). The resulting deep organic soils, or peat, sequester and store vast amounts of carbon and support unique hydrologic functions, including flood attenuation, water filtration, and baseflow regulation (read more about DNR's [Peatland Resilience Initiative](#)). Minnesota's peatlands are broadly categorized into bogs and fens, distinguished primarily by their hydrology and nutrient availability.

Bogs receive water only from precipitation. Over time, accumulating sphagnum moss raises the peat surface above the surrounding groundwater table, resulting in acidic (pH < 5.5), nutrient-poor conditions (see also [Acid Peatland System](#)). These environments limit species diversity and are often dominated by black spruce and acid-tolerant sphagnum species. Bogs are most prevalent in northern Minnesota, particularly within the Laurentian Mixed Forest Province, where cool temperatures and high precipitation enable extensive peat development (see also [Laurentian Mixed Forest Province](#)).

Fens receive mineral-rich groundwater, resulting in more neutral pH conditions (5.6–7.5) and higher concentrations of nutrients such as calcium and magnesium. Fens are divided into forested rich peatlands and open rich peatlands (see also [Open Rich Peatland System](#)), depending on their vegetation structure. Open rich peatlands are graminoid- or low-shrub dominated communities characterized by sedges (*Carex* spp.), brown mosses, and minerotrophic *Sphagnum* species. These environments favor sun-loving, shade-intolerant species that thrive in wet, nutrient-poor but mineral-rich conditions. Species such as buckbean (*Menyanthes trifoliata*), tufted bulrush (*Scirpus cespitosus*), and Kalm's lobelia (*Lobelia kalmii*) are commonly found in these systems. Adaptations such as aerenchyma tissue allow these plants to survive in saturated soils, while others, like tufted bulrush, form hummocks that lift the plant above anaerobic conditions. Open rich peatland communities are widespread in the Laurentian Mixed Forest Province, where abundant precipitation, cool climate, and poorly drained glacial basins support sustained peat development. However, these communities also extend into the Eastern Broadleaf Forest, Prairie Parkland, and Tallgrass Aspen Parklands provinces, where they are typically found near the southern and western margins of the peatland range. In these warmer, more evaporative regions, open rich peatlands are restricted to floating mats or areas with sufficient groundwater input to maintain saturation throughout the growing season.

Calcareous fens are a rare type of fen that develops in areas with particularly high calcium content, often where calcareous glacial till or limestone bedrock is present. These systems support a number of plant species in Greatest Conservation Need, including highly specialized and rare species such as [twig rush](#) (*Cladium mariscoides*) and [hairlike beak rush](#) (*Rhynchospora capillacea*), as well as grass of Parnassus (*Parnassia* spp.), a calciphilic (calcium loving) indicator species. Calcareous fens are predominantly located in western and southern Minnesota, including the Minnesota River Valley and parts of the southeastern karst region, where conditions favor calcium-rich groundwater discharge (see also [Calcareous Fens](#)).

Habitat Map

To depict non-forested wetland habitat (see Figure 3.15), we compiled spatial data from several sources: DNR's Native Plant Communities, National Wetland Inventory for Minnesota, and Midwest Terrestrial Habitat System created by the USFWS Midwest Landscape Initiative (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Marsh (MR)

[MRn83 Northern Mixed Cattail Marsh \(PDF\)](#)

[MRn93 Northern Bulrush-Spikerush Marsh \(PDF\)](#)

[MRu94 Lake Superior Coastal Marsh \(PDF\)](#)

[MRp83 Prairie Mixed Cattail Marsh \(PDF\)](#)

[MRp93 Prairie Bulrush-Arrowhead Marsh \(PDF\)](#)

Wet Meadow / Carr (WM)

[WMn82 Northern Wet Meadow/Carr \(PDF\)](#)

[WMs83 Southern Seepage Meadow/Carr \(PDF\)](#)

[WMs92 Southern Basin Wet Meadow/Carr \(PDF\)](#)

[WMP73 Prairie Wet Meadow/Carr \(PDF\)](#)

Open Rich Peatland (OP)

[OPn81 Northern Shrub Shore Fen \(PDF\)](#)

[OPn91 Northern Rich Fen \(Water Track\) \(PDF\)](#)

[OPn92 Northern Rich Fen \(Basin\) \(PDF\)](#)

[OPn93 Northern Extremely Rich Fen \(PDF\)](#)

[OPp91 Prairie Rich Fen \(PDF\)](#)

[OPp93 Prairie Extremely Rich Fen \(PDF\)](#)

Acid Peatland (AP)

[APn90 Northern Open Bog \(PDF\)](#)

[APn91 Northern Poor Fen \(PDF\)](#)

National Wetlands Inventory

From the [National Wetland Inventory for Minnesota \(NWI\) Layer](#) we included these Simplified Plant Community Class types: Shallow Marsh, Deep Marsh, Seasonally Flooded/Saturated Emergent Wetland, Shrub Wetland, Open Bog, and Shallow Open Water Aquatic Community.

Midwest Terrestrial Habitat System

From the [Midwest Terrestrial Habitat System](#) we included these groups: Eastern North American Boreal-Subboreal Alkaline Fen, Midwest Prairie Alkaline Fen, Eastern North American Freshwater Marsh, Eastern North American Ruderal Meadow & Shrubland, and Eastern Ruderal Wet Meadow & Marsh.




Figure 3.15. A map depicting Non-forested Wetland habitat in Minnesota, including DNR Native Plant Community Classes in the Ecological Systems of Marsh, Wet Meadow/Carr, Open Rich Peatland, and Acid Peatland as well as several types from the National Wetland Inventory for Minnesota and Midwest Terrestrial Habitat System (see narrative).

Conservation Overview

Wetlands provide a number of ecological benefits to the landscapes in which they are found including erosion control, flood control, groundwater recharge and discharge, water quality, recreation, habitat for rare plant and animal species, and economic and cultural value.

Prior to European settlement, Minnesota's wetlands covered about 18.5 million acres, accounting for nearly 35 percent of the state's landscape (Phillips 1997). Since then, approximately half of those wetlands have been lost, primarily through conversion and drainage for agriculture, urban development, and other land use changes, leaving just over 10 million acres of wetland today (Haga 2015). Losses have been most severe in southern and western Minnesota, where up to 95 percent of original wetlands have vanished, whereas northern wetland complexes remain relatively intact and high-quality (Minnesota Pollution Control Agency 2025). Although Minnesota's 1991 Wetland Conservation Act has largely limited acreage losses, overall wetland quality continues to face threats from invasive species, altered hydrology, and nutrient loading (Marohn 2024). While there are new additions to wetland acreage in Minnesota, replacement wetlands can lack the diversity and complexity of natural wetlands (Whigham 1999). Of particular note are peatlands, one of Minnesota's most ecologically significant landscapes. Historically, peatlands were considered as unproductive and unusable land, and European settlers drained them for conversion to agriculture and forestry uses (The Nature Conservancy 2025). Peat was also harvested for fuel and horticultural use, and in the 1970s and 1980s, proposals emerged to develop peat as an energy resource (Keirstead 1992). However, most of these plans were abandoned due to economic, environmental, and regulatory concerns, coupled with increasing recognition of the ecological importance of peatlands.

Today, peatlands are valued for the many ecosystem services they provide. They are globally significant carbon sinks, storing vast amounts of carbon and helping to regulate greenhouse gases. When disturbed or drained, however, peatlands can become major sources of carbon dioxide and methane emissions. Ecologically, they support unique plant and animal communities adapted to wet, acidic, nutrient-poor conditions, including carnivorous plants, rare orchids, boreal birds, and several SGCN. Peatlands also play an important role in hydrology, storing and slowly releasing water to moderate flood risks and maintain base flows, while filtering sediments and nutrients to improve water quality (Bonn et al. 2016). Their relative intactness also makes them likely refuges for species as climate conditions shift.

Recognizing their ecological value, Minnesota has taken steps to conserve peatlands through legislation and land protection. The Minnesota Peatland Protection Act of 1991 established a system for identifying and safeguarding high-quality peatland ecosystems, including the designation of Scientific and Natural Areas (SNAs) such as the Red Lake and Lost River peatlands. Many peatlands are now managed under state or federal ownership, with careful consideration given to land use, restoration potential, and climate resilience. These efforts reflect the growing understanding that peatlands are not only biologically rich and hydrologically important but also essential in mitigating climate change.

Calcareous fens are one of Minnesota's rarest and most ecologically unique wetland types, which supports an assemblage of specialized and often endangered plant and animal species, many of which are found nowhere else in the state. Historically, calcareous fens were more common across Minnesota's landscape, particularly in the western and southern regions where glacial activity and underlying calcareous bedrock created suitable conditions (see also [Calcareous Fens](#)). However, their extent has been dramatically reduced since European settlement due to agricultural drainage, peat mining, groundwater pumping, and development (Prince 1997).

Their location on flat or gently sloping lands made them attractive for conversion to cropland, and many were drained or filled before their ecological importance was fully understood. In the 20th century, some fens were also impacted by excavation for peat or marl and by groundwater withdrawals that altered the delicate hydrology required to sustain fen ecosystems (Prince 1997).

Today, calcareous fens are recognized as exceptionally valuable and fragile ecosystems. They are home to a diversity of rare and state-endangered plant species, such as the flowering plant [valerian \(*Valeriana edulis*\)](#), [sterile sedge \(*Carex sterilis*\)](#), and slender reedgrass (*Calamagrostis stricta*), as well as rare orchids, mosses, and invertebrates including many SGCN (see [Calcareous Fens](#)). Because they are groundwater-fed, calcareous fens are especially sensitive to changes in hydrology and land use, including groundwater withdrawals from nearby municipal, industrial, or agricultural wells (Prince 1997). Their ecological function depends on maintaining stable, clean, and abundant groundwater flow, which not only sustains fen vegetation but also buffers downstream water bodies against sedimentation and nutrient overload. The Minnesota Department of Natural Resources (DNR) maintains a registry of over 200 documented calcareous fens, and active efforts are in place to monitor, research, and protect these critical habitats (see also [Calcareous Fens](#)).

Climate Profile for Non-forested Wetlands

We provide information anticipated future climate impacts in non-forested wetland habitat types from two reports created by the US Geological Survey Midwest Climate Adaptation Science Center in support of State Wildlife Action Plans. All content in this section is derived from these reports: “Effects of Climate Change on Midwestern Ecosystems: Eastern North American Temperate Freshwater Marsh, Wet Meadow and Shrubland” (Ratcliffe et al. 2025a) and “Effects of Climate Change on Midwestern Ecosystems: Bogs and Fens” (Ratcliffe et al. 2025b).

The primary stressor for marsh and wet meadow/carr wetlands is fluctuating hydrology. There is anticipated more frequent and severe flooding driven by increased spring precipitation in more frequent extreme events. In addition, summer and fall droughts will likely be more frequent, driven by higher temperatures and evapotranspiration demands.

Projected changes in annual temperatures in the Mixed Wood Plains ecoregion of central Minnesota for the late century (2070-2099) are 6 – 11 degrees F over historical (1971-2000), depending on emissions and using the mean of 20 climate models (Table 2 in Ratcliffe et al. 2025a). Spring precipitation is projected to rise 25% over historical conditions by end of century (Table 4 in Ratcliffe et al. 2025a). The frequency of 30-yr droughts is projected to be as much as 917% over historical rates by end of century (Table 5 in Ratcliffe et al 2025a).

Marshes (from the North American Freshwater Marsh section of Ratcliffe et al. 2025a)

Marshes are adapted to fluctuating hydrology, typically with plants suited for various inundation levels in zones within them. However, shallow marshes may be sensitive to water deficits.

- **Habitat Structure** - Flooding may increase nutrient inputs and sediment deposition from adjacent developed or agricultural areas, increasing eutrophication, the distribution of invasive species, and altering habitat structure through scouring and deposition. Increased drought phases may accelerate decomposition, carbon dioxide release, and mineralization during dry periods.
- **Community Composition** – Due to the increased incidence of flooding as well as drought, the particulars of which species will be favored under future climate conditions is likely quite site-specific.

More frequent, severe drought may favor facultative wetland species rather than obligate and promote meadow and shrub species more typical of drier-adapted communities. Wild rice (*Zizania palustris*) marshes may be particularly vulnerable to hydrological change, because as annuals they can be easily uprooted during flooding or outcompeted under both flooding and drought.

- **Problematic Species** – Prolonged flooding and increased nutrient loads may favor species such as reed canarygrass (*Phalaris arundinacea*), European common reed (*Phragmites australis* ssp. *australis*), and cattails (*Typha* spp.).

Wet Meadow/Carr (from the Midwest Wet Prairie, Wet Meadow, and Shrub Swamp of Ratcliffe et al. 2025a)

These wetlands are typically shallow, seasonally or intermittently flooded wetlands. Their hydrology is largely reliant on precipitation and snowmelt, and increased evapotranspiration rates driven by higher temperatures may exacerbate their vulnerability to drying.

- **Habitat Structure** - Flooding may increase nutrient inputs and sediment deposition from adjacent developed or agricultural areas, increasing eutrophication, the distribution of invasive species, and terrestrialization. Increased drought phases may reduce wetland habitat extent through drying soils. Frequent shifting of soil between aerobic and anaerobic states may slow decomposition, denitrification, and plant nutrient uptake during saturation and accelerate decomposition, carbon dioxide release, and mineralization during drought.
- **Community Composition** – Given the projected increase in both flooding and drought, the plants that will be favored under future climate conditions will likely vary considerably from site to site. Flood-tolerant species may become more dominant. Drought may facilitate dominance by annuals and rhizomatous graminoids, shifting communities toward shrubs and graminoids typical of drier-adapted communities.
- **Problematic Species** – Prolonged flooding and increased nutrient loads may favor species such as reed canarygrass, European common reed, and cattails. Warming trends may also create conditions favorable to expansion of invasive taxa from more southern regions.

Bogs (acid peatlands) – (from the Eastern North American Boreal-Subboreal Bog and Acidic Fen section of Ratcliffe et al. 2025b)

In the Northern Forest ecoregion of northeastern Minnesota, projected changes in annual temperature by end of century (2070-2099) are 6 – 11 degrees higher than historic (1970-2000), depending on the carbon emissions level and using the mean of 20 climate models (see Table 9 in Ratcliffe et al. 2025b). Rainfall projections include greater expected winter and spring rain and more summer drought. Less summer rainfall paired with higher temperatures is projected to raise the vapor pressure deficit (VPD) 32-66% by end of century (2070-2099) over historic (1970-2000), depending on emissions levels, using the mean of models (Table 11 in Ratcliffe et al. 2025b).

Anticipated climate change impacts:

- **Habitat Structure** - Higher VPD corresponds with rapid evaporation and water loss from plants and soils and reduced water table levels. When water levels drop, peat becomes more exposed to oxygen, and this leads to accelerated aerobic decomposition and release of nutrients including carbon. Sustained heat and lack of moisture can hurt the productivity of Sphagnum moss (*Sphagnum* spp.) and diminish its dominance (Bu and others, 2011; Norby and others, 2019 in Ratcliffe et al. 2025b). Drier

conditions lead to peat subsidence, compaction, and cracking, which reduces water retention, limits hydraulic conductivity, and increases probability of wildfire (Ratcliffe et al. 2025b).

- **Community Composition** – climate changes, particularly increased temperatures that enhanced nutrient availability and extended growing seasons, are anticipated to be unfavorable for black spruce, tamarack, and sphagnum moss species, while being more favorable for ericaceous shrubs such as highbush blueberry (*Vaccinium corymbosum*), bog rosemary (*Andromeda polifolia*) and small cranberry (*Vaccinium oxycoccos*) (Ratcliffe et al. 2025b).
- **Problematic Species** – Increased nutrient availability and longer growing seasons create more favorable conditions for species such as glossy buckthorn (*Frangula alnus*), reed canarygrass, common reed, and cattails.
- **Pests and pathogens** – In general, climate change may increase populations and facilitate spread of pests and pathogens, as well as decreasing the fitness of dominant species, making them more susceptible to outbreaks, such as tamarack (*Larix laricina*) being vulnerable to eastern larch beetle (*Dendroctonus simplex*) and black spruce (*Picea mariana*) to spruce budworm (*Choristoneura fumiferana*) (Bellemin-Noël and others, 2021; McKee and others, 2022; Ren and others, 2020 in Ratcliffe et al. 2025b).

Fens – (from Midwest Prairie Alkaline Fen section in Ratcliffe et al. 2025b)

Anticipated climate change impacts:

- **Habitat Structure** – warming paired with reduced summer rainfall creates water deficits that can lower water tables and expose peat to oxygen, increasing decomposition and releasing nutrients and carbon. Drier conditions lead to peat subsidence, compaction, and cracking, which reduces water retention, limits hydraulic conductivity, and increases probability of wildfire. These changes could reduce conditions for fen-adapted specialists while making these habitats more available to more generalist forbs and graminoids.
- **Community Composition** – more frequent water deficits from drought may create difficult growing conditions for some species such as spikerushes (*Eleocharis* species) and grass of parnassus (*Parnassia glauca*), while favoring more deep-rooted plants and facilitate encroachment by woody shrubs and trees. Longer growing seasons and enhanced nutrient availability will shift competitive dominance by fen specialist forbs and mosses while creating conditions more favorable to ericaceous shrubs and graminoids.
- **Problematic Species** – Longer growing seasons along with increased nutrient availability may facilitate conditions favorable for species such as glossy buckthorn (*Frangula alnus*), reed canarygrass, common reed, and cattails.
- **Pests and pathogens** – Pest and pathogen pressure is anticipated to increase directly through higher pest and pathogen populations and indirectly through heightened host susceptibility due to temperature and water deficit related stress.

Species in Greatest Conservation Need

Non-forested wetlands provide primary or secondary habitat for 97 animal Species in Greatest Conservation Need (SGCN) and 52 plant SGCN (see Table 3.13.). Primary habitats are those that species rely on and use most consistently, and the loss or degradation of which would most greatly impact the species. Secondary habitats are used by species less frequently.

Table 3.13. Numbers of SGCN associated with Non-forested Wetlands.

Species Group	Primary Habitat*	Secondary Habitat	Total
Amphibians	4	1	5
Birds	18	12	30
Mammals	3	3	6
Reptiles	1	5	6
Aquatic Invertebrates			
Mussels	0	1	1
Dragonflies and damselflies			7
Terrestrial Invertebrates			
Bees			20
Beetles (terrestrial)			1
Butterflies			6
Moths			14
Spiders	1	0	1
Plants	52		52
Total	126	22	149

*Primary and secondary habitat associations were distinguished for vertebrate and some invertebrate animals but not for insects, which are shown in 'total'. Plants were associated with only one primary habitat each.

Animals with more general habitat requirements were associated with multiple habitats, while specialists were associated with fewer. Plants were only associated with a single primary habitat for each species. Detailed tables associating each of the SGCN with each of the 15 habitats can be found in Appendix C for animals and Appendix D for plants.

Non-forested wetlands provide habitat for one-third of the 131 terrestrial vertebrate SGCN identified in the 2025-2035 SWAP (36%; 47 species). Thirty of these SGCN are bird species, and the 30 birds utilizing these wetlands comprise 38% of the 79 avian SGCN. The 52 plant SGCN found in this habitat constitute 12% of the 447 that are designated as SGCN. Examples of some of the SGCN and their significance are described below.

Amphibians

Non-forested Wetlands are a rich and important habitat for amphibian SGCN. Five of the seven amphibian SGCN are found in this habitat. It is especially critical for four species: [*Blanchard's cricket frog \(Acris*](#)

[blanchardi](#)), [spotted salamander \(*Ambystoma maculatum*\)](#), pickerel frog (*Lithobates palustris*), and eastern newt (*Notophthalmus viridescens*). As noted in the Riparian and Floodplain Forest chapter which also provides critical habitat, the cricket frog is found right along the water's edge; open areas with muddy shorelines and dense emergent vegetation are preferred. They reside near water throughout the summer but, during dry conditions, they may seek out wetter conditions elsewhere. Crayfish burrows or small openings along the bank of a wetland or river provide sites for hibernation. Less than 2 inches in size, their breeding call resembles that of some birds, making visual identification essential. A state endangered species in Minnesota, currently populations are limited to sites in Hennepin and Winona counties.

[Spotted salamanders \(*Ambystoma maculatum*\)](#), a state Special Concern species, also have a very limited distribution in the state, restricted to two east-central counties: Carlton and Pine. Found in forests with seasonal ponds or in shallow emergent wetlands that do not contain predatory fish to ensure successful reproduction. Mature, close-canopy forests that provide sufficient shade and organic soils with woody debris for foraging are important habitat features.

Pickerel frogs are also found in Riparian Forests and Floodplains as well as open non-forested wetlands, preferring the edges of ponds, small rivers, and streams that flow through wooded areas or open habitats (Moriarty and Hall 2014). The wetlands they use are typically on a valley floor near a river or stream, are permanent, and are never far from forested habitats. They rarely venture far from the water's edge or from nearby wet meadows. Winter months are spent hibernating in their water habitats. Pickerel frogs are restricted to the far southeastern corner of the state; county records are available from Goodhue, Fillmore, Houston, Mower, Olmsted, and Winona counties (Moriarty and Hall 2014). Eastern newts have a much wider state distribution in Minnesota than other amphibian SGCN. They can be found temporary or permanent wetlands throughout deciduous and coniferous forests across the eastern half of the state. A rich, organic understory with abundant leaf litter and woody debris is required (Moriarty and Hall 2014). They are known to be very active during warm, rainy days and hide under the leaf litter when it is drier.

The fifth amphibian found in open wetlands is the [Great Plains Toad \(*Anaxyrus cognatus*\)](#). It is more prominent in prairies and other grasslands but can also be found in shallow wetlands during the breeding season. They are not especially selective about their wetland breeding sites; they also utilize any shallow pool of water, including low, ephemeral sites in agricultural fields that form following a heavy rainstorm.

Birds

[Yellow rails \(*Coturnicops noveboracensis*\)](#) and [Nelson's sparrows \(*Ammodramus nelsoni*\)](#) are almost entirely restricted to **rich fens** and **wet meadows**. Permanent water depth during the nesting season ranging from a few centimeters to about 25-30 centimeters seems critical to occupancy. A canopy of dead sedges (previous years' growth) is a key habitat feature. Nelson's sparrows often utilize isolated shrubs or small clumps of cattails as singing perches. However, neither species is found in cattail-dominated wetlands, or wetlands with dense shrub cover. Sedge wetlands of acid peatlands, with dense sphagnum mosses or ericaceous shrubs, are also avoided.

Sedge wrens (*Cistothorus stellaris*) and LeConte's sparrows (*Ammodramus leconteii*) are also found in **wet meadows** and **rich fens**, but will utilize mesic upland grasslands with dense, tall grasses. [Wilson's phalaropes \(*Phalaropus tricolor*\)](#) have similar habitat requirements, but typically require at least small patches of shallow, open water within fens, wet meadows, or upland grasslands. Northern harriers (*Circus hudsonius*) and [short-](#)

[eared owls \(*Asio flammeus*\)](#) may occupy the same sedge-dominated habitats as well as open acid peatlands, marshes, shrub swamps, and upland grasslands.

Emergent marshes provide key habitat for [king rails \(*Rallus elegans*\)](#), [common gallinules \(*Gallinula chloropus*\)](#), [Franklin's gulls \(*Leucophaeus pipixcan*\)](#), black terns (*Chlidonias niger*), [Forster's terns \(*Sterna forsteri*\)](#), least bitterns (*Botaurus exilis*), and American bitterns (*Botaurus lentiginosus*). Each of these species have slightly different microhabitat requirements. In general, a vital habitat characteristic is a moderate to high degree of interspersed of emergent vegetation and open water. Dense cattail stands without accessible open water are avoided by all of these species. Exposed bare soil, floating mats of vegetation, or muskrat house provide important nesting and loafing sites, particularly for Franklin's gulls, black terns, and Forster's terns. Common gallinules often utilize areas of open water with water lilies within hemi-marshes. Least and American bitterns are restricted to marsh vegetation, typically foraging at the vegetation-open water interface. While least bitterns are largely restricted to cattails (as well as patches of phragmites), American bitterns will also occupy shrub swamps, wet meadows, and sometimes acid peatlands.

Another group of wetland-dependent SGCN birds utilize the open water and/or shorelines of non-forested wetlands, particularly emergent marshes, as well as lakes (also see Lakes Chapter). This includes [trumpeter swan \(*Cygnus buccinator*\)](#), ducks, grebes, and common loon (*Gavia immer*). [Common terns \(*Sterna hirundo*\)](#) and [American white pelicans \(*Pelecanus erythrorhynchos*\)](#) forage in/over open water but nest on shorelines (typically islands). Black-crowned night herons (*Nycticorax nycticorax*) forage at wetland edges or shoreline, but nest colonially in trees or sometimes marshes. Buff-breasted sandpipers (*Calidris subruficollis*) utilize very short vegetation, often associated with wetlands.

Mammals

[Northern bog lemmings \(*Synaptomys borealis*\)](#) are limited to lowland conifer forests and open peatlands in extreme northern Minnesota; they have been shown to disappear from peatlands altered by human activities. Little is known about the species. Current populations are believed to be isolated relics of a larger population that occurred much farther south when glaciers advanced during the Pleistocene epoch. Fewer than 12 records have been documented in Minnesota since the species was first found in Lake County in 1932. Those records have been limited to open bogs, shrub carrs, and black spruce swamps. Since the [Eastern heather vole \(*Phenacomys ungava*\)](#) was first documented in Lake County in 1940, fewer than 20 specimens have been collected in Cook, Lake, and St. Louis Counties. Like the northern bog lemming it also prefers lowland conifer forests and open peatlands but may be found in upland fire-dependent habitats as well. Proximity to water and the presence of boulders, coarse woody debris, and ericaceous plants appear to be important habitat components. Larger species typical of conifer wetlands include the SGCN snowshoe hare (*Lepus americanus*), [Canada lynx \(*Lynx canadensis*\)](#).

Franklin's ground squirrel (*Poliocitellus franklinii*) is strongly associated with prairie, other grasslands, and savanna habitats. However, the grasslands it selects usually include wooded banks and gullies for its underground burrows ([Missouri DNR](#)). The species is declining throughout its range and is under consideration for federal status.

Reptiles

Where they have access to them, [Blanding's turtles \(*Emydoidea blandingii*\)](#) have been documented using bogs and fens as well as open wetland complexes that are adjacent to sandy uplands for nesting. For example, adult females have been tracked to fens prior to and following nesting; both to hydrate and to forage. Potentially, hatchlings may seek the deep muck for protection. Shallow waters with abundant aquatic vegetation are preferred. Ephemeral wetlands are utilized in spring and early summer, while deeper marshes and backwater pools are utilized in both the summer and winter. In southwestern Minnesota, meandering streams and rivers, fens, prairie marshes, backwaters, and oxbows are important aquatic habitats, and upland habitats include adjacent agricultural lands. In addition to Blanding's turtle, four snakes may also be found in or near open non-forested wetlands: [plain hog-nosed snake \(*Heterodon nasicus*\)](#); Eastern hog-nosed snake (*Heterodon platirhinos*); smooth greensnake (*Opheodrys vernalis*); and [gopher snake \(*Pituophis catenifer*\)](#).

Invertebrates

Many insects both terrestrial and aquatic are associated with Non-forested Wetlands, including 50 SGCN. Please see the Aquatic Invertebrates and Terrestrial Invertebrates sub-chapters for more information.

Climate Profile: Zigzag Darner

This profile is excerpted from "Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest" published by the Midwest Climate Adaptation Science Center (Ratcliffe et al. 2025c).

The [zigzag darner \(*Aeshna sitchensis*\)](#) is a boreal species of Odonata [dragonfly] found in northern Minnesota, Wisconsin, and Michigan (DuBois and others, 2004). While little is known about the direct effects of climate change on zigzag darners, research on other Odonata (e.g., dragonfly and damselfly) species has documented climate-driven changes such as advanced phenology, altered development time, pigmentation changes, behavioral adjustments in time spent basking and other activities, and shifts in geographic distribution (Hassall and Thompson, 2008; Zeuss and others, 2014). Additionally, higher temperatures may increase Odonata sensitivity to environmental stressors, including insecticides and other pollutants (Ishiwaka and others, 2024; Sirois-Delisle and Kerr, 2022). As generalist aquatic predators, larval Odonata are unlikely to be directly affected by changes in prey phenology (Hassall and Thompson, 2008). However, rising temperatures may alter their ability to capture prey. One study found that at higher temperatures, zigzag darner larvae showed decreased activity, while their prey, wood frog (*Lithobates sylvaticus*) tadpoles, became more active (Hayden and others, 2015). However, this relationship is complex, as the number of larval attacks also increased with temperature.

Zigzag darners breed in cool-water bogs and fens with little to no emergent vegetation. In the Great Lakes region, these habitats are vulnerable to climate-driven changes in water temperature, ice-cover, and water-level fluctuations (DuBois and others, 2004; DuBois and others, 2018; Magee and others, 2021). This species prefers small pools that often dry during summer (DuBois and others, 2004), but increasing rainfall variability, with cycles of drought and deluge, may lead to more frequent early drying or pool connections to larger water bodies, reducing suitable breeding habitat. Maintaining and restoring boreal, cool-water bogs and fens will be essential to zigzag darner persistence under warming temperatures and variable precipitation patterns. Additionally, because warming temperatures often amplify other environmental stressors, management strategies may include reducing exposure to toxins, such as insecticides (Ishiwaka and others, 2024) and copper (Hayden and others, 2015).

Plants

Minnesota's peatlands provide refuges for several [endangered, threatened, or special concern](#) plant species in the state, including [linear-leaved sundew \(*Drosera linearis*\)](#), [English sundew \(*Drosera anglica*\)](#), [coastal sedge \(*Carex exilis*\)](#), [twig rush \(*Cladium mariscoides*\)](#), [bog rush \(*Juncus stygius* var. *americanus*\)](#), [sooty colored beak-rush \(*Rhynchospora fusca*\)](#), and [montane yellow-eyed grass \(*Xyris montana*\)](#) (see also [Minnesota Scientific and Natural Areas Patterned Peatlands](#)).

Marshes and wet meadow are habitat for rare species including: sweet-smelling [Indian plantain \(*Hasteola suaveolens*\)](#), [marginated rush \(*Juncus marginatus*\)](#), [auricled twayblade \(*Listera auriculata*\)](#), [sea milkwort \(*Lysimachia maritima*\)](#), [autumn fimbry \(*Fimbristylis autumnalis*\)](#), [neat spikerush \(*Eleocharis nitida*\)](#), [smooth-sheathed sedge \(*Carex laevivaginata*\)](#), [upswept moonwort \(*Botrychium ascendens*\)](#), [stream parsnip \(*Berula erecta*\)](#), [yellow bartonia \(*Bartonia virginica*\)](#), and [great Indian plantain \(*Arnoglossum reniforme*\)](#).

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts (e.g. catastrophic wildfire following a prolonged period of fire suppression) while serving as valuable conservation tools in others (e.g., appropriately planned and applied prescribed fire).

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Pressure for urban and exurban development puts pressure on wetlands, both directly, such as in the Anoka Sandplain, and indirectly in the use and alteration of groundwater hydrology, which is particularly detrimental to groundwater-fed wetlands such as calcareous fens.



Crop Production

Wetlands in general and calcareous fens in particular are affected by agricultural practices either by direct use, chemical inputs such as fertilizers or pesticides, or by changes to the flow of groundwater to the wetlands, such as by installation of wells or drain tile, which once disrupted, is essentially not possible to reinstate and causes extirpation of the wetland.

Open wetlands, particularly seasonal and temporary wetlands, have been lost from the landscape due to being able to be farmed in some years, in some cases through the placement of drain tile, conversion of these to farmland reduces rich, seasonally available habitat for migratory waterfowl and shorebirds.



Mining and Quarrying

Mining and quarrying operations can threaten wetlands directly with insufficient buffers and indirectly through changes in hydrology, including impacts to the level or flow of water in the water table.

Peat mining remains a significant stressor on Minnesota’s peatlands, impacting ecosystem structure, water quality, and greenhouse gas emissions (Winkler and DeWitt 1985, Jones 2021). There are at least seven identified major environmental impacts associated with large-scale peat mining including potential release of toxic metals, eutrophication of surface waters, increased surface runoff, release of organic pollutants,

alterations to freshwater-saltwater balance, changes in groundwater supply, and air pollution and increased fire risk (Winkler and DeWitt 1985).



Wind and Solar Energy Infrastructure

Siting of solar development can impact wetland hydrology and associated communities.



Roads, Trails, and Railroads

The development of roads and trails can disrupt wetlands directly and indirectly by changing hydrology, both above and below ground in the aquifer, including in some cases the flow of cold groundwater into wetlands.



Utility Corridors

Powerlines present a risk to waterfowl when situated between feeding and roosting areas. This is especially true for lines near lakes, rivers and wetlands. Less agile birds, such as trumpeter swans are particularly susceptible to collision with overhead lines located near waterbodies (APLIC, 2012).

Gas pipelines can fragment wetlands habitats and also disrupt groundwater hydrology.



Recreation

Recreational use can contribute to altered hydrology through soil compaction as well as the introduction of non-native invasive species to recreational sites with wetlands by inadvertently bringing in seeds on footwear or gear.



Dams and Water Management

Water impoundments and surface water diversions can disrupt the upwelling of groundwater that maintains fen and open rich peatlands unique chemistry and hydrology. For wet meadows and sedge-dominated systems, dams and irrigation can elevate water levels which favor invasive or more water-tolerant species like cattails (*Typha* spp.) and reed canary grass (*Phalaris arundinacea*) (Apfelbaum and Sams 1987, Boers et al. 2007). Water impoundments reduce the natural variability in flow and moisture regimes, potentially degrading habitat for species dependent on specific hydrologic conditions such as nesting birds, amphibians, and rare wetland invertebrates (Winter and Woo 1990).



Invasive Species (Problematic Non-native Species)

Numerous invasive plant species with detrimental effects are present in Minnesota wetlands. For example, purple loosestrife (*Lythrum salicaria* L.), invasive cattail (*Typha angustifolia* L., *Typha x glauca*), and invasive common reed (*Phragmites australis australis* (Cav.) Trin. ex Steud.) are all associated with habitat declines for birds, homogenization and dominance of wetland plant communities, and modification of wetland soil and nutrient cycling. In addition to these emergent plants, multiple submerged species including curlyleaf

pondweed (*Potamogeton crispus*) and Eurasian watermilfoil (*Myriophyllum spicatum*) have impacts including increasing internal phosphorus loading and outcompeting native plant species, respectively. Human mediated movement of species means that in addition to the current distribution of invasive species, there is a persistent risk of additional introductions of invasive species. Hybridization of native and invasive species is an additional concern, with hybrid cattail and hybrid Eurasian watermilfoil both wetland examples of species with documented hybrid vigor (hybrids are superior competitors relative to parent strains) and hybrid Eurasian watermilfoil having documented herbicide tolerance.

The invasive subspecies of Phragmites (*Phragmites australis subsp. australis*) is a tall densely growing perennial grass that can be found in wetlands, shorelines, wet ditches, and other wet areas. The species can rapidly spread through seed, stolons, rhizomes, and stem fragments, forming dense stands that alter wetland hydrology, lower plant diversity, and reduce habitat quality for invertebrates, fish, and waterbirds. There is a native subspecies of Phragmites that can be difficult to distinguish from its invasive counterpart. This makes it difficult to estimate the distribution of the invasive species throughout the state. Source: Minnesota Aquatic Invasive Species Research Center (MAISRC) [Invasive Phragmites | Minnesota Aquatic Invasive Species Research Center \(MAISRC\)](#)



Problematic Native Species

Reed canary grass (*Phalaris arundinacea*) is a hardy, aggressive, cool-season grass that can form dense, monotypic stands in wetlands. Despite recent research suggesting that reed canary grass found in Minnesota is predominantly native, it can outcompete other native wetland species, posing challenges for wetland restoration and mitigation efforts. Source: University of Minnesota Extension ([Reed canary grass](#)).



Water-borne Pollution

Nonpoint source pollution, including phosphorus, nitrogen, sediment, bacteria, and other contaminants generally results from land runoff, precipitation, atmospheric deposition, drainage, seepage or hydrologic modification. As runoff flows, it picks up and carries away natural and human-made pollutants, depositing them into lakes, rivers, wetlands, and groundwater, and estuaries ([EPA NPS Webpage](#)). These nonpoint source pollutants represent the largest combined contribution (an estimated 86%) to the state's water pollution. ([MPCA Water Quality Initiatives](#)) Water quality measured by nutrients and contaminants has been shown to decline with higher agricultural or urban development in the watershed. Some key impacts include excess algal growth, reduced plant diversity, and reduced diversity in fish and invertebrate communities. ([DNR Water quality non-point sources](#))



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control

Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). Also, projected temperature increases are likely to affect ecosystems unevenly. For example, peatlands are sensitive to changes in temperature and hydrology and may experience reduced capacity to provide critical ecosystem services such as carbon sequestration and water storage. For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al. 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Examples of conservation actions are grouped below under these five strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Also note that some strategies, such as Strategy 3, collaborating with partners, could truly be applied to all actions to most broadly and effectively implement them. Some actions might not relate to specific stressors, in which case we indicate that with a 'n/a' in the stressor column.

Potential Conservation Actions for Non-forested Wetlands



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Protections for rare calcareous fens are in place. However, early project planning is needed to limit the installation of new wells in aquifers that support them and to site projects such as mining far enough from fens so that groundwater is not impacted. Consider acquiring lands where calcareous fens occur, along with a buffer area, to protect them and their hydrology.

	Preserve and prevent loss or degradation of all types of nonforested wetlands, especially in the Eastern Broadleaf Forest and Prairie Parkland provinces. Focus on protecting wetlands larger than 10 hectares (25 acres) and wetland complexes. Enforce wetland protection regulations (“no-net loss”).
	To reduce the impacts of pipelines, consider policies that increase the fees to cross state land and public waters. Also develop clearer rules around pipelines for human and environmental safety, including for CO2 pipelines.
	Ensure mining operations stay above and do not impact the water table, to protect the hydrology of calcareous fens.
	Ensure mining and quarrying operations and solar and wind development operate with sufficient buffers and provide sufficient mitigation for any damage to wetlands. Increase incentives to avoid all wetlands, including wet meadows.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

Stressor	Action
	Restore large wetland complexes, with attention to the habitat features required by SGCN.
	Control non-native and hybrid cattails to promote a diversity of plants. Minimize the spread of invasive plants in nonforested wetlands (e.g., purple loosestrife, Phragmites). Mitigate impacts and manage potential invasive species dominance in affected areas.
	Reverse ditching and try to restore hydrology to drained peatlands.
	Implement bridges and boardwalks to prevent damage to wetlands from recreational use.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

Stressor	Action
	Avoid creating impoundments that flood wet meadows, except during wetland restoration projects.
	Farmed wetlands can be restored, with resources such as provided through the Natural Resource Conservation Service , or in cooperation with Soil and Water Conservation Districts or Minnesota Board of Water and Soil Resources . This can be particularly helpful when fields are too wet for planting or for crops to come in properly, which is happening more frequently with heavier and more prolonged spring rain events and with projections of even greater spring rainfall in the coming years.
	Increase early coordination with counties and local governmental units in the early pre-planning stages to ensure avoidance of wetlands. Review and comment on plant seed mixes for restoration activities.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions

n/a	Wetlands identification, mapping, and monitoring are ongoing in several projects, with current focus on less well-known wetlands such as vernal pools (see Vernal Pools sub-chapter) and calcareous fens. Also see Chapter 4. Monitoring.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Stressor	Action
	To reduce the impacts of agriculture, mining, and development on wetlands, in particular, calcareous fens, several actions are needed: technical support to identify and study hydrological contexts, information about tiling and groundwater depletions that may affect calcareous fen hydrology, sharing of information with local governments and

	entities including counties and Soil and Water Conservation Districts, and political support to address tough conservation issues.
	Education and outreach to promote soil and hydrological management among agricultural producers and other water consumers to reduce soil and chemical runoff, noting buffer rules, listening to barriers and finding solutions, and taking advantage of financial incentives, including discussions of hydrology associated with drain tiles.
	Increase awareness of importance of peatlands and recognize relative abundance in Minnesota compared to other states, regulate and consider reducing peat mining in the state, increase awareness about anticipated stress from warming and impacts it could have to our states ability to store carbon and changes that may need to be made in the timing of management activities and the amount of area needed to be set-aside.

Case Study: Peatlands Resilience Initiative

Minnesota's [Peatland Resilience Initiative](#), led by the DNR, aims to sustain and restore the state's vast peatland ecosystems, over 7 million acres or roughly 10% of Minnesota's land area, recognizing their critical role in carbon storage, water regulation, and wildlife habitat. Peatlands accumulate partially decomposed plant material under saturated conditions, creating some of the deepest organic soils in the world and storing carbon for millennia. As the second most peat-rich state in the continental U.S. after Alaska, Minnesota's peatlands are vital for both regional biodiversity and global climate mitigation.

Under the initiative, the DNR is advancing a science-based strategy to protect peatlands within Minnesota's Climate Action Framework. Central actions include mapping deep organic soils and wetlands to identify priority areas, developing criteria for conserving or restoring peatland sites, and executing demonstration projects. One such effort evaluates restoration at Winter Road Lake Peatland SNA to quantify changes in greenhouse gas fluxes and mercury export, thereby measuring ecological as well as climate benefits.

Beyond carbon sequestration, the initiative supports additional co-benefits, including improved flood water storage and filtration, reduced mercury and sediment loading in water bodies, and enhanced wildlife habitat for peatland-dependent species. Restoration work focuses on increasing hydration to drained peatlands on state lands, often by raising water levels, to reestablish saturated conditions crucial for peat accumulation and ecosystem resilience.

By integrating mapping and prioritization, cross-agency collaboration, targeted restoration, and rigorous monitoring, the Peatland Resilience Initiative provides a strategic, climate-forward model for peatland conservation. It aligns ecological restoration with climate mitigation goals, establishing Minnesota as a leader in leveraging natural landscapes to address the challenges of a warming world.

References for Non-forested Wetlands

- Apfelbaum, S. I., & Sams, C. E. (1987). Ecology and control of reed canary grass (*Phalaris arundinacea*). *Natural Areas Journal*, 7, 69–74.
- APLIC. (2012). *Reducing Avian Collisions with Power Lines: The State of the Art in 2012* (p. 184). Edison Electric Institute and APLIC.
https://www.aplic.org/uploads/files/11218/Reducing_Avian_Collisions_2012watermarkLR.pdf
- Boers, A. M., Veltman, R. L. D., & Zedler, J. B. (2007). *Typha x glauca* dominance and extended hydroperiod constrain restoration of wetland diversity. *Ecological Engineering*, 29(3), 232–244.
<https://doi.org/10.1016/j.ecoleng.2006.04.011>
- Bonn, A., Allott, T., Evans, M., Joosten, H., & Stoneman, R. (Eds.). (2016). *Peatland restoration and ecosystem services: science, policy, and practice*. Cambridge University Press.
- Finch, D. M., Butler, J. L., Runyon, J. B., Fetting, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing.
https://doi.org/10.1007/978-3-030-45367-1_4
- Haga, C. (2015). *Wetlands 101: What to Know Before You Work*. <https://Bwsr.State.Mn.Us/>.
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Jones, M. W. (2021). *Not so Renewable: Implications for continued peat mining in Minnesota*. University of Minnesota. Retrieved from the University Digital Conservancy, <https://hdl.handle.net/11299/229964>.
- Keirstead, M. E. (1992). Management of Minnesota's Peatlands and Their Economic Uses. In H. E.
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Marohn, K. (2024). Report shows gain in Minnesota's wetland acres. *MPR News*.
- Minnesota Pollution Control Agency. (2025). *Wetland Quality*. <https://www.pca.state.mn.us/air-water-land-climate/wetland-quality> Minnesota. University of Minnesota Press.
- Moriarty, J. J. & Hall, C. D. 2014. Amphibians and Reptiles
- Phillips, M. J. (1997). Forestry best management practices for wetlands in Minnesota. In C. C. Trettin (Ed.), *Northern forested wetlands: Ecology and Management* (1st ed., pp. 403–410). CRC Press.

Prince, H. C. (1997). *Wetlands of the American Midwest: a historical geography of changing attitudes*. The University of Chicago Press.

Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. 2025a. Effects of Climate Change on Midwestern Ecosystems: Eastern North American Temperate Freshwater Marsh, Wet Meadow and Shrubland. Midwest Climate Adaptation Science Center. (64 pages) Available: <https://mwasc.umn.edu/node/821>

Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. 2025b. Effects of Climate Change on Midwestern Ecosystems: North American Bog and Fen. Midwest Climate Adaptation Science Center. (67 pages) Available: <https://mwasc.umn.edu/node/821>

Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. 2025c. Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwasc.umn.edu/node/821>

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.
<https://statesummaries.ncics.org/chapter/mn/>

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434.
<https://doi.org/10.1111/cobi.14434>

The Nature Conservancy. (2025). *Restoring Minnesota's Peatlands for Climate & Water*.
<https://www.nature.org/en-us/about-us/where-we-work/united-states/minnesota/stories-in-minnesota/peatland-restoration-study>

Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report].
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Whigham, D. F. (1999). Ecological issues related to wetland preservation, restoration, creation and assessment. *Science of The Total Environment*, 240(1–3), 31–40. [https://doi.org/10.1016/S0048-9697\(99\)00321-6](https://doi.org/10.1016/S0048-9697(99)00321-6)

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in

Winkler, M. G., & DeWitt, C. B. (1985). Environmental Impacts of Peat Mining in the United States: Documentation for Wetland Conservation. *Environmental Conservation*, 12(4), 317–330.
<https://doi.org/10.1017/S0376892900034433>

Winter, T. A., & Woo, M. K. (1990). Hydrology of lakes and wetlands. In M. G. Wolman & H. C. Riggs (Eds.), *Surface water hydrology. The Geology of North America* (Vol. 1, pp. 159–187). Geological Society of America.

Rivers and Streams

Habitat Description

This chapter focuses on the aquatic habitat within streams and rivers and the shore directly adjacent. Please see the Riparian and Floodplain Forest Chapter for information about forested riverine habitat. Minnesota contains streams and rivers representing a continuum of size and volume ranging from headwaters to joining with tributaries all the way to the great rivers of the Minnesota, Mississippi, Red River of the North, and the St. Croix. Rivers and streams are commonly categorized by size using the Strahler Stream Order classification system (SSO; Strahler 1957). This system ranks streams based on the hierarchy of their tributaries, with each increase in order reflecting an additional level of branching, including small rivers (i.e., ‘streams’; SSO 1-4), medium-large rivers (i.e., ‘rivers’; SSO 5-6), and large rivers (i.e., SSO 7-8). Within each size class, rivers and streams differ considerably in their hydrology, geomorphology, water quality, connectivity, and biological communities—variations that are often shaped by their geographic location or ecoregion. For example, headwater streams are usually characterized by cooler temperatures, fast flow velocities, and shallower pools. In contrast, larger streams and rivers generally exhibit warmer water, slower flow rates, and deeper pools. Smaller streams, owing to their limited watershed size, often maintain higher water quality and experience less exposure to contaminants, excess nutrients, and sediment loads.

The [River Continuum Concept](#) describes a river system as a continuous gradient of physical, chemical, and biological changes from headwater to mouth (Vannote et al. 1980). Along this gradient, longitudinal connectivity, refers to the uninterrupted pathways that link the entire length of a stream. As the water flows downstream, shifts in physical conditions drive corresponding changes in chemical dynamics and biological communities, reflecting the natural integration of upstream and downstream processes.

Streams

Streams in Minnesota exhibit considerable variation in hydrology, geomorphology, temperature, and water quality, shaped largely by regional geology, land use, and climate. For example, prairie streams in western and southern parts of the state tend to have slower velocities, higher turbidity, and greater lateral movement within broad floodplains. Prairie streams are also highly productive and often hotspots of biodiversity (American Prairie Foundation 2025). In contrast, streams in the eastern portion of the state are typically more constrained by rocky substrates and steeper gradients, which result in faster flows and clearer water. Coldwater streams, primarily located in the southeastern karst regions and northeastern forested areas (see Figure 3.16), are strongly influenced by consistent groundwater discharge. These inputs help maintain cool, stable water temperatures throughout the year—conditions that are critical for supporting cold-water biota, including native brook trout (*Salvelinus fontinalis*) and other species of special concern. Many of these streams

are formally designated as trout streams due to their ecological and recreational value (see also [Trout fishing in southeastern Minnesota](#)).

In heavily agricultural regions of Minnesota, especially those shaped by glacial activity, hydrological conditions have been profoundly altered by human intervention. Thousands of miles of artificial surface and subsurface drainage systems—such as ditches and tile drains—have been installed to increase the arability of poorly drained soils. As a result, the density of stream-like channels on the landscape has dramatically increased. In some cases, it can be difficult to distinguish natural streams from constructed ditches due to similar morphology and vegetation, though they often differ in ecological function and stability (Schottler et al., 2014). These modifications accelerate runoff, increase sediment and nutrient loads, and disrupt natural hydrologic regimes, all of which can degrade aquatic habitat and reduce biodiversity.

Smaller streams are particularly vulnerable to anthropogenic stressors and extreme weather events due to their proximity to surrounding landscapes, limited capacity to buffer impacts, and diminished regulatory protection. Because of their smaller size, agricultural or urban development can have immediate and pronounced effects on water quality, sedimentation, and temperature regimes (Freeman et al., 2007). Their small volume and low dilution capacity makes them vulnerable to significant ecological harm from even modest pollutant inputs such as nutrients, pesticides, or heavy metals (Alexander et al., 2007). These streams also respond more dramatically to hydrologic extremes, such as flash floods or droughts, which are increasing in frequency and intensity with climate change (Jackson et al., 2001; Wohl, 2017). Despite their ecological importance, many small streams remain underregulated or unprotected, particularly intermittent or ephemeral channels, leaving them especially susceptible to anthropogenic disturbance and extreme weather events (Leibowitz et al., 2008).

Rivers

As rivers progress from headwaters to larger downstream segments, their geomorphology typically becomes more complex, with wider channels, more developed floodplains, and increasingly intricate channel features such as meanders, oxbows, and backwaters (Leopold, Wolman, & Miller, 1995). These changes in physical structure also correspond with a rise in anthropogenic modifications, particularly the prevalence of dams, levees, and other impoundments that disrupt the longitudinal connectivity of river systems. Dams can fragment habitats, alter sediment and nutrient transport, and affect flow regimes, which in turn disrupt the life cycles of many aquatic organisms, including migratory fish and invertebrates that depend on free-flowing conditions (Poff et al., 1997).

As rivers enlarge and receive flow from multiple tributaries, their pollutant loads tend to increase in both quantity and diversity. Larger rivers collect contaminants from across their watershed, including industrial effluents, municipal wastewater discharges, and nonpoint source runoff, leading to cumulative impacts on water quality (Allan, 2004).

Additionally, the natural increase in flow volume downstream tends to elevate sediment transport capacity, particularly in prairie and agricultural landscapes where erosion rates are often higher. Many large river species have evolved to tolerate or even require these sediment-rich environments. However, excessive sedimentation, especially when combined with nutrient loading, can smother benthic habitats, reduce water clarity, and fuel harmful algal blooms, disrupting aquatic ecosystems (Waters, 1995). In contrast, many species in headwaters are dependent on clear, cool waters with stable substrates and minimal pollution, underscoring the importance of protecting these upstream habitats for overall watershed health.

Large Rivers

Minnesota's large rivers—the Red River, the Minnesota River, the Lower St. Croix River, and the Mississippi River south of St. Anthony Falls—represent some of the state's most ecologically and hydrologically significant waterways. These wide river systems carry large volumes of water and are characterized by slower velocities, meanders and oxbows, numerous islands, and significant backwater systems. These rivers periodically experience significant flooding events that shape the physical landscape and sustain the biological integrity of associated floodplain forests, wetlands, and aquatic habitats (Naiman, Décamps, & McClain, 2005). Notably, several segments of these large rivers widen into lake-like expanses such as Lake Pepin on the Mississippi River and Lake St. Croix on the Lower St. Croix River

These large rivers are also among the most biologically diverse freshwater ecosystems in Minnesota. The Mississippi River, in particular, supports more than 120 species of fish, along with a rich assemblage of mussels, aquatic invertebrates, amphibians, waterbirds, and floodplain plant communities (Houser, 2022). The dynamic hydrology and variety of habitat types, ranging from main channels and side channels to sloughs and marshes, create a mosaic of ecological niches that support high biodiversity. Similarly, the Minnesota and St. Croix rivers serve as important migration corridors and spawning grounds, linking upstream and downstream habitats essential for the life cycles of many aquatic and semi-aquatic species (see also [2005 Minnesota Wildlife Action Plan](#); [St. Croix National Scenic Riverway: Natural Features & Ecosystems](#)).

Habitat Map

To depict Rivers and Stream habitat (see Figure 3.16), we compiled spatial data from several sources: DNR's Native Plant Communities, the National Wetland Inventory for Minnesota, and the Stream Routes with Strahler Stream Order (for more information, see *Habitat Map Methods in Chapter 3. Habitats*). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

River shore (RV)

[RVx32 Sand/Gravel/Cobble River Shore \(PDF\)](#)

[RVx43 Rocky River Shore \(PDF\)](#)

[RVx54 Clay/Mud River Shore \(PDF\)](#)

National Wetlands Inventory

From the [National Wetland Inventory for Minnesota \(NWI\) Layer](#) we included the 'Riverine' Wetland Type.

Stream Routes with Strahler Stream Order

From the Stream Routes with Strahler Stream Order, we classified stream orders 1-4 as small, 5-6 as medium large, and 7-8 as large.




Figure 3.16. A map depicting River and Stream habitat in Minnesota, including the Wetland Type, Riverine, from the National Wetland Inventory for Minnesota and the Stream Order Field from the Stream Routes with Strahler Stream order Layer. River shore habitats were not depicted in this map but are included in the data.

Conservation Overview

The rivers and streams of Minnesota provide important aquatic habitat as a connected system of flowing water. In addition to in-stream habitat, the adjacent shoreline and riparian habitats provide essential interactions for a wide diversity of plant and animal communities.

In the western prairie region, healthy riparian vegetation along prairie streams historically and today consists mostly of sedges, grasses and at times shrubs, such as willows or dogwood (Thornton and Sather 2007; Dodd et al. 2004). Historically, these western riverine landscapes included more trees. They were in locations where prairie fires did not reach them regularly, areas considered a "fire shadow" based on topography and proximity to large rivers or lakes.

Streams in other areas of the state developed diverse riparian vegetative corridors over time, such as overhanging grasses along cold-water streams, water resistant tree species and forbs along expansive riverine floodplains, and forests of hardwoods, pines and shrubs along more topographically dissected stream and river corridors in forested regions of the state.

Backwater areas are another important type of habitat found adjacent to flowing water in small streams to large rivers. These areas have little current and provide important spawning areas for several species of fish. They are also important refuge habitat for many other animal species. For instance, the [Blanding's turtles \(*Emydoidea blandingii*\)](#), [Topeka shiner \(*Notropis topeka*\)](#), [plains topminnow \(*Fundulus sciadicus*\)](#), and [Blanchard's cricket frogs \(*Acris blanchardi*\)](#) are frequently associated with oxbows in the Prairie Coteau. Expansive backwater areas adjacent to large rivers hold a mosaic of land and water that provide important habitat for many species and contributes greatly to flood abatement by regulating high and low flows.

Connectivity of these rivers (the free-flowing condition that maintains pathways that move organisms, energy, and matter throughout the watershed) is an important ecological feature. Historically, unconstrained flowages influenced the distribution of fish populations and likely other taxa as well. Some natural barriers such as St. Anthony Falls located in what is now Minneapolis, limited the range of Minnesota fish species, such as [yellow bass \(*Morone mississippiensis*\)](#), [pirate perch \(*Aphredoderus sayanus*\)](#), pugnose minnow (*Opsopoeodus emiliae*), and [warmouth \(*Lepomis gulosus*\)](#), limiting those species to waters downstream of the falls.

Today, many important features of these river systems, such as connectivity, balanced sediment loads, and water quality, are jeopardized by multiple human activities. In particular, the construction of dams and the placement of culverts has restricted movement of aquatic species that otherwise would migrate significant distances. For example, restricting fish motility has consequences for the distribution of freshwater mussels. Mussels often rely on a specific host fish and when the movement the host fish is blocked it will limit the viability of the mussel population. Barriers that limit the range of the [skipjack herring \(*Alosa chrysochloris*\)](#) as the host fish for the state endangered [elephant-ear \(*Elliptio crassidens*\)](#) and [ebonyshell \(*Reginaia ebenus*\)](#) mussels is an example that has been well documented. (Watters 1994). Dams and hydropower plants also change the natural variability in water flows, causing sedimentation in some areas and dewatering in others, and preventing seasonal flooding, which maintains the health of backwater areas and low flow conditions that allow germination of aquatic plants. Dredging to maintain the 9-foot navigation channel in the Mississippi River for commercial barge traffic presents other challenges including directly destroying mussel beds or burying beds by increased sedimentation or turbidity downstream of the dredging. Water levels in the vicinity of the

dredging may lower and river channels erode; sites to place the removed material must also be found (Pinter et al. 2003). Many of Minnesota's smaller-order streams carry flow laden with heavy amounts of sediments, nutrients, and pollutants from upland activities. As these streams flow together, these contaminants continue downstream. This problem is most pronounced in the agricultural areas where streams and ditches carry pollutants to the Minnesota and Red Rivers. A map of Minnesota's impaired waters illustrates the wide extent that smaller rivers and streams throughout the state have been impacted, especially in southern watersheds ([Minnesota Pollution Control Agency Impaired Waters Viewer 2024](#)). Many aquatic species in greatest conservation need have been extirpated from these two rivers, including the [Higgin's eye mussel \(*Lampsilis higgins*\)](#), while relatively stable populations still occur in the St. Croix and Mississippi Rivers.

The Watershed Health Assessment Framework (WHAF), developed by the DNR, uses five core components to assess the condition of natural systems across the state's watersheds: biology, connectivity, geomorphology, hydrology, and water quality. Together, these components provide a holistic view of watershed function and resilience, emphasizing how natural processes and human actions interact within a watershed context (for details see [The Five Component Framework](#)).

- **Biology** reflects the living systems within a watershed, focusing on the diversity, abundance, and composition of species, particularly aquatic organisms like fish and macroinvertebrates. Biological communities integrate and respond to physical, chemical, and ecological changes over time, making them strong indicators of overall watershed health.
- **Connectivity** examines the degree to which natural systems such as rivers, wetlands, and uplands remain physically and ecologically linked across the landscape. This includes both longitudinal connectivity (upstream to downstream) and lateral connectivity (stream to floodplain), which are essential for species movement, gene flow, sediment transport, and nutrient cycling. Fragmentation by roads, dams, and land development can disrupt these vital connections.
- **Geomorphology** addresses the shape and structure of the landscape, focusing on how sediment moves through stream systems and how stream channels respond to flow and land use changes. Stable stream systems maintain a balance between erosion and deposition. Altered geomorphology can indicate instability, leading to degraded habitat and increased sediment loading.
- **Hydrology** evaluates the timing, magnitude, and duration of water movement through a watershed. It reflects both natural precipitation and groundwater contributions, as well as alterations from impervious surfaces, drainage systems, and water withdrawals. Changes in hydrology can influence flooding, baseflow conditions, and the ecological integrity of aquatic habitats.
- **Water Quality** assesses the chemical, physical, and biological characteristics of water in relation to ecosystem and human health. It includes measures such as nutrient levels, temperature, clarity, and contaminants. Degraded water quality can result from runoff, point source pollution, or landscape disturbances and is a key factor affecting both biodiversity and watershed function.

Species in Greatest Conservation Need

Rivers and streams provide habitat for 133 animal and 11 plant SGCN as primary or secondary habitat (see Table 3.14.) Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants). Examples of selected SGCN are described below.

Table 3.14. Numbers of Species in Greatest Conservation Need associated with Rivers and Streams as either primary or secondary habitat¹.

Species Group	Primary Habitat	Secondary Habitat	Total
Amphibians	3	0	3
Birds	1	0	1
Fish	31	0	31
Mammals	0	9	9
Reptiles	3	0	3
Aquatic Invertebrates			
Bees			1
Beetles (aquatic)			1
Caddisflies (Trichoptera)	26	1	27
Dragonflies and Damselflies (Odonata)			12
Mayflies (Ephemeroptera)			8
Mussels			32
Stoneflies (Plecoptera)			5
Plants	11		11
Total	75	10	144

1. Habitat associations for some insects were not differentiated into primary and secondary habitats and are shown only in the total column. For plants, only a single primary habitat was identified per species.

Rivers and streams provide primary habitat for 31 species, or 86%, of SGCN fish. Fish dominate the vertebrate SGCN in rivers and streams, followed by a number of aquatic invertebrates. All but one of the 31 SGCN mussels are restricted to rivers and 52% of all the caddisfly SGCN inhabit rivers and streams. A more detailed breakdown of what size of stream and river stretches these species utilize can be found in the Fish and in the Aquatic Invertebrate SGCN chapters. The 10 vascular plant SGCN found in these flowing waters represent only 2% of all the SGCN vascular plant flora.

Amphibians

There are three SGCN amphibians who are primarily dependent on streams and rivers in Minnesota.

[Mudpuppies \(*Necturus maculosus*\)](#) are fully aquatic salamanders that typically prefer rocky substrates and structures in medium and large rivers. This is their only habitat and is therefore critical to their survival. Blanchard's frogs are typically found along the waters edge in wetlands, lakes streams, and smaller rivers in Southeastern and Southwestern Minnesota; rivers and streams serve as travel corridors allowing this rare frog to colonize new wetland habitat. Pickerel frogs (*Lithobates palustris*) breed in the backwaters of rivers and streams, and spend summer in clear, cool streams under dense forest canopies in Southeastern Minnesota.

Birds

[Louisiana Waterthrush \(*Parkesia motacilla*\)](#) is closely tied to small streams, or less often stream channels along larger rivers, but only in mature, closed canopy forest (with a strong preference for Mesic Hardwood Forest). Louisiana waterthrushes nest in eroded stream banks and forage on small, aquatic invertebrates. Because they are a small songbird, suitable foraging microhabitat must have a shallow water interface such as small, exposed rocks, woody vegetation (e.g., small logs or dead branches), or in-stream leaf litter. Optimal streams are typically rocky, with rapids and riffles, and have good water-quality that supports the aquatic invertebrates that waterthrushes eat.

The vast majority of SGCN birds associate with rivers and streams are dependent on emergent wetlands, floodplain forests or other riparian habitats. Though not formally listed as using rivers and streams as primary or secondary habitat, many birds do use rivers as habitat. For example, marshes with emergent and submergent vegetation, particularly those in large river backwaters, provide important habitat for several bird SGCN. American bitterns (*Botaurus lentiginosus*), least bitterns (*Botaurus exilis*), and [common gallinules \(*Gallinula galeata*\)](#) require relatively dense emergent vegetation. Black terns (*Chlidonias niger*) build floating nests anchored to emergent vegetation or on piles of rotting plant matter, but require shallow open water for foraging. Birds typical of sedge wetlands, including [Yellow Rail \(*Coturnicops noveboracensis*\)](#), [Wilson's phalarope \(*Phalaropus tricolor*\)](#), and [Nelson's Sparrow \(*Ammospiza nelsoni*\)](#), can nest on relatively broad, floating sedge mats associated with the shorelines of shallow, slow-moving rivers.

Prothonotary warblers (*Protonotaria citrea*) are a floodplain forest specialist, nesting in tree cavities. They have a strong preference for nesting over water, so suitable nest sites are typically in dead trees or snags in forested river backwaters. The [red-shouldered hawks \(*Buteo lineatus*\)](#) breed in mature forests with wetland openings, and may be found in both upland and floodplain forests. Suitable floodplain forest habitat typically occurs along medium to large rivers that have associated wetland openings such as marshes, side channels, or oxbows lakes.

Fish

The [redside dace \(*Clinostomus elongatus*\)](#), a fish species in greatest conservation need, is most abundant in clear, spring-fed, coldwater streams. It typically occurs in pools with moderate current and overhanging vegetation and spawns in riffles or shallow flowing pools. Another fish SGCN, the plains topminnow, is also most common in headwater streams of high water quality but does not require cold water.

The largescale stoneroller(*Camptostoma oligolepis*), [black redhorse \(*Moxostoma duquesni*\)](#), and greater redhorse (*Moxostoma valenciennesi*) are all found in moderate-sized warmwater streams. Largescale stonerollers require clear water with moderate to swift current, often in deep, fast riffles. They require gravel bottoms and are intolerant of siltation, as they feed by overturning small stones; hence the name stoneroller. Black redhorses prefer pools with gravelly to rocky, occasionally sandy and silty, bottoms. They spawn in gravel

and fine rubble runs and riffles in water about 8 - 24 inches (20- 60 centimeters) deep. The greater redhorse also requires clear water with moderate to fast-flowing currents, and clean sand or gravel substrates.

The [least darter \(*Etheostoma microperca*\)](#) occurs in moderate to large warmwater streams, as well as large, deep lakes. It prefers heavily vegetated areas with sluggish flow, immediately downstream of pools in the spawning and growing season, and overwinters in the deep water of pools.

The [crystal darter \(*Crystallaria asprella*\)](#) occurs in large to very large, warmwater rivers. It is usually in water more than 24 inches (60 centimeters) deep with strong current, and along expanses of clean sand and gravel, where it buries itself in sand with only its eyes protruding.

Case Study: Lower Mound Creek Restoration

The [Lower Mound Creek Restoration at Blue Mounds State Park](#) entailed removal of an old dam after it had failed and restoring the stream and adjacent prairie. It has been a successful multi-disciplinary approach to improve [watershed health](#) including biology, hydrology, connectivity, water quality and geomorphology. Species in Greatest Conservation were a key consideration, including Topeka shiner and plains topminnow which have returned to this section. When their requirements are incorporated, similar restorations can benefit other SGCN of the Prairie Coteau that co-occur in streams such as Blanding's turtle, Blanchard's cricket frog, and [pondmussel \(*Sagittunio subrostratus*\)](#). There are many other "stackable" benefits for the ecosystem and for humans. This DNR [video](#) gives an excellent overview from several perspectives.

Mammals

No SGCN mammals are primarily or secondarily listed as using rivers and streams as habitat.

Mussels

Twenty-two of Minnesota's 48 native freshwater mussel species only or primarily occur in large river systems. Many of these mussel species are presently restricted to the lower St. Croix River and the Mississippi River below St. Anthony Falls, where water quality, flow regimes, and/or substrates such as boulders or gravel beds are present in sufficient quality and quantity to allow for their persistence. Freshwater mussel diversity typically declines as streams become smaller due to mussel life history characteristics like host fishes utilized. Only three mussel species are typically strongly associated with small streams. One example is the [creek heelsplitter \(*Lasmigona compressa*\)](#) that is found in headwater to moderate sized, warmwater streams. It is usually found in swift currents with a substrate of sand, fine gravel, and mud, often downstream of riffles in small pools. Most others are generalists, occurring across stream sizes with only a few occurring predominantly in medium sized rivers (Hornbach et al. 2024).

Case Study: Reconnecting Pomme de Terre

The Appleton Milldam, built in 1872, cut off connectivity within the Pomme de Terre river, and the Marsh Lake Dam separated the lower Pomme De Terre river from the Minnesota River. Over a century later, these dams were no longer providing the benefits they once had. A series of dam removal, channel rerouting, and dam modification projects between 1999 and 2018 reconnected these divided waterways. Ten native mussel species have recolonized the historical river channel and abundances of five species appear to be increasing. An earlier restoration project on the Pomme de Terre River in 2001-2005 involved channel stabilization after the Appleton Dam washed out. Since that time, state threatened [elktoe mussels \(*Alasmidonta marginata*\)](#) were documented for the first time upstream of the dam and have expanded their known distribution in the Pomme de Terre River some 30 miles upstream to the next dam near the town of Morris. This further highlights the importance of river connectivity to aquatic animals. [Reconnecting the lower Pomme de Terre River | Minnesota DNR](#)

Reptiles

Many reptiles rely on a combination of upland and stream habitats to complete their life cycles. Blanding's turtles refer calm, shallow wetlands associated with rivers and streams as their primary habitats. Fluvial outwash plains, such as those in Weaver Bottoms along the Mississippi River, provide nesting habitat for Blanding's turtles while the flowing waters are important travel corridors and, where open wetlands are scarce, may be used as overwintering sites. [Wood turtles \(*Glyptemys insculpta*\)](#) hibernate in small to medium fast-moving rivers and streams with sand and gravel substrates, nest on undisturbed sandy banks, and spends much of their time in nearby upland forests. [Smooth softshell turtles \(*Apalone mutica*\)](#) require large, unpolluted rivers with sand substrates for their complete life cycle. Sandbars within or along the river's edge provide essential nesting habitat. Lakes are avoided, as are rivers with muddy substrates.

Climate Profile: Blanding's Turtle

This profile is excerpted from "Effects of Climate Change on Midwestern Ecosystems: Eastern North American Temperate Freshwater Marsh, Wet Meadow and Shrubland" published by the USGS Midwest Climate Adaptation Science Center (Ratcliffe et al. 2025).

Blanding's turtles are long-lived, semiaquatic freshwater turtles found across the northern and eastern United States and southern Canada. In the Midwest, their reliance on a variety of aquatic and upland habitats during their annual activity cycle, combined with low thermal tolerance, make them potentially vulnerable to both the direct and indirect impacts of climate change (Lyons and others, 2023, p. 202). Of all North American freshwater turtles, Blanding's turtles have the lowest critical thermal threshold, and summer maximum temperatures are increasingly likely to exceed this threshold, reducing foraging time (Lyons and others, 2023). Warmer winters and increasingly variable freeze-thaw conditions may also increase mid-winter movements, raising energy demands during a time when food is scarce (Newton and Herman, 2009). Additionally, temperature fluctuations may prompt premature emergence from overwintering sites (Markle and Chow-Fraser, 2014).

Climate change also poses risks to Blanding's turtle reproduction. Higher temperatures during incubation are likely to skew hatchling sex ratios toward females if not compensated by selecting cooler, shaded nest sites or nesting earlier in the season (Gutzke and Packard, 1987). More frequent extreme precipitation events may increase nest failure due to flooding, while drought may hinder nest construction, cause nest collapse, or increase embryo and hatchling mortality through desiccation (Congdon and others, 2000).

Blanding's turtles depend on freshwater ponds, bogs, and marshes for overwintering and foraging and upland, well-drained sites with sparse vegetation for nesting. In the Midwest, these aquatic habitats are likely to face growing impacts from more frequent and intense cycles of drought and deluge. Prolonged summer dry periods may reduce foraging habitat, leading to food shortages and increased movement in search of suitable conditions (Hall and Cuthbert, 2000). These movements during low water periods may increase mortality risk due to road collisions (Ruane and others, 2008) and heightened exposure to predators (Gasbarrini and others, 2021). While some degree of behavioral plasticity, such as earlier nesting or shorter incubation periods, may buffer against these risks, additional adaptive management strategies will likely be necessary, including the installation of turtle road crossings and the implementation of head-starting programs, which raise hatchlings in captivity through their most vulnerable early stages before releasing them into the wild.

Plants

There are a wide variety of plant SGCN dependent on rivers and streams. For example, [small white waterlily \(*Nymphaea leiberghii*\)](#) is found in slow-moving streams, especially those impounded by beaver dams, and is susceptible to being choked out by invasive aquatic plants. [Glade mallow \(*Napaea dioica*\)](#) found in forests on the banks of small to medium-sized streams, is also directly tied to stream health. Encroachment of its stream-side habitat by reed canary grass is a major threat to this species, and natural flooding processes seem to be necessary for seedling establishment. Similarly, [sessile-flowered Yellow Cress \(*Rorippa sessiliflora*\)](#) is a very

rare annual plant dependent on the natural rising and falling of the Minnesota and Mississippi rivers to expose and maintain its very specialized mud flat habitat.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience. Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Straightening and ditching rivers and streams has been done to try to contain the footprint of riverine systems to make space for urban and other forms of development (see also Dams and Water Management and Use). Built infrastructure, such as rip-rap, can impede movement by animals such as turtles, even trapping them, and also can diminish the extent and replenishment of sandy areas used by nesting turtles. Development results in habitat loss and habitat fragmentation along rivers, as well as an increase in roads and trails.



Crop Production

Runoff from cropland can bring sediments, excess nutrients, and chemicals into streams and rivers. Drainage from fields reduces the natural absorption of the landscape for heavy rain events and increases sheet flow. Agricultural practices such as the installation of wells or drain tile can also change the flow of groundwater to streams and are often associated with straightening of the channel (Blann et al. 2009). These changes result in flashy hydrology and incised channels that cause streams to function poorly (Blann et al. 2009). Smaller headwater streams are disproportionately affected by agricultural practices (Wohl 2017).



Livestock Management

Cattle grazing in streams and rivers can cause erosion, reduce habitat quality by trampling and grazing, and degrade water quality (O’Callaghan et al. 2019). Livestock effluent in proximity to streams can introduce *E. coli* into the water.



Mining and Quarrying

Potential mines near and potentially within rivers pose pollution and sedimentation risks to water quality within rivers. There are several proposed mines that could pose a risk to Minnesota rivers, including an in-riverbed sand and gravel mine in the Mississippi River and sulfide-ore mines in northern Minnesota.



Roads, Trails, and Railroads

Roads, trails, and railroads can be a source of erosion and pollution. In addition, they can reduce connectivity with crossing structures such as culverts that are insufficiently sized or due to placement or sedimentation prevent regular access and movement for aquatic species such as fish. Many riverine wildlife species, especially turtles, are often compelled to cross roads, trails, and railroad tracks which interrupt their natural movement paths in the proximity of river corridors. Nesting females are particularly vulnerable during these movements; they are frequently drawn to road edges and adjacent areas for nesting, significantly increasing their risk of fatal encounters with vehicles (Steen et al., 2006). Studies have found that adult females suffer disproportionately from road mortality, leading to skewed sex ratios and diminished reproductive output in turtle populations (Steen et al., 2006; Gibbs & Shriver, 2002). Because female turtles are long-lived and mature slowly, even modest increases in adult mortality can have long-lasting negative effects on population viability (Steen et al., 2006; Piczak, Markle, & ChowFraser, 2019). De-icing salts dispersed on roadways increase the salinization of rivers [Chloride | Minnesota Pollution Control Agency](#)



Shipping Lanes

The maintenance of shipping lanes, including locks and dams, create stressors on the natural movement of sediments, nutrients, and organisms in the Mississippi River. Dredging of the channel creates a constant disturbance that prevents natural channel formation, movement, and sediment transport. Dredging activities can actively remove and prevent freshwater mussels from colonizing an area and impede the presence of dense mussel beds that aid in stabilizing the river bottom, provide habitat for macroinvertebrates and fish, and filter the water (Vaughn 2018). Nutrient cycling is reduced because levees keeping the river in the navigation channel under most flow conditions limits access to floodplain habitats that offer an exchange of terrestrial and aquatic nutrients (Schramm et al. 2015). Impounded sections of river upstream of locks and dams trap sediment and create warmer water temperatures (Poff and Zimmerman 2010). Wakes and turbulence from barges can cause excess sedimentation (Spear et al. 2024).



Hunting and Collecting Animals

Illegal collection of threatened and endangered species, such as wood turtles, is a concern along rivers, particularly as roads, trails, and development increase access to the rivers creating the opportunity for illegal collection to occur.



Timber Harvest

Runoff from timber harvest sites may introduce sediments into waterways including streams and rivers, particularly if riparian buffer areas are not scaled appropriately. Loss of riparian canopy along streams can result in increased stream temperatures. Managing for early-successional habitat like aspen can increase

the presence of beavers in rivers and streams. In northern MN, timber harvest has shifted the forest composition along rivers from later successional, older pine forest to early successional, younger aspen forest in many areas. Downed large diameter trees in rivers are very important for providing habitat for aquatic wildlife. These trees can be lacking if adequate riparian buffers are not implemented.



Fishing

The potential for ingestion of lead fishing tackle ([Getting lead out of fishing tackle | Minnesota Pollution Control Agency](#)), and entanglement in fishing line and lures poses a threat to wildlife in Minnesota rivers and streams.



Recreation

Recreational use on sandbars and beaches can disturb sand-nesting SGCN (i.e., ATVs, picnicking, bonfires, party beaches). These activities can destroy nests and attract predators to nest sites. Increased recreation also increases the chances of illegal collection of listed species.

Boat wakes from recreational boaters have the potential to create large waves that can erode shorelines and re-suspend sediments that increase turbidity (Lorenz et al. 2013, Marr et al. 2022, Spear et al. 2024). Recreational use can contribute to the introduction of invasive species by seeds of problematic non-native plant being inadvertently brought in on footwear or gear including boats.



Fire Management

Historically, fires may have helped create openings in forests and riparian areas along rivers, creating nesting habitat for turtles.



Dams and Water Management

Dams, culverts, bridges and levees impede the free flow and hydrologic connectivity of riverine habitats. Hydrologic connectivity refers to the flow, exchange and pathways that carry water and move organisms, energy and matter throughout the watershed system. These interactions create complex, interdependent processes that vary over time. When these stream processes are interrupted, the impact to aquatic biota is far-reaching. Examples include the complex relationship between fish migrations and the mussel populations that are dependent on the presence of a specific host fish. When fish migrations are limited, mussel dispersal and persistence can be negatively impacted.

Straightening and ditching rivers and streams has been done to try to contain the footprint of riverine systems to make space for development. Around 50% of Minnesota's streams and rivers have been altered from free-flowing systems into ditched, straightened or drained systems. This extreme level of alteration changed these river systems from natural meandering streams with complex habitats into simplified channels that move water quickly off the landscape, removing aquatic habitats, limiting water storage and increasing erosion. (More about altered watercourses)

Large rainfall events cause large flows of water to enter the river and spill out over the riverbanks and into the floodplain. These pulses of water are important for creating sand/gravel points, maintaining riparian habitat,

and terrestrial-aquatic nutrient exchange. When too much of a watershed is developed or managed as young forest, it changes river dynamics. The amount of water running into the river increases. This can cause the riverbed and banks to erode, and banks can become steep enough to disconnect the river from the floodplain. The energy of floodwaters then becomes concentrated within the river's banks, causing sand points and gravel bars to wash away. All of these effects are worsened when stream channels are artificially straightened. These changes can fundamentally change the river and its habitat (Blann et al. 2009).



Invasive Species (Problematic Non-native Species)

Invasive carp (bighead *Hypophthalmichthys nobilis*, silver *H. molitrix*, black *Mylopharyngodon piceus*, and grass carp *Ctenopharyngodon idella*) are a threat to native ecosystems, local economies, and recreation. Minnesota has acted as a sink for the invasive carp populations that are reproducing further downstream, and the movement of invasive carp into Minnesota appears to be dependent on river connectivity (Fritts et al. 2024). When dams are open or are passable, invasive carp move further upstream. While reproduction has not yet been documented in Minnesota, captures and other monitoring methods suggest that invasive carp abundance has increased in Minnesota in recent years. Minnesota DNR has recently updated its statewide Invasive Carp Action Plan (DNR 2024) to identify priority actions that DNR will use to prevent and manage invasive carp.

Sea lamprey are an invasive species that reproduce in fast moving waters, specifically tributaries to Lake Superior. The Great Lakes Fishery Commission and the U.S. Fish and Wildlife Service maintain a sea lamprey control program to reduce impacts on native and desirable fish such as lake trout (GLFC 2025). The program is focused around treating streams where lamprey reproduce to kill larval sea lamprey, and around maintaining strategic barriers to prevent sea lamprey from expanding their spawning grounds. Whether a dam or other structure is acting as a barrier to movement for sea lamprey should be considered before removal.



Problematic Native Species

Reed canary grass (*Phalaris arundinacea*) is a hardy, aggressive, cool-season grass that can form dense, monotypic stands in wetlands. Previously considered an invasive species, recent research suggests that reed canary grass found in Minnesota is predominantly native. It can outcompete other native wetland species, posing challenges for wetland restoration and mitigation efforts. Source: University of Minnesota Extension ([Reed canary grass](#)).



Water-borne Pollution

Nonpoint source pollution is pollution that comes from many diffuse sources (like pollution from many agricultural fields). Nonpoint source pollution includes phosphorus, nitrogen, sediment, bacteria, and other contaminants generally results from land runoff, precipitation, atmospheric deposition, drainage, seepage or hydrologic modification. As runoff flows, it picks up and carries away natural and human-made pollutants, depositing them into lakes, rivers, wetlands, and groundwater, and estuaries. ([EPA NPS Webpage](#)) These nonpoint source pollutants represent the largest combined contribution (an estimated 86%) to the state's water pollution. ([MPCA Water Quality Initiatives](#)) Water quality measured by nutrients and contaminants has been shown to decline with higher agricultural or urban development in the watershed.

Some key impacts include excess algal growth, reduced plant diversity, and reduced diversity in fish and invertebrate communities. ([DNR Water quality non-point sources](#)).

Polyfluoroalkyl Substances (PFAS) and Polychlorinated biphenyls (PCBs) are human made chemicals that are both banned in Minnesota, but persistent in the environment. PFAS may get into rivers from industrial discharge, leaching from landfills into groundwater, and from firefighting foam products. The specific effects of these pollutants on wildlife are poorly known, but their presence is documented in many riverine fish and the Minnesota Department of Health has fish consumption guidelines due to contaminated fish ([Rivers Waterbody Specific Guidelines April 2025](#)) Air-borne Pollutants

Mercury is emitted into the atmosphere and moves long distances with wind and weather and can deposit in aquatic systems like rivers. Mercury pollution is common in aquatic food webs and bioaccumulates, such that species higher on the food chain, like large fish, become contaminated. Fish in many of Minnesota's rivers have concerning levels of mercury ([Rivers Waterbody Specific Guidelines April 2025](#)).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Low water levels due to frequent and prolonged droughts are forecast to impact Minnesota streams as the climate continues to change. Warming summer air temperatures will likely cause many streams to become warmer, especially during times of low water. Impounded stream reaches associated with dams may be disproportionately affected by these warming temperatures ([Climate trends affecting lakes and rivers | Minnesota DNR](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a

corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. Fish spawning habitat can be disturbed by the sedimentation caused by such high flow events and species like turtles may be killed, washed downstream, buried under sand, or experience nest failure ([Climate trends affecting lakes and rivers | Minnesota DNR](#)). The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al. 2022). For more information and resources for climate-adapted management strategies, see the Climate subchapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Examples of conservation actions are grouped below under these five strategies and tagged with icons for the stressor(s) that they address. Some actions will combine multiple strategies, in which case we present it under the one it fits best. Also note that some strategies, such as Strategy 3, collaborating with partners, could truly be applied to all actions to most broadly and effectively implement them. Some actions might not relate to specific stressors, in which case we indicate that with a ‘n/a’ in the stressor column.

Potential Conservation Actions for Rivers and Streams



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Look for opportunities to restore or protect (acquisition, conservation easements, use designations) key river or stream reaches. Consider the following criteria in prioritizing project areas: rivers and streams that lack or have limited barriers to aquatic organism passage or with high potential for restoration; Rivers/streams with significant groundwater inputs; Rivers/streams with intact floodplains, meanders, and potentially associated oxbows; Rivers and streams with documented SGCN are necessary for SGCN population connectivity, genetic exchange, and climate resiliency.
	Since many animals require both upland and stream habitats for their life cycles, it is essential to keep these physical habitats connected. Protect habitats along river corridors to allow movement of wildlife and facilitate movement of nutrients and energy between the stream and surrounding landscape. Management options include near shore riparian corridor conservation and habitat restoration to improve longitudinal as well as lateral connectivity.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

Stressor	Action
	Support the removal of dams or replacing or upgrading with structures that allow better passage of aquatic organisms such as fish, turtles, amphibians, and mussels will assist in the health and connectivity of free-flowing rivers.
	To protect and enhance water quality parameters such as sediment loads and chemical pollution, options include improved stormwater systems, soil erosion control techniques, maintaining vegetated buffers, and point - nonpoint source pollution abatement.
	Culvert replacement with larger capacities and naturalized bottoms will maximize aquatic habitat connectivity and can hold up with resilience under future climatic conditions that entail heavier spring rain events. Where indicated, include other tested means to improve wildlife connectivity such as passage benches, J-fencing, wing walls, and other measures (Hernick et al. 2019, DNR/MNDOT 2014).
	Restore sinuosity to rivers and streams that have been previously straightened with considerations for SGCN in both restoration design and during the construction process to avoid or minimize adverse effects. Advocate for maintenance of natural flow regimes.
	Map drain tile and wells to better understand and document hydrological changes in groundwater levels, flow, and inputs to streams. Consider new agricultural practices

	such as irrigating from below with agri-drains installed on drain tiles to reduce hydrological impacts on streams.
	Develop and implement better stream management and restoration designs that will be resilient under the higher flows from heavier and more prolonged rainfall events as well as maintain flow during late summer droughts.
	Implement riparian restoration that increases shade on streams to ameliorate the effects of ambient temperature increases.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

Stressor	Action
	Watershed scale restorations involving multiple landowners are complex and expensive but necessary.
	Incentive programs, technical assistance, and outreach to livestock producers to reduce impacts of cattle in stream and river corridors, such as providing alternative water access and partnering on riparian habitat restoration. Successful programs will entail long-term relationships among farmers and cattle producers, increased technical guidance at the community level, and a support network that goes beyond governmental participants.
	Best management practices, mitigation, and cross-training and increased collaboration within and among agencies can normalize expectations of aquatic habitat connectivity and reduce the impacts of roads and road crossings on riverine water flow and quality.

(No actions were articulated for Strategy 4.)



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCM

Stressor	Action
	Outreach, education, and sharing success stories about dam removals or upgrades can help overcome the fear of change and the sense that the dams cannot be removed, upgraded, or enhanced.
	Provide technical assistance and incentives to support best management practices and the maintenance of native vegetation in riparian areas.
	Education and outreach campaigns encouraging fewer chemical inputs to yards and lawns and other municipal practices to minimize pollutants are warranted.

	Incentivize agricultural practices that create and maintain buffers around streams, reduce erosion and runoff, and increase soil organic carbon to reduce sedimentation into streams.
	Promote best management practices for timber management near riparian zones, assuring sufficient buffer areas and that woody debris is left intact in riparian and in-stream locations. Contribute to MFRC Forest Management Guideline revision processes by providing up-to-date literature to support science-based guideline revisions. Consider prioritizing forests near rivers and streams as places to manage for older forest and older successional forest.
	Develop and share educational materials to alert recreational users of sandbars and beaches when there may be sand-nesting SGCN present, such as shorebirds and turtles.

References for Rivers and Streams

Alexander, R. B., Boyer, E. W., Smith, R. A., Schwarz, G. E., & Moore, R. B. (2007). The Role of Headwater Streams in Downstream Water Quality. *JAWRA Journal of the American Water Resources Association*, 43(1), 41–59. <https://doi.org/10.1111/j.1752-1688.2007.00005.x>

Allan, J. D. (2004). Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems. *Annual Review of Ecology, Evolution, and Systematics*, 35(1), 257–284. <https://doi.org/10.1146/annurev.ecolsys.35.120202.110122>

American Prairie Foundation. (2025). *Streams and Wetlands*. <https://americanprairie.org/project/streams-and-wetlands/>

AvERY, Har. W., van Loben Sels, R., & Tinkle, D. (2000). Nesting ecology and embryo mortality: implications for hatchling success and demography of Blanding's turtles (*Emydoidea blandingii*). *Chelonian Conservation and Biology*, 3(4), 569–579.

Blann, K. L., Anderson, J. L., Sands, G. R., & Vondracek, B. (2009). Effects of Agricultural Drainage on Aquatic Ecosystems: A Review. *Critical Reviews in Environmental Science and Technology*, 39(11), 909–1001. <https://doi.org/10.1080/10643380801977966>

Dodds, W. K., Gido, K., Whiles, M. R., Fritz, K. M., & Matthews, W. J. (2004). Life on the Edge: The Ecology of Great Plains Prairie Streams. *BioScience*, 54(3), 205. [https://doi.org/10.1641/0006-3568\(2004\)054\[0205:LOTETE\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0205:LOTETE]2.0.CO;2)

Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4

Freeman, M. C., Pringle, C. M., & Jackson, C. R. (2007). Hydrologic Connectivity and the Contribution of Stream Headwaters to Ecological Integrity at Regional Scales¹. *JAWRA Journal of the American Water Resources Association*, 43(1), 5–14. <https://doi.org/10.1111/j.1752-1688.2007.00002.x>

- Fritts, M., Gibson-Reinemer, D., Appel, D., Lieder, K., Henderson, C., Milde, A., Brey, M., Lamer, J., Turney, D., & Witzel, Z. (2024). Flooding and dam operations facilitate rapid upstream migrations of native and invasive fish species on a regulated large river. *Scientific Reports*, 14(1), 20609.
- Gasbarrini, D., Lesbarrères, D., Sheppard, A., & Litzgus, J. (2021). An enigmatic mass mortality event of Blanding's Turtles (*Emydoidea blandingii*) in a protected area. *Canadian Journal of Zoology*, 99(6), 470–479.
- Gibbs, J. P., & Shriver, W. G. (2002). Estimating the effects of road mortality on turtle populations. *Conservation Biology*, 16(6), 1647–1652.
- Gutzke, W. H. N., & Packard, G. C. (1987). The Influence of Temperature on Eggs and Hatchlings of Blanding's Turtles, *Emydoidea blandingii*. *Journal of Herpetology*, 21(2), 161. <https://doi.org/10.2307/1564476>
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Hernick, M., Lenhart, C., Kozarek, J., & Nieber, J. (2019). *Minnesota guide for stream connectivity and aquatic organism passage through culverts*.
- Holland, J. S. (2018). *Deadly Crossings*. National Wildlife Federation. <https://www.nwf.org/Magazines/National-Wildlife/2018/June-July/Conservation/Turtle-Crossings>
- Hornbach, D. J., Sietman, B. E., & William Bouchard, R. (2024). The relationship between stream size and life-history traits in freshwater mussels: an examination of the Host-Habitat Continuum Concept. *Hydrobiologia*, 851(18), 4419–4437. <https://doi.org/10.1007/s10750-024-05595-2>
- Houser, J. N., ed. (2022). *Ecological status and trends of the Upper Mississippi and Illinois Rivers (ver. 1.1, July 2022): U.S. Geological Survey Open-File Report* [Open-File Report]. U.S. Geological Survey. <https://doi.org/10.3133/ofr20221039>
- Jackson, C. Rhett., Sturm, C. A., & Ward, J. M. (2001). TIMBER HARVEST IMPACTS ON SMALL HEADWATER STREAM CHANNELS IN THE COAST RANGES OF WASHINGTON. *JAWRA Journal of the American Water Resources Association*, 37(6), 1533–1549. <https://doi.org/10.1111/j.1752-1688.2001.tb03658.x>
- Leibowitz, S. G., Wigington, P. J., Schofield, K. A., Alexander, L. C., Vanderhoof, M. K., & Golden, H. E. (2018). Connectivity of Streams and Wetlands to Downstream Waters: An Integrated Systems Framework. *JAWRA Journal of the American Water Resources Association*, 54(2), 298–322. <https://doi.org/10.1111/1752-1688.12631>
- Leopold, L. B., Wolman, M. G., & Miller, J. P. (1995). *Fluvial processes in geomorphology* (Dover ed). Dover Publications.
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>

Lorenz, S., Gabel, F., Dobra, N., & Pusch, M. T. (2013). Modelling the effects of recreational boating on self-purification activity provided by bivalve mollusks in a lowland river. *Freshwater Science*, 32(1), 82–93.

Marr, J., Riesgraf, A., Herb, W., Lueker, M., Kozarek, J., & Hill, K. (2022). *A Field Study of Maximum Wave Height, Total Wave Energy, and Maximum Wave Power Produced by Four Recreational Boats on a Freshwater Lake*. <https://hdl.handle.net/11299/226190>

Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). (2025). *Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state's greenhouse gas emissions contributing to climate change* [Report to Legislature]. Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>

Naiman, R. J., Décamps, H., & McClain, M. E. (2005). *Riparia: ecology, conservation, and management of streamside communities*. Elsevier Academic.

National Park Service. (2023). *St. Croix National Scenic Riverway: Natural Features & Ecosystems*. <https://www.nps.gov/sacn/learn/nature/naturalfeaturesandecosystems.htm>

Newton, E. J., & Herman, T. B. (2009). Habitat, movements, and behaviour of overwintering Blanding's turtles (*Emydoidea blandingii*) in Nova Scotia. *Canadian Journal of Zoology*, 87(4), 299–309. <https://doi.org/10.1139/Z09-014>

O'Callaghan, P., Kelly-Quinn, M., Jennings, E., Antunes, P., O'Sullivan, M., Fenton, O., & hUallacháin, D. Ó. (2019). The Environmental Impact of Cattle Access to Watercourses: A Review. *Journal of Environmental Quality*, 48(2), 340–351. <https://doi.org/10.2134/jeq2018.04.0167>

Open-File Report. (2023). [Open-File Report].

Piczak, M. L., Markle, C. E., & Chow-Fraser, P. (2019a). Decades of Road Mortality Cause Severe Decline in a Common Snapping Turtle (*Chelydra serpentina*) Population from an Urbanized Wetland. *Chelonian Conservation and Biology*, 18(2), 231. <https://doi.org/10.2744/CCB-1345.1>

Piczak, M. L., Markle, C. E., & Chow-Fraser, P. (2019b). Decades of Road Mortality Cause Severe Decline in a Common Snapping Turtle (*Chelydra serpentina*) Population from an Urbanized Wetland. *Chelonian Conservation and Biology*, 18(2), 231. <https://doi.org/10.2744/CCB-1345.1>

Pinter, N., Miller, K., Wlosinski, J. H., & Van Der Ploeg, R. R. (2004). Recurrent shoaling and channel dredging, Middle and Upper Mississippi River, USA. *Journal of Hydrology*, 290(3–4), 275–296. <https://doi.org/10.1016/j.jhydrol.2003.12.021>

Poff, N. L., Allan, J. D., Bain, M. B., Karr, J. R., Prestegard, K. L., Richter, B. D., Sparks, R. E., & Stromberg, J. C. (1997). The Natural Flow Regime. *BioScience*, 47(11), 769–784. <https://doi.org/10.2307/1313099>

Poff, N. L., & Zimmerman, J. K. (2010). Ecological responses to altered flow regimes: a literature review to inform the science and management of environmental flows. *Freshwater Biology*, 55(1), 194–205.

- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. P., & LeDee, O. E. (2025). *Effects of climate change on midwestern ecosystems: Temperate flooded and swamp forest* (p. 110pp). Midwest Climate Adaptation Science Center. Available: <https://mwccasc.umn.edu/node/821>
- Ruane, S., Dinkelacker, S. A., & Iverson, J. B. (2008). Demographic and reproductive traits of Blanding's turtles, *Emydoidea blandingii*, at the western edge of the species' range. *Copeia*, 2008(4), 771–779.
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Rupasinghe, P. A., & Chow-Fraser, P. (2024). Change detection of wetland vegetation under contrasting water-level scenarios in coastal marshes of eastern Georgian Bay. *Landscape Ecology*, 39(3), 44. <https://doi.org/10.1007/s10980-024-01829-9>
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>
- Schottler, S. P., Ulrich, J., Belmont, P., Moore, R., Lauer, J. W., Engstrom, D. R., & Almendinger, J. E. (2014). Twentieth century agricultural drainage creates more erosive rivers. *Hydrological Processes*, 28(4), 1951–1961. <https://doi.org/10.1002/hyp.9738>
- Schramm, H. L., Richardson, W. B., & Knights, B. C. (2015). Managing the Mississippi River Floodplain: Achieving Ecological Benefits Requires More Than Hydrological Connection to the River. In P. F. Hudson & H. Middelkoop (Eds.), *Geomorphic Approaches to Integrated Floodplain Management of Lowland Fluvial Systems in North America and Europe* (pp. 171–201). Springer New York. https://doi.org/10.1007/978-1-4939-2380-9_8
- Spear, M. J., Harris, B. S., Bookout, T. A., Ickes, B. S., Jankowski, K. J., Solomon, L. E., Maxson, K. A., Harris, A. L. W., Mathis, A. T., & Schaick, S. J. (2024). Reduction of large vessel traffic improves water quality and alters fish habitat-use throughout a large river. *Science of the Total Environment*, 946, 172705.
- Steen, D. A., Aresco, M. J., Beilke, S. G., Compton, B. W., Condon, E. P., Kenneth Dodd, C., Forrester, H., Gibbons, J. W., Greene, J. L., Johnson, G., Langen, T. A., Oldham, M. J., Oxier, D. N., Saumure, R. A., Schueler, F. W., Sleeman, J. M., Smith, L. L., Tucker, J. K., & Gibbs, J. P. (2006). Relative vulnerability of female turtles to road mortality. *Animal Conservation*, 9(3), 269–273. <https://doi.org/10.1111/j.1469-1795.2006.00032.x>
- Strahler, A. N. (1957). Quantitative analysis of watershed geomorphology. *Eos, Transactions American Geophysical Union*, 38(6), 913–920. <https://doi.org/10.1029/TR038i006p00913>
- Thornton, R., & Sather, N. (2007). *Analysis and narrative: a complementary approach to understanding the 19th century landscape and forests of Minnesota's Prairie Coteau*. Symposium of the Minnesota Native Plant Society.
- Vannote, R. L., Minshall, G. W., Cummins, K. W., Sedell, J. R., & Cushing, C. E. (1980). The River Continuum Concept. *Canadian Journal of Fisheries and Aquatic Sciences*, 37(1), 130–137. <https://doi.org/10.1139/f80-017>

Vaughn, C. C. (2018). Ecosystem services provided by freshwater mussels. *Hydrobiologia*, 810(1), 15–27.
<https://doi.org/10.1007/s10750-017-3139-x>

Waters, T. F. (1995). *Sediment in streams: sources, biological effects, and control*. American Fisheries Society.

Watters, G. (n.d.). *An annotated bibliography of the reproduction and propagation of the Unionoidea (Primarily of North America)* (No. 1; Miscellaneous Contributions, p. 158). Ohio Biological Surveys.

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report].
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Wills, M. (2021). *Road density threatens turtle populations*. J Stor Daily. <https://daily.jstor.org/road-density-threatens-turtle-populations/>

Wohl, E. (2017). The significance of small streams. *Frontiers of Earth Science*, 11(3), 447–456.
<https://doi.org/10.1007/s11707-017-0647-y>

Lakes

Habitat Description

Lakes are iconic and ecologically important in Minnesota, numbering over 11,800 at least 10 acres in size. Glacial activity formed most of Minnesota lakes (Schwartz and Thiel 1954) with the physical characteristics of each individual lake differing based on factors including the local geology and hydrology at each site. Lakes vary from shallow to 280 feet deep for Lake Saganaga—Minnesota’s deepest inland natural lake. Lake Superior is 1,290 feet deep. Minnesota has several other very large lakes: Lake of the Woods (950,400 ac, shared with Canada), Upper and Lower Red Lake (288,800 ac), Mille Lacs (132,516 ac), and Leech Lake (111,527 ac) ([Lakes, Rivers, and Wetlands Facts](#)). There are many ways to classify lakes but for simplicity, here we categorize lakes based on depth.

Deep Lakes

Here, we use ‘deep lakes’ to refer to water bodies greater than 15 feet (5 meters) deep. Deep lakes vary in productivity, ranging from low productivity northeastern Minnesota lakes with clear water, cold temperatures, and minimal aquatic vegetation to high productivity southern Minnesota lakes with high nutrient concentrations, low water clarity, and lesser amounts of rooted aquatic vegetation or dominated by filamentous or planktonic algae. A combination of factors related to climate, geology, hydrology, land use, and land cover results in the general trend of lakes increasing in water clarity and decreasing in productivity and alkalinity from southwestern to northeastern Minnesota. Communities of fish and other aquatic organisms also vary along this distribution of lake types in Minnesota (Schupp 1992).

The close association between aquatic plant communities and fish communities is well recognized. Aquatic plants provide food and oxygen, shelter and spawning areas, and improved water quality for fish. Plants also provide excellent food and habitat for zooplankton, insects, waterfowl, and other wildlife. Aquatic vegetation also anchors sediments, maintaining water clarity. Radomski et al., (2019) provides a short synthesis of the scientific studies and management of aquatic plants to highlight lake vegetation management actions that may benefit fisheries.

Deep, thermally stratified lakes with abundant dissolved oxygen throughout the water column are especially threatened in Minnesota. These lakes are commonly referred to as Coldwater Lakes. They support cold-adapted fish species, including lake trout (*Salvelinus namaycush*), lake whitefish (*Coregonus clupeaformis*), cisco (*C. artedii*), and burbot (*Lota lota*). In Minnesota, there are about 1000 lakes that support these cold-water fish communities. In northeast Minnesota, these lakes also support a variety of isoetids plant species, which are submerged aquatic plants that are slow-growing, evergreen, and forming basal rosettes, such as quillworts (*Isocetes* spp.), which are unique to these low nutrient cold-water lakes. Eutrophication (excessive nutrients) and climate change is reducing the number of lakes that support these two communities.

Shallow Lakes

There are an estimated 4000 shallow lakes in Minnesota, permanent or semi-permanent water bodies less than 15 feet (5 meters) deep and over 50 acres (20 hectares) in size (see DNR’s [Shallow Lakes Program](#)). These

lakes are typically dominated by wetland habitat (see also Non-forested Wetlands Chapter). While shallow lakes can support fish populations, often low levels of dissolved oxygen in winter limit their numbers.

Shallow lakes may have abundant aquatic plant growth due to sunlight reaching much of the lake bottom in these basins. Stands of emergent and floating-leaved aquatic plants such as cattails (*Typha* spp.), bulrush (*Schoenoplectus* spp.), bur-reeds (*Sparganium* spp.), water lily (Nymphaeaceae), as well as submerged plants, such as stoneworts (Characeae), coontail (*Ceratophyllum* spp.) and pondweeds (*Potamogeton* spp.), may be present throughout the entire basin. These plants provide excellent food and habitat for zooplankton, insects, fish, waterfowl, and other wildlife. Aquatic vegetation also anchors sediments, maintaining water clarity.

Shallow lakes are dynamic ecosystems that often express considerable ecological variability. Nutrients in shallow lakes, unlike in deeper lakes, are regularly mixing. Shallow lakes do not thermally stratify for long periods of time so there is a constant exchange of nutrients between the sediments and water column. Wave action also can often reach the bottom of the lake, suspending sediments in the water. Droughts lower water levels in shallow lakes resulting in exposed sediments. This exposure stimulates beneficial aquatic plant growth. Low water levels can produce conditions for fish winterkills that can decrease or eliminate fish populations. Such low water cycles have been impacted by higher levels of precipitation and watershed alterations including draining of wetlands into larger water bodies. Persistent and high water levels restrict plant growth and reduce water quality, allowing significant algal growth. Milder winters also favor fish such as bullheads (*Ameiurus* spp.) and common carp (*Cyprinus carpio*), which in shallow lakes can reduce water quality and diminish the abundance of rooted aquatic plants.

Some shallow lakes can support wild rice (*Zizania palustris*, also known as *manoomin*), designated as a SGCN plant in this Plan due to cultural significance to many of the Tribal Nations in Minnesota. Manoomin is harvested traditionally each fall. Minnesota supports more acres of wild rice than any other state in the country. Wild rice has been historically documented in 45 of Minnesota's 87 counties and is thought to have been even more broadly distributed prior to European settlement (see [Wild Rice Management](#)).

Habitat Map

To depict Lake and Lakeshore habitat (see Figure 3.17), we compiled spatial data from several sources: Shallow Lakes Identified by DNR Wildlife, Lake Basin Morphology, the National Wetland Inventory, and the DNR Native Plan Community Layer (for more information, see Habitat Map Methods in Chapter 3. Habitats). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Lakeshore (LK)

[LK_i32 Inland Lake Sand/Gravel/Cobble Shore \(PDF\)](#)

[LK_i43 Inland Lake Rocky Shore \(PDF\)](#)

[LK_i54 Inland Lake Clay/Mud Shore \(PDF\)](#)

[LK_u32 Lake Superior Sand/Gravel/Cobble Shore \(PDF\)](#)

[LKu43 Lake Superior Rocky Shore \(PDF\)](#)

National Wetlands Inventory

From the [National Wetland Inventory for Minnesota \(NWI\) Layer](#) we included only the Non-Vegetated Aquatic Community type from the Simplified Plant Community Classification attribute field.

Lake Basin Morphology

We included all lakes identified by the [Lake Basin Morphology layer](#).

Shallow Lakes Identified by DNR Wildlife

We included all Lakes identified in the [Shallow Lakes Identified by DNR Wildlife layer](#).




Figure 3.17. A map depicting Lake habitat in Minnesota including Non-Vegetated Aquatic Communities from the National Wetland Inventory for Minnesota as well as all lakes from the Shallow Lakes Identified by DNR Wildlife and the Lake Basin Morphology layers. Lakeshore habitat is not depicted in the map but is included in the data.

Conservation Overview

Minnesota's lakes receive considerable recreation and development, and are faced with threats from invasive species, shoreline alteration, altered hydrology, eutrophication and other pollution, and climate change.

Recreationalists transport aquatic species to lakes, such as on their boats, and the arriving species can alter lake habitat. Several invasive aquatic species are now present in many Minnesota lakes. The invasive aquatic plants [Eurasian watermilfoil \(*Myriophyllum spicatum*\)](#) and [curly-leaf pondweed \(*Potamogeton crispus*\)](#) have been present in Minnesota lakes for decades. Eurasian watermilfoil can grow to the water surface and compete with native plants. Several invasive aquatic animals also occur in many Minnesota lakes, including [common carp](#) and [zebra mussels \(*Dreissena polymorpha*\)](#). Common carp is widely distributed in central and southern Minnesota, and this fish severely impacts shallow lakes as their feeding habits uproot aquatic plants and suspends fine sediments into the water. Zebra mussels filter the water, which reduces food for larval fish and other animals and increases water clarity. So, zebra mussels can change the food web and reduce the carrying capacity for popular fish such as walleye (*Sander vitreus*). [Infested Waters List | Minnesota DNR](#)

Minnesota lakes face serious changes due to human alteration of land, water, and air. Shorelines are altered by lakeshore property owners. Minnesota has currently lost 40 to 50% of its natural shorelines, and the rate of loss is estimated to be 1 to 2% per decade (Minnesota Natural Shoreline Partnership 2023). As a lake's watershed becomes more developed, nutrient levels increase and water quality decreases due to pollution runoff, poor stormwater management, and shoreline phosphorus pollution from shoreland sewer systems and lawns down to the lake. Nutrients reaching the lake from this pollution result in eutrophication where lakes receive surplus nutrients which can result in excessive plant and algal growth. Agricultural lands shed large amounts of sediment and pollution and is one of the leading causes of lake water quality impairments. These lands are also drained with ditches and tiling, and thousands of acres of Minnesota's shallow lakes have been ditched and drained.

Recreation can be problematic in lake ecosystems. Boating in or near shallow water suspends lake sediment, destroys aquatic plants, and erodes shoreline, especially when shoreline vegetation is removed and replaced with lawn down to the lake. Boat strikes kill wildlife and species like turtles and loon chicks are particularly vulnerable. Fish stocking has altered the native aquatic faunal communities, and homogenized fish assemblages in Minnesota lakes (Radomski and Goeman 1996). Aquatic vegetation can be impacted by docks and shallow water disturbance (Radomski et al., 2010). Aquatic vegetation loss for northcentral Minnesota lake was estimated at 15% (Radomski 2006). Lead, a toxic element, is extensively used in fishing tackle. In popular angling lakes, tackle loss results in many metric tons of lead across lake bottoms annually (Radomski et al., 2006). Waterbirds ingest this lead, which is often lethal. Additionally, other tackle and discarded fishing line can entrap wildlife.

Air pollution is degrading Minnesota lakes. This pollution includes a large range of chemicals, from mercury via primarily from coal burning, pesticides via drift from farm and residential lands, microplastics and nanoplastics via wind and waves moving broken down plastic waste fragments, to greenhouse gases via combustion of coal and gas. Mercury bioaccumulates and biomagnifies through lake food webs, which affects the wildlife and human health as it is a potent neurotoxicant [Mercury | Minnesota Pollution Control Agency](#). Fish from lakes in the northeast generally have the higher mercury concentrations (Radomski et al., 2024).

Of particular concern, due to the magnitude of change, is climate change from burning fossil fuels. In Minnesota, lakes now have warmer water and shorter ice cover seasons. Fish communities are changing, and those changes and other widespread environment changes will accelerate. [Climate trends affecting lakes and rivers | Minnesota DNR](#)

Species in Greatest Conservation Need

Lakes provide habitat for 93 animal and 36 plant SGCN as primary or secondary habitat (see Table 3.15.) Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants). Examples of selected SGCN are described below.

Table 3.15. Numbers of Species in Greatest Conservation Need associated with lakes as either primary or secondary habitat¹.

Species Groups	Primary Habitat	Secondary Habitat	Total
Amphibians	2	1	3
Birds	18	6	24
Fish	8	3	11
Mammals	0	9	9
Reptiles	1	0	1
Aquatic Invertebrates			40
Beetles (aquatic)			1
Caddisflies	18	3	21
Dragonflies and Damselflies			13
Mayflies			2
Mussels	1		1
Snails (aquatic)	1		1
Terrestrial Invertebrates			5
Bees			4
Spiders			1
Plants	36		36
Total	84	23	129

1. Habitat associations for most insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Amphibians

The [mudpuppy \(*Necturus maculosus*\)](#) is Minnesota's only fully aquatic salamander and is restricted to medium or large rivers and lakes (primarily in deep lakes but secondarily with shallow lakes). Lakes may be used more often in western Minnesota than farther east. Submerged rocks or logs provide cover and important breeding habitat. Eastern newt (*Notophthalmus viridescens*) occupies shallow lakes and will be threatened by a pathogenic fungus if or when it arrives to North America. [Blanchard's cricket frog \(*Acris blanchardi*\)](#) may utilize lakes, particularly when they are fed, or are near, rivers and streams. Large, deep lakes, even with predatory fish, may be used but generally only if they are associated with a river or stream.

Birds

Of 24 SGCN bird species using lakes, 8 are primarily limited to shallow lakes such as the American bittern (*Botaurus lentiginosus*), [common gallinule \(*Gallinula galeata*\)](#) and [Wilson's phalarope \(*Phalaropus tricolor*\)](#). Together, the 22 bird SGCN utilizing lake habitats represent 28% of all bird SGCN.

Both shallow and deep lakes are well recognized for their importance to waterfowl and other waterbirds. Many of these species will use open water during migration, as well as for foraging and loafing during the breeding season. However, most SGCN waterbirds require emergent vegetation, or in some cases islands or other shoreline habitat, for nesting.

Waterfowl species, such as the American black duck (*Asas rubripes*), canvasback (*Aythya valisineria*), and northern pintail (*Anas acuta*), nest in emergent marsh or grassland vegetation adjacent to lake shores or open water in marshes. Buffleheads (*Bucephala albeola*), restricted to far northwestern Minnesota during the breeding season, are cavity-nesters. American bitterns, least bitterns (*Ixobrychus exilis*), and common gallinules require relatively dense emergent vegetation, so only nest on portions of shallow lakes with suitable marsh vegetation. Four SGCN grebe species (Horned (*Podiceps auritus*), red-necked (*Podiceps grisegena*), eared (*Podiceps nigricollis*), and western (*Aechmophorus occidentalis*)), plus [Franklin's gulls \(*Leucophaeus pipixcan*\)](#), black terns (*Chlidonias niger*), and [Forster's terns \(*Sterna forsteri*\)](#), build floating nests anchored to emergent vegetation, or on platforms or piles of vegetation. Franklin's gulls are commonly observed when migrating and during the breeding season, but only nest in very large colonies on one or two large, marshy lakes in northwestern Minnesota.

[American White Pelicans \(*Pelecanus erythrorhynchos*\)](#) nest on higher portions of shore habitats, typically on treeless islands, but feed or loaf on a wide variety of lakes and marshes. Black-crowned Night Herons (*Nycticorax nycticorax*) are a colonial-nesting species that forages along shorelines and in marshes. In Minnesota, they usually nest either in trees on islands, or (less commonly) on floating platforms of marsh vegetation.

During migration, many shorebird species use lake shoreline communities as resting and feeding sites. Two SGCN shorebirds do not nest in Minnesota but solely occur in the state as migrants. The Red Knot (*Calidris canutus*) is a relatively rare migrant, most often found on sand beaches of Lake Superior (particularly those on Minnesota Point). They can also be found, very rarely, on beaches or other types of shorelines away from Lake Superior. Buff-breasted Sandpipers (*Tryngites subruficollis*) are a migrant most often found in open uplands with very short or sparse vegetation. When they are in areas associated with lake shores they do not occur at the water-shoreline interface (wave zone), but may use higher, drier portions of shore habitats.

The Wilson's phalarope is a breeding shorebird species that nests in Minnesota, preferring sedge or grass dominated wetlands such as rich fen and wet prairie. Suitable nesting habitat includes relatively wide, floating sedge mats associated with the shorelines of lakes and rivers. Although Wilson's phalaropes are a species of sedge wetlands, breeding sites can border the shores of large, deep lakes (e.g., Lake-of-the-Woods and Leech Lake). Phalaropes are more aquatic than most shorebirds, frequently feeding while swimming in open water. However, they usually stay in relatively shallow water close to shore, particularly during the nesting season.

[Piping plovers \(*Charadrius melodus*\)](#) now nest sporadically in only one location in Minnesota, along sand beaches on the south shores of Lake of the Woods, although they once nested along the dunes of Lake Superior. Critical habitat for this species has been designated by the U.S. Fish and Wildlife Service on Pine and Curry islands and also in the Duluth Harbor area. [Common terns \(*Sterna hirundo*\)](#) nest in a handful of colonies on islands with rocky or sandy shorelines in large lakes such as Leech Lake, Mille Lacs Lake, Lake of the Woods, and Lake Superior (Duluth Harbor).

Fish

Minnesota lake dwelling fish species of greatest conservation need include several species. Pugnose shiner (*Notropis anogenus*), [Northern sunfish \(*Lepomis peltastes*\)](#), and [least darter \(*Etheostoma microperca*\)](#) are associated with or depend on aquatic vegetation. These fish species are adversely impacted by alterations to natural shorelines and increased nutrient loads. Lake sturgeon, a fish of large rivers and lakes, is impacted by loss of connectivity due to dams (see River subchapter). Deepwater sculpin (*Myoxocephalus thompsonii*) inhabit deep lakes with cold waters, which are threatened by eutrophication and climate change. See also the Fish sub-chapter.

Mammals

Perhaps the mammal SGCM most closely identified with lake habitats is the [moose \(*Alces alces*\)](#). Moose depend on lakes for drinking water. During the spring and summer they also are attracted to lakes and their shoreline habitats for an abundance of favorite foods including sedges, horsetails, and aquatic plants such as pondweeds and water lilies. SGCM bat species not only utilize lakes for drinking but are also attracted to lakes and their riparian shorelines during the summer months for the aquatic insects that emerge from the water's surface.

Reptiles

[Blanding's turtle \(*Emydoidea blandingii*\)](#) use a variety of aquatic habitat, including wetlands and lakes. Deep lakes are usually avoided when shallow, well-vegetated wetlands are nearby. Populations in the southern portion of their range are threatened by warming water temperature due to climate change.

Insects

Lakes provide primary habitat for 18 species of caddisflies (*Trichoptera*) as well as secondary habitat for an additional 3 species, as well as habitat (unspecified as to primary or secondary) for 13 species of dragonflies and damselflies (*Odonata*). Please see the Aquatic Invertebrates sub-chapter for more information.

Mussels

Minnesota is home to the only known lake endemic freshwater mussel species, the Lake Floater (*Pyganodon lacustris*), which requires silt or soft mud bottoms, and low to non-existent flow. This species historically occupied lakes as far south as the Iowa/Minnesota border but now is only known from central to northern Minnesota.

Plants

The 36 vascular plant SGCN found associated with lake habitat represent 8% of all the SGCN plants.

Moyle (1945) suggested three aquatic floral regions of the state based on water chemistry data, which included soft-water, hard-water, and alkaline (Table 3.16).

Table 3.16. Limnologists and biologists have classified midwestern lakes into biologically meaningful categories, often using a combination of hydrology and water chemistry (Sather 1991).

Aquatic Flora	Region	Alkalinity	Sulphate	pH
Soft-water	northeast	<40 ppm	<5 ppm	6.8-7.5
Hard-water	central	90-250 ppm	<50 ppm	8.0-8.8
Alkaline	western	>150 ppm	>125 ppm	8.4-9.2

A relatively small group of plants have evolved to live in the Minnesota's aquatic environments with special adaptations to survive unique habitat conditions across the state. In soft-water lakes of the northeast, [*American awlwort* \(*Subularia aquatica* ssp. *americana*\)](#), [*algae-like pondweed* \(*Potamogeton confervoides*\)](#), and [*lavender bladderwort* \(*Utricularia resupinata*\)](#) occur in the shallow sandy sites of low nutrient, clear water lakes. Human-caused nutrient increases in these lakes can result in lower clarity and a shift to more leafy aquatic plants at the detriment of the small, low-nutrient adapted, rare species. In central Minnesota, [*purple-flowered bladderwort* \(*Utricularia purpurea*\)](#) and [*spotted pondweed* \(*Potamogeton pulcher*\)](#) appear restricted to very few lakes but may grow locally abundantly within sites. Direct removal of vegetation for recreational use as well as declines in water clarity threaten these species. Widgeon grass (*Ruppia maritima*) and [*spiny naiad* \(*Najas marina*\)](#) are "salt-loving" submerged plants that are adapted to the naturally higher alkalinity of western Minnesota lakes. Drainage and upland development have degraded many of these habitats.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. Species in Greatest Conservation Need.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors. See also: [Stressors to Biological Communities in Minnesota’s Lakes](#).



Development

Shorelines are altered by lakeshore property owners. Minnesota has currently lost 40 to 50% of its natural shorelines, and the rate of loss is estimated to be 1 to 2% per decade (Minnesota Natural Shoreline Partnership 2023). As a lake’s watershed becomes more developed, nutrient levels increase and water quality decreases due pollution runoff, poor stormwater management, and shoreline phosphorus pollution from shoreland sewer systems and lawns down to the lake.

Both permitted and illegal removal of aquatic vegetation (via mechanical or herbicide) is a threat to lake habitats, as aquatic vegetation is important for many species including fish and loons. Common Loons (*Gavia immer*) nest near the water edge on lakes and use mud and aquatic vegetation to create the nest. Shoreline development and property management that includes the clearing of natural vegetation along the shoreline are threats to loon nesting success. Minnesota is losing natural shorelines due to development and land management (i.e. mowing) [Minnesota’s Vanishing Natural Shorelines: A Loss that Contributes to Degraded Lake Quality](#).

Artificial shoreline stabilization structures are built to protect human infrastructure and prevent erosion. Riprap, bulkheads, jetties and other shoreline protection strategies are designed to alter the behavior and movement of sand and sediments. These strategies can effectively retain or increase sediment in select locations while eroding shoreline down shore. Artificial structures can eliminate access to beaches for some shore nesting species and beach erosion can degrade or eliminate nesting and foraging habitat for shore nesting SGCN (Prosser et al., 2018).



Crop Production

Drainage from agricultural fields within the watershed of a lake, either by surface flow, or by drain tiles, can affect downslope hydrology and water quality through the transmission of sediments and chemicals.

Agricultural lands contribute as a leading cause of lake water quality impairments and contribute substantially to lake eutrophication. Over half of Minnesota's wetlands and shallow lakes have been ditched and drained. Wetland losses have been the greatest in the Prairie Parkland ecoregion where well over 90% of the wetlands have been drained or filled. Forty to 50% of remaining wetlands have altered hydrology from storm water inputs, roads, or drain tiles (Genet et al., 2019).

Direct wetland loss is lost habitat but also the water conveyed through drainage systems has increased levels of sediments, nutrients and chemicals that further impair the receiving water bodies. Excess phosphorus and silt remain in lakes and wetlands and result in degraded water quality and wildlife habitat. Tile drainage bypasses natural riparian areas that would normally serve to filter some of the nutrients and sediments so many of the wetlands remaining are in poor condition. Blann et al. (2009) is a thorough review of impacts of agricultural drainage on aquatic ecosystems.



Mining and Quarrying

Mining or quarrying operations may affect hydrology or produce sediments that can impact waters with sediments and chemicals. Proposed sulfide ore mines in northern Minnesota could produce pollution issues.



Roads, Trails, and Railroads

Roads and railways can disrupt connectivity to lake habitats. In general, increased impervious surfaces adjacent to lakes and in watershed lead to increased sedimentation, turbidity, and chemical levels. De-icing salts dispersed on roadways increase the salinization of the lakes. [Chloride | Minnesota Pollution Control Agency](#)



Utility Corridors

Powerlines present a risk to waterfowl when situated between feeding and roosting areas. This is especially true for lines near lakes, rivers and wetlands. Less agile birds, such as trumpeter swans are particularly susceptible to collision with overhead lines located near waterbodies (APLIC, 2012).



Fishing

A leading cause of mortality in the common loon and other waterbirds is ingestion of lead fishing tackle. In popular angling lakes, tackle loss results in many metric tons of lead across lake bottoms annually (Radomski et al., 2006). This can occur when loons eat fish that have ingested lead tackle, eating bait with lead tackle attached, or from eating tackle directly off the lake bottom. Loons may also get entangled in loose fishing line or injured from ingested hooks. Lead is particularly toxic when ingested as the grinding mechanisms and acidic environment of the gizzard cause the lead to be transported into the blood and organs. A small lead fishing split shot weight can kill a loon in a matter of weeks. [Get the lead out | Minnesota DNR](#) [Lead Poisoning in Loons | U.S. Fish & Wildlife Service](#).



Recreation

Large powerful boats, and particularly the use of wake boats (designed to throw a big wake for watersports) can disrupt vegetation and nesting waterbirds including loons and leads to erosion.

Boat strikes commonly kill wildlife like loon chicks and turtles during the summer.

Human disturbance on beaches and islands serving as nesting and foraging areas for birds such as piping plovers and common terns disrupts nesting behavior and can limit reproductive success (DeRose-Wilson et al., 2018; Stantial et al., 2021). Beach access at times is a threat and temporary beach closures are needed.

Boats, docks, and fishing bait that are moved among lakes and not properly decontaminated contribute to the spread of invasive species (see below).



Dams and Water Management

Drainage of wetlands and fields in watersheds of lakes can impacts seasonal hydrology of the remaining water bodies by delivering water at a faster rate through drainage networks. Dissolved nutrients and sediments can also be elevated in this runoff, leading to excess nutrients in the receiving water bodies.

Natural connections between water bodies have been influenced by dams, bridges and culverts and impede species movement and alters habitat. See Rivers subchapter for

Loons nest near the water's edge, and waterbodies that undergo large fluctuations in water levels may flood nests and reduce reproductive success of loons and other shore nesting species. (Windels et al. 2013). Water level fluctuations impact nesting and foraging habitat conditions for shore nesting bird species. High water levels erode or eliminate nesting sites (Maxson S.J., Haws K.V, 2000) Water level fluctuations also impact the nesting habitat of overwater nesters such as western grebes and red-necked grebes, black and Forster's terns. Water level management that results in rapid water level changes during the nesting season can destroy nests (Bracey A.M., et al., 2024).



Invasive Species (Problematic Non-native Species)

Several invasive aquatic species are present in Minnesota lakes. [Eurasian watermilfoil](#) and [curly-leaf pondweed](#) have been present in Minnesota lakes for a century and decades, respectively. Eurasian milfoil can grow to the water surface and compete with native plants. Curly-leaf pondweed is well adapted to nutrient-rich or nutrient polluted lakes. A newer invasive species, [starry stonewort \(*Nitellopsis obtusa*\)](#), is associated with native species declines and drastic changes in the structural complexity of plant communities.

Invasive aquatic animals present in many Minnesota lakes include [common carp](#), spiny water flea, rusty crayfish, and zebra mussels. Common carp is widely distributed in central and southern Minnesota, disrupting shallow lakes as their feeding habits uproot aquatic plants and suspends fine sediments into the water. Zebra mussels filter the water, which reduces food for larval fish and other animals and increases water clarity. So, zebra mussels can change the food web and reduce the carrying capacity for popular fish such as walleye.

Shipping in the Great Lakes does not pose a direct threat to Lake Superior's fish but the ships themselves are a vector for the introduction of invasive species when they release their ballast water. As of April 2010, 89 non-native aquatic species had already been introduced into Lake Superior. Among the long-list of non-native

species are the following fish: Eurasian ruffe (*Gymnocephalus cernuus*), round goby (*Apollonia melanostomus*), sea lamprey (*Petromyzon marinus*), alewife (*Alosa pseudoharengus*), and rainbow smelt (*Osmerus ordax*). Changes to ballast water regulations have reduced the rate of invasive species by an estimated 85% since 2006 (Ricciardi and MacIsaac 2022). Many biologists consider aquatic invasive species the biggest threat to the Great Lakes (Lake Superior Binational Program 2014; see also Non-native Invasive Species and [Infested Waters List | Minnesota DNR](#)).



Water-borne Pollution

Eutrophication occurs when a lake becomes enriched in nutrients which leads to excessive plant and algal growth, and is a major threat to Minnesota lakes. Nonpoint source pollution (pollution originating from a broad area) includes phosphorus, nitrogen, sediment, bacteria, and other contaminants generally results from land runoff, precipitation, atmospheric deposition, drainage, seepage or hydrologic modification. As runoff flows, it picks up and carries away natural and human-made pollutants, depositing them into lakes, rivers, wetlands, and groundwater, and estuaries. ([EPA NPS Webpage](#)) These nonpoint source pollutants represent the largest combined contribution (an estimated 86%) to the state's water pollution. ([MPCA Water Quality Initiatives](#)) Water quality measured by nutrients and contaminants has been shown to decline with higher agricultural or urban development in the watershed. Some key impacts include excess algal growth, reduced plant diversity, and reduced diversity in fish and invertebrate communities. ([DNR Water quality non-point sources](#)).

Polyfluoroalkyl Substances (PFAS) and Polychlorinated biphenyls (PCBs) are human made chemicals that are both banned in Minnesota, but persistent in the environment. PFAS may get into lakes from industrial discharge, leaching from landfills into groundwater, and from firefighting foam products. The specific effects of these pollutants on wildlife are poorly known, but their presence is documented in many fish and is reported on the DNR Lakefinder website for specific lakes [LakeFinder | Minnesota DNR](#).

Chemicals used in proposed sulfide mines may leach into watersheds, causing water acidification and leaching of heavy metals.



Air-borne Pollution

Air pollution includes a large range of chemicals, from mercury via primarily from coal burning, pesticides via drift from farm and residential lands ([Introduction to Pesticide Drift | US EPA](#)), microplastics and nanoplastics via wind and waves moving broken down plastic waste fragments, to greenhouse gases via combustion of coal and gas. Mercury bioaccumulates and biomagnifies through lake food webs, which affects the wildlife and human health as it is a potent neurotoxicant. Fish from lakes in the northeast generally have the higher mercury concentrations (Radomski et al. 2024).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter

temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

Minnesota's thousands of lakes, including coldwater and small shallow lakes, are increasingly vulnerable to rising temperatures driven by climate change. Coldwater lakes capable of supporting species like lake trout (*Salvelinus namaycush*) and cisco (*Coregonus artedii*) require cool, oxygen-rich waters to sustain their sensitive aquatic communities. However, as average temperatures rise, particularly in winter and summer, these lakes are experiencing thermal stress that threatens the long-term viability of cold-adapted species (Jacobson et al., 2010; Fang et al., 2012). Warming water reduces the volume of suitable cold, oxygenated habitat, often referred to as an "oxythermal squeeze", which can lead to population declines or local extirpations (Jane et al., 2024).

Rising air temperatures have also disrupted the natural freeze-thaw cycles of Minnesota's lakes. Ice-in now occurs later in the fall and ice-out earlier in the spring, significantly shortening the duration of seasonal ice cover. These shifts increase the risk of winterkill in shallow lakes, alter stratification and mixing patterns, and affect the timing and productivity of aquatic food webs (Magnuson et al., 2000; Sharma et al., 2019).

The ecological and recreational consequences of these changes may be severe. Angling opportunities for coldwater species may diminish, while lakes become more susceptible to algal blooms and invasive species like [common carp \(*Cyprinus carpio*\)](#) and [Eurasian watermilfoil \(*Myriophyllum spicatum*\)](#), which thrive in warmer, nutrient-rich conditions.



Changes in Precipitation and Hydrology related to Climate


From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

More specifically in lake systems across Minnesota, an anticipated higher incidence of heavy storms may increase flooding, runoff, affect water level fluctuations in lakes and alter hydroperiods in shallow lakes. Also, high wind events on large lakes can generate intense wave action which threatens shorebird nesting habitat and aquatic plant establishment. Snow on frozen lakes acts as an insulating barrier, and influences ice formation and ice melt, and light penetration during winter. Changing patterns of snowfall will accordingly affect lakes as well. [Climate trends affecting lakes and rivers | Minnesota DNR](#)

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, to protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN, five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Examples of conservation actions are presented below grouped by the primary strategy that they implement and tagged with icons for the stressor(s) that they address. Some of the strategies, such as Strategy 3, collaborating with partners, could truly be applied to most broadly and effectively implement all actions. In this case, under Strategy 3 we highlight actions for which collaboration is particularly vital.

Potential Conservation Actions for Lakes



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Utilize Aquatic Management Areas to protect shorelines and wetlands. Aquatic Management Areas Minnesota DNR
	Prevent loss or degradation of all types of shallow lakes. Preserve shallow lakes and wetlands, especially in the Eastern Broadleaf Forest and Prairie Parkland provinces. Focus on protecting larger shallow lakes (> 4 acres (10 hectares)) and shallow lake and wetland complexes. Protect known nesting areas of Black terns and Forster's terns.
	Enforce wetland protection regulations ("no-net loss") as they pertain to shallow lakes and wetlands.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.

Stressor	Action
	Restore large complexes of shallow lakes and wetlands, with attention to the habitat features required by SGCN.
	Enforce shoreline development and aquatic vegetation removal restrictions.
	Manage the spread of new invasive species, particularly in rare very deep-water habitats. Manage the encroachment of invasive plants in shallow lakes. See DNR's Aquatic Invasive Species webpage for up-to-date information on DNR's suggested actions regarding reducing the spread.
	To address altered hydrology of shallow lakes, encourage restoration of wetlands and grasslands and natural hydrological function in the watershed.

	Manage for natural water regimes in shallow lakes.
	Prevent problematic practices in key watersheds to reduce external nutrient loading in shallow lakes.




Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.

Stressor	Action
	To address boat traffic and wake boats disturbing nesting habitat and causing erosion, ensure appropriate provisions are in place to limit the use of wake boats to protect loon nesting habitat. Work with lake associations and county governments. Ensure there is sufficient enforcement. Invest in research and education on the topic.
	When SGCN shorebirds are nesting on lakeshores, temporary beach restrictions can be beneficial to protect the nests from recreational disturbance. Participation by agency, conservation partners, and volunteers can help to implement, explain, and enforce the closures. Seasonal beach restrictions protect nesting and foraging areas of birds such as piping plovers and common terns (Hillman et al., 2015). Wildlife sanctuary signage placed around breeding habitat reduces recreational disturbance to these sites. Local support from landowners, conservation partners and enforcement officers is important in supporting these beach closures.
	Encourage anglers to voluntarily phase out lead tackle through education and outreach. Provide a directory of non-toxic tackle manufacturers. Encourage tackle retailers to stock non-toxic lead alternatives for anglers. There are several programs underway with partnerships with the Minnesota Pollution Control Agency aimed at these goals, linked under key partnerships below.
	Promote early coordination on utility corridor design and siting. Avoid routing lines near or across lakes where feasible.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.

Stressor	Action
	Examine efficacy of floating artificial nest platforms for common loons. Floating platforms may improve reproductive success on waterbodies with considerable water-level fluctuations. Artificial Nesting Platforms (ANPs) for common loons Minnesota DNR
	Long term monitoring efforts are underway for loons. Two different volunteer programs, the Minnesota Loon Monitoring Program and the Volunteer LoonWatcher Survey continue to provide useful data about the status of loons and will be critical to keep track of long-term population trends. The Minnesota Loon Restoration Project partnership continues to protect habitat, deploy and measure the efficacy of Artificial Nesting Platforms, support non-lead tackle initiatives, and monitor loon populations.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Stressor	Action
	To protect lake shorelines from development pressure, promote initiatives to preserve sensitive shorelines, promote shoreline restorations, and implement shoreline setbacks. Lack of education and public buy-in can be addressed by education at local level within communities (e.g. lake associations, see Loon Restoration case study), governments, and managers. Develop BMPs for technical staff. In addition, more funding is needed for lakeshore preservation and restorations, including funding to support research to inform decision making.
	Continue outreach, education and enforcement regarding movement of invasive species, including Watercraft Inspection Program, mobile invasive species decontamination sites, and “Clean, Drain, Dispose” laws that reduce invasive species spread.
	Consider changes in regulation of ditching, tiling, and erosion to prevent shallow lake losses due to sedimentation. Support agricultural policies for crop production incentives to further develop and deliver programs that are lucrative for landowners to maintain working lands that protect and restore habitat.

Case Study: Monitoring Minnesota's Lakes

In the Land of 10,000 Lakes, a suite of complementary lake monitoring programs provides necessary information for proper lake management. Below several of our lake monitoring programs are highlighted.

Lake Index of Biotic Integrity: The DNR and Minnesota Pollution Control Agency (MPCA) monitor the health of many lakes by tracking the fish and plants with indices of biological integrity ([Lake Index of Biotic Integrity](#)). Indices of biotic integrity (IBI) use the diversity and species composition of community to describe the relative health or impairment of the ecosystem as compared to what would be expected if the ecosystem was undisturbed by anthropogenic activities. When applied to plants these are termed 'floristic quality indices' (FQI). For Minnesota lakes, aquatic ecosystem health is described by the fish community with a fish IBI, supplemented by a plant FQI. The fish IBI was developed for four different lake types to accurately evaluate different expectations for their associated fish communities: deep lakes with complex shaped shorelines, moderately deep lakes with rounder shorelines, moderately shallow, heavily vegetated lakes, and shallow lakes (where more than 80% of the lake is 15 feet or less deep). Fish-based IBI scores are used by DNR staff to place lakes into one of six assessment categories: exceptional, fully supporting, vulnerable, not supporting (impaired), inconclusive information and insufficient information.

Lake Water Quality Monitoring: All lakes over 500 acres and a subset of smaller lakes are assessed for water quality by MPCA ([Lake Water Quality Monitoring](#)). The goal of this monitoring is to determine if water quality supports recreation, compare regional differences in water quality, and identify long term trends. Based on these results, a total maximum daily load (quantity of pollutants a watershed can receive without violation quality standards) and a watershed and restoration and protect strategy are completed for each watershed. These reports are the foundation for restoration and protection projects in each watershed, and acknowledge various unique needs of different waters (e.g. oxygen and temperature standards for cold water lakes) [Lake aquatic life and recreation | Minnesota Pollution Control Agency](#)

Fisheries Surveys: Routine lake surveys by the DNR Section of Fisheries include monitoring of fish populations, water chemistry, and fish habitat to guide fisheries management. Lake survey data is used to track fish population trends, evaluate the effectiveness of management actions such as stocking, and establish management goals for a given lake. On average, the Section of Fisheries conducts 650 lake surveys each year. Most of the important fishing lakes are surveyed once every 3-5 years. The 11 largest lakes, including Mille Lacs, Superior, and Lake of the Woods, are surveyed annually.

Sentinel Lakes Program: This program is an intensive, long-term lake ecosystem monitoring program created to detect and understand the physical, chemical and biological changes occurring in Minnesota's lakes. Long-term monitoring on a set of 25 Sentinel Lakes includes water chemistry, fisheries, zooplankton, phytoplankton, habitat, groundwater as well as watershed development (see [Monitoring Minnesota's Changing Lakes](#)). This program documents biological, physical and chemical trends, identifies mechanisms behind them, and directs management solutions for long-term sustainability of lakes.

Watershed Health Assessment Framework: Assesses three components of aquatic ecosystem health: biology, hydrology, and water quality and integrates numerous metrics to derive lake health scores for about 4000 Minnesota lakes [Watershed Health Assessment Framework - Lakes](#).

Lakes of Biological Significance: High-quality lakes based on biological sampling are listed and updated every five years by DNR. The list is used to focus lake protection efforts. Unique plant and animal presence is the primary measure of a lake's biological significance. Lakes are rated and grouped for each of the following communities: aquatic plants, fish, birds, and herpetofauna (amphibians and a turtle species). Lakes are assigned to one of three biological significant classes (Outstanding, High, or Moderate); see [Lakes of Biological Significance - Resources - Minnesota Geospatial Commons](#).

Case Study: Minnesota Common Loon Restoration Project

In 2010, the Deepwater Horizon oil spill negatively impacted the wildlife and ecosystems in the Gulf of America, and an environmental damage settlement of over \$20 billion was approved. Minnesota's state bird, the common loon, winters on coastal waters. Radio telemetry work and chemical analysis of eggs indicated that loons overwintering in the areas impacted by the spill breed in Minnesota. The Open Ocean Trustee Implementation Group provided funds to Minnesota DNR to restore loons lost from the spill in a project spanning 2021-2025.

A partnership between the DNR, United States Geological Survey, US Fish and Wildlife Service, and MPCA was established to accomplish the following goals: 1) acquire and protect lakeshore nesting and foraging habitat, 2) enhance habitat and increase lake stewardship, and 3) reduce lead exposure through advocacy of non-toxic tackle.

Progress on goals:

1. As of July 2025, over 6.5 miles of lakeshore habitat across four parcels has been purchased and protected perpetually as DNR Aquatic Management Areas or Wildlife Management Areas. Two more acquisitions are planned for later in 2025. Extensive surveys confirm nesting and foraging use of the protected habitats by loons.
2. Artificial Nesting Platforms (ANPs) are floating platforms covered in natural vegetation that loons use to nest. They reduce nest failure from predation and water fluctuation. Volunteers were recruited to build and deploy 44 artificial loon nesting platforms to enhance nesting success of loons on lakes with low reproductive output, and the DNR and USGS has engaged in a research study to assess the efficacy of these nesting platforms as conservation tools. Teams of researchers monitored loons on these lakes throughout the reproductive season and provided detailed data on habitat use, nest success/failure, and productivity from nesting platforms. Platform use increased each consecutive year of deployment (1% in 2023, 18% in 2024, and 34% in 2025). Efforts are currently underway to continue the nest platforms project beyond 2025. Additionally, substantial outreach efforts have increased public engagement and interest in loon conservation and shoreline preservation and restoration (see Public Engagement Chapter)
3. The MN PCA led a non-toxic fishing advocacy effort that has included 213 public engagement events and has provided 33,700 free non-toxic tackle sample packs between 2022-2024.

This loon conservation work will continue. By leveraging partnerships with local sports teams (MN United Football Club; The Loons) the DNR has created a new permanent position, the Loon Program Coordinator, to continue and develop loon conservation, research, and outreach efforts across the state and to coordinate citizen science loon monitoring efforts. For more information on the Loon Restoration Project, see annual reports: [2021](#), [2022](#), [2023](#), [2024](#).

Also see related: Minnesota Pollution Control Agency Get the Lead Out [Getting lead out of fishing tackle | Minnesota Pollution Control Agency](#) Minnesota Loon Restoration Project: Partners include Department of Interior, US Fish and Wildlife Service, US Geological Survey, MN Pollution Control Agency: Minnesota Loon Restoration Project | Minnesota DNR Case Studies

References for Lakes

- APLIC. (2012). *Reducing Avian Collisions with Power Lines: The State of the Art in 2012* (p. 184). Edison Electric Institute and APLIC.
https://www.aplic.org/uploads/files/11218/Reducing_Avian_Collisions_2012watermarkLR.pdf
- Blann, K. L., Anderson, J. L., Sands, G. R., & Vondracek, B. (2009). Effects of Agricultural Drainage on Aquatic Ecosystems: A Review. *Critical Reviews in Environmental Science and Technology*, 39(11), 909–1001.
<https://doi.org/10.1080/10643380801977966>
- Bracey, A., Bednar, J., Kolbe, S., & Grinde, A. (2024). *Conserving Black Terns and Forster's Terns in Minnesota*. University of Minnesota Duluth.
- DeRose-Wilson, A. L., Hunt, K. L., Monk, J. D., Catlin, D. H., Karpanty, S. M., & Fraser, J. D. (2018). Piping plover chick survival negatively correlated with beach recreation. *The Journal of Wildlife Management*, 82(8), 1608–1616. <https://doi.org/10.1002/jwmg.21552>
- Fang, X., Alam, S. R., Stefan, H. G., Jiang, L., Jacobson, P. C., & Pereira, D. L. (2012). Simulations of water quality and oxythermal cisco habitat in Minnesota lakes under past and future climate scenarios. *Water Quality Research Journal*, 47(3–4), 375–388. <https://doi.org/10.2166/wqrjc.2012.031>
- Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing.
https://doi.org/10.1007/978-3-030-45367-1_4
- Genet, J., Bourdaghs, M., & Gernes, M. (2019). *Status and trends of wetlands in Minnesota: Depressional Wetland Quality Assessment (2007 – 2017)* (p. 61). MN PCA.
<https://www.pca.state.mn.us/sites/default/files/wq-bwm1-12.pdf>
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Hillman, M. D., Karpanty, S. M., Fraser, J. D., & Derose-Wilson, A. (2015). Effects of aircraft and recreation on colonial waterbird nesting behavior. *The Journal of Wildlife Management*, 79(7), 1192–1198.
<https://doi.org/10.1002/jwmg.925>
- Jacobson, P. C., Stefan, H. G., & Pereira, D. L. (2010). Coldwater fish oxythermal habitat in Minnesota lakes: influence of total phosphorus, July air temperature, and relative depth. *Canadian Journal of Fisheries and Aquatic Sciences*, 67(12), 2002–2013. <https://doi.org/10.1139/F10-115>
- Jane, S. F., Detmer, T. M., Larrick, S. L., Rose, K. C., Randall, E. A., Jirka, K. J., & McIntyre, P. B. (2024). Concurrent warming and browning eliminate cold-water fish habitat in many temperate lakes. *Proceedings of the National Academy of Sciences*, 121(2), e2306906120. <https://doi.org/10.1073/pnas.2306906120>

Lake Superior Binational Program. (2014). *Lake Superior Aquatic Invasive Species Complete Prevention Plan* (p. 92). <https://conservancy.umn.edu/items/dd691b08-029b-435b-be5f-efcb45533c36>

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>

Magnuson, J. J., Robertson, D. M., Benson, B. J., Wynne, R. H., Livingstone, D. M., Arai, T., Assel, R. A., Barry, R. G., Card, V., Kuusisto, E., Granin, N. G., Prowse, T. D., Stewart, K. M., & Vuglinski, V. S. (2000). Historical Trends in Lake and River Ice Cover in the Northern Hemisphere. *Science*, 289(5485), 1743–1746. <https://doi.org/10.1126/science.289.5485.1743>

Maxson, S. J., & Haws, K. V. (2000). Population Studies of Piping Plovers at Lake of the Woods, Minnesota: 19 Year History of a Declining Population. *Waterbirds: The International Journal of Waterbird Biology*, 23(3), 475. <https://doi.org/10.2307/1522185>

Minnesota Natural Shoreline Partnership. (2023). *Minnesota's Vanishing Natural Shorelines: A Loss that Contributes to Degraded Lake Quality* (p. 30). <https://files.dnr.state.mn.us/assistance/backyard/shorelandmgmt/minnesota-vanishing-natural-shorelines.pdf>

Moyle, J. B. (1945). Some chemical factors influencing the distribution of aquatic plants in Minnesota. *The American Midland Naturalist*, 34(2), 402–420.

Prosser, D. J., Nagel, J. L., Howlin, S., Marbán, P. R., Day, D. D., & Michael Erwin, R. (2018). Effects of Local Shoreline and Subestuary Watershed Condition on Waterbird Community Integrity: Influences of Geospatial Scale and Season in the Chesapeake Bay. *Estuaries and Coasts*, 41(S1), 207–222. <https://doi.org/10.1007/s12237-017-0288-0>

Radomski, P. (2006). Historical changes in abundance of floating-leaf and emergent vegetation in Minnesota lakes. *North American Journal of Fisheries Management*, 26(4), 932–940.

Radomski, P., Bergquist, L. A., Duval, M., & Williquett, A. (2010). Potential impacts of docks on littoral habitats in Minnesota lakes. *Fisheries*, 35(10), 489–495.

Radomski, P., Brentrup, J. A., Acquah, S., Preimesberger, A. L., & Kvit, A. (2024). Spatiotemporal variation in fish mercury contamination: Comparing approaches for crafting consumption advisories and assessing impairment. *Journal of Environmental Management*, 370, 122992.

Radomski, P., Carlson, K., & Perleberg, D. (2019). Advancing aquatic vegetation management for fish in north temperate lakes. *Lake and Reservoir Management*, 35(4), 355–363.

Radomski, P., & Goeman, T. J. (2001). Consequences of human lakeshore development on emergent and floating-leaf vegetation abundance. *North American Journal of Fisheries Management*, 21(1), 46–61.

Radomski, P., Heinrich, T., Jones, T. S., Rivers, P., & Talmage, P. (2006). Estimates of tackle loss for five Minnesota walleye fisheries. *North American Journal of Fisheries Management*, 26(1), 206–212.

- Radomski, P. J., & Goeman, T. J. (1996). Decision making and modeling in freshwater sport-fisheries management. *Fisheries*, 21(12), 14–21.
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. P., & LeDee, O. E. (2025). *Effects of climate change on midwestern ecosystems: Temperate flooded and swamp forest* (p. 110pp). Midwest Climate Adaptation Science Center. Available: <https://mwcas.umn.edu/node/821>
- Ricciardi, A., & MacIsaac, H. J. (2022). Vector control reduces the rate of species invasion in the world's largest freshwater ecosystem. *Conservation Letters*, 15(2), e12866. <https://doi.org/10.1111/conl.12866>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>
- Sather, N. (1991a). *A proposed aquatic classification for use by the Minnesota Natural Heritage Program*. Minnesota Department of Natural Resources.
- Sather, N. (1991b). *A proposed aquatic classification for use by the Minnesota Natural Heritage Program*. Minnesota Department of Natural Resources.
- Schupp, D. H. (1992). *An ecological classification of Minnesota lakes with associated fish communities*. Minnesota Department of Natural Resources, Section of Fisheries St. Paul.
- Schwartz, G., & Thiel, G. (1963). *Minnesota's Rocks and Waters: A Geological Story*. University of Minnesota Press.
- Sharma, S., Blagrove, K., Magnuson, J. J., O'Reilly, C. M., Oliver, S., Batt, R. D., Magee, M. R., Straile, D., Weyhenmeyer, G. A., Winslow, L., & Woolway, R. I. (2019). Widespread loss of lake ice around the Northern Hemisphere in a warming world. *Nature Climate Change*, 9(3), 227–231. <https://doi.org/10.1038/s41558-018-0393-5>
- Stantial, M. L., Cohen, J. B., Darrah, A. J., Farrell, S., & Maslo, B. (2021). Habitat-specific behavior, growth rate, and survival of piping plover chicks in New Jersey, USA. *Ecosphere*, 12(10), e03782. <https://doi.org/10.1002/ecs2.3782>
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Windels, S. K., Beever, E. A., Paruk, J. D., Brinkman, A. R., Fox, J. E., Macnulty, C. C., Evers, D. C., Siegel, L. S., & Osborne, D. C. (2013). Effects of water-level management on nesting success of common loons: Water-Level Management and Common Loons. *The Journal of Wildlife Management*, 77(8), 1626–1638.
<https://doi.org/10.1002/jwmg.608>

Vernal Pools

Habitat Description

Vernal pools are naturally occurring ephemeral wetlands generally found in depressions embedded within upland forests of the Eastern Broadleaf Forest and Laurentian Mixed Forest Province within Minnesota. Within these ecological provinces, the Western Superior Uplands, Northern Minnesota Drift and Lake Plains, and Minnesota and Northeast Iowa Morainal ecological sections host higher concentrations of vernal pools due to topographical features driven by historical glacial ice movement patterns. At a more local scale, the presence of vernal pools is strongly influenced by the surrounding forested environment including elements of topography, canopy shading, and leaf litter.

Vernal pools are typically small, shallow, fishless, confined basin wetlands less than 2.4 ac/1 ha in size. They are considered hydrologically 'isolated' wetlands because there is no continuous surface water connection to permanently flooded water bodies. Pools typically receive most of their water from snow melt runoff or rain, however, in settings where pools intersect the water table, groundwater inputs may be important. Based on local topography and surrounding forest characteristics, vernal pools can also be part of larger intricate pool complexes and can be connected to other wetlands or experience overflow from other wetlands/streams, still, any flow connections to permanently flooded water bodies remain temporary.

Vernal pools are characterized by seasonal water fluctuations and temporary flooding. They experience alternating periods of flooding and drying and must go through a drying phase annually or every few years (e.g., 3 out of 5 years). Pools typically fill with water in the spring and dry down by later in the summer, however, some pools also fill again after fall or winter rains or after large precipitation events. Flood duration and timing, as well as the frequency of flood events, can vary widely from pool basin to pool basin and even from year to year given both temporal and spatial differences in precipitation amounts. The hydroperiod, or length of time that the pool is inundated (as well as the amount of water held), has a strong effect on the type of vegetation and wildlife that occupy and utilize vernal pool habitats. Additionally, maintenance of surrounding forest conditions for canopy shading is important for minimizing impacts to wetland hydrology, plant communities, nutrient cycles, and thermal regimes.

These wetlands are not well captured by Minnesota's or the national wetlands inventory and therefore not yet comprehensively mapped for the state of Minnesota. Efforts are currently underway to better understand their distribution. They are thought to be distributed mostly within Mesic Hardwood Forests and Deciduous Wet Forests (and to a lesser extent in Upland Deciduous Forest and Woodland; see those sub-chapters).

Soils are typically hydric and often covered by leaf litter. Large coarse woody debris is common and contributes to the microsite heterogeneity and complexity of vernal pools. Other variables that influence conditions within the wetland include basin (depression) configuration and topography, light availability (canopy cover), composition of surrounding forest (and litter quality and availability), landscape setting, infiltration or drainage capacity of soil, and water inputs (which can vary spatially and temporally).

Habitat Map

To depict Vernal Pool habitat (see Figure 3.18), we compiled spatial data from several sources: The National Land Cover Database (NLCD) and the National Wetland Inventory (for more information, see [Habitat Map Methods](#) in Chapter 3. Habitats). The map in Figure 3.18 depicts a model of where vernal pools may exist based on the parameters outlined below.

National Wetlands Inventory

From the [National Wetland Inventory for Minnesota \(NWI\) Layer](#) we used the Circular 39 attribute field to include Seasonally Flooded Basins or Flat, Shrub Swamp, and Wooded Swamp that were in forested areas and 1 hectare in size or smaller.

National Land Cover Database

From the [National Land Cover Database](#), we narrowed our search for vernal pools by including only those in Mixed or Deciduous Forest parcels that were larger than 10 hectares in size.

Native Plant Communities

Vernal pools have not been characterized in Minnesota's Ecological Classification system, so we cannot classify them here by a particular ecological system or native plant community. At the time of this publication, DNR is actively investigating the locations and ecological characteristics of vernal pools. Emerging information about the vegetation composition of vernal pools is based on survey work in the northcentral part of the state within the range of the [four-toed salamander \(*Hemidactylium scutatum*\)](#). In the paragraphs below we share what preliminary information has been gathered to date.

Vegetation composition of sampled vernal pools varies widely across pools and even within pools. As noted above, hydroperiod can also vary dramatically between pools (and within a pool if basin topography is complex) and has a direct influence on the vegetative response.

Vegetation in pools (or portions of pools) may be dominated by trees like black ash (*Fraxinus nigra*) or yellow birch (*Betula alleghaniensis*) or other woody species like the shrub speckled alder (*Alnus incana*). Pools may also be dominated by herbaceous vegetation or be nearly devoid of vegetation altogether. Vegetation of deeper and/or more open pools can include plants often observed in wet meadow or marsh/aquatic systems like Canada bluejoint (*Calamagrostis canadensis*), water parsnip (*Sium suave*), northern water plantain (*Alisma triviale*), and duckweed (*Lemna* spp.). Species that were encountered more regularly across a variety of pool types include: brownish sedge (*Carex brunnescens*), tuckerman's sedge (*C. tuckermanii*), bladder sedge (*C. intumescens*), spinulose woodfern (*Dryopteris carthusiana*), and marsh marigold (*Caltha palustris*). Mosses are often present in abundance, sometimes forming larger hummocks (*Sphagnum* species) and growing on leaf litter substrate (non-sphagnum species).




Figure 3.18. A map depicting potential (unverified) Vernal Pool habitat in Minnesota, modeled from three wetland types from the National Wetland Inventory for Minnesota, limited to those found within Mixed or Deciduous Forest parcels larger than 10 hectares in size from the National Land Cover Database. Investigations of the locations of vernal pools are ongoing.

Conservation Overview

Vernal pools are a unique and dynamic habitat feature that many animals and plants depend upon for reproduction and survival. Due to their ephemeral nature, these wetlands act as a resource pulse that is available for only a few months of the year and often become a biodiversity hotspot during this time. Several SGCN animals and plants utilize these wetlands; therefore, there is a strong need to understand and protect these sensitive habitats.

Conversion of forested lands to development, agriculture, and other land-uses may result in the degradation and/or loss of vernal pools. Unlike wetlands that hold permanent water, vernal pools are not protected under Minnesota's Wetland Conservation Act (WCA) and often only receive limited protection during forest management activities. Additionally, climate change presents new challenges and many unknowns for the future of vernal pools and the organisms that rely on them. These wetlands are not a readily replaceable resources on the landscape and artificial vernal pools have been met with mixed success. Thus, our primary conservation approach is the protection and enhancement of existing vernal pools.

With the aforementioned reasons, vernal pools are in need of additional research and monitoring in Minnesota. Efforts are currently underway to classify vegetation composition within these wetlands to categorize different types of vernal pools and understand these wetlands connection to the surrounding forest. Furthermore, research is being conducted to understand how vernal pools, and the SGCN that rely upon them, respond to forest disturbance and to identify strategies to minimize detrimental impacts. Through these efforts, our hope is to raise awareness of these important forested wetlands, identify effectiveness techniques to protect the plants and animals using these habitats, and ensure that vernal pools continue to thrive in Minnesota.

Species in Greatest Conservation Need

Vernal pools, despite being ephemeral, very small, and poorly documented, provide primary habitat for 10 animal Species in Greatest Conservation Need (SGCN) and secondary habitat for two others (see Table 3.17). Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently.

Table 3.17. Numbers of Species in Greatest Conservation Need associated with vernal pools as primary or secondary habitat¹.

Species Group	Primary Habitat	Secondary Habitat	Total
Amphibians	3	0	3
Birds	0	1	1
Reptiles	0	1	1
Invertebrates			
Dragonflies and Damselflies			6
Snails (aquatic)	1	0	1
Total	4	2	12

1. Habitat associations for dragonflies and damselflies were not differentiated into primary and secondary habitats and are shown in the total column.

A detailed table associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found for animals in Appendix C. Plants were associated only with a single primary habitat type, based on Minnesota's native plant classification. Because vernal pools are not yet incorporated into that system, none were formally associated to the vernal pool type. However, some SGCN plants that may be found in vernal pools are discussed in the section below.

It should be noted that although the Eastern fairy shrimp (*Eubrachipus vernalis*) is endemic to vernal pools, they are not represented in the table above because habitats were not delineated for any of the SGCN Branchiopoda (shrimp). Also, given the ephemeral nature of this habitat, some additional species mentioned in the text below, while not associated formally with the habitat, opportunistically use these pools.

Vernal pools provide critical habitat for a whole suite of animal species in Minnesota. Due to the periods of prolonged drawdown in vernal pools, permanent fish populations are not able to survive. This provides habitat for organisms that require or benefit from fishless conditions for successful reproduction. Some of these are obligate species and include organisms that use vernal pools almost exclusively for breeding and larval development. Many of the species that utilize vernal pools also rely on the integrity of the upland forest surrounding the pool for things like cover, foraging sites, and overwintering habitat.

Amphibians

Three SGCN salamanders associate with vernal pools as their primary habitat. As a quintessential vernal pool obligate species, [spotted salamanders \(*Ambystoma maculatum*\)](#) depend upon these fishless, seasonal wetlands for successful reproduction. As soil and air temperatures increase and spring rains fall, adults migrate

from upland forests to vernal pools to breed in late March - April (Sexton et al., 1990, Moriarty and Hall, 2014). Additionally, four-toed salamanders also use these forested wetlands for reproduction and seek out vernal pools with abundant sphagnum moss hummocks as their preferred nesting habitat (Chalmers and Loftin, 2006; Moriarty and Hall, 2014). While adult spotted salamanders generally leave the vernal pools soon after breeding, female four-toed salamanders will continue to guard their eggs for approximately one month after oviposition (Banning et al., 2008). Larval development for both of these species occurs within these pools and can last until late summer. In general, these species need these pools to hold water (or ample moisture in leaf litter) until approximately mid-July for successful reproduction in Minnesota (Moriarty and Hall, 2014). In some rare cases, spotted salamander larvae may overwinter in the vernal pools and complete their development the following spring (Whitford and Vinegar, 1966). Both of these species eventually migrate to adjacent upland forests for the remainder of the active season. Eastern newts (*Notophthalmus viridescens*), another SGCN salamander, may also breed within vernal pools, yet they are not as dependent on these specific wetland features as the other two SGCN salamanders. Adult eastern newts will also frequent vernal pool in the spring to forage on other amphibian eggs.

Birds

Several SGCN birds breed and forage within mature upland hardwood forests with interspersed vernal pools. Wetland openings such as vernal pools are a key component of forests occupied by [Red-shouldered hawks](#) (*Buteo lineatus*), which often prey on amphibians and reptiles that inhabit vernal pools (King et al. 2011). While not formally associated with this habitat, [acadian flycatchers](#) (*Empidonax virescens*), and [cerulean warblers](#) (*Setophaga cerulea*) that use mature deciduous forest are often loosely associated with vernal pools. They may be responding to foraging opportunities provided by invertebrate prey linked to vernal pools, or simply the forest structural diversity associated with small canopy gaps.

Mammals

Although not formally associated with vernal pools, several forest bat SGCN, such as [northern long-eared bat](#) (*Myotis septentrionalis*), [little brown myotis](#) (*Myotis lucifugus*), and [big brown bat](#) (*Eptesicus fuscus*), may forage in forest openings above vernal pools (Francl, 2008).

Reptiles

[Blanding's turtles](#) (*Emydoidea blandingii*) use vernal pool habitat early the spring for foraging, thermoregulation, and breeding opportunities (Beaudry et al., 2009). These turtles capitalize on highly abundant and concentrated food resources like amphibian eggs, tadpoles, and various invertebrates. These wetlands are also some of the first water features to ice out in the spring and provide opportunities to thermoregulate via basking and water submersion. Also, vernal pools may attract numerous individuals at once; thus, providing breeding opportunities for adult turtles.

Invertebrates

Some invertebrate species, such as the crustacean Eastern fairy shrimp, are endemic to vernal pools and live out their entire life cycle within these wetlands. Over the course of approximately three weeks, fairy shrimp larvae hatch from dormant cysts, develop into adults, reproduce, and die off (usually by mid-to late May; McCafferty et al., 2010).

Other aquatic invertebrates that utilize vernal pools to forage, reproduce, and undergo larval development include numerous insect species such as caddisflies (*Trichoptera*), dragonflies and damselflies (*Odonata*), water bugs (*Hemiptera*), midges (*Chaoboridae* and *Chironomidae*), beetles (*Coleoptera*), as well as some mollusk species like snails (*Lymnaeidae*, *Physidae*, and *Planorbidae*), and clams (*Sphaeriidae*) (Colburn, 2004; Batzer et al., 2004; Bischof et al., 2013). Dragonflies SGCN associated with vernal pools include the following six species: springwater dancer (*Argia funebris*); arrowhead spiketail (*Zoraena obliqua*), azure bluet (*Enallagma aspersum*), elfin skimmer (*Nannothemis bella*), [ocellated darner \(*Boyeria grafiana*\)](#), and [Quebec emerald \(*Somatochlora brevicincta*\)](#).

Plants

The intersection of vernal pools and the upland forest that surrounds them can provide a key habitat for a number of SGCN plant species. When the transition zone or ecotone is gradual and rich and mesic or wet-mesic in nature, it can provide habitat conditions that support some of MN's most rare interior forest species of moonworts (*Botrychium*) and grapeferns (*Sceptridium*) as well as the mycorrhizal fungi associates that they rely upon. This includes SGCN species such as [narrow triangle moonwort \(*Botrychium angustisegmentum*\)](#), swamp moonwort (*B. tenebrosum*), and [blunt-lobed grapefern \(*Sceptridium oneidense*\)](#).

Shallow, sparsely vegetated vernal pools with shorter hydroperiods also provide key habitats for grape ferns, including blunt-lobed grapefern, a SGCN species with a state conservation status of imperiled (S2).

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “Stressors” section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Conversion of forests to developed landscapes can result in the direct loss of vernal pools, degradation of vernal pools and their surrounding terrestrial habitat, increased road mortality of vernal pool wildlife, inhibited animal movement due to impervious surfaces, as well as reduced gene flow between wetlands (Windmiller and Calhoun, 2007; Hamer and McDonnell, 2008; Beaudry and deMaynadier, 2009; Evans et al., 2017).



Crop Production

Forest conversion to agriculture may result in the loss and degradation of vernal pools. Additionally, proximity to agricultural lands can increase the likelihood of herbicide and pesticide introductions into vernal pools, which can have profound impacts on plant and invertebrate communities (Battaglin et al., 2009).



Roads, Trails, and Railroads

Roads and trails can produce hazardous run-off (such as salt, fertilizer, and pesticide/herbicide) that may alter water quality and chemistry within vernal pools. For example, salinization of these freshwater pools may occur from road deicers found in spring snowmelt. This has been shown to negatively impact zooplankton and cyanobacteria communities, as well as a reduction of dissolved organic matter, creating unsuitable conditions for various invertebrates and amphibians (Eakin et al., 2019; Vigil and Schuler, 2024). Additionally, roads may act as a movement barrier and/or source of mortality (via car strikes) as herpetofauna migrate to and from vernal pools (Colino-Rabanal and Lizana 2012).



Timber Harvest

Forest management can alter forest composition and cover (shade), thereby affecting, water levels and temperature, litter source, quality, and quantity, vegetation composition within pools, and

seasonal food availability with an overall impact on pool food webs. This can impact both plant and animal SGCNs that utilize the pools directly as habitat for at least part of their life history as well as species utilizing the transition zone of the pools and the surrounding upland forest with close proximity to pools.

Several studies have examined the impacts forest management on hydrology, plant and animal communities of vernal pools. One study suggested that riparian buffers around vernal pools help maintain similar hydrology to those of undisturbed sites, whereas thinning or clearcutting up to the pool boundary resulted in greater water depths and took several years to return to pre-harvest water depths (Kolka et al., 2011). Palik and Kastendick (2010) determined that harvesting up to the margins of the vernal pool resulted in significant changes in plant communities in the ground layer and shrub/tree regeneration within the wetland. Furthermore, this study demonstrated that a 15.25-m (50 ft) uncut buffer around the vernal pool helped mitigate changes in the plant community. The utilization of buffers around vernal pools generated mixed results when attempting to mitigate impacts to invertebrate communities (Batzer et al., 2004, Bischof et al., 2013). More specifically, particular species (such as Eastern fairy shrimp) appear to be more sensitive to changes in the surrounding forests and the subsequent alterations to hydrology and sediment load (Hanson et al., 2010). Another study examined the long-term effects that timber harvest had on vernal pool obligate species that also utilize the adjacent upland forests: spotted salamanders and wood frogs (*Lithobates sylvaticus*). Within vernal pools that experienced timber harvests, researchers found that occupancy of breeding adults and egg mass abundance significantly decreased over time for both species, with wood frogs experiencing a more drastic decline (Wiewel et al., 2023). Furthermore, it was also noted that many of the vernal pools within harvested forests were smaller in size and held less water than those in unharvested forests. While these results contradict Kolka et al.'s (2011) assessment that timber harvests resulted in greater water depths, it should be noted that there are many other habitat factors that influence vernal pool depth and size such as precipitation frequency and amount, previous winter's snow depth and surrounding topography.



Recreation

Recreation (both foot and OHV) within a vernal pool may significantly degrade sensitive features, especially with repeated disturbance. The bottom of these pools are often mucky and soft; therefore, human activity within the wetland may impact the topography and alter how water is distributed within the pool. Additionally, if recreation activity occurs during a sensitive period, such a breeding and larval development of amphibians, this can result in mortality of breeding individuals and loss of reproductive potential. Finally, recreational traffic may act as a vector for disease and invasive species if care is not taken to disinfect and clean equipment.



Invasive Species (Problematic Non-native Species)

Increasing density and species diversity of invasive European earthworms can drastically alter the leaf litter/humus, as well as soil horizon composition, within the adjacent forest around a vernal pool (Hale et al., 2005). In turn, this impacts soil nutrient availability and ultimately plant species composition (Alexander et al. 2022). Areas with high earthworm invasion may result in a loss of SGCN plants (such as rare ferns of *Botrychium* and *Sceptridium*; Gundale, 2002; Alexander et al., 2022), as well as a loss of vital cover for forest-floor dwelling animals.

Emerald ash borer (*Agilus planipennis*) poses a risk to vernal pools since black ash is a common species within vernal pools. The mortality of ash within these wetlands may impact the hydrology, canopy cover, wetland

depth, and plant species composition. This may result in the vernal pool becoming unsuitable for various species. Grinde et al. (2022) conducted experimental manipulations of black ash wetlands to mimic EAB damage and examined the response of bird, amphibian, and small mammal communities. While there was an initial increase in species richness attributed to the change in forest structure, researchers suggested that there would be a shift from forest-dependent wildlife to more open-canopy, wetland-associated species within these wetlands.



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years (Minnesota Pollution Control Agency and Minnesota Department of Commerce 2025). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al., 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al., 2022; [Climate Change in Minnesota](#)). Warmer temperatures may result in shorter hydroperiods and create challenges for larval development of some animals that require water availability (Montrone et al., 2019). However, amphibian eggs and larva may develop faster under warmer water conditions, thus, some animals may be able to adapt better than others. Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter,

drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

More specifically for vernal pools, increased frequency, duration, and intensity of droughts may drastically reduce hydroperiods. In some extreme cases, this may result in water shortages for an entire season, which may inhibit some animals’ ability to successfully reproduce (Wiewel et al., 2023). Alternatively, large increases in precipitation will increase the hydroperiod, but may also flood previously suitable habitat. For example, sphagnum moss hummocks used for nesting by four-toed salamanders may become inundated with water after extreme rainfall events, which may impact the viability of deposited egg masses.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, to protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN, five strategies were identified:

-  Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.
-  Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience
-  Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas
-  Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions
-  Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Examples of conservation actions are presented below grouped by the primary strategy that they implement and tagged with icons for the stressor(s) that they address. Some of the strategies, such as Strategy 3, collaborating with partners, could truly be applied to most broadly and effectively implement all actions. In this case, under Strategy 3 we highlight actions for which collaboration is particularly vital. If an action does not relate to a particular stressor, ‘n/a’ is marked in the stressor column.

Potential Conservation Actions for Vernal Pools



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	A minimum of a 50 ft “no impact” buffer should be maintained year-round around the whole perimeter of the vernal pool. This includes no removal of overstory and understory vegetation, as well as no use of heavy equipment.
	If habitat disturbance surrounding a vernal pool is unavoidable, plan work from October 1 through March 15 and conduct work under frozen soil conditions only.
	Maintain habitat corridors between adjacent vernal pools (especially those within 300 ft of each other) to enable wildlife movement and promote genetic diversity.
	Avoid development of roads or trails in proximity to vernal pools to minimize an alteration of hydrology. For existing or new roads/trails, ensure run-off cannot enter the vernal pool.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.

Stressor	Action
	Avoid herbicide/pesticide application within forested habitat surrounding vernal pools.
	Avoid or minimize soil compaction and rutting if operating heavy equipment in proximity to vernal pool.
	In the case of forest management, ensure the slash is distributed evenly and not piled into or surrounding a vernal pool.
	Avoid ditching or draining of vernal pools during any land conversion process.

Stressor	Action
	Follow proper disinfecting protocols if traveling between isolated vernal pools to prevent the movement of invasive species or diseases that may affect salamanders (see <i>Amphibian</i> sub-chapter within <i>Chapter 2. SGCN</i>).

(no actions were identified under Strategy 3)



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.

Stressor	Action
n/a	Given that this habitat is not yet well-mapped and understood in Minnesota, continue studies to identify and map the habitat and learn more about the plant communities and animals reliant on the habitat. Continue building and sharing knowledge about managing for healthy vernal pool systems.
	Prior to any work in forested habitat that may contain vernal pools, locate and assess the condition of these wetland features in the spring/early summer to ensure proper delineation of vernal pool boundary.

(no actions were identified under Strategy 5)

Case Study: Monitoring the Response of Rare Forest Dependent Wildlife Populations to Forest Management: Four-toed Salamander Monitoring

Understanding how a species, and their habitat, responds to forest management is a vital piece to conserving rare forest-dependent wildlife. Initiated in 2022, the DNR Nongame Wildlife Program began a study to explore how four-toed salamander populations respond to forest management activities in proximity to their preferred nesting habitat: vernal pools. This study utilizes a Before-After-Control-Impact (BACI) study design in an effort to tease apart natural annual variation in salamander abundance from the impact that forest management may have on these populations. Within four-toed salamander range in Minnesota, 45 vernal pools were identified for long-term monitoring with approximately half of these vernal pools scheduled for forest management activities within 350 meters of the pool boundary (i.e., treatment vernal pools). The other half of the survey pools are identified as controls and are not planned for forest management. Survey pools were selected based on historical and recent observations of four-toed salamanders or pools that consisted of suitable nesting habitat. Survey work involves conducting nesting surveys that consist of 30 minutes of "active" search time or until all suitable nesting habitat within a vernal pool has been searched, whichever comes first. "Active" search time is defined as any time spent searching suitable nesting habitat (moss hummocks, woody debris, sedges clumps, etc.) within the highwater mark of a vernal pool for female four-toed salamanders and/or eggs. Relative abundance is determined by counting the occurrence of nesting activity (i.e., presence of salamander guarding eggs, gravid female salamanders, or unattended egg masses) within each wetland. All wetlands are surveyed at least once during the nesting season (May 1 – June 15).

Started in 2024, vegetation surveys and trail cameras deployment occurred within several survey pools (both treatments and controls) to better understand vegetation composition and hydroperiod of vernal pools, respectively. Vegetation surveys were conducted in August after the drying period within these vernal pools to allow for ample growing season and to minimize impacts to salamander larval development. These surveys employed a relevé sampling structure and documented species abundance, diversity, and other vernal pool characteristics. The goals of these surveys are to document vegetation composition/diversity within vernal pools and to develop a classification structure for this understudied wetland type. Additionally, these data establish a baseline of vegetation composition and provide the opportunity to examine changes in vegetation after forest management has occurred. The objectives of the trail camera deployment are to record spatial and temporal variation in hydroperiod across most of the study vernal pools, and to determine whether hydroperiod significantly changed after forest management operations. These cameras are programmed to record timelapse photos of the wetlands and identify how many days standing water was present/visible. The deepest location of the wetlands was located during nesting surveys and a measuring post and camera was installed to monitor this location.

As of 2024, four-toed salamanders were detected in approximately 90% of the study vernal pools. Over 500 nests have been detected with over 290 female four-toed salamanders observed during nesting surveys. The mean egg mass size is 39.5 eggs (StDev: 53.2). The majority of vernal pool depths ranged from 24 – 35 cm at the time of nesting surveys. With May 1 as the start day for hydroperiod monitoring, most vernal pools held water for 74 to 152 day (July 14 and September 30, respectively) with a mean hydroperiod of 103 days (August 12). To date, few vernal pools have experienced the planned forest management activities; thus, not enough data exists to make any inference on the impacts of forest management on four-toed salamander abundance. As this study continues for the next several years, additional data will yield valuable information on how best to protect these sensitive vernal pools and the species that rely on them.

References for Vernal Pools

- Alexander, G., Almendinger, J. and White, P., 2022. The long-term effects of invasive earthworms on plant community composition and diversity in a hardwood forest in northern Minnesota. *Plant-Environment Interactions*, 3(2), pp.89-102.
- Banning, J.L., Weddle, A.L., Wahl lli, G.W., Simon, M.A., Lauer, A., Walters, R.L. and Harris, R.N., 2008. Antifungal skin bacteria, embryonic survival, and communal nesting in four-toed salamanders, *Hemidactylium scutatum*. *Oecologia*, 156, pp.423-429.
- Battaglin, W.A., Rice, K.C., Focazio, M.J., Salmons, S. and Barry, R.X., 2009. The occurrence of glyphosate, atrazine, and other pesticides in vernal pools and adjacent streams in Washington, DC, Maryland, Iowa, and Wyoming, 2005–2006. *Environmental Monitoring and Assessment*, 155, pp.281-307.
- Batzer, D. P., B. Palik, and R. Buech. 2004. Relationships between environmental characteristics and macroinvertebrate assemblages in seasonal woodland pools of Minnesota, U.S.A. *Journal of the North American Benthological Society* 23: 50–68.
- Beaudry, F., P.G. deMaynadier, and M.L. Hunter. 2009. Seasonally dynamic habitat use by spotted (Clemmys guttata) and Blanding's turtles (Emydoidea blandingii) in Maine. *Journal of Herpetology*, 43(4), pp.636-645.
- Bischof, M.M., M.A. Hanson, M.R. Fulton, R.K. Kolka, S.D. Sebestyen, and M.G. Butler. 2013. Invertebrate community patterns in response to hydrology and environmental variables in seasonal ponds in Minnesota, USA. *Wetlands*, 33(2): 245-256.
- Chalmers, R.J. and Loftin, C.S., 2006. Wetland and microhabitat use by nesting four-toed salamanders in Maine. *Journal of Herpetology*, 40(4), pp.478-485.
- Colburn, E.A., 2004. Vernal Pools: Natural History and Conservation. McNaughton & Gunn, Inc., Saline, MI, p.426.
- Colino-Rabanal, V.J. and M. Lizana. 2012. Herpetofauna and roads: a review. *Basic and Applied Herpetology*, 26, pp.5-31.
- Eakin, C.J., A.J. Calhoun, and M.L. Hunter. 2019. Effects of suburbanizing landscapes on reproductive effort of vernal pool-breeding amphibians. *Herpetological Conservation and Biology*, 14(2), pp.515-532.
- Evans, J.P., Cecala, K.K., Scheffers, B.R., Oldfield, C.A., Hollingshead, N.A., Haskell, D.G. and McKenzie, B.A., 2017. Widespread degradation of a vernal pool network in the southeastern United States: challenges to current and future management. *Wetlands*, 37, pp.1093-1103.
- Francl, K.E., 2008. Summer bat activity at woodland seasonal pools in the northern Great Lakes region. *Wetlands*, 28, pp.117-124.
- Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in*

Forests and Rangelands of the United States (pp. 57–83). Springer International Publishing.

https://doi.org/10.1007/978-3-030-45367-1_4

Grinde, A., B Youngquist, M., A Slesak, R., R Kolbe, S., D Bednar, J., J Palik, B. and W D'Amato, A., 2022. Potential impacts of emerald ash borer and adaptation strategies on wildlife communities in black ash wetlands. *Ecological Applications*, 32(4), p.e2567.

Gundale, M. J. (2002). Influence of Exotic Earthworms on the Soil Organic Horizon and the Rare Fern *Botrychium mormo*. *Conservation Biology*, 16(6), 1555–1561. <https://doi.org/10.1046/j.1523-1739.2002.01229.x>

Hale, C.M., Frelich, L.E., Reich, P.B. and Pastor, J., 2005. Effects of European earthworm invasion on soil characteristics in northern hardwood forests of Minnesota, USA. *Ecosystems*, 8, pp.911-927.

Hamer, A.J. and McDonnell, M.J., 2008. Amphibian ecology and conservation in the urbanising world: a review. *Biological conservation*, 141(10), pp.2432-2449.

Hanson, M. A., Palik, B. J., Church, J. O., & Miller, A. T. (2010). Influences of upland timber harvest on aquatic invertebrate communities in seasonal ponds: efficacy of forested buffers. *Wetlands ecology and management*, 18(3), 255-267.

He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>

King, J.C., Dubay, S.A. and Woodford, J.E., 2011. Distribution and nest site selection of red-shouldered hawks (*Buteo lineatus*) in forests of northeastern Wisconsin (USA). *Forest ecology and management*, 261(1), pp.169-177.

Kolka, Randall K.; Palik, Brian J.; Tersteeg, Daniel P.; Bell, James C. 2011. Effects of riparian buffers on hydrology of northern seasonal ponds. *Transactions of the American Society of Agricultural and Biological Engineers (ASABE)*. 54(5):2111-2116.

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>

McCafferty, S.S., Warren, N., Wilbur, C. and Shumway, S., 2010. Genetic divergence among Massachusetts populations of the vernal pool fairy shrimp *Eubranchipus vernalis* (Crustacea: Anostraca). *Northeastern Naturalist*, 17(2), pp.285-304.

Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). 2025. Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state's greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/lraq-3sy25.pdf>

- Montrone, A., Saito, L., Weisberg, P.J., Gosejohan, M., Merriam, K. and Mejia, J.F., 2019. Climate change impacts on vernal pool hydrology and vegetation in northern California. *Journal of Hydrology*, 574, pp.1003-1013.
- Moriarty, J.J. and C.D. Hall. 2014. *Amphibians and Reptiles in Minnesota*. U of Minnesota Press., Minneapolis, MN., p.370
- Palik, B. J., & Kastendick, D. (2010). Response of seasonal pond plant communities to upland forest harvest in northern Minnesota forests, USA. *Forest ecology and management*, 260(5), 628-637.
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. 2025. Effects of Climate Change on Midwestern Ecosystems: North American Bog and Fen. Midwest Climate Adaptation Science Center. (67 pages) Available: <https://mwasc.umn.edu/node/821>
- Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>
- Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434. <https://doi.org/10.1111/cobi.14434>
- Sexton, O. J., Phillips, C., & Bramble, J. E. (1990). The Effects of Temperature and Precipitation on the Breeding Migration of the Spotted Salamander (*Ambystoma maculatum*). *Copeia*, 1990(3), 781–787. <https://doi.org/10.2307/1446443>
- Vigil, J.P. and M.S. Schuler. 2024. Salt pollution reduces turbidity, dissolved organic matter, and cyanobacteria in experimental vernal pool communities. *Science of The Total Environment*, 931, p.172948.
- Wiewel, A. N. M., Brand, A. B., & Grant, E. H. C. (2023). Long-Term Effects of Timber Harvest on Ephemeral Pools and Occupancy of Spotted Salamanders (*Ambystoma maculatum*) and Wood Frogs (*Lithobates sylvaticus*). <https://doi.org/10.1670/21-059>
- Whitford, W.G. and Vinegar, A., 1966. Homing, survivorship, and overwintering of larvae in spotted salamanders, *Ambystoma maculatum*. *Copeia*, pp.515-519.
- Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report]. <https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>
- Windmiller, B., Calhoun, A.J.K., 2007. Conserving vernal pool wildlife in urbanizing landscapes. In: Calhoun, A.J.K., deMaynadier, P.G. (Eds.), *Science and Conservation of Vernal Pools in Northeastern North America*. CRC Press, Boca Raton, Florida, pp. 233–251.

Cliff, Talus, and Rock Outcrop

Habitat Description

This chapter features several types of habitat that are composed of sparsely vegetated native plant community types. These habitats, which include cliffs, talus slopes, and rock outcrops, all have extensive areas of exposed substrate. They occur as small patches or long linear strips throughout the state (see Figure 3.19).

Cliff and talus communities are present on cliffs or talus slopes on steep-sided bluffs, along lakes and streams, on margins of bedrock ridges, and in other settings with sheer bedrock exposures. Cliffs and talus slopes are often associated with one another because talus slopes are composed of rock fractured either from cliffs or smaller areas of exposed bedrock on steep hillsides. The vegetation of these communities is generally open. Lichens and mosses are often the dominant life forms, and vascular plants are sparse or patchy because of scarcity of soil.

In the Laurentian Mixed Forest Province, cliffs and talus communities are mostly restricted to the North Shore Highlands and Border Lakes subsections, where the Precambrian bedrock is frequently at or just below the surface and rugged topography is common. In the Eastern Broadleaf Forest Province, these communities are abundant in the Blufflands Subsection of southeastern Minnesota, where sedimentary bedrock is typically at or near the surface and topography is rugged. Scattered cliffs are present on bedrock formations elsewhere in the state.

Rock outcrop communities are open plant communities on horizontal or sloping bedrock exposures. Some outcrops are relatively level whereas others are very rugged. They are common in landscapes with thin soils over bedrock and are typically, although not always, small in size (i.e., less than 25 acres (10 hectares)). Crustose and foliose lichens typically cover exposed rock surfaces, and fruticose lichens are also common. Vascular plant cover is sparse to patchy, depending on the amount of fracturing of the bedrock surface and accumulation of soil in cracks, crevices, and shallow depressions. Outcrops that have minimal fracturing and little accumulation of soil are dominated by lichens, with scattered shrubs and herbaceous plants. Outcrops with deep crevices provide vital year-round habitat, including for overwintering below the frostline, for some SGCN reptiles, like the [common five-lined skink \(*Plestiodon fasciatus*\)](#).

Many plants on bedrock outcrops are adapted to frequent desiccation because of the low moisture-holding capacities of substrates, and exposure to direct sunlight and strong winds. Plants must also withstand rapid fluctuations in substrate temperatures, which are significantly colder at night than in surrounding forests and much warmer during midafternoon on sunny days. Limited availability of nutrients in outcrop communities strongly influences community composition and limits growth rates of plants. Fire, as well as frequent drought and scarce soils, play a role in maintaining the open vegetation characteristic of these communities. Rock outcrop communities are most common in the Border Lakes and North Shore Highlands subsections and in the upper Minnesota River valley

Habitat Map

To depict Cliff, Talus, and Rock Outcrop habitat (see Figure 3.19), we compiled spatial data from several sources: DNR's Natural Plant Community layer and the Minnesota Geological Survey. (for more information, see Habitat Map Methods in Chapter 3. Habitats). We note included sub-types below; underlined items have links to online descriptions.

Associated Native Plant Community Classes by Ecological Systems

Cliff/Talus

[CTn11 Northern Dry Cliff \(PDF\)](#)

[CTn12 Northern Open Talus \(PDF\)](#)

[CTn24 Northern Scrub Talus \(PDF\)](#)

[CTn32 Northern Mesic Cliff \(PDF\)](#)

[CTn42 Northern Wet Cliff \(PDF\)](#)

[CTu22 Lake Superior Cliff \(PDF\)](#)

[CTs12 Southern Dry Cliff \(PDF\)](#)

[CTs23 Southern Open Talus \(PDF\)](#)

[CTs33 Southern Mesic Cliff \(PDF\)](#)

[CTs43 Southern Maderate Cliff \(PDF\)](#)

[CTs46 Southern Algific Talus \(PDF\)](#)

[CTs53 Southern Wet Cliff \(PDF\)](#)

Rock Outcrop (RO)

[ROn12 Northern Bedrock Outcrop \(PDF\)](#)

[ROn23 Northern Bedrock Shrubland \(PDF\)](#)

[ROs12 Southern Bedrock Outcrop \(PDF\)](#)

Minnesota Geological Survey

From the [Minnesota Geological Survey](#) we used the Minnesota Bedrock Outcrop polygons.




Figure 3.19. A map depicting Cliff, Talus, and Rock Outcrop habitat in Minnesota including DNR Native Plant Community Classes in the Ecological System of Rock Outcrop as well as rock outcrop data from the Minnesota Geological Survey.

Conservation Overview

Moderate cliffs and algific talus slopes in the Blufflands Subsection of southeastern Minnesota provide the only habitat for several rare SGCN land snails, some of which only occur a couple of locations. These cliff and talus communities provide a cool, moist, equable climate that allows the snail species to persist. A variety of human activities, either within these communities (such as rock climbing) or affecting the sinkholes on lands above these cliffs, may threaten these communities.

Rocky outcrops are critically important for protecting specialized flora and fauna and for maintaining biodiversity at multiple spatial scales (Fitzsimons and Michael, 2017). Their locations as dispersed across the landscape make designing conservation strategies challenging. Rock outcrop communities along the Minnesota River are threatened by woody encroachment (e.g eastern red cedar (*Juniperus virginiana*) and common buckthorn (*Rhamnus cathartica*), mineral extraction, and residential and recreational development.

Species in Greatest Conservation Need

Despite taking up a small proportion of the land in terms of acres, cliffs, talus and rock outcrops provide habitat for 76 animal, and 69 plant SGCN as primary or secondary habitat (see Table 3.18). Primary habitats are those that species rely on and use most consistently; if these habitats were degraded or lost, it would have the greatest effect on the species. Secondary habitats are those in which a species can be found some of the time, but less frequently than a primary habitat.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Plant species were only associated with their single most primary habitat. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C (animals) and Appendix D (plants).

Table 3.18. Numbers of Species in Greatest Conservation Need associated with the Cliff, Talus, and Rock Outcrops as either primary or secondary habitat¹.

Species Group	Primary Habitat	Secondary Habitat	Total
Birds	4	0	4
Mammals	2	12	14
Reptiles	8	0	8
Terrestrial Invertebrates			50
Bees			15
Beetles (terrestrial)			4
Butterflies			8
Moths			14
Snails (terrestrial)	8	0	8
Spiders	0	1	1
Plants	69		69
Total	91	13	145

1. Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column. For plants, only a single primary habitat was identified per species.

Examples of SGCN are described below. Please note that species when underlined contain a clickable link to their profile in Minnesota's [Rare Species Guide](#).

Birds

Common nighthawks (*Chordeiles minor*) associate with rock outcrops as a primary habitat. They nest on sparsely vegetated sites, especially rock outcrops in the upper Minnesota River valley and the Border Lakes Subsection, as well as sites in prairies and oak savannas, and in recently burned or logged areas. The [peregrine falcon \(*Falco peregrinus*\)](#) is an SGCN whose primary habitat association is with cliffs. This species had been extirpated in Minnesota due to DDT poisoning by the 1960's, and after the ban of DDT and active reintroduction efforts, a breeding population has been re-established in Minnesota and their range in the United States continues to expand (see [Rare Species Guide profile](#) for more details). Both of these species also nest on buildings and are associated therefore with urban and other developed spaces.

Mammals

Although found in a variety of habitats in northeastern Minnesota, rock voles (*Microtus chrotorrhinus*) are most often associated with frost-fractured rock outcrops and rocky streambeds. Moist conditions and boulders or crevices seem to be important habitat features. The [smoky shrew \(*Sorex fumeus*\)](#) is also found in a variety of northern forest habitats but its preferred microhabitat includes cool, damp forests with mossy covered rocks and decaying debris. They also have been documented in glacial boulder streams and within talus slopes. Although these two SGCN closely associated with talus slopes, rock outcrops and boulder piles, many other mammal SGCN may be observed using these habitats, including the Eastern Heather Vole (*Phenacomys ungava*), Western harvest mouse (*Reithrodontomys megalotis*), and American Badger (*Taxidea taxus*).

Reptiles

In Minnesota, common five-lined skink is found in or near [granite outcrops \(PDF\)](#) in the Minnesota River Valley ([Lang 1982 \(PDF\)](#)), and in exposed limestone and sandstone outcrops and [bluff prairies \(PDF\)](#) in the eastern part of the state (Oldfield and Moriarty 1994). The areas inhabited by skinks generally have abundant cover and basking areas. The presence of rocks or bedrock outcrops is a very important habitat component for this species ([Lang 1982 \(PDF\)](#); Howes and Lougheed 2004; Gelvin-Innvaer et al., in press).

The timber rattlesnake (*Crotalus horridus*) occurs in the eastern and central United States. Minnesota is on the northwestern periphery of its range, which extends north along the Mississippi River from Illinois to Minnesota. Timber rattlesnake specimens have been collected from 8 counties in southeastern Minnesota, but survey efforts in the late 1990s and early 2000s found that populations were substantially reduced or extirpated from several areas where they occurred historically ([Keyler and Fuller 1999 \(PDF\)](#); Keyler and Oldfield 2003). In Minnesota, the ideal habitat for timber rattlesnakes includes forested bluffs, south-facing rock outcrops ([rock outcrops PDF](#)), and bluff prairies ([bluff prairies \(PDF\)](#)), particularly in the Mississippi River valley. Bluff prairies located on steep, south or west-facing hillsides, with rock outcroppings and ledges, are essential habitat components because over-wintering dens are often located in these areas. Surrounding forests, prairies, and agricultural lands are used as summer feeding grounds. Two necessary habitat components for this species are open areas for thermoregulation and dens for over-wintering.

Case Study: Common Five-Lined Skink Microhabitat Research

Gelvin-Innvaer et al. (2025) describe an initiative by DNR Nongame Wildlife Program and the Minnesota Biological Survey, in cooperation with other partners, to assess microhabitat use of and response by the common five-lined skink, a SGCN, to outcrop management along the Upper Minnesota River Valley. This resulted in recommendations for management thresholds for this population. These recommendations help to update the DNR's Rare Species Guide Account for this species. See also Gelvin-Innvaer, 2018.

See also the draft chapter for Shore, Cliff, Talus, [Rock Outcrop](#), and Dune under Case Studies which includes companion initiatives for protection, restoration, and management of rock outcrops in the Upper Minnesota River Valley. Collectively, these constitute multi-faceted, integrated initiatives that involved many, diverse partners and funding sources.

See also Gelvin-Innvaer et al. 2025 which describes a companion initiative by DNR Nongame Wildlife Program and MBS – in cooperation with partners to assess microhabitat use of and response to outcrop management by a common five-lined skink, a SGCN. See also [Reptile Renaissance | July–August 2018 | Minnesota Conservation Volunteer](#)

Invertebrates

Several species and subspecies of rare tiger beetles of the genus *Cicindela* are found in the shoreline-dunes-cliff/talus habitat. [The laurentian tiger beetle \(*Cicindela denikei*\)](#) is limited to rock outcrops and sandy openings in hardwood forests in the Border Lakes Subsection.

The [bluff vertigo \(*Vertigo meramecensis*\)](#) and other species of Pleistocene land snails are found on steep, moist, shaded, cool north-facing slopes and cliff faces in the Blufflands Subsection. This small land snail also occurs on algific slopes and moderate cliffs but generally avoids areas with continuous cooling from cold air or water discharge. Although researchers have raised questions recently about the taxonomic distinctness of some of the SGCN land snails, all are rare and associated with the same habitats.

Plants

In western Minnesota, [waterhyssop \(*Bacopa rotundifolia*\)](#) is found primarily in small rainwater pools on [bedrock outcrops \(PDF\)](#) and occasionally along the [margins of shallow prairie ponds \(PDF\)](#). The water is rarely more than inches deep and typically evaporates by July, leaving the sediments at the bottom of the pool moist or sometimes dry and cracked. *Bacopa rotundifolia* is generally able to persist for a couple more weeks after the pools dry up and then completely disappears from view.

The known habitats of [wolf's spikerush \(*Eleocharis wolfii*\)](#) in Minnesota are primarily associated with level outcrops in the southern part of the state ([Minnesota River Prairie](#) and [Inner Coteau](#) subsections) of [bedrock \(PDF\)](#). Specifically, they grow in the margins of shallow rain water pools that form in depressions on the bedrock or in thin sediments over the [bedrock](#) within a [prairie or savanna \(PDF\)](#) setting.

While associated primarily with the muddy margins of prairie ponds in non-forested wetlands, [Short pointed umbrella-sedge \(*Cyperus acuminatus*\)](#) also occurs at the edge of [shallow rock pools \(PDF\)](#). The shallow rock pools where *C. acuminatus* has been found are mostly level with shallow depressions where rainwater accumulates. The pools are typically only a few meters across and 5-7 cm (2-3 in.) deep. Plants are typically rooted in a thin layer of organic sediment that accumulates at the bottom. This is a sparsely vegetated,

ephemeral habitat that may support only a few scattered plants or several hundred, depending on habitat conditions.

Another extremely rare plant in Minnesota is the [ball cactus \(*Coryphantha vivipara*\)](#), which occurs in crevices on granite outcrops. New occurrences of these and several other SGCN plant species were documented in a systematic survey of ephemeral rock pools by Harris (2010) during the summers of 2006-2008.

In southeastern Minnesota [Leedy's Roseroot \(*Rhodiola integrifolia* ssp. *leedyi*\)](#), a SGCN that is also federally threatened, is restricted to shallow ledges on north-facing dolomite cliffs that can be nearly 100 feet high. It is further limited to sites where groundwater seeps through the sinkholes and openings amidst the region's karst landscape, releasing cool air through subsurface vents along the cliff face. Dependent on these cool, moist conditions, the species is considered a glacial relic, "a plant that was more widespread at the end of the last glaciation but which has since become isolated because of the loss of appropriate habitat in intervening areas as the climate has warmed" (Sather 1993). The four populations known in Minnesota are among just six that exist; the remaining two sites are in New York, one of which supports only a single plant. As the climate continues to warm, the unique habitat where this species is found faces another threat. Increased precipitation, especially several storm events, can disturb the cliff environment, dislodging vegetation and substrate materials. Indeed, in recent years plants were found in the talus at the base of cliffs at three sites and, at another site, the entire cliff face had been dislodged.

Lichens

Rock outcrops are home to many unique lichen species. Lichen 'species' are a complex that consists of two or more, typically obligate, symbionts. The dominant partner is a fungus, which creates the main body of the organism. The endosymbiont is either a green alga or a cyanobacteria that lives within the fungal tissue. [Usnea mutabilis \(Bloody Beard Lichen\)](#) is a Minnesota State Threatened species that occurs primarily on sandstone outcrops near cool rivers associated with a variety of other lichen and moss species. It has been documented living on wood in other parts of its global range, but we know very little about why it is restricted to outcrops in Minnesota.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key 'stressors' that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the "Stressors" section in Chapter 1. *Species in Greatest Conservation Need*.

It is important to note that some of the factors listed as 'stressors' can also be used to advance conservation goals. Broad terms such as 'fire management' reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Mining and Quarrying

Pit quarries and dimensional stone quarries occur on mapped bedrock sources with visible rock outcrops in the Minnesota River Valley. Some of these are adjacent to protected conservation lands with documented or potential for rare features associated with rock outcrops. Some of these conservation lands exhibit evidence of quarrying prior to protection. It is unclear from available GIS data layers, how many of these quarries are still active, although some appear to be. Rock outcrop habitats are not a renewable resource, therefore quarrying would result in habitat loss and fragmentation for species dependent on rock outcrops which can exacerbate geographic isolation and limit gene flow. Active quarrying also increases the potential for direct injury or mortality.

Arguably one of the rarest plant species in Minnesota, *Coryphantha vivipara* ([ball cactus](#)), occurs in crevices on granite outcrops. Historically, it also occurred in thin-soiled prairies which were largely converted to agriculture. Today, its remaining populations are threatened by granite quarrying. Much of its potential habitat in Minnesota has already been destroyed by mining.



Recreation

Rocky habitats are often not considered fragile when in fact they can be very sensitive to disturbance. Surface rocks that are even slightly displaced (e.g. rock collecting, moved for trails, etc.) can change thermal conditions considerably and impact their value as refugia for reptiles (Pike et al., 2010). Foot traffic, mountain bikes, OHVs, and horses can damage fragile and endemic plants. Increased recreational trails and human presence are often associated with the spread of invasive species and increases in commensal predators (domestic and wild). Rock climbing may impact sensitive vegetation (Clark and Hessler 2015; March-Salas et al. 2023; Johanson et al. 2024)



Fire Management

With fire suppression since Euro-American settlement, woody plant cover is increasing in many rock outcrop communities. Prescribed fire is a valuable tool for managing woody encroachment on outcrops (Johanson et al. 2024). However, often fires do not “carry” well on outcrops when there are limited natural fuels. In fact, large rock outcrops can act as natural fire breaks. High intensity fires on open rock outcrops typically are naturally rare (Fitzsimons et al. 2017). Further, burning large piles of down woody debris atop outcrops (such as those that remain from large woody removal projects) can result in intense flames that can sterilize thin soils hosting rare endemic plants.



Invasive Species (Problematic Non-native Species)

European and glossy buckthorn impact rock outcrop ecosystems in a variety of ways. These include reducing the availability of open habitats (sunlight and thermal effects) and altering soil and litter properties which in turn can impact native vegetation and invertebrates which may have broader trophic effects (Gelvin-Innvaer et al. in press).



Problematic Native Species

Eastern red cedar and sumacs encroaching on rock outcrops reduce the suitability of the habitat for some SGCN, such as the five-lined skink (Gelvin-Innvaer et al. in press).



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al. 2022). For more information and resources for climate-adapted management strategies, see the Climate sub-chapter in Chapter 6. Implementation.

Increases in precipitation, especially heavy rainfall events, may impact some of the more geologically unstable (i.e., friable) cliff features in the southeast, causing steep rock faces or cliffs to crumble more easily when disturbed. Some increases in cliff sloughing and crumbling at sites have been detected where the federally threatened [Leedy's roseroot \(*Rhodiola integrifolia* subsp. *leedyi*\)](#) is found.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, to protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN, five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Examples of conservation actions are presented below grouped by the primary strategy that they implement and tagged with icons for the stressor(s) that they address. Some of the strategies, such as Strategy 3, collaborating with partners, could truly be applied to most broadly and effectively implement all actions. In this case, under Strategy 3 we highlight actions for which collaboration is particularly vital.

Potential Conservation Actions for Cliff, Talus, and Rock Outcrop



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Protect algal talus and moderate cliffs, as well as rock outcrops from mining and other development activities.
	Restrict recreational access to and use of cliffs, talus slopes and rock outcrops, including but not limited to hiking, rock climbing, rock collecting and mountain biking.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.

Stressor	Action
	Manage rock outcrops to prevent woody encroachment that may be advancing due to a history of fire suppression, the presence of invasive species like European and glossy buckthorn, the presence of problematic native species like red cedar and sumac, and climatic factors favorable to woody species. In the Minnesota River valley, work to prevent encroachment of junipers. Consider the application of grazing, especially by goats, to help reduce woody encroachment on rock outcrops.
	When using fire to manage invasive species or woody encroachment, consult habitat specialists to ensure that fires are not so intense as to negatively impact the habitat.

(no actions were noted for Strategies 4 – 6)

Case Study: Protection of Granite Rock Outcrop Ecosystems on the Upper Minnesota River

Granite rock outcrops along the Upper Minnesota River are among the oldest exposed rock in North America, dating back approximately 3.6 billion years. These outcrops are also home to rare and specialized plant and animal communities rarely found elsewhere in Minnesota, including several types of cactus and one of Minnesota's only three lizard species, the five-lined skink. However, these rock outcrops are increasingly threatened by mining, overgrazing, and development. With funding from the Minnesota Environment & Natural Resources Trust Fund, the Renville County Soil and Water Conservation District worked with Minnesota's Board of Water and Soil Resources to acquire conservation easements that will permanently preserve portions of this endangered habitat.

Soil & Water Conservation District (SWCD) employees saw a need to protect the natural environment and to provide economically viable choices for the landowners. The Minnesota River Valley contains exposed ancient granite rock outcrops that provide unique landscape features and habitat for specialized plant and animal communities rarely found elsewhere in Minnesota. No programs existed that would give landowners a payment if they chose to protect the area from development by mining, overgrazing and other development interests. Rock outcrops are a component of the Minnesota River's riparian zone, and destruction of this unique habitat degrades water quality and wildlife habitat in the Minnesota River and its tributaries. Removal of the rock results in severe degradation and permanent loss of these unique landscape features. The Minnesota River Corridor is easily susceptible to fragmentation because it comprises such a small percentage of the Minnesota River Watershed. Past development activities and mining operations have already fragmented large areas of the fragile Minnesota River Corridor.

A total of 560.4 acres of rare and unique Minnesota River Valley landscape were permanently protected and sixteen landowners were paid \$1,379,814 for voluntarily placing perpetual conservation easements on those acres. Five counties participated in the project including Lac qui Parle, Chippewa, Yellow Medicine, Redwood and Renville. Easement applications were scored by resource professional teams and funding was based on those scores. For more information, see: [Protection of Granite Rock Outcrop Ecosystem](#).

References for Cliff, Talus, and Rock Outcrop Habitat

- Clark P. and A Hessler. 2015. The effects of rock climbing on cliff-face vegetation. *Applied Vegetation Science* 18:705-715. <https://cdnsiencepub.com/doi/abs/10.1139/b98-178>
- Fitzsimons, James A. and Damian R. Michael. 2017. Rocky outcrops: A hard road in the conservation of critical habitats. *Biological Conservation* 211:36-44.
- Gelvin-Innvaer, Lisa A., Jeffrey B. LeClere, Mike Worland, and Carol D. Hall. In press. Microhabitat use by disjunct Common Five-lined Skink (*Plestiodon fasciatus*) populations at their northwestern range extent. *Herpetologica*.
- Harris, F. 2009. Rock Pools on the Prairie. *Minnesota Conservation Volunteer* Mar-April p.30-39
- Harris, F. 2010. Recent Rediscovery of rare plants of ephemeral rainwater pools in Sioux quartzite prairies in Southwestern Minnesota. *Proceedings of the 22nd North American Prairie Conference*. Pp. 105 –109. <https://publications.iowa.gov/50658/>
- Howes, B. J., & Loughheed, S. C. (2004). The importance of cover rock in northern populations of the five-lined skink (*Eumeces fasciatus*). *Herpetologica*, 60(3), 287–294. <https://doi.org/10.1655/03-85>
- Johanson, Connor J. , Samuel A. Schmid, and Neal D. Mundahl. 2024. Fire History and Woody Encroachment Influence Saxicolous Macrolichen Community Composition on Dolomite Outcrops in Dry Bluff Prairies of the Minnesota Driftless Area. *Prairie Naturalist* 56:56-70 https://www.gri.msstate.edu/publications/docs/2024/11/169582024_johanson_PNAT_lichen_ecology.pdf
- Keyler, D. E., and B. Oldfield. 2003. Timber Rattlesnake field survey. Report to the Minnesota Department of Natural Resources. 17 pp
- Lang, J.W. 1982. Distribution and abundance of the Five-lined Skink (*Eumeces fasciatus*) in Minnesota. Final report submitted to Nongame Wildlife Program, Minnesota Department of Natural Resources. Minnesota Department of Natural Resources, USA. Available at https://files.dnr.state.mn.us/eco/nongame/projects/consgrant_reports/1982/1982_lang.pdf. Accessed on 24 March 2022.
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- March-Salas, M., Morales-Armijo, F., Hernández-Agüero, J. *et al.* Rock climbing affects cliff-plant communities by reducing species diversity and altering species coexistence patterns. *Biodivers Conserv* **32**, 1617–1638 (2023). <https://doi.org/10.1007/s10531-023-02567-1>
- Oldfield and Moriarty. (1994). *Amphibians and Reptiles Native to Minnesota*. University of Minnesota Press.

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.
<https://statesummaries.ncics.org/chapter/mn/>

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434.
<https://doi.org/10.1111/cobi.14434>

Sather, Nancy. 1993. Leedy's Roseroot: A cliffside glacial plant. Biological Report #42 of the Minnesota Natural Heritage and Nongame Wildlife Programs. Minnesota Department of Natural Resources.

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S.

Caves

Habitat Description

Karst landscape features are formed by the dissolution of soluble rocks like limestone and dolomite. The resulting porous and soluble rock in the Upper Mississippi Valley has been recognized as one of the most “important karst regions” (Herak and Stringfield 1972, p. 470-471). Key features of karst landscapes include springs, disappearing rivers, sinkholes, cool and stable temperatures, high humidity, and caves and it is in the southeast corner of Minnesota where over 400 caves are found.

Cave and karst features are concentrated in the southeast corner of Minnesota where thick units of limestone and dolomite rock exist (Alexander and Brick, 2021). Fillmore County has the highest concentration of caves in southeast Minnesota, where caves are primarily formed in the Dubuque and Galena formations (Alexander, 1980). Caves exist in other counties of southeastern Minnesota including those formed in the Prairie du Chien Group, St. Peter Sandstone, and Platteville limestone.

The largest cave in Minnesota, with over 13 miles of cave passage, is Mystery Cave which is located in Forestville/Mystery Cave State Park. Mystery Cave is an important bat hibernaculum and provides habitat for many cave invertebrates. Water from the South Branch Root River sinks underground at Mystery Cave and emerges at Seven Springs Wildlife Management Area, approximately 1.5 miles east of the cave ([Forestville/Mystery Cave State Park](#)).

In Fillmore County another significant area of caves is Cherry Grove Blind Valley Scientific and Natural Area (SNA). Cherry Grove SNA protects a large cave system including Goliath Cave as well as several hydrologically connected smaller caves. Water that sinks into the Cherry Grove Blind Valley travels underground and reemerges approximately 2.7 miles to the northeast at springs on Canfield Creek in Forestville/Mystery Cave State Park ([Cherry Grove Blind Valley SNA](#)).

Spring Valley Wildlife Management Area, also in Fillmore County, includes a sinkhole that has been hydrologically linked to Spring Valley Caverns and nearby Bly Spring. Other sinkholes on the property are likely hydrologically connected to Spring Valley Caverns as well. Spring Valley Caverns is privately owned with nearly 6 miles of mapped passages. Bat River Cave, also in Fillmore County and privately owned, is ecologically significant because it once supported the largest hibernating colony of bats in Minnesota.

Additional important sites for bats in Minnesota include sandstone caves in Pine and Washington counties, including Robinson’s Ice Cave and Leslie Cave. In addition, in Lake County, two natural caves, Hole-in-the-Head and Gnomen, provide habitat for hibernating bats (Knowles, 1992). Due to the geology in the northeast region, natural caves are uncommon, and these sites are likely of regional importance to hibernating bat populations. Other geologic features in the north that can offer alternative overwintering habitat for bats include talus slopes and deep vertical rhyolite crevices (Gerda Nordquist, DNR, pers. comm.).

We created a map of important caves from Open Street Map and DNR Natural Heritage Information System database of caves hosting bat colonies (see Figure 3.20).

Habitat Map

To depict Cave habitat (see Figure 3.20), we compiled spatial data from DNR's Natural Heritage Information System and Open Street Map. For more information, see Habitat Map Methods in Chapter 3. Habitats.

Natural Heritage Information System

From the [Natural](#) Heritage Information System we used data relating to the location of bat colonies.

Open Street Map

From [Open Street Map](#), we digitized Public/semi-public caves.




Figure 3.20. A map depicting Cave habitat in Minnesota including DNR Natural Heritage Information System data regarding caves used as bat hibernacula and digitized public/semi-public caves from Open Street Map.

Conservation Overview

Caves provide habitat for rare and unique species of animals, the majority of which in Minnesota are invertebrates specially adapted to living in subterranean environments. Many of these species are troglobites, or organisms that spend their whole life in caves; many of these species have evolved specialized traits such as the loss of eyes and pigmentation. Minnesota cave ecology is only partially understood and the animals living there are exposed to extreme conditions with few food sources, no light, yet a stable and consistent temperature of approximately 8.8°C and nearly 100% humidity. Other surface-dwelling animals such as bats, flies, moths, raccoons, rodents, and snakes will use caves to overwinter.

Groundwater within the cave forming karst landscape of southeastern Minnesota is critical to maintaining healthy cold-water stream habitats for trout in southeast Minnesota. Water traveling through bedrock aquifers typically cools as it travels underground. When groundwater emerges at a spring, it provides cool isothermal temperatures critical for cold-water trout species. Native brook trout (*Salvelinus fontinalis*) are found in some of these stream environments. Recreation opportunities provided by trout streams are of great economic importance to communities throughout southeast Minnesota (Anderson, 2016). These fisheries, like the caves themselves, are affected by [nitrate contamination](#) and pesticide run-off (also see Water-borne Pollutants below).

Species in Greatest Conservation Need

Bats

Four SGCN species in Minnesota rely on caves and mines for winter survival: [little brown bats \(*Myotis lucifugus*\)](#), [big brown bats \(*Eptesicus fuscus*\)](#), [tricolored bats \(*Perimyotis subflavus*\)](#), and [northern long-eared bats \(*Myotis septentrionalis*\)](#).

Caves and mines have been an important habitat for early documentation of bat species in Minnesota. The first account of winter ecology for bats was published in 1934 by Charles A. Evans (Evans, 1934), which was followed by a period of scientific intrigue for bats in the state into the 1950's. Studies over the years added more sites with known hibernacula, more recently well described in the Distribution and Status of Bats in Minnesota report (Nordquist and Birney, 1985).

Caves provide the narrow range of ideal conditions for bats to hibernate for extended periods (Kunz, 1982). To hibernate, bats seek out subterranean environments that do not fluctuate with above-ground weather conditions, where cool temperatures and high humidity remains stable year-round. This stability provides ideal conditions for bats to enter torpor, a physiological state characterized by a controlled reduction in body temperature, metabolic rate, and overall energy expenditure (Barclay et al., 2001). Torpor is a critical mechanism that allows bats to maintain fat reserves when ambient temperatures are unfavorably cool, and food availability is low (Geiser, 1988). During the long harsh winters in Minnesota, caves provide protective and stable environments where bats can use extended bouts of torpor with sparse active periods to survive without foraging for up to eight months underground.

While there are 490 listed caves described in Minnesota (Alexander and Brick, 2021), not all are suitable for bats. Suitable hibernating conditions within caves are primarily found in large cavities which have small entrances that reduce fluctuation with outside air temperatures, Nordquist and Birney 1985 describes “cull-de-sac or labyrinth passages that protect bats from drafty corridors.” Stable caves in Minnesota generally maintain a constant year-round temperature of 8°C or 47°F, and variation in temperature and humidity levels can be found based on the site entrance exposure, level of water seepage, and site size (Nordquist and Birney, 1985). This variation results in differing levels of importance for each cave in supporting bat populations.

The majority of the important bat hibernaculum in Minnesota are localized to southeastern Minnesota where the karst region is found (Nordquist and Birney, 1985). Located in Fillmore County, Mystery Cave is the largest limestone cave system in Minnesota. Thus, this site supports generally constant temperature and humidity levels beneficial to bat populations, particularly species like little brown bats. Bat counts and banding efforts were conducted by the Minnesota Speleological Survey beginning in the late 1970s. Other important sites were mentioned above.

Considering the limited number of suitable hibernacula across Minnesota, combined with concentrations of vulnerable bat populations, and the narrow range of conditions that make caves irreplaceable habitat for bats, caves are extremely important environments to protect for the health and persistence of bat populations in Minnesota. White-nose syndrome threatens cave-wintering bats in Minnesota (see Diseases and Pathogens section below).

Species in Need of Information

Three species of springtails (arthropods in the class *Collembola*) were designated as Species in Need of Information (SNI). These were highlighted for inclusion due to having global conservation status ranks of vulnerable or imperiled (*Arrhopalites dubius* and *Tullbergia hades*, G2, imperiled; *Onychiurus obesus*, G3, vulnerable). These organisms are completely confined to cave systems and adapted to subterranean life. As detritivores, these springtails feed on microbial films, fungi, and decomposing organic matter lining cave pools, playing a vital role in nutrient cycling and supporting food webs in the nutrient-poor conditions of cave ecosystems. Their troglotactic (i.e., obligated to underground living) lifestyle makes them excellent bioindicators of cave habitat integrity as they depend completely on stable subterranean conditions (Christiansen, 1964). They are also likely to be very sensitive to water quality, particularly pollutants from agricultural runoff, which also makes them bioindicators for assessing cave ecosystem health. These species are found in Mystery Cave and associated caves as well as Goliath cave and associated caves at Cherry Grove Blind Valley Scientific and Natural Area. Due to a lack of information about these species, they were designated SNI rather than SGCN.

Invertebrates

Caves provide a unique environment for the evolution of specially adapted species, and due to the disparate distribution of caves across the landscape, many fauna from caves are likely to be significantly rare. However, the taxonomy for many invertebrate groups remains poorly understood, and there is a shortage of active taxonomists. As a result, even though some collections exist, such as a 2022 survey of Forestville/Mystery Cave State Park (by [Zara Environmental LLC](#); Krejca and McDermid, 2022), our understanding of representative

invertebrate communities remains limited and there is not enough information to determine if any cave-dwelling species should be designated SGCN.

Several species of mites in the family Rhagidiidae (Acari), including some troglobitic forms have been reported from caves across the United States (Elliott & Strandtmann, 1971; Zacharda, 1985). These tiny arachnids, which are relatives of spiders and ticks, are a common but often overlooked component of cave ecosystems. In the 2022 invertebrate inventory of Mystery Cave, a member of the Rhagidiidae was recorded for the first time, marking a new addition to the known fauna of the cave and contributing to the broader understanding of subterranean biodiversity in the region (Krejca and McDermid 2022). Like springtails, these animals play vital roles in cave ecosystems as predators, nutrient recyclers, and indicators of environmental health. Their presence likely helps maintain ecological balance in sensitive cave environments, and their loss could disrupt entire subterranean food webs. Additionally, many Rhagidiid mites remain poorly studied, so their conservation protects not only biodiversity but also future opportunities for scientific discovery.

Small, shrimp-like crustaceans known as amphipods (Order *Amphipoda*) were recorded for the first time in Mystery Cave during a 2021 biological survey (Krejca and McDermid 2022). This finding is noteworthy, as a previous study by Montz (1993) specifically highlighted the absence of amphipods in this cave. The newly collected specimens may represent a range extension of known Driftless Area species or possibly a previously undescribed species. Either scenario highlights the ecological significance of the discovery and underscores the need for continued biological exploration and conservation of subterranean habitats like Mystery Cave.

Spiders (*Araneae: Nesticidae*) are often among the most diverse and ecologically important invertebrates found in cave ecosystems, including species that are either fully adapted to life underground (troglobites) or partially so (troglophiles). While it is not yet clear how specialized the spiders found in Mystery Cave are, or whether they represent unique endemic species, the individuals collected during recent surveys are notable and merit further investigation (Krejca and McDermid 2022). Continued research could reveal important insights into the cave's biodiversity, the evolutionary adaptations of subterranean spiders, and the role that spiders play in subterranean food webs.

Bathynellacea are an order of minute, groundwater-dwelling crustaceans typically found in caves, wells, and sometimes within the interstitial spaces of other aquatic environments (Schminke & Noodt, 1988). Despite their global distribution, occurring on every continent except Antarctica, they are notably absent from the fossil record. Their presence is also generally limited in regions previously covered by glacial ice, with only sparse recolonization documented in such areas (Grzimek, 2004). A 2021 collection in Mystery Cave marked the first recorded occurrence of *Bathynellacea* within the park, representing a species adapted specifically to groundwater environments (Krejca and McDermid, 2022). As such, it is restricted to subterranean habitats like caves and aquifers, highlighting the unique and specialized biodiversity found in Minnesota's karst systems.

Between 1989 and 1993, Montz (1993) documented the presence of several copepod and cladoceran species in the Mystery Cave system. Three copepod species representing two orders were identified from the Disappearing River: *Acanthocyclops vernalis* (tentatively identified) and *Eucyclops speratus* from the order Cyclopoida, and *Phyllognathopus viguieri* from the order Harpacticoida. More recently, in 2021, copepods were collected from several locations within the cave system, including Coon (Goliath's) Cave, Old Mystery Cave, Woodchuck Cave, the Big Spring complex, and Seven Springs (Krejca and McDermid, 2022).

Water fleas (order *Cladocera*) were also found in the system. Montz (1993) reported six cladoceran species from the Disappearing River and one additional species from Blue Lake within Mystery Cave. In a 2021 survey, specimens resembling cladocerans were collected from Coon (Goliath's) Cave, Old Mystery Cave, Big Spring, and Seven Springs (Krejca and McDermid, 2022). However, these were not identified to species level. These findings suggest that both copepods and cladocerans are relatively widespread in the Mystery Cave hydrological network and may play an important role in the cave's aquatic food web.

In summary, cave-dwelling invertebrates are highly specialized organisms that play essential roles in nutrient cycling and food web dynamics within subterranean ecosystems. Despite their ecological importance, little is known about their population status, distribution, or specific ecological functions. Many species are adapted to stable, low-light environments and are highly sensitive to environmental changes, making them valuable indicators of groundwater and habitat health. Improved surveillance and monitoring of these invertebrates are critical for understanding and conserving the unique biodiversity of cave ecosystems in Minnesota.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key 'stressors' that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the "Stressors" section in Chapter 1. Species in Greatest Conservation Need.

It is important to note that some of the factors listed as 'stressors' can also be used to advance conservation goals. Broad terms such as 'fire management' reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Crop Production

Commercial agriculture dominates southeast Minnesota where Minnesota has its highest concentration of caves. The use of nitrogen-based fertilizers, herbicides, pesticides, and other chemicals can degrade water quality in cave environments. In karst areas of southeast Minnesota, human influences on groundwater and surface water quality has been documented for decades (Mohring, 1983; Jameson and Alexander, 1994 Kuehner et al., 2025).

The main Goliath Cave sinkhole entrance has been completely filled with sediment from run-off. The SNA is surrounded by farm fields that are tilled. When sediments, including agricultural plant material (like corn

stalks), and manure run off farm fields, it can significantly alter the cave's hydrology and introduce pollutants that threaten the unique underground ecosystem. A recent inventory noted approximately 20 different species of invertebrates found in Goliath Cave (Krejca and McDermid, 2022). These highly specialized species, such as blind isopods, cave beetles, or springtails, have evolved in stable, low-nutrient, low-light environments with little tolerance for change and are therefore threatened by a sudden influx of sediments and pollutants.



Livestock Management

Livestock operations, particularly concentrated feed lots, can threaten caves through pollution. Manure and agricultural runoff can contaminate water sources, which can enter caves through sinkholes, sinking streams, and fractures in bedrock. Caves act as conduits for groundwater and can transport pollutants very quickly downstream. For instance, Mystery Cave has an underground stream that is a meander cut-off of the South Branch Root River. Agriculture and grazing operations are active on the surface above Mystery Cave. The caves' river flows underground two miles and resurfaces at Seven Spring Wildlife Management Area, which is an important trout habitat. Goliath's Cave in Cherry Grove Blind Valley has its' own underground river which flows five miles and resurfaces in Canfield Creek in Forestville/Mystery Cave State Park another important trout habitat. Goliath Cave is surrounded by agriculture and livestock operations.



Mining and Quarrying

Mining, including limestone extraction, can directly destroy caves, eliminating habitat for cave adapted species. Mining operations can also impact cave and karst systems in other ways such as thermal impacts to springs and surface-water streams, changes to water turbidity, and lowering of groundwater and surface-water levels from mine dewatering (Green et al., 2005).



Recreation

During the fall and winter months (September 1 – May 1) when bats are likely to be present in caves, human presence can disturb and wake bats similar to white-nose syndrome symptoms that result in mortality. In addition, recreators can inadvertently spread the causative fungus of white-nose syndrome if shoes and gear taken underground are not sufficiently decontaminated. For these reasons, major bat hibernacula in Minnesota are protected with gated entrances and restricted access for safety of bats and people.

The presence of visitors in caves can also elevate air temperatures. In Mystery Cave, tours cause an elevation in temperature of about 1.5 degrees F every time a group of visitors passes by a set of data loggers over the past five years (Dawn Ryan, DNR, pers. Comm.).



Diseases and Pathogens

As critical habitat for overwintering bats, caves have also been at the heart of white-nose syndrome (WNS), an epidemic that has devastated hibernating bat populations across North America. This disease impacts hibernating bats by disturbing the delicate hibernation cycle, resulting in bats waking from torpor more frequently and burning the fat reserves needed to survive the winter with no insects (Frick et al., 2010). The causative fungus, *Psuedogymnoascus destructans*, is a non-native, subterranean, cold-loving fungus that thrives in the same areas of caves where bats hibernate. The fungus infects tissues of hibernating bats, vulnerable with no immune response while in torpor, and becomes visible white patches on the forearms, ears

or nose of the bat, giving the disease the WNS namesake. The confirmation of WNS in Minnesota during the winter of 2015-2016 has been followed by declines averaging greater than 90% for hibernating bats across the state, particularly for the little brown bat, Northern long-eared bat, tricolored bat.



Water-borne Pollution

One of the most frequently detected contaminants in groundwater and surface waters of southeastern Minnesota is nitrate-nitrogen (commonly referred to as nitrate). Nitrate concentration is commonly elevated in the root zone underlying row-crop agriculture in Minnesota, where concentrations collected from lysimeters average greater than 20 ppm (Kuehner et al., 2020). Nitrate is leached in cultivated row-crop settings following precipitation and snowmelt and moves to shallow groundwater where it migrates to underlying aquifers and streams. Depending on the timing of recharge, nitrate concentrations in shallow karst may rapidly increase or decrease (Barry et al., 2020). In southeastern Minnesota, trout-stream baseflow nitrate concentration positively correlates to the percentage of row crop agriculture in the watershed (Watkins et al., 2013).

Elevated levels of nitrate may indicate that other surface contaminants have the potential to reach an aquifer. Pesticides frequently co-occur with nitrate, with the likelihood of detecting at least one pesticide compound increasing as the concentration of nitrate increases (MDA, 2019; Goedjen et al., 2024). Trout fisheries are also impacted by [nitrate contamination](#) directly as well as fed by underwater sources from caves as noted above.

Southeastern Minnesota has several karst aquifers with different properties. In general, the shallow Galena karst aquifer (which is present across Fillmore County) has high nitrate levels and short groundwater residence time (Kuehner et. al, 2025). Groundwater residence time is the time it takes water to travel from the surface to the point it was sampled; residence time is related to the amount of nutrient removal it provides. The Prairie du Chien aquifer, more common across counties east of Fillmore, has longer residence times and lower nitrate levels.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. Increased flooding events carry pollutants from agricultural run-off, faulty septic systems, and pesticide, which can impact springtails and other animals who live in caves. While many cave-dwelling animals are aquatic, they are adapted to near constant temperatures. In addition, flooding events cause increases in cave temperatures. In Mystery Cave, increases in temperature of close to 10 degrees °F have been detected during floods, and the cave takes months to return to normal temperatures (Dawn Ryan, DNR, pers. comm.).

The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al., 2022; Climate Change in Minnesota). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al., 2022). For more information and resources for climate-adapted management strategies, see the Climate subchapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, to protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN, five strategies were identified:



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.



Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas.



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Examples of conservation actions are presented below grouped by the primary strategy that they implement and tagged with icons for the stressor(s) that they address. Some of the strategies, such as Strategy 3, collaborating with partners, could truly be applied to most broadly and effectively implement all actions. In this case, under Strategy 3 we highlight actions for which collaboration is particularly vital.

Potential Conservation Actions for Caves



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.

Stressor	Action
	Purchase land easements above known caves to help reduce run-off into caves and ground water.
	Increase protection of important cave ecosystems from mining and quarrying.



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience.

Stressor	Action
	Maintain stock limits and reduce capacity of confined feedlots in sensitive karst habitats where caves receive runoff than can contain contaminants.

(no activities were articulated for Strategy 3)



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions.

Stressor	Action
n/a	Encourage cave-related research, such as through the Cave Research Foundation



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN.

Stressor	Action
	Encourage and reward landowners who properly buffer sinkholes, swallets, and other karst features that have direct relationships to groundwater. Promote best practices regarding runoff including: properly seal wells and maintain septic systems; reduce the use of nitrates, pesticides, and fertilizers; increase cover crops, and develop denitrification wetlands.
	Encourage adoption of best practices for cavers and spelunkers to reduce impacts when exploring caves, such as in guidelines set forth by the National Speleological Society .
	Follow decontamination procedures when visiting caves to avoid spreading the causative fungus of White-Nose Syndrome. While effective treatments for widespread containment of the fungus causing white-nose syndrome or prevention of disease have yet to be developed, there are actions that can be taken to protect bats. Preventing the spread of fungal spores between caves and avoiding disturbance to hibernating bats are key methods to reduce further harm. Spores are not visible to the human eye and readily attach to clothing or gear carried into caves. Follow protocol for decontaminating gear after visiting any cave or underground feature, particularly if you travel to multiple caves.

Case Study: Goliath Cave and Cherry Grove Blind Valley SNA

Goliath Cave and Cherry Grove Blind Valley Scientific and Natural Area (SNA) was established in March 2000 after the owner applied for a quarry permit to mine the limestone. Goliath Cave and several other small caves in the SNA would have been mined away and lost, had the state not purchased the property. Goliath Cave includes ancient rock formations dating back to the Ordovician Period (443.7—488.3 million years before present), when what is now Minnesota was located south of the equator and covered in shallow seas. The cave system also provides cold water to trout streams in the area. See [Cherry Grove Blind Valley Scientific and Natural Area](#) for more information.

References for Caves

Alexander Jr, E.C. (1980). An Introduction to Caves of Minnesota, Iowa, and Wisconsin, Guidebook for the 1980 National Speleological Society Convention.

Alexander, E.C., Brick, G.A. (2021). Minnesota Caves and Karst. In: Brick, G.A., Alexander Jr., E.C. (eds) Caves and Karst of the Upper Midwest, USA. Cave and Karst Systems of the World. Springer, Cham.
https://doi.org/10.1007/978-3-030-54633-5_3

Anderson D. (2016). Economic impact of recreational trout angling in the Driftless Area. Report to Driftless Area Restoration Effort. Trout Unlimited.

Barclay, R. M., Lausen, C. L. and Hollis, L. (2001). What's hot and what's not: defining torpor in free-ranging birds and mammals. – Can. J. Zool. 79: 1885–1890.

Barry JD, Miller TP, Steenberg JR, Runkel AC, Kuehner KJ, Alexander EC Jr (2020). Combining high resolution spring monitoring, dye tracing, watershed analysis, and outcrop and borehole observations to characterize the Galena Karst, southeast Minnesota, USA.

Christiansen, K. (1964). Bionomics of Collembola. Annual Review of Entomology. Vol. 9:147-178.
<https://doi.org/10.1146/annurev.en.09.010164.001051>

Elliott, W. R., & Strandtmann, R. W. (1971). New Locality Records for Rhagidia from Mexican and American Caves. *Journal of the Kansas Entomological Society*, 44(4), 468–475. <http://www.jstor.org/stable/25082450>

Evans C.A. (1934). Hibernating Bats in Minnesota. *Journal of Mammalogy* 15:240-241

Frick et al. (2010). An Emerging Disease Causes Regional Population Collapse of a Common North American Bat Species. *Science* 329(1) 679-682

Geiser, F. (1998). Evolution of daily torpor and hibernation in birds and mammals: importance of body size. *Clin. Exp. Pharmacol. Physiol.* 25: 736–739.

Goedjen GJ, Capel PD, Barry JD, Arnold WA. Occurrence and distribution of neonicotinoids and fiproles within groundwater in Minnesota: Effects of lithology, land use and geography. *Sci Total Environ.* (2024) Dec 1;954:176411. doi: 10.1016/j.scitotenv.2024.176411. Epub 2024 Sep 22. PMID: 39317252.

Green, J.A., Pavlish, R.G., and Leete, J.L., (2005). Hydraulic Impacts of Quarries and Gravel Pits, Prepared for the Legislative Commission on Minnesota Resources.

Herak, M and VT Stringfield. (Eds.). (1972). KARST Important Karst Regions of the Northern Hemisphere. Elsevier publishing company.

Jameson, R.A. and Alexander Jr, E.C., (1994). The Waters of Mystery Cave, Forestville State Park, Minnesota, Mystery Cave Resource Evaluation (Groundwater) Interpretive Report.

Knowles B. (1992). Bat Hibernacula on Lake Superior's North Shore, Minnesota. *The Canadian Field-naturalist.* Vol 106(2) 252-254.

Krejca, J., McDermid, K. (2022). Invertebrate inventories in four caves at Forestville/Mystery Cave State Park, Minnesota. Report prepared by Zara Environmental, LLC, for DNR. 51 pages.

Kuehner KJ, Dogwiler TJ, Kjaersgaard J (2020) Examination of soil water nitrate: N concentrations from common land covers and cropping systems in southeast Minnesota karst. Minnesota Department of Agriculture, 31 pp.

Kuehner, K.J., Runkel, A.C. & Barry, J.D. Informing nitrate concentration trends: estimating groundwater residence time in a karstic, multiaquifer system using anthropogenic tracers (Minnesota, USA). *Hydrogeol J* **33**, 167–192 (2025). <https://doi.org/10.1007/s10040-024-02871-2>

Kunz 1982 - Kunz TH (1982) Roosting ecology. In: Kunz TH (ed) *Ecology of bats*. Plenum, New York, pp 1–55

Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>

MDA (Minnesota Department of Agriculture) (2019) Final township testing nitrate report–Fillmore County 2017–2018. Minnesota Department of Agriculture, St Paul, MN, 80 pp.

Mohring EH (1983) A study of subsurface water flow in a southeastern Minnesota karst drainage basin. MSc Thesis, University of Minnesota, Minneapolis, MN, 99 pp

Nordquist, G.E. and Birney E.C. (1985). Distribution and Status of Bats in Minnesota. Final Report Nongame Wildlife Program. Minnesota Department of Natural Resources.

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS. <https://statesummaries.ncics.org/chapter/mn/>

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification

of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434.
<https://doi.org/10.1111/cobi.14434>

Schminke, H. K. and Noodt W. (1988). Groundwater Crustacea of the Order Bathynellacea (Malacostraca) from North America. *Journal of Crustacean Biology*, 8(2), 290–299. <https://doi.org/10.2307/1548321>

Watkins J, Rasmussen N, Streitz A (2013) Nitrate-nitrogen in the springs and trout streams of Minnesota. *Minnesota Groundwater Association Newsletter* 32(3).

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report].
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Zacharda, M. (1985). New Rhagidiidae (Acarina: Prostigmata) from caves of the U.S.A., *Veštno Ak C I eskoslovenskeA Společnosti Zoologické A*, 49, 67-80.

Urban and Other Developed Lands

Habitat Description

This chapter is devoted to the places where we live and work. Developed lands include those urban, exurban, or rural areas for housing, business, and municipal or community uses. These developed areas contain parklands with a variety of land cover types including grasslands, woodlands, forests, lakes, and rivers as well as managed wetlands such as water treatment facilities. Road rights-of-way extend beyond typically developed areas forming a network throughout the state. Areas developed for energy generation and transmission include mines, oil and gas wells and pipelines, nuclear, wind, and solar power generation facilities, and transmission lines. Working lands refers to agricultural lands managed for crop and livestock production, and forest land cultivated for timber harvest (see grassland and forest chapters).

These developed and working lands can also provide habitat for a subset of SGCN that inhabit, visit, or fly through. These are also important places where many actions may be applied to assist SGCN because of the number of willing and interested community members who can contribute.

Habitat Map

For many of the other habitat sub-chapters, we referenced Minnesota's Ecological Classification System, noting Native Plant Community (NPC) Classes or Types in each. However, other developed lands are not represented within the NPC system but are labeled as Developed and Ruderal types within the Midwest Landscape Initiative Midwest Landcover map. Developed is used for areas with less permeable surfaces. Ruderal references vegetative communities that are dominated by non-natives or are otherwise heavily human-influenced. Agricultural lands are also included in this map (see Figure 3.21).

Associated Habitat Types

Developed

Urban – residential, corporate, municipal, manufacturing, etc.

Developed grasslands (examples: urban parks, airstrips, solar arrays)

Developed aquatic habitats (examples: water treatment ponds)

Rights-of-ways (the areas directly adjacent to roads, railroads, and trails)

Mining and Energy

- Sand and gravel mines

- Solar farms

- Wind farms

- Power and pipeline corridors

Data Centers

Agricultural

Cropland

Hayfields (cross-list to Prairies and Other Grasslands)

Pasture and rangelands (cross-list to Prairies and Other Grasslands)

Animal rearing and livestock production facilities

CRP and other restoration lands (cross-list to Prairies and Other Grasslands)

Woodlots and Tree Plantations




Figure 3.21. A map depicting Urban and Agricultural landcover in Minnesota from the Midwest Terrestrial Habitat System (see narrative).

Conservation Context

Conservation within urban and developed landscapes is unique. These landscapes are typically fragmented due to development for housing, business, energy, industry, and transportation or in the case of working lands, agricultural, livestock, or timber production. These are the places that provide services to our communities. Because of their proximity to where we live and work, these lands also represent excellent opportunity locations to engage community members in conservation actions.

Urban conservation efforts over the last 30 years have been successful at identifying tracts of land that contain unique habitat types or those with high ecosystem function and proactively protecting them from development through conservation easements or fee title purchase as open space, natural areas or parks. These lands have become focal points for current conservation efforts as well as drawing the attention of residents to the benefits of those habitats to wildlife, watershed health, and proper ecosystem function. Through local outreach efforts, public engagement, and conservation programs, urban and suburban landowners have grown to understand the benefits of these habitats and implement small scale practices such as pollinator plantings, prairie restoration and the use of native species in landscaping and lawns.

Landowners that manage working lands have access to federal, state, local, and private conservation resources focused on helping them implement grazing management, soil health, and conservation practices that minimize or mitigate potential negative environmental impacts. Agricultural producers have access to and incentives to implement conservation practices such as cover cropping, precision agricultural practices, and various crop management techniques. Livestock managers can implement rotational grazing systems that can enhance soil and ecosystem health and may be tailored to certain habitat types and long-term conservation goals.

Even with all the progress mentioned above, pressure from urbanization continues to threaten working lands, and remaining wildlife habitats alike. Once these lands have been developed, they will never return to their previous state. Future conservation efforts need to capitalize on past and current successes while focusing on partnerships between private landowners, non-governmental conservation organizations, state, federal and local governments that proactively coordinate conservation efforts that aim to mitigate the effects of development on wildlife habitats.

Species in Greatest Conservation Need

Urban and Other Developed Lands provide primary or secondary habitat for 36 animal SGCN (see Table 3.19.) Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently. Plant SGCN were not associated with this habitat type.

Table 3.19. Numbers of Species in Greatest Conservation Need associated with Urban and Other Developed Habitat as either primary or secondary habitat¹.

Species Group	Primary Habitat	Secondary Habitat	Total
Amphibians	0	1	1
Birds	3	6	9
Mammals	2	15	17
Reptiles	0	3	3
Terrestrial Invertebrates			
Butterflies			1
Bees			2
Beetles (terrestrial)			3
Total	5	25	36

1. Habitat associations for insects were not differentiated into primary and secondary habitats and are shown in the total column.

Animals with more general habitat requirements are associated with multiple habitat types, while specialists are associated with one or few. Detailed tables associating each SGCN with the 15 habitats identified in the 2025-2035 SWAP can be found in Appendix C for animals and Appendix D for plants. Examples of selected SGCN are described below. SGCN that use urban or other developed habitats should not be interpreted as requiring this habitat type, rather we acknowledge developed land uses are part of the landscape where SGCN can persist, and conservation opportunities can be highlighted. For example, urban and other developed lands may serve an important role in providing stopover habitat for highly mobile species and migrating birds.

Amphibians

Minnesota's only endangered amphibian, the [Blanchard's cricket frog \(*Acris blanchardi*\)](#), readily exploits aquatic habitats that have been or are significantly altered by human actions. This species appears to express some preference for aquatic habitats with limited aquatic vegetation which are sometimes created by sand and gravel mining and artificial water level manipulation. Once nearly lost from Minnesota, some of the habitats this species has persisted in and recolonized as they recover, include active sand and gravel mines, river barge docking areas, streams in agricultural landscapes and urban parks.

Birds

Several bird SGCN are almost entirely dependent on human-created nest structures. Formerly nesting in large, hollow trees or other natural cavities, chimney swifts (*Chaetura pelagica*) now build their nests almost

exclusively inside of chimneys. Suitable chimneys usually occur in urban areas, often in either large cities or small-town centers. Chimney swifts may also use artificial nest structures created specifically for them. [Purple martins \(*Progne subis*\)](#) are dependent on artificial nest structures built, placed, and maintained by humans. Nest structures must be built following specific guidelines to minimize predation and competition with other bird species. Suitable purple martin nest structures are usually close to water, often placed near lakeshores.

Common nighthawks (*Chordeiles minor*) and [peregrine falcons \(*Falco peregrinus*\)](#) often nest on artificial structures in developed habitats, but both species also nest in native habitats. Common nighthawks will nest on gravel rooftops on buildings, as well as on native rock outcrops and other barren microhabitats. Changes in roofing practices, eliminating many gravel rooftops, may be one factor in common nighthawk population declines. Historically, peregrine falcons nested on native cliff habitats, but during reintroduction efforts were largely dependent on artificial nest boxes placed on bridges, buildings, or other human-created structures, as well as on cliff faces. As the population has increased, some peregrines have returned to native cliffs, but artificial structures continue to be very important nest sites.

Red-headed Woodpeckers (*Melanerpes erythrocephalus*) nest in native habitats such as oak savanna and floodplain forest, but in rural, agricultural regions they commonly nest in small woodlots at rural farmsteads. These farmsteads, particularly in southwestern Minnesota, support a significant red-headed woodpecker population. However, woodpeckers nesting in these areas often forage along roads, where they are susceptible to mortality from being struck by motor vehicles.

Various bird SGCN use urban areas as stopover habitat during migration. Migration has been identified as the primary period determining survival (or mortality) of adult migrating birds (Hutto 1998; Sillett and Holmes 2002). There may be untapped potential for urban habitat restorations (particularly replacement of turfgrass with native plantings) to support migrating birds as stopover habitat.

Mammals

Two SGCN mammals are associated with now human dominated habitat types of grazed pastures, developed grasslands, and road rights-of-way: the [Richardson's ground squirrel \(*Urocitellus richardsonii*\)](#) and [Northern pocket gopher \(*Thomomys talpoides*\)](#). Both can be found in open habitats preferring dry well-drained soils for burrowing. Richardson's ground squirrels, a species of tallgrass and short grass upland prairies, prefer short vegetation as high visibility is important for detecting potential predators and can also be found in grazed pastures and open urban areas in western Minnesota. Broadly distributed across western North America, the Northern pocket gopher can be found in a range of habitats. In Minnesota, however, it is limited to suboptimal habitat associated with agricultural fields and roadways in the extreme northwestern part of the state. Competition with the larger and more numerous plains pocket gopher (*Geomys bursarius*), combined with bounties for the species, may be responsible for its restricted distribution here.

All seven SGCN bat species frequent urban and other human-altered habitats. They may be observed foraging in parks, along boulevards and under streetlights. Cave bats, such as [little brown myotis \(*Myotis lucifugus*\)](#) and [big brown bat \(*Eptesicus fuscus*\)](#), also roost in developed spaces such as bridges and attics during summer and hibernate in tunnels, mines and buildings during winter.

Reptiles

In the upper Midwest, [timber rattlesnakes \(*Crotalus horridus*\)](#) live in the hills and valleys of the Mississippi River drainage. South to west-facing bluffs or hillsides with rock outcrops are used as overwintering sites. Due to cold winter weather, most snakes are inactive and in their dens from mid-October to mid-April. Nearby forests and grasslands are used as summer feeding grounds where snakes eat mostly small mammals. In summer, snakes like to be anywhere there is a food source for them. In some cases, this can include residential yards with bushes, rocks, mulch, a water source, bird feeders or any other feature that attracts rodents; all these features can also attract snakes. Another SGCN snake, the [North American racer \(*Coluber constrictor*\)](#), also occurs in the southeastern region of MN and uses woodland margins and field edges as the preferred summer habitat.

Many turtles in MN including the [wood turtle \(*Clemmys insculpta*\)](#) and [Blanding's turtle \(*Emydoidea blandingii*\)](#), can be found using developed land types, especially with respect to nesting or dispersal between wetlands. Roadways, gardens and yards often provide a loose dirt substrate suitable for digging their nests.

Invertebrates

The [ghost tiger beetle \(*Cicindela lepida*\)](#) and two other SGCN tiger beetles persist and even thrive in human dominated habitats that create or maintain open soil like sand and gravel mines. Tiger beetles are dependent on bare or sparsely vegetated soil for the development of their larva and to facilitate their hunting strategies. Soil stabilization, invasion of invasive plants (which may readily exploit bare soil habitats) and artificial water level manipulation can limit the availability of the habitats this group of insects require.

The [rusty patched bumble bee \(*Bombus affinis*\)](#) is a generalist forager and relies on consistent food sources throughout the growing season from a wide array of flowers. Worker bees have been observed in a variety of habitats, including forests, wetlands, grasslands, roadsides, agricultural fields, and residential lawns and parks (Colla and Packer 2008). Rusty-patched bumble bee occupancy is positively associated with developed land uses (Boone 2023), and this bee can be found foraging on native and non-native flowering plants in urban and suburban gardens and parks within its range.

The monarch butterfly (*Danaus plexippus*) is another insect SGCN that can thrive in developed habitat types such as roadsides, fields, gardens, wetlands and parklike settings, if milkweed is present for the caterpillar phase of their life cycle. During migration, adult monarch butterflies forage for nectar on a wide range of blooming plants. Minnesota is within the reproductive and migratory ranges for the eastern migratory monarch population that overwinters in Mexico.

Plants

Although none of the plant SGCN were associated with Urban and Other Developed Lands as their single primary habitat, open spaces within developed and working lands can provide valuable habitat for SGCN plant species and act as connections to more intact habitat. Even altered or fragmented habitat can retain the necessary habitat characteristics required by SGCN to persist. These areas are often under surveyed, and the value of the habitat is not recognized until a development action is proposed.

[Lance-leafed violet \(*Viola lanceolata*\)](#) is a prime example of a species that is frequently found persisting in altered or fragmented habitat. Lance-leafed violet is a pioneer species that does not compete well with larger, later successional species. It requires some disturbance to its habitat to be successful. In some cases, anthropogenic disturbance that removes competing vegetation and releases the seed bank can be beneficial to the species. However, development remains the biggest threat to the species and was the basis for listing the species as State Threatened in 1995. A large percentage of Minnesota's *Viola lanceolata* element occurrences (EOs) are found in the Anoka Sand Plain in the north Minneapolis-St Paul metro area. The north metro has and likely will continue to have a great amount of development pressure which results in the destruction of lance-leafed violet habitat and the take of individual plants. The loss of lance-leafed violet from this part of the state would greatly reduce its range and ability to survive in Minnesota.

Projects such as pollinator gardens, rain gardens and conservation plantings often use Minnesota plant SGCNs. Examples include [rattlesnake master \(*Eryngium yuccifolium*\)](#), [beachgrass \(*Ammophila breviliqulata*\)](#) and [plains wild indigo \(*Baptisia leucophaea*\)](#). These species are SGCNs in Minnesota because they are believed to possess unique genetic composition that allows them to survive at the extreme edge of their natural range. Knowing the origin of seeds purchased commercially is important as there is concern that plants imported from other regions may hybridize with native strains.

Primary Stressors in this Habitat

Throughout Minnesota, habitats have been lost and degraded due to pressures associated with human settlement, subsistence, livelihoods, and recreation. Indeed, habitat loss or alteration remains the primary threat to most, if not all, SGCN. In this section, we identify key ‘stressors’ that may continue to contribute to habitat degradation and loss. The list is adapted from a globally recognized threats lexicon developed by the International Union for the Conservation of Nature (Salafsky et al., 2024). For additional details, see the “*Stressors*” section in Chapter 1. Species in Greatest Conservation Need.

It is important to note that some of the factors listed as ‘stressors’ can also be used to advance conservation goals. Broad terms such as ‘fire management’ reflect the dual nature of these factors as they may function as stressors in some contexts while serving as valuable conservation tools in others. For example, an intense wildfire following prolonged fire suppression may cause significant stress for the habitat and species affected, while prescribed fire, when planned appropriately, can enhance ecosystem health and resilience.

Information about a subset of primary stressors specifically impacting this habitat is included below, followed by a set of conservation actions addressing those stressors.



Development

Habitat loss and fragmentation, loss of movement or connectivity corridors and increased competition from invasive species are more prominent in developed landscapes. Yet the remaining patches of wildlife habitat or natural areas within developed infrastructure can provide benefits for SGCN. There are 3 sub-categories of development: commercial and industrial areas, housing and urban areas, tourism and recreation areas.

Bird collisions with glass claims the lives of hundreds of millions of birds each year in the United States. It is second only to domestic cats as a source of mortality linked directly to human action. Glass is dangerous for strong, healthy, breeding adults, as well as sick or young birds; it can have a particularly serious impact on populations (Sheppard and Phillips, 2015). Bird glass strikes often happen in the day and are due to reflective glass mimicking sky or clear glass not being recognized as an obstacle. While these collisions are often documented in more urban settings and associated with large glass skyscrapers, even small windows in residential settings can be high strike hazards.

A separate but related issue is artificial lighting which can exacerbate the threat of collisions. Bright lights at night have been documented as causing mortality events of nocturnal migrants, especially during storms or low ceiling events. Becoming disorienting, the birds either collide with lighted structures or circle for hours leading to exhaustion and fall out.

A known threat associated with developed areas is unsupervised domestic and feral cats which can pose a serious threat to local wildlife populations. According to Loss et al. (2013) an estimated 164 million cats reside in the United States, with 30-80 million unowned (barn cats, human-fed strays, feral). The impact of free-ranging cats has often been downplayed with comparison to anthropogenic threats, like habitat destruction,

pollution, and direct collisions with man-made structures (Loss et al. 2013). Yet, studies have found that cat-induced wildlife mortality, especially for birds, is comparable or can exceed additional mortality factors. It's been estimated that just in the contiguous United States cats kill between 1.3 and 4 billion birds annually, with most mortality (69%) caused by un-owned cats (Loss et al. 2013). For mammals, Loss et al. (2013) estimated that cats are responsible for the mortality of 6.3-22.3 billion mammals annually, with 89% caused by un-owned cats. Although there is no statewide estimate for domestic cat populations in Minnesota, Lepczyk et al. (2010) highlighted the steady increase in domestic house cat populations in the United States. Current research is limited in the United States, however there may also be significant impacts on reptiles and amphibians. An Australian study, for example, found that 609 million reptiles were killed annually by cats (Stobo-Wilson et al. 2022). Generally, free-ranging cats can cause mortality, and population declines of native wildlife through direct predation, injury, nest failure, and indirect mortality through spreading disease and harassment which may lead to behavioral changes that can reduce prey species fitness.



Crop Production

In the Midwest there is increasing pressure to produce greater quantities of food, livestock feed and fuel, while better protecting environmental quality. Key environmental issues in this region include water contamination from the runoff of nutrients, soil degradation from the aerial and ground applications of herbicides and pesticides, increasingly intensive agricultural practices, and a lack of non-agricultural habitat to support diverse communities of native plants and animals. Projected changes in climate, which include increases in the proportion of precipitation coming from extreme events, are an increasing concern, making soil and water conservation in existing cropping systems more difficult (Liebman et. al. 2013).

According to the Minnesota Department of Agriculture's [2023 Minnesota Agricultural Facts and Stats](#), MN is ranked 6th in the nation for agricultural production, 5th for crop production and 8th in livestock production. MN is also ranked 10th in the nation for the number of organic farms. Clearly crop production, and agriculture in general, is foundational to Minnesota's economy. In the action portion of this chapter, we highlight some opportunities to focus on the environmental quality opportunities within the agricultural landscape.



Tree Plantations

Tree plantations convert native landscapes to lands planted for the main purpose of timber harvest. The threats tree plantations pose vary depending on how different the tree plantation is from the native landscape. The threat of tree plantations to SGCNs is especially apparent in the Anoka Sand Plain Subsection. Prior to European settlement the Anoka Sand Plain was primarily an assemblage of fire dependent native plant communities including oak savanna and prairie, both of which included large openings and barrens. These open spaces have been lost due to urban development, agriculture, and tree plantations. The Anoka Sand Plain Subsection Forest Resource Management Plan ([Anoka Sand Plain SFRMP](#)) highlights the threats tree plantations pose to the area including the loss biodiversity and loss of ecologically intact landscapes. The Anoka Sand Plain SFRMP recognizes the need for the removal of tree plantations and restoration of the subsection to oak savanna and prairie.



Livestock Management

In MN livestock production includes poultry/eggs, dairy, cattle/calves, and hogs as described in the [Minnesota Livestock Profile: 2023](#). [The key issues of concern regarding the environmental impacts of livestock](#)

[production are overgrazing, and runoff from livestock impoundment and production facilities. Both concerns can affect soil health and water quality, as well as watershed function.](#) Overgrazing, if left unchecked, can reduce diversity in vegetative cover and structure, and have severe negative impacts on biodiversity, ecosystem function, soil health, and forage production.



Mining and Quarrying

Mining and quarrying can permanently destroy large areas of key habitats, including aquatic habitats as sand mining expands to the beds of public waters.



Wind and Solar Energy Infrastructure

According to [Our Minnesota Climate](#) “The amount of energy our state produces from renewable resources like solar power and wind has risen 60% over the past decade. Today, renewable energy accounts for 28% of Minnesota’s electricity generation, with 52% of our energy coming from carbon-free sources like renewables, nuclear, and hydropower.” These successes in renewable energy production can be complex in the context of wildlife and habitat impacts. Direct impacts to birds and bats are a particular concern, especially during migration, with wind turbine collisions documented as a high source of mortality for both groups. Loss and his colleagues (2013b) estimated between 140,000 and 328,000 birds are killed annually at wind turbines across the contiguous United States; mortality increased with the height of the turbine hub. Foliage and tree-roosting bats, such as hoary bats and eastern red bats, also are frequently found dead below wind turbines (Arnett et al. 2008). Although variable in timing, most mortality occurs during late summer and fall, corresponding with migration and mating seasons. Estimates place the loss at approximately 888,000 bats per year (Smallwood 2013). To date, lighting and noise appear to not influence the level of mortality.

Similarly land-use impacts in and around solar arrays can negatively impact native plant communities (predominantly prairie) and rare features. Taking lessons from traditional power infrastructure and development, we can be mindful to wildlife considerations in the siting and design of emerging alternative energy resources.



Roads, Trails, and Railroads

Animals being struck and killed by vehicles has been recognized as a global threat to wildlife and human safety and presents significant economic and conservation challenges (Balčiauskas 2025). Road mortality due to collisions with cars is one of the leading contributors to decline for some species groups. According to a study synthesis and review in 2009, amphibians and reptiles tended to show negative effects in association to roads. Birds showed mainly negative or no effects, with a few positive effects for some small birds (e.g. seeking food, water, grit or roadside perches along utility lines and fences) and for vultures (feeding on roadkilled animals). Small mammals generally showed either positive effects or no effect, mid-sized mammals showed either negative effects or no effect, and large mammals showed predominantly negative effects (Fahrig & Rytwinski, 2009). Animal life history and behavior plays a role in determining the impacts of roads and transportation corridors on their survivorship. Some species, for example, are attracted to roadways for access to resources, such as carrion or minerals, therefore increasing their potential for direct collisions (Jacobson et. al., 2016).

In addition to direct mortality, roads, trails and railways can have many impacts which include the creation of barriers, habitat fragmentation, isolation of local populations, noise pollution that interferes with behavior and reproduction (Parris, 2015), altered hydrology (runoff), and water pollution. These corridors also can facilitate the establishment and spread of invasive species, including noxious weeds; this, in turn, can lead to the increased application or roadside spraying of herbicides (Forman and Alexander 1998). Pollutants from de-icing salts, vehicle wear and tears, and the legacy of leaded gasoline are present in Minnesota roadside soils and plants (Mitchell et al. 2020, Shephard et al. 2022).

Roadside mowing is necessary for maintaining sight lines for safety. However, frequent mowing reduces the ability for monarch butterflies to successfully reproduce on milkweed that is often present in that habitat (Hopwood et al. 2015). When road rights-of-way are planted in nonnative plants, there is a missed opportunity to provide native plants on the landscape, which can be used by a wide variety of wildlife, particularly insects (Mitchell et al. 2024).



Utility Corridors

Utility corridors contain both above ground electricity transmission lines as well as pipelines for oil and gas. These corridors and their associated infrastructure can cause local-scale and landscape-scale disturbances. Locally, there can be soil disturbance, compaction, displacement of rare plant species, introduction of invasive species, disturbance to groundwater and surface water flows, and vegetation management activities including use of herbicide, mowing, and woody removal. Bird collisions are problematic, particularly for some transmission line designs, and in wetlands that support waterfowl and other waterbirds. More broadly, utility corridors fragment habitat of all types which can be an added stressor for reptiles and amphibians. Among mammals, DNR staff have found that pine martens require narrow corridors, with few disturbances. On the other hand, grassy, open corridors can attract predators that find hunting prey in open areas can be more efficient.



Hunting and Collecting Animals

Unlicensed hunting or poaching is illegal and a threat to some species groups more than others. Amphibians, turtles and other reptiles are subject to collection more prominently than other animal groups due to their predictable life-history and slow movements. More education is needed to foster appreciation and connection with wildlife without the need to take them out of their environment and into our homes.



Gathering Plants and Fungi

Commercial harvest of ginseng (*Panax quinquefolius*) and goldenseal (*Hydrastis canadensis*) is controlled by legislation, but foraging for personal use is not. Guidelines for where and how to forage responsibly are needed. Several native plant SGCNs are also known to be taken directly from the wild for use in urban gardens. This is especially true for orchids in the genera *Platanthera* and *Cypripedium* which are very difficult to grow from seed and are not available commercially. These species often occur in small, fragile, remnant habitats that cannot withstand the removal of large, reproducing individuals.



Timber Harvest

Timber harvest in fragmented landscapes such as those dominated by human use can further fragment, eliminate or reduce the availability of required habitats for SGCN. Management such as timber harvest must account for a modern landscape often dominated by islands of habitat among a matrix of inhospitable human systems. While timber harvest is often a temporary loss of habitat until trees re-grow, islands of trees in an urban matrix leave displaced wildlife nowhere to persist for the 40-100+ years most forests take to recover. On the other hand, urban areas often lack timber harvest which can affect the populations of early successional species, such as the chestnut-sided warbler (*Setophaga pensylvanica*). Where there are opportunities, homeowners, local governments, and commercial landscape operations should be encouraged to plant native tree species adapted to local environments.



Fishing

Fishing creates and fosters a connection to nature which will be required for its persistence. Fishing pressure can be high in urban and developed areas just based on sheer number people and easy access. However, anglers should employ good stewardship practices to avoid harming the resources they love. Numerous studies have documented the impacts to waterbirds and fish-eating raptors of consuming lead-based fishing tackle and lead sinkers.



Recreation

Minnesotans are known for their love of the outdoors. However, recreational overuse can occur in urban and developed habitats. High recreational use can threaten sensitive plant communities and the wildlife they support. Intense use of popular hiking areas may reach a threshold where certain rare plants are trampled, and the level of foot traffic is high enough to degrade the plant community. Foot traffic also introduces the risk of spreading invasive species. The greater recreational use in each area, the higher the likelihood of invasive species introduction on the site.

Moving wood for campfires can inadvertently move pests such as emerald ash borer in the firewood, so firewood is not to be moved across the state.

Boaters and beach goers can limit turtle nesting by compacting and overusing sandy soil habitat that turtles require for nesting. Boats can also impact waterbirds such as loons and eagles and cause erosion of lakeshore habitats through their wake action.



Fire Management

Fire management in urban and other developed settings is not like fire management in other non-developed habitat types. Human health and safety are the top priority and fire prevention practices are paramount. Impacts of fire to SGCN in this landscape focus more on controlling unintended fires due to high human activity (accidental, electrical, arson) that may impact SGCN that have adapted to human structures such as chimney swifts or bats; these species can be directly impacted by catastrophic or accidental fires in developed settings. The use of prescribed fire as a management tool does occur in agricultural and forested lands, more information can be found in those habitat sub-chapters.



Dams and Water Management

Ditching, channeling and draining wetlands to increase the amount of arable land was a major contributor to the loss of Minnesota's wetlands and wet prairies over one hundred years ago.

Today, the loss continues at a smaller scale and is often impacted directly by commodity prices. Farmers are more likely to make such conversions when commodity prices are high and the potential return on their investment in the cost of tiling is much higher.

Water level manipulation (draw downs, flooding, water level maintenance) are common management tools in human dominated landscapes. Wide varieties of SGCN depend on these human dominated aquatic habitats and can be killed or harmed by water level manipulation that does not account for their persistence.

The maintenance of navigable channels in navigable waterways in Minnesota results in the disruption of benthic aquatic habitats as well as the upland habitats upon which the dredged material is stored and moved. While this activity is necessary to maintain commerce along Minnesota's navigable waterways plans should always account for and seek to minimize effects on SGCN and their habitats through informed, long-range planning.



Invasive Species (Problematic Non-native Species)

Changes in habitat can give invasive species an advantage. Moving soil and equipment around can spread invasive species such as invasive plants and earthworms. Some invasive plant species have been introduced purposefully as ornamental plants for yards and gardens. They then spread from the urban and developed areas to natural areas. Jumping worms (*Amyntas* and *Metaphire* species) can have impacts to yards and gardens as well as to natural areas.



Problematic Native Species

Residents in urban, suburban and even rural areas can easily identify a list of problematic native species that regularly visit their properties and can cause some damage, especially to landscape plantings, vegetable gardens, and building structures. The list includes such species as white-tailed deer, coyotes, raccoons, rabbits, squirrels, chipmunks and woodpeckers. Once the culprit is accurately identified, the majority of these problems can be dealt with by implementing non-lethal actions that exclude, repel or deter wildlife. Numerous websites provide ample advice to homeowners.



Water-borne Pollution

Nonpoint source pollution, including phosphorus, nitrogen, sediment, bacteria, and other contaminants generally results from land runoff, precipitation, atmospheric deposition, drainage, seepage or hydrologic modification. As runoff flows, it picks up and carries away natural and human-made pollutants, depositing them into lakes, rivers, wetlands, and groundwater, and estuaries. ([EPA NPS Webpage](#)) Water quality measured by nutrients and contaminants has been shown to decline with higher agricultural or urban development in the watershed. Some key impacts include excess algal growth, reduced plant diversity, and reduced diversity in fish and invertebrate communities. ([DNR Water quality non-point sources](#))

A significant example of non-point source pollution is chloride. It enters Minnesota's lakes, streams, wetlands, and groundwater from a variety of sources, predominantly road salt use, fertilizers, and wastewater

treatment. Chloride is toxic to aquatic organisms and its presence in high concentrations can alter the function of aquatic communities. In turn, individual populations and communities may become too small or impaired to perform an essential ecosystem function. Up to 70% of road salt applied on Minneapolis–St. Paul metro roads ends up in groundwater aquifers and nearby lakes, many of which exceed regulatory limits for chloride concentrations. Potassium acetate, a de-icing alternative poses its own issues (Gulliver et al. 2022). MPCA (2020a, b) discusses impacts to plants, wildlife, and ecosystems.

Lead in fishing tackle and ammunition can become toxic to wildlife species through eating contaminated fish or wildlife, such as has been well documented in some species such as common loons and bald eagles.



Air-borne Pollution

Experiments under controlled greenhouse conditions have shown that increased levels of atmospheric carbon dioxide may favor the competitiveness of woody plants and cool season invasives by stimulating photosynthesis and growth. It may also reduce mild to moderate abiotic stress in woody plants if root growth and nutrients are not limited.

Nitrogen levels in the atmosphere are driven by natural processes as well as transportation and agricultural sources, primarily fertilizer (Burke et al., 2002; Wall and Pearson, 2013). A study indicates that increased atmospheric deposition of nitrogen reduced plant species numbers by 17% by increasing soil acidity and increasing above ground production of certain plants that reduces light availability to other plant species. This can lead to competitive exclusion. This suggests that chronic but low-level nitrogen deposition may have a greater impact on diversity than previously thought (Clark, Tilman, 2008).



Light and Noise Pollution

Nearly half of the United States land surface has light-polluted skies (Falchi et al. 2016). Sources of light pollution at night include interior and exterior lights on homes, buildings, billboards, parking lots, and streets as well as lighted sports fields, industrial plants, and airports (DarkSky International 2025a). Studies also point to the ecological impacts created during the daytime by polarized light that reflects off surfaces such as asphalt roads, glass panes, and solar panels (Chock et al. 2020; Horvath et al. 2009). Because most of the nation's large urban areas (with multiple artificial light sources every night), are embedded in the Temperate Deciduous Forest biome, biota occupying temperate forests are the most vulnerable to these impacts (Aubrecht et al. 2010).

Impacts of light pollution include disrupting the behavior of amphibians whose breeding calls are delivered at night, drawing nocturnal insects to artificial lights that increase their predation risk, and disorienting birds that use environmental cues to guide their nighttime migration (DarkSky International 2025a, 2025b). In addition, the phenology, growth rates and biomass of plants are impacted (Bucher et al. 2023). Bright day and dark night cycles synchronize the internal clocks of plants and animals, and alterations to these cycles can impact hormone production, reproductive behavior, neural activity, and metabolic functions (Bumgarner and Nelson 2021).

Noise pollution from aircraft, vehicles, boats, and other human causes is pervasive. Anthropogenic noise was detected in 36% of national parks surveyed (Buxton et al. 2019), and 12% of wilderness areas had anthropogenic noise levels 3 decibels higher than levels predicted to occur naturally (Buxton et al.

2017). Exposure to increasing noise levels can change the spatial distribution and movement patterns of wildlife, cause avoidance of feeding and nesting areas, and interrupt sleeping patterns and communications (Kok et al. 2021). For plants, noise can reduce seedling recruitment (Phillips et al. 2021).



Changes in Temperature related to Climate

Minnesota has experienced a clear warming trend over the past century. Between 1895 and 2020, average statewide temperatures increased by 3.0 degrees Fahrenheit (°F; [Climate Trends](#)). This warming has become more pronounced in recent decades and during the winter months. Since 1985, average winter temperatures in Minnesota have risen by 5.4°F, with average winter low temperatures increasing even more significantly by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shortened season of snow cover and a reduction in lake ice duration by 10-14 days over the past 50 years ([MPCA 2025](#)). Furthermore, these shifts in thermal regimes are ecologically significant. Many species are adapted to narrow temperature ranges, and such rapid changes can result in increased thermal stress, the spread of invasive species, and heightened disease and pathogen risks (Ratcliffe et al. 2025).

Looking ahead, this warming trend is expected to continue. By mid-century (2040-2059), Minnesota's average annual temperature is projected to rise by an additional 3.8 - 4.5 °F, depending on future greenhouse gas emissions scenarios (Liess et al. 2022; [Climate Change in Minnesota](#)). Climate change does not act in isolation, interacting with invasive species dynamics, land-use change, and shifts in water quality and quantity, compounding ecological impacts (He et al., 2019; Finch et al., 2021). For additional context and resources, refer to the Climate sub-chapter in Chapter 6. Implementation.

Green spaces utilized by SGCNs can help mitigate the urban heat island effect, which is intensified by the high density of impervious surfaces in urban environments (Purohit, 2025). A Heat Watch report conducted in July 2024 for the Twin Cities modeled the distribution of urban temperatures, confirming patterns observed in previous studies: the hottest areas contain more impervious surfaces, while the coolest areas benefit from shade, vegetation, or nearby water bodies ([Urban Heat Island Study | Ramsey County](#)). Preserving and enhancing green spaces for SGCNs not only supports biodiversity but also provides co-benefits for human health and well-being as urban temperatures continue to rise.



Changes in Precipitation and Hydrology related to Climate

From 1895 to 2020, Minnesota's average annual precipitation increased by 3.4 inches ([Climate Trends](#)). The state has also seen a notable rise in the frequency and intensity of heavy precipitation events. Since 2000, very heavy rains (6 inches or more in a single day) have occurred two to three times more frequently than during the 20th century (Williams-Sether & Sanocki, 2025; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). These extreme events have led to a corresponding increase in flooding, which can disrupt ecosystems, human infrastructure, and water quality (Williams-Sether & Sanocki, 2025).

Future projections indicate continued increases in annual precipitation, especially during the winter and spring months, which are likely to exacerbate flooding risks. Increased frequency of heavy rainstorms leads to more frequent high-flow events in streams, often carrying elevated sediment loads from sheet flow and erosive forces. These processes can alter channel morphology, including downcutting or channel incision. During flooding, streams passing under roads may overtop and deposit sediments that fill culverts. Subsequently, as

channels downcut and flows decrease, these culverts can become perched above the water level, creating barriers that impede the passage of fish and other aquatic wildlife.

The same climate models also forecast an increase in late summer drought events, underscoring the variability and unpredictability of hydrologic patterns under a changing climate ([Climate Change in Minnesota](#)). By mid-century (2040-2059), average annual precipitation is projected to increase by up to 1.2 inches, depending on emissions scenario (Liess et al. 2022; [Climate Change in Minnesota](#)). This seemingly counterintuitive pattern—wetter winters and springs, punctuated by hotter, drier late summers—has profound implications for water availability, wetland health, soil stability, and species dependent on seasonal hydrologic cycles (Runkle et al. 2022). For more information and resources for climate-adapted management strategies, see the Climate subchapter in Chapter 6. Implementation.

Priority Conservation Strategies

To implement the Habitat Goal of this Plan, to protect and enhance the resilience, function, and ability of habitats to support biodiversity, especially for SGCN, five strategies were identified:

-  Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.
-  Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience
-  Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas
-  Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions
-  Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Examples of conservation actions are presented below grouped by the primary strategy that they implement and tagged with icons for the stressor(s) that they address. Some of the strategies, such as Strategy 3, collaborating with partners, could truly be applied to most broadly and effectively implement all actions. In this case, under Strategy 3 we highlight actions for which collaboration is particularly vital. If an action does not relate to a stressor, we mark this with a ‘n/a’ in the stressor column.

Potential Conservation Actions for Urban and Other Developed Lands



Strategy 1. Protect, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits, particularly across the Conservation Action Network.


Stressor	Action
	Promote comprehensive plans that prioritize and protect open space, habitat connectivity, and higher density developments. New development projects should plan for the maintenance of urban wildlife habitat.
	Conservation partners can collaborate to provide landowners information regarding best management practices for green/open space in urban and suburban areas. Urban/municipal planners can focus on developing plans that will protect sites that are important to SGCN plants. Management actions might include prescribed burning, invasive species removal, population augmentation, etc.
	To minimize the reduction of natural lands as habitat, encourage siting of solar arrays on top of buildings and parking areas. Support the co-development of land, such as wind turbines sitting on cropland.
	Consider wildlife-friendly solutions such as building new lines within the same right of ways as existing lines to minimize overall impact footprint, orient transmission lines horizontally to limit transmission line heights and minimize collision potential, and/or add lighted markings or ultra-violet lighting to make transmission lines more visible to birds– which has found to lower collisions by 88% .



Strategy 2. Restore, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience

Stressor	Action
	Encourage conversion of turfgrass to more diverse plantings that benefit butterflies, bumble bees, migrating birds, and other wildlife. Monitor SGCN use and study how effectiveness varies with size, floral composition, landscape context, and other variables. Evaluate benefits while considering threats such as window collisions, pesticides, and light pollution.

Stressor	Action
	Minimize outdoor use of insecticides (including those for home and garden use), especially those marketed as long-lasting treatments and those with broad spectrum (nontargeted) effects on many insects. Minimize the use of herbicides that reduce native forb abundance.
	Use native seed mixes to restore areas after improvement work along roadside corridors; include blooming plants to benefit pollinators, such as milkweed for monarchs (Mitchell et al. 2024; Darst et al. 2024). Pollinators and birds will benefit by managing the area to produce wildflower blooms by less frequently mowing the backside of the right-of-way.
	Mowing road rights-of-ways is critical to maintain sight lines for safety. Yet, beyond these zones less frequent mowing (once per summer optimally) enables better outcomes for blooming flowers to provide nectar to pollinators and for butterfly eggs and larva to survive.
	Broader, natural-bottomed stream crossings are a climate-smart solution when replacing culverts that have become filled with sediments or are no longer aligned with stream flows. Identify culverts where larger capacity, well-signed road crossings can provide greater resilience with future fluctuations in rainfall events and stream levels.
	Avoid bisecting wetland or aquatic habitats with roads without including barrier fencing to keep wildlife off roads. Wildlife mortality occurs most often where roads bisect aquatic habitats.
	Advocate for non-reflective coating on solar farms to reduce waterfowl collisions and insect reproductive sinks when reflective panels appear as water to passing wildlife.
	To minimize bird SGCN collisions into utility lines, participate in site selection to ensure wildlife collision concerns are represented, including siting away from areas where birds are accessing wetlands. Where bird strikes have occurred implement diverters. Reduce chemical vegetation control methods such as broadcast use of herbicides.
	Move control of vegetation, such as mowing and brush removal, to outside the bird breeding season (May-July), turtle nesting season (June), and migratory seasons (May/June, Sept/Oct) to minimize impacts to SGCN.
	Provide recreational users information on potential boating disturbances to loons and eagles, and suggest mitigation measures such as buoys to delineate nesting areas and no-wake zones.
	Avoid stormwater maintenance (drawdown and in-water work) in winter; these are critical winter habitats for a variety of aquatic species (turtles, frogs, fish, waterfowl). Winter maintenance work generally precludes wildlife use of these sites.

Stressor	Action
	Design invasive or nuisance species treatment approaches with protective measures for native species and SGCN. For example: avoid drawing down aquatic habitats in the winter to temporarily reduce curly-leaf pondweed without planning for avoidance of native wildlife (turtle, fish, etc.) mortality. Reduce chemical vegetation control methods such as broadcast use of herbicides. Time control of vegetation such as mowing and brush removal to outside the bird breeding season (May-July), turtle nesting season (June) and migratory seasons (May/June, Sept/Oct) to minimize impacts to SGCN.
	Encourage the use of lead-free fishing tackle. Fish, waterfowl, and waterbirds can consume lead tackle that can lead to impaired immunity and neurological function, bioaccumulation, and death. Support policies to reinforce the importance of adopting lead-free tackle.
	Suggest wildlife friendly erosion control products. Plastic erosion control blankets should specifically be avoided as they entangle and kill wildlife and introduce microplastics into the environment. See DNR Erosion Control .
	Support maintenance of vegetative buffers around riparian and drainage areas to reduce sedimentation and pollutants making their way into streams and lakes.




Strategy 3. Collaborate with conservation partners and landowners to enhance conservation delivery, particularly in the Conservation Action Network and Conservation Opportunity Areas

Stressor	Action
	Collaborate with MnDOT, Counties, Townships, and Cities to integrate road crossings for vulnerable wildlife (such as natural-bottomed underpasses for turtles and fish). Study use patterns to determine the most effective placement, work collaboratively to write funding requests, and plan adequate timeframes for project permitting and delivery. Advocate for wildlife passage to be an integral part of building and maintaining roads and for reduced application of de-icing salts to reduce salinization of watersheds. The design and maintenance of bridges should be done with wildlife in mind (e.g. bats, cliff swallows).
	Engage in transportation corridor planning projects to minimize road and motorized trail building in key movement areas for wildlife such as wood turtles and American goshawks.
	Implement best practices for wildlife-friendly outdoor lighting, such as turning off lights when they are not needed and employing lights in the warmer part of the spectrum. For more detailed information, refer to solutions advised by Darksky , the Xerces Society , and National Audubon's Lights Out Program .



Strategy 4. Monitor SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to natural disturbances, conservation actions, and climatic conditions

Stressor	Action
	Monitor the movement of invasive plants through road, trail and utility corridors to rapidly address problem areas as needed (see EdMaps). Research best practices regarding time of year for various maintenance activities to support native plants and reduce the spread of invasive plants.
	Continue research to help develop effective strategies to improve wildlife passage/connectivity and reduce wildlife-vehicle collisions.



Strategy 5. Connect to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

Stressor	Action
	Advocate and disseminate information regarding the safe and healthy option of maintaining domestic animals inside the home or in controlled settings. Feral and outdoor cats have a measurable impact on some vertebrate populations (songbirds, small mammals). Existing management strategies such as trap-neuter-release have unfortunately proven unsuccessful in controlling cat populations and their impacts on wildlife species. Additional steps to minimize domestic cat impacts include the disbandment of cat colonies through adoption, eliminate feeding of free-ranging cats, pet cat sterilization, and public education. All pigs should remain under containment to prevent the establishment of feral swine in Minnesota.
	Provide materials regarding the appropriate disposal of unwanted pets and responsible pet ownership. Each year there are dozens of abandoned pet reports made with the DNR – from red-eared slider turtles dumped in state parks to pet alligators released into the Mississippi river. No unwanted pets, plants, or bait may be released into Minnesota waters.
	Work with developers of private and commercial properties to implement actions that address window-strike bird collisions. Increase public outreach/education on collision

Stressor	Action
	prevention. The American Bird Conservancy and Audubon's Bird Friendly Buildings initiative provide more details.
	Collaborate with counties, BSWR, NRCS, and other farm conservation organizations to share information regarding SGCN and encourage the use of rowcrop conservation measures that benefit wildlife. Valuable resources include: Science-Based Trials of Rowcrops Integrated with Prairie Strips , Best Management Practices (BMPs) Minnesota Department of Agriculture , Guides to Prairie Restoration in Minnesota TNC , Restoration-Guide-Crop-to-Utility-Meadow.pdf , Restoration-Guide-Crop-to-Conservation-Meadow.pdf , and Restoration-Guide-Crop-to-Conservation-Prairie.pdf . Incentivize innovation and implementation of conservation measures
	Communicate with the public, using public service announcements, educational materials, radio, and social media during seasons when turtles are most likely crossing roads, requesting motorists to slow down and be on the lookout for turtles on roads.
	Provide lake shore and lake owner associations with information regarding the natural systems and communities found in their water bodies and the protective management options to ensure both recreational opportunities as well as wildlife habitat on human dominated water bodies.
	Where people recreate, work with partners to provide outreach and education on cleaning gear and equipment to reduce the introduction and spread of invasive species. DNR's Prevent the spread webpage details actions for different land-based activities like biking, hiking and off-highway vehicle riding. At trailheads, add invasive species prevention messages, such as those on boot brush kiosks developed by the PlayCleanGo: Stop Invasive Species in Your Tracks program. DNR's What You Should Do webpage details actions for different water activities like kayaking and waterfowl hunting. At water accesses, share Clean in, Clean Out messaging to clean, drain, and dispose to reduce the spread of aquatic invasive species.
	To reduce impacts of chloride, increase long-range planning, communicate more with zoning officials, facilitate enforcement of rules and ordinances, publicize best practices such as smart salting, and reducing chloride in water softeners.

Case Study: Rare Plant Rescue Project

Anoka Sand Plain Rare Plant Rescue Program is a collaborative partnership led by the UMN Landscape Arboretum and focused on protecting rare plants in the Anoka Sand Plain (ASP) region. The ASP spans across several metro counties and supports 59 of MN's state-listed species (or nearly 20%) while representing only 2.2% of MN's land area. Our team collaborates with the DNR, landowners and developers to rescue rare plants from permitted development project areas before construction begins. We accomplish this with a team of ecologists and dedicated volunteers who help us to quickly mobilize and implement rescues. Rescued plants are transplanted into permanently protected natural areas with ecologically appropriate habitat and are monitored overtime.

Since the 2019 inception of the ASP Rare Plant Rescue Program, 10,384 rare plants have been rescued and transplanted into permanently protected habitats. Plants have been rescued from 11 development sites and transplanted into 230 monitored plots on 13 recipient sites. Rescues have involved 10 different species including: 3 state endangered, 5 threatened, 1 special concern, and 1 watch list.

Case Study: Bird City Program by American Bird Conservancy

Bird City started with a single program in Wisconsin in 2009 and expanded to almost a dozen programs recognizing communities across the US and Canada by 2021. Bird City is a conservation program that brings people together to create healthier communities benefiting both people and birds. The program serves as a guide - individuals working together in the community learn about the problems birds face and about potential solutions. They view a list of specific, locally appropriate, bird-friendly actions and select a subset they want to complete or in many cases, are already in motion. Once the required number of actions are complete, the community can apply and be designated as a Bird City. Actions are focused on improving habitat, addressing threats, engaging people, and promoting sustainability. American Bird Conservancy and Environment for the Americas have joined forces to help deliver solutions on the ground through the Bird City Network with over 200 communities across 4 countries currently working to become healthier for birds and people.

Case Study: Optimal Bat Box Design and Placement

Artificial roost structures, also called bat boxes, provide vertical crevices for bat colonies to roost during the daytime during summer months. These structures typically house maternity colonies, or groups of female bats gathered to give birth and raise pups. Bat boxes are often used to compensate for roost loss by increasing roosting opportunities for building-roosting bats in Minnesota, specifically SGCN such as the little brown bat (*Myotis lucifugus*) and the big brown bat (*Eptesicus fuscus*). These structures are relatively easy to build and can be deployed anywhere on posts or buildings. As a result, bat boxes have emerged as a popular conservation tool for supporting bat conservation efforts among the public.

Until recently, bat box guidelines emphasized installing a single bat box, painting it a dark color, and placing it in an area with full sun to create conditions attractive for bats. While these efforts can provide favorable structures for bats, occupancy rates are often low and highly variable, ranging from 1-48% of installations (Fontaine et al. 2021). In Minnesota, mixed success for bat boxes has been reported with many observations of boxes that failed to attract bats. In 2020, the Minnesota Biological Survey and the Parks and Trails Division (PAT) initiated a study to investigate what box and installation characteristics were associated with use by maternity colonies to improve guidelines for using bat boxes in

Minnesota. Initial project goals focused on creating an inventory of bat boxes installed across parks and trails lands, documenting box characteristics, and observing evidence of bat use.

In 2021, with the help of PAT staff across 28 state parks and trails lands, the project documented 138 bat boxes with maternity colonies using 36% of inventoried boxes. With the idea that certain box characteristics would be favored by maternity colonies, we collected information such as box design type, color, dimensions of the box, condition, solar aspect, entrance height, and mounting method. From these data, it was evident that maternity colonies were found in larger volume boxes with multiple roosting chambers. Small, single chamber boxes were never used by maternity colonies and were rarely used by single bats. Maternity colonies were also primarily found where 3 or more boxes were installed within sight distance, providing multiple roost choices.

After the inventory was complete, a subset of 55 boxes was selected for monitoring from 2022 – 2024 to understand colony preferences throughout the summer months. These boxes were primarily found in sites that had multiple boxes clustered together, ranging from 3 – 15 bat boxes available in each study site. Monitoring efforts estimated bat counts, or group sizes, in each box nearly daily, while temperature loggers installed inside boxes monitored temperatures from April - October. Colonies were composed primarily of little brown bats, and total colony sizes at sites ranged from 103 – 539 bats. The data from the monitored boxes showed that bat counts in boxes changed throughout the season, larger groups tended to gather in certain boxes in the spring and early summer, later dispersing to smaller groups spread out among available boxes in the late summer. Temperatures inside the boxes were extremely variable, changing with time of day, ambient temperature, solar exposure, and structural characteristics such as box dimension and design.

The effectiveness of boxes to provide suitable and safe roost conditions, or microclimates (temperature and humidity) for bat occupants has emerged as an important component of guidelines for using bat boxes (Holroyd et al. 2023). Roost microclimate plays a critical role in bat energy budgets, reproductive fitness of mothers, and pup growth rates (Kunz and Lumsden 2003; Tuttle 1976). Reproductive females select roosts with suitable microclimates based on the reproductive period (pregnancy, lactation, post-lactation) to maximize energetic savings for mothers and increased growth rates for pups (Lausen and Barclay 2003). Buildings are attractive to bats because these roosts offer warm, stable temperatures, and ample space with a range of microclimates that bats need throughout reproductive periods, providing energetic benefits that allow bats to maximize reproductive success and avoid switching roosts through the season (Voigt et al. 2016, Lausen and Barclay 2006). Conversely, bat boxes do not provide the same range of conditions or microclimate stability found in building roosts, meaning strategies that increase microclimate choices for bats is an important component of bat box guidelines.

As observed in our study, the temperatures inside bat boxes vary widely, influenced by roost characteristics such as dimensions, design, materials used for construction, and environmental conditions such as air temperatures, solar exposure, and season (Doty et al. 2016; Martin Bideguren et al. 2019; Rueegger 2019). In addition, we observed a pattern of fluctuating daily bat counts, shifting group sizes through the season, and the association of maternity colonies with sites where multiple boxes were available. These observations suggest maternity colonies using bat boxes are likely to use roost-switching and social thermoregulation strategies to find suitable microclimates throughout the reproductive periods, like bat colonies roosting in tree roosts (Ruczyński and Bartoń 2020; Russo et al. 2017).

Temperature variation is also observed in cavities of tree roosts, with tree characteristics explaining less than 28% of variation and ambient conditions explaining over 80% of variation (Clement and Castleberry 2013). Roosting space within tree roosts is also limited as bats rely on pre-existing cracks or crevices, loose bark, and cavities. As a result, reproductive females roosting under loose bark or in tree cavities maintain a network of roost trees, switching roosts almost nightly to find appropriate roost microclimates (Swingen et al. 2018; Sedgely 2001). Social thermoregulation is also an important strategy to help facilitate suitable microclimates in tree roosts, with group sizes changing nightly based on the selected roost, ambient conditions, and reproductive period (Russo et al. 2017).

Our project recommends installing multiple large volume (>3,000 in³) multi-chamber designs, with varying solar aspects nearby, to provide multiple options for maternity colonies. This framework allows bats to change group sizes and select roosts that facilitate suitable microclimates given the reproductive period and ambient conditions to increase effectiveness of boxes to provide quality reproductive habitat for bats. Our findings align with key recommendations recently published in Holroyd et al. 2023, Best Management Practices for Bat Houses in the USA and Canada. Our recommendations for using bat boxes in Minnesota have been summarized to an [online PDF guidance document](#) available to the public, with links to bat box designs that were used by maternity colonies across our study and more.

A major accomplishment from this study was the documentation of significant little brown bat colonies relying on these bat boxes. As one of the most heavily impacted species by White-Nose Syndrome in Minnesota, it is apparent that artificial roosts can play an important role supporting these declining populations. Across our monitored boxes at six state parks, we found a total of at least 2,068 bats relying on these artificial roosts during the reproductive season. We also recommend the consideration of longevity of bat boxes when conducting installations. Little brown bats are long-lived and will return every year to the bat boxes once colonies establish, so it is important to consider monitoring and maintenance plans through time. Ideally, with suitable habitat to support summer colonies, these bats could become common once again.

References for Urban and Other Developed Lands

Aubrecht, C., M.S. Jaiteh, A. de Sherbinin. 2010. Global Assessment of Light Pollution Impact on Protected Areas. Research Gate.

https://www.researchgate.net/publication/238704417_Global_Assessment_of_Light_Pollution_Impact_on_Protected_Areas

Balčiauskas, L., Kučas, A., & Balčiauskienė, L. (2025). A Review of Wildlife–Vehicle Collisions: A Multidisciplinary Path to Sustainable Transportation and Wildlife Protection. *Sustainability*, 17(10), 4644.

<https://doi.org/10.3390/su17104644>

Boone, Michelle L., Zachary M. Portman, Ian Lane, Sujaya Rao. 2023. Occupancy of *Bombus affinis* (Hymenoptera: Apidae) in Minnesota is highest in developed areas when standardized surveys are employed. *Environmental Entomology* 52: 918–938, <https://doi.org/10.1093/ee/nvad088>

Bucher, S.F., L. Uhde, A. Weigelt, S. Cesarz, N. Eisenhauer, A. Geler, C. Kyba, C. Romermann, T. Shatwell and J. Hines. 2023. Artificial light at night decreases plant diversity and performance in experimental grassland communities. <https://doi.org/10.1098/rstb.2022.0358>.

Bumgarner, J.R. and R. Nelson. 2021. Light at Night and Disrupted Circadian Rhythms Alter Physiology and Behavior. *Integrative and Comparative Biology* 61 (3); 1160–1169.

<https://academic.oup.com/icb/article/61/3/1160/6206372>.

Buxton R.T., McKenna M.F., Mennitt D.J., et al. 2017a. Noise pollution is pervasive in US protected areas. *Science* 356: 531–33.

- Buxton, R.T., M.F. McKenna, D.Mennitt, E. Brown, K. Kristrup, K.R. Crooks L.A. Angeloni, and G. Wittemyer. 2019. Anthropogenic noise in US national parks – sources and spatial extent. *Frontiers in Ecology and the Environment* (17) 559-564. <https://doi.org/10.1002/fee.2112>.
- Chock, R.Y., B. Clucas, E.K. Peterson, B. F. Blackwell, D. T. Blumstein, K. Church, E. Fernandex-Jurcic, G. Francescoli, A. L. Greggor, P. Kemp, G.M. Pinho, P. M. Sanzenbacher, B. A. Schulte, P. Toni. 2020. Evaluating potential effects of solar power facilities on wildlife from an animal behavior perspective. *Conservation Science and Practice*. <https://doi.org/10.1111/csp2.319>.
- Clement, M.J., Castleberry S.B. 2013. Tree structure and cavity microclimate: implications for bats and birds. *Int J Biometeorol* 57,437-450.
- DarkSky International. 2025a. [DarkSky International | Protecting the night skies for present and future generations](https://darksky.org). Accessed 2025. <https://darksky.org>
- DarkSky International. 2025b. Artificial Light at Night: State of the Science 2025. DOI:10.5281/zeno.
- Darst, A.L., Mitchell, T.S., Verhoeven, M.R., Evans, E., Tonsfeldt, L., Kjaer, S. et al. (2024) Diversity of bumble bees and butterflies in Minnesota roadsides depends on floral diversity and abundance but not floral native status. *Insect Conservation and Diversity*, 17(4), 690–701. Available from: <https://doi.org/10.1111/icad.12739>
- Doty, A.C., Stawski, C., Currie, S.E., Geiser, F. 2016. Black or white? Physiological implications of roost colour and choice in a microbat. *Journal of Thermal Biology* 60, 162–170.
- Dutcher, A., K. Pias, G.C. Sizemore, and S.M. Vantassel. 2021. Free-ranging and Feral Cats. *Wildlife Damage Management Technical Series*. USDA, APHIS, WS National Wildlife Research Center. Fort Collins, Colorado. 25p.
- Falchi, Fabio, Cinzano, Pierantonio, Duriscoe, Dan, Christopher C. M. Kyba, Christopher C. M., Elvidge, Dhristopher D., Daugh, Kimberly, Portnov, Boris A., Rybnikova, Nataliya A., and Furgoni, Riccard. 2016. The new world atlas of artificial night sky brightness. *Science Advances*. 10 June 2016. Vol 2(6): DOI: [10.1126/sciadv.1600377](https://doi.org/10.1126/sciadv.1600377)
- Fahrig, L., & Rytwinski, T. (2009). Effects of Roads on Animal Abundance: an Empirical Review and Synthesis. *Ecology and Society*, 14(1). <http://www.jstor.org/stable/26268057>
- Finch, D. M., Butler, J. L., Runyon, J. B., Fettig, C. J., Kilkenny, F. F., Jose, S., Frankel, S. J., Cushman, S. A., Cobb, R. C., Dukes, J. S., Hicke, J. A., & Amelon, S. K. (2021). Effects of Climate Change on Invasive Species. In T. M. Poland, T. Patel-Weynand, D. M. Finch, C. F. Miniati, D. C. Hayes, & V. M. Lopez (Eds.), *Invasive Species in Forests and Rangelands of the United States* (pp. 57–83). Springer International Publishing. https://doi.org/10.1007/978-3-030-45367-1_4
- Fontaine, A., Simard, A., Dubois, B., Dutel, J., Elliott, K.H., 2021. Using mounting, orientation, and design to improve bat box thermodynamics in a northern temperate environment. *Sci Rep* 11, 7728. <https://doi.org/10.1038/s41598-021-87327-3>
- Forman, Richard T. T. & Alexander, L. E. (1998). Roads and Their Major Ecological Effects. *Annual Review of Ecology and Systematics*, 29, 207-C2. <http://www.jstor.org/stable/221707>

- Gulliver, John, Chan Lan Chun, Peter Weiss, Andrew Erickson, William Herb, Jerry Henneck, and Kathryn Cassidy. 2022. [Environmental Impacts of Potassium Acetate as a Road Salt Alternative](#) (University of Minnesota evaluation) Report no. MnDOT 2022-27A
- He, X., Liang, J., Zeng, G., Yuan, Y., & Li, X. (2019). The Effects of Interaction between Climate Change and Land-Use/Cover Change on Biodiversity-Related Ecosystem Services. *Global Challenges (Hoboken, NJ)*, 3(9), 1800095. <https://doi.org/10.1002/gch2.201800095>
- Holroyd, S., Lausen, C.L., Dulc, S., de Freitas, E., Crawford, R., O’Keefe, J., Boothe, C., Segers, J. and Reichard, J. 2023. Best Management Practices for the Use of Bat Houses in US and Canada --with focus on summer habitat mitigation for Little Brown Myotis, Yuma Myotis, and Big Brown Bat. Wildlife Conservation Society Canada, produced in cooperation with US Fish and Wildlife Service and Canadian Wildlife Health Cooperative for the WNS Conservation and Recovery Working Group. Version (last updated): 20 Sept. 2023. <https://doi.org/10.7944/P99K4BF5>
- Hopwood, J., S. H. Black, E. Lee-Mader, A. Charlap, R. Preston, K. Mozumder, and S. Fleury. 2015. “Literature Review: Pollinator Habitat Enhancement and Best Management Practices in Highway Rights-of-Way.” Prepared by The Xerces Society for Invertebrate Conservation in collaboration with ICF International. 68 pp. Washington, D.C.: Federal Highway Administration.
- Horvath, G., G. Kriska, P. Malik, and B. Robertson. 2009. Polarized light pollution: a new kind of ecological photopollution. *Frontiers in Ecology and the Environment* 7 (6): 317-325. <https://doi.org/10.1890/080129>.
- Hutto, Richard L. 1998. On the importance of stopover sites to migrating birds. *The Auk* 115: 823 – 825.
- Jacobson, S. L., L. L. Bliss-Ketchum, C. E. de Rivera, and W. P. Smith. 2016. A behavior-based framework for assessing barrier effects to wildlife from vehicle traffic volume. *Ecosphere* 7(4):e01345. 10.1002/ecs2.1345
- Kok A.C.M, Boris W. Berkhout, Nora V. Carlson, Neil P. Evans, Nicola Khan, Dominique A. Potvin, Andrew N. Radford, Marion Sebire, Saeed Shafiei Sabet, Graeme Shannon, and Claudia A. F. Wascher. 2023. [How chronic anthropogenic noise can affect wildlife communities](#). *Frontiers in Ecology and Evolution* 04 (11) <https://doi.org/10.3389/fevo.2023.1130075>.
- Kunz, T.H., and Lumsden L.F. 2003. Ecology of cavity and foliage roosting bats. In: *Bat ecology* (T.H. Kunz and M.B. Fenton, eds.), pp 3-89.
- Lausen, C.L., Barclay, R.M.R., 2003. Thermoregulation and roost selection by reproductive female big brown bats (*Eptesicus fuscus*) roosting in rock crevices. *Journal of Zoology* 260, 235–244. <https://doi.org/10.1017/S0952836903003686>
- Lausen, C.L., Barclay, R.M.R., 2006. Benefits of living in a building: Big brown bats (*Eptesicus fuscus*) in rocks versus buildings. *Journal of Mammalogy* 87, 362–370. <https://doi.org/10.1644/05-MAMM-A-127R1.1>
- Lepczyk, C. A., Dauphiné, N., Bird, D. M., Conant, S., Cooper, R. J., Duffy, D. C., Hatley, P. J., Marra, P. P., Stone, E., & Temple, S. A. (2010). What Conservation Biologists can do to Counter Trap-Neuter-Return: Response to Longcore et al.. *Conservation Biology*, 24(2), 627–629. <https://doi.org/10.1111/j.1523-1739.2009.01426.x>

- Liebman, M., Helmers, M. J., Schulte, L. A., & Chase, C. A. (2013). Using biodiversity to link agricultural productivity with environmental quality: Results from three field experiments in Iowa. *Renewable Agriculture and Food Systems*, 28(2), 115–128. doi:10.1017/S1742170512000300++++
- Liess, S., Twine, T. E., Snyder, P. K., Hutchison, W. D., Konar-Steenberg, G., Keeler, B. L., & Brauman, K. A. (2022). High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*, 9(3), e2021EA001893. <https://doi.org/10.1029/2021EA001893>
- Loss, S., Will, T. & Marra, P. The impact of free-ranging domestic cats on wildlife of the United States. *Nat Commun* 4, 1396 (2013). <https://doi.org/10.1038/ncomms2380>
- Martin Bideguren, G., López-Baucells, A., Puig-Montserrat, X., Mas, M., Porres, X., Flaquer, C., 2019. Bat boxes and climate change: testing the risk of over-heating in the Mediterranean region. *Biodivers Conserv* 28, 21–35. <https://doi.org/10.1007/s10531-018-1634-7>
- Minnesota Pollution Control Agency. 2020. [Statewide Chloride Management Plan](#) and [Summary](#).
- Minnesota Prairie Plan Working Group. 2018. Minnesota Prairie Conservation Plan, 2nd ed. Minneapolis, MN. 69 pp.
- MN Pollution Control Agency. [Getting Lead out of Fishing Tackle](#)
- DNR and MNDOT. 2014. [Best Practices for Meeting DNR General Public Waters Work Permit GP 2004-0001](#)
- Mitchell, T. S., Agnew, L., Meyer, R., Sikkink, K. L., Oberhauser, K. S., Borer, E. T., & Snell-Rood, E. C. (2020). Traffic influences nutritional quality of roadside plants for monarch caterpillars. *Science of The Total Environment*, 724, 138045. <https://doi.org/10.1016/j.scitotenv.2020.138045>
- Mitchell, T. S., Verhoeven, M. R., Darst, A. L., Patterson, C. H., & Snell-Rood, E. C. (2024). Seeding roadsides is necessary but not sufficient for restoring native floral communities. *Ecological Solutions and Evidence*, 5, e12364. <https://doi.org/10.1002/2688-8319.12364>
- Parris, K.M. (2015). Ecological Impacts of Road Noise and Options for Mitigation. In *Handbook of Road Ecology* (eds R. van der Ree, D.J. Smith and C. Grilo). <https://doi.org/10.1002/9781118568170.ch19>
- Phillips, J.N., S. E. Termondt, and C. D. Francis. 2021. Long-term noise pollution affects seedling recruitment and community composition, with negative effects persisting after removal. *Proceeding of the Royal Society*. <https://doi.org/10.1098/rspb.2020.2906>.
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. P., & LeDee, O. E. (2025). *Effects of climate change on midwestern ecosystems: Temperate flooded and swamp forest* (p. 110pp). Midwest Climate Adaptation Science Center. Available: <https://mwccasc.umn.edu/node/821>
- Ruczyński I., Bartoń K.A. 2020. Seasonal changes and the influence of tree species and ambient temperature on the fission-fusion dynamics of tree-roosting bats. *Behavioral Ecology and Sociobiology* 74:5, 1-8.

Ruegger, N., 2019. Variation in Summer and Winter Microclimate in Multi-Chambered Bat Boxes in Eastern Australia: Potential Eco-Physiological Implications for Bats. *Environments* 6, 13.

<https://doi.org/10.3390/environments6020013>

Runkle, J., Kunkel, K. E., Frankson, R., Easterling, D. R., & Champion, S. M. (2022). *Minnesota State Climate Summary 2022* (NOAA Technical Report NESDIS 150-MN, p. 4 pp.). NOAA/NESDIS.

<https://statesummaries.ncics.org/chapter/mn/>

Russo, D. Cistrone L., Budinski I., Console G., Corte M.D., Milighetti C., Di Salvo I., Nardone V., Brigham R. M., Ancillotto L. 2017. Sociality influences thermoregulation and roost switching in a forest bat using ephemeral roosts. *Ecology and Evolution* 7, 5310-5321.

Salafsky, N., Relton, C., Young, B. E., Lamarre, P., Böhm, M., Chénier, M., Cochrane, E., Dionne, M., He, K. K., Hilton-Taylor, C., Latrémouille, C., Morrison, J., Raymond, C. V., Seddon, M., & Suresh, V. (2024). Classification of direct threats to the conservation of ecosystems and species 4.0. *Conservation Biology*, 39(3), e14434.

<https://doi.org/10.1111/cobi.14434>

Sedgeley, J.A., 2001. Quality of cavity microclimate as a factor influencing selection of maternity roosts by a tree-dwelling bat, *Chalinobus tuberculatus*, in New Zealand: Roost microclimate of long-tailed bats. *Journal of Applied Ecology* 38, 425–438. <https://doi.org/10.1046/j.1365-2664.2001.00607.x>

Shephard, A. M., Agnew, L., Herdtle, A., Mitchell, T. S., Borer, E. T., & Snell-Rood, E. C. (2022). Traffic patterns, more than adjacent land use, influence element content of roadside forbs for insect pollinators. *Ecological Solutions and Evidence*, 3, e12195. <https://doi.org/10.1002/2688-8319.12195>

Sheppard, Christine and Phillips, Glenn. Bird-Friendly Building Design, 2nd Ed. (The Plains, VA: American Bird Conservancy, 2015)

Sillett, Scott T. and Richard T. Holmes. 2002. Variation in survivorship of a migratory songbird throughout its annual cycle. *Journal of Animal Ecology* 71: 296 – 308.

Stobo-Wilson, A. M., Murphy, B. P., Legge, S. M., Caceres-Escobar, H., Chapple, D. G., Crawford, H. M., Dawson, S. J., Dickman, C. R., Doherty, T. S., Fleming, P. A., Garnett, S. T., Gentle, M., Newsome, T. M., Palmer, R., Rees, M. W., Ritchie, E. G., Speed, J., Stuart, J., Suarez-Castro, A. F., ... Woinarski, J. C. Z. (2022). Counting the bodies: Estimating the numbers and spatial variation of Australian reptiles, birds and mammals killed by two invasive mesopredators. *Diversity and Distributions*, 28(5), 976–991. <https://doi.org/10.1111/ddi.13497>

Swingen, Morgan; Moen, Ronald; Walker, Marcus; Baker, Richard; Nordquist, Gerda; Catton, Timothy; Kirschbaum, Kari; Dirks, Brian; Dietz, Nancy. (2018). Northern Long-Eared Bat Roost Tree Characteristics 2015-2017. Retrieved from the University Digital Conservancy, <https://hdl.handle.net/11299/204334>.

Tuttle, M.D., 1976. Population Ecology of the Gray Bat (*Myotis grisescens*): Factors Influencing Growth and Survival of Newly Volant Young. *Ecology* 57, 587–595. <https://doi.org/10.2307/1936443>

U.S. Fish and Wildlife Service. 2023. Species Status Assessment Report for the Blanding's turtle (*Emydoidea blandingii*). Version 1.1. Midwest Region, Bloomington, Minnesota.

Voigt, C.C., Phelps, K.L., Aguirre, L.F., Corrie Schoeman, M., Vanitharani, J., Zubaid, A., 2016. Bats and Buildings: The Conservation of Synanthropic Bats, in: Voigt, C.C., Kingston, T. (Eds.), Bats in the Anthropocene: Conservation of Bats in a Changing World. Springer International Publishing, Cham, pp. 427–462.
https://doi.org/10.1007/978-3-319-25220-9_14

White, Lauren. 2024. Mitigation of Road Mortality of Freshwater Turtles in New Hampshire with an Emphasis on Blanding's Turtles (*Emydoidea blandingii*). Master's Theses and Capstones. 1905.
<https://scholars.unh.edu/thesis/1905>

White, Lauren. 2024. Mitigation of Road Mortality of Freshwater Turtles in New Hampshire with an Emphasis on Blanding's Turtles (*Emydoidea blandingii*). Master's Theses and Capstones. 1905.
<https://scholars.unh.edu/thesis/1905>

Williams-Sether, T., & Sanocki, C. (2025). *Peak streamflow trends in Minnesota and their relation to changes in climate, water years 1921–2020, chap. E* (Peak Streamflow Trends and Their Relation to Changes in Climate in Illinois, Iowa, Michigan, Minnesota, Missouri, Montana, North Dakota, South Dakota, and Wisconsin: U.S. Geological Survey Scientific Investigations Report 2023–5064, p. 55) [Scientific Investigations Report].
<https://pubs.usgs.gov/sir/2023/5064/e/sir20235064e.pdf>

Chapter 4. Public Engagement

Introduction

The conservation of Minnesota's Species in Greatest Conservation Need (SGCN) requires actions by all Minnesotans. If actively engaged, the people of Minnesota can make important contributions to wildlife conservation wherever they live, from rural to urban settings. Effective engagement of our state's residents begins with an understanding of their demographics and values towards wildlife. As of 2016, 81% of Minnesota residents lived in urban areas (Minnesota Department of Transportation [MnDOT], 2022a), with over half of the state's population residing in the seven-county metro area (MnDOT, 2022b). Recent trends show that growth will steadily continue in urban areas with the highest growth rates in suburban edges (Metropolitan Council, 2024). For these residents living in more highly developed areas, connecting to rare and declining wildlife species and their habitats might pose a particularly difficult challenge. Additionally, the America's Wildlife Values Study (Dietsch et al., 2018), which sought to explore the values, attitudes, and beliefs of residents across the U.S. in relation to wildlife, found that the following breakdown of wildlife value orientations for Minnesota respondents: 37% traditionalist (believe wildlife should be used and managed for human benefit), 27% mutualist (believe people should live in harmony with wildlife), 24% pluralist (their wildlife values vary depending on the context), and 12% distanced (wildlife issues are less salient to them). While participating in the conservation of rare and declining wildlife may be most intuitive for just under a third of Minnesota's residents (the mutualists), we have the potential to engage any Minnesotan in SGCN conservation. However, our strategies will need to be varied and tailored to our residents' differing values and interests. Given the mix of wildlife value orientations in Minnesota, and the state's ratio of urban to rural residents, it is vital to the conservation of SGCN that we invest in connecting the people of Minnesota to its rare and vulnerable wildlife. By actively engaging the public through multi-faceted approaches, we can work towards reducing human-caused threats to SGCNs and increase capacity and funding for their conservation. This chapter provides a strategic guide for engaging Minnesotans to become a part of the solution.

With a deep passion and appreciation for the outdoors, the people of Minnesota are primed to engage in conservation action, setting this plan up for success. This passion was evident in November 2024, when in remarkable agreement, over 77 percent of the state's voters approved a renewal of the Environment and Natural Resources Trust Fund. This vote assured that a portion of Minnesota state lottery monies would fund environmental projects for another 25 years.

To develop the recommendations in this chapter, the Minnesota Department of Natural Resource's Nongame Wildlife Program (NWP) utilized a participatory process. NWP staff reviewed the input of 161 conservation professionals from 28 organizations that attended a series of seven statewide SWAP workshops in 2023. To obtain further input on content specifically relevant to the SWAP's public engagement strategies, the NWP also consulted with 90 conservation educators, outreach program leaders, and communication professionals from 29 organizations across the state in two additional workshops and two working group meetings. In total, this chapter is based upon input from over 250 individuals, representing multiple DNR divisions and over 50 other agencies, organizations, and academic institutions.

Building on the two Prior Plans

2005-2015 Tomorrow's Habitat for the Wild and Rare

One of the original Plan's three strategic goals was to: **Enhance people's appreciation and enjoyment of Species in Greatest Conservation Need (SGCN)**. The actions associated with this goal were included within each of the profiles for Minnesota's 25 Ecological Subsections:

1. Create new information and communicate with people to enhance their appreciation of SGCN
2. Create opportunities for people to appropriately enjoy SGCN-based recreation.

2015-2025 Minnesota's Wildlife Action Plan

In the 10-year revision of the 2005-2015 State Wildlife Action Plan, public engagement remained one of the three primary goals, to: Enhance opportunities to enjoy Species in Greatest Conservation Need and other wildlife and to participate in conservation. The goal was revised from the 2005 Plan to emphasize creating opportunities for Minnesotans to actively participate in SGCN conservation efforts. Two objectives to reach these goals were delineated:

1. Increase the number and diversity of people participating in wildlife-related outdoor recreation.
2. Increase the number and diversity of people directly engaged in conservation of SGCN.

Our recommendations for public engagement take a multi-faceted approach, from sparking inspiration to motivating and empowering conservation action. We seek to first connect Minnesotans to our state's SGCN and habitats by using varied educational approaches that build knowledge, relevancy and positive attitudes around these species. These approaches include information sharing, environmental education, and wildlife viewing recreation. From there, with that essential groundwork in place, we lead Minnesotans to take the following conservation actions: recreating outdoors responsibly, creating and caring for habitat, volunteering to collect wildlife data, and donating to wildlife conservation. Environmental education practices utilizing this continuum from inspiring connections to motivating and empowering action have been found to develop pro-environmental behaviors, build community capacity, support conservation action projects and contribute to long-term conservation goals (Ardoin et al., 2020).

To further increase the efficacy of these strategies for all communities in Minnesota and to address potential historical, cultural and societal barriers, our public engagement recommendations follow insights gained from a study the DNR's NWP contracted from Virginia Tech, which sought to learn how to expand Minnesota's audience for public conservation efforts (Jennings et al., 2024a). Accordingly, some themes throughout our recommended actions are meeting people where they are, working collaboratively with partners and local groups, and developing programs relevant to communities.

Our public engagement recommendations for 2025-2035 are organized into seven overarching strategies, each supported by a suite of actions, flowing from inspirational tactics to action-oriented ones. These strategies include information sharing, environmental education, wildlife viewing, responsible recreation, habitat stewardship, participatory science, and funding support.

Goal: *Inspire, motivate, and empower Minnesotans to engage in conservation actions that support Species in Greatest Conservation Need and their habitats.*

Recommended Strategies for Public Engagement

1. Develop and distribute informational resources and media on SGCN, their habitats, and threats.
2. Support environmental education on SGCN and their habitats on a statewide level.
3. Provide and promote wildlife viewing opportunities and programs that inspire curiosity, caring, and connection with Minnesota's SGCN.
4. Foster responsible recreation practices that reduce recreational impacts to SGCN and their habitats while allowing for safe and satisfying access to the outdoors.
5. Engage and empower communities to improve wildlife habitats where they live, work, and recreate.
6. Offer participatory science programming for Minnesotans to take part in conservation volunteerism that addresses SGCN data needs.
7. Cultivate financial support for the conservation of SGCN through the development and promotion of public funding opportunities and public-private partnerships.

Information Sharing

Strategy 1: Develop and distribute informational resources and media on SGCN, including their habitats and threats.

One of the first steps towards inspiring and motivating people towards conservation action is raising awareness that there is a need for conservation action. The people of Minnesota cannot aid in the conservation of SGCN if they do not know these species exist in our state or are unaware of the issues impacting them. Minnesotans are passionate about conserving our state's natural heritage, so if conservation professionals provide the public more information about SGCN, their threats and their needs, we have an opportunity to foster changes in human attitudes and behaviors to benefit these species in need. To be effective, this information must be provided in easily accessed formats and in a relevant manner.

Research has shown that social media can increase pro-conservation behaviors among the public, increase conservation funding, and inspire policy change (Bergman et al., 2022). In addition, sharing information about wildlife conservation, in tandem with efforts like increasing environmental education, can help shift social norms and have a positive impact on rare wildlife species (Salazar et al., 2018).

The species groups that may benefit most from information campaigns include native fish, pollinating insects, mussels, amphibians, reptiles, mammals, and birds. Within these groups are Minnesota SGCN that are threatened by human behaviors such as illegal take (wanton killing/poaching), accidental take (e.g. road mortality, bird building collisions), illegal possession, pollution, climate change, and the spread of invasive species. These are also groups of wildlife that are interesting and/or relevant to Minnesotans (Sinkular et al., 2022; Duda, 2021), in which it may be easier to capture people's attention and foster behavior changes.

Sharing more information about the SGCN listed above will enable Minnesotans to make more informed decisions about their actions and interactions with nature. For example, informing Minnesotans when turtles are more likely to be crossing roads and how to help them safely cross the road may help reduce road mortality. Sharing the important role of native mussels in our water ecosystems is a gateway to discussing pollution, climate change, invasive species and how Minnesotans can make a difference for our waters.

By providing communication products that increase Minnesotans' knowledge about SGCN, the stressors they face, and what people can do to alleviate these stressors, we can foster a deeper appreciation for SGCN, the desire to help them, and motivate people towards the conservation actions outlined in Strategies 4-7.

Stressors Addressed: Development (e.g. bird building collisions), Roads, Trails and Railroads (e.g. road mortality), Hunting and Collecting Animals (wanton killing, poaching, illegal possession), Invasive Species (Problematic Non-Native Species), Diseases and Pathogens, Water-borne Pollution, Air-borne Pollution, Changes in Temperature related to Climate, and Changes in Precipitation and Hydrology due to Climate.

Recommended Actions for Information Sharing:

- 1. Identify specific SGCN as ambassador species (i.e. charismatic champions that represent a suite of other SGCNs or habitats) to help build a broader audience to receive wildlife information by streamlining information and making it appealing. These ambassador SGCN could be used for leveraging popular interests, communicators, communication channels, and events to reach new audiences.**

Work with wildlife biologists to identify specific SGCN to utilize as ambassador species—charismatic champions representing a suite of other taxa (or habitats)— in which by garnering public familiarity, increasing knowledge, and leveraging support for that specific SGCN, a greater suite of species will benefit. These ambassador species can be used in a variety of ways, such as in communication campaigns around calls to action for certain stewardship practices, in an “Adopt-a-SGCN” program to engage schools and municipalities in selecting a species to represent their community, or as features in outreach products like education posters, coloring books, etc. Ambassador species could also be utilized to connect with unique, less traditional audiences, such as reaching sports fans by partnering with Minnesota sports teams that feature SGCN as their mascot (e.g. the Minnesota Lynx), artists through art contests/exhibitions themed around these species, and social media users with the use of influencers and short videos providing educational content on these species. Holding festivals to celebrate these species, serves as an attractive, fun event that also engages people in learning.

- 2. Promote and support an update to the DNR's Rare Species Guide (RSG) as a means for providing an information hub on Minnesota's SGCN and their habitats and range.**

The RSG is a valuable information resource on state listed animals and plants (see case study), offering an area of opportunity to present information on more SGCN. It is also in need of updates given that it launched in 2008. In 2023, a MNIT Business Value Assessment showed the RSG as having high business value and moderate to low technical health with multiple modernization efforts needed. A RSG Modernization and Enhancement IT Project has been proposed to address these needs and a survey

was implemented in 2024 to better understand how people use the RSG and how they would like to see it improved. By increasing the capacity to help with these needed technical, design, and content updates (including developing new RSG profiles for SGCN animal species that don't currently have a profile), we can deliver a modern product that will better inform Minnesotans about the needs and threats of SGCN.

3. Provide solutions for DNR staff, conservation organizations, and municipalities to address common wildlife questions and concerns.

- a. Collaborate with the DNR Information Center to develop FAQs for the public that may be disseminated through communications media and materials by varied agencies and organizations. These FAQs could cover topics like keeping wild species wild, Threatened and Endangered Species poaching, and illegal take/possession penalties.
- b. Enhance internal DNR collaboration between divisions, Listed Species Program, and the Nongame Wildlife Program to increase and support knowledge of SGCN regulations to help prevent wanton killing, illegal take, or illegal possession of SGCN. Species that are protected by state and federal laws, such as Blanding's and wood turtles, migratory birds, native fish, and timber rattlesnakes, would all benefit from this action. Some examples of work to be done in this arena include, providing informational resources and trainings to DNR employees that will support their protection efforts and public communications work of these legally protected SGCN. Trainings and resources could include summarizing SGCN-related rules and regulations, species identification, and locations of important populations that could be exploited.

Example SGCN: Black buffalo

The [black buffalo \(*Ictiobus niger*\)](#) is a State Threatened, long-lived native fish species that inhabits large river systems in Minnesota including the Mississippi and Minnesota and their tributaries. Minnesota is on the Northern edge of the species' continental range and is an important area for conservation. While the black buffalo is protected from any harvest or take due to its status as a state threatened species, it is part of a group native fish including gar, buffaloes, suckers, and redhorse that have historically been classified as "rough fish" and lacked the same protections as game fish, such as legal harvest methods and bag limits. Further, confusion about native fishes' role in the ecosystem and a lack of public identification resources to help people differentiate native fish from common carp often led to unregulated overharvest. Fortunately, recent conservation and legislative efforts have resulted in these native fish being reclassified and receiving the same protections as game fish (Winter, 2024). Specific regulations for these species are relatively new and still being developed. Many people may be unaware of changing regulations and related laws regarding native fish species, and lack of resources for species identification remains an issue. Ongoing communication, education, and outreach will benefit the black buffalo and other native fish by raising awareness, reducing unintentional take, empowering conservation officers to enforce regulations, and filling gaps in scientific knowledge about native fish in Minnesota.

4. Increase understanding of misunderstood SGCN and foster tolerance for them in human spaces through informational products, trainings, and programs.

- a. Create outreach products for SGCN that may be perceived as frightening or as a nuisance (e.g. snakes, bats, hawks, wolves, mudpuppies, northern pocket gophers and insects), so that people may learn to appreciate these interesting animals. An example of this work for snakes could include updating the DNR's "Living with Snakes" informational brochure and webpage.

- b. Create outreach products that clarify common and problematic species misidentifications and distribute/promote these to target audiences. For example, a species identification guide helps anglers understand differences among native SGCN fish, common carp, and invasive carp ([Have you seen an invasive carp?](#); also available in Hmong and Spanish on the DNR's [Invasive Carp](#) webpage). Similarly, work with herpetologists to develop an updated field guide and/or infographics on Minnesota's snakes, so that people can learn to discern the many nonvenomous snakes in Minnesota from our one venomous snake, the state threatened timber rattlesnake.
- c. Partner with pest control companies to provide professional development on species identification and best practices.
- d. Connect with [Advancing Conservation through Empathy for Wildlife Network \(ACE\)](#) to use and share their resources.
- e. Wildlife viewing engagement programs would also help to support tolerance and understanding (see Strategy 4) for SGCN.

Example SGCN: Timber rattlesnake

[Timber rattlesnakes \(*Crotalus horridus*\)](#) are generally docile creatures. When threatened, their initial reaction is to hide or flee. They are rarely observed, as they have a localized population in the bluffs of southeastern Minnesota. However, because they are a venomous snake, many Minnesotans fear them and regard them as dangerous. This fear is passed onto many other snake species, as there are several rattlesnake mimics in the state. While they are not venomous themselves, they utilize the rattlesnake's behavior to protect themselves. Unfortunately, this perpetuates the perception that snakes are dangerous and results in wanton killings, which in the case of timber rattlesnake and five other snake SGCN that are state threatened, is also illegal. By sharing more information about the rarity of timber rattlesnakes in our state, their status as a state threatened species, how to distinguish them from other snake species, and educating Minnesotans about their behaviors and importance to balancing our ecosystem, we can hopefully reduce the killings that they and other snakes face and begin to correct the perception and misinformation that snakes are dangerous. By simply keeping their distance, leaving snakes alone and understanding their fascinating behaviors and important role, Minnesotans can hopefully learn to tolerate and perhaps even enjoy a rare reptile sighting.

5. Increase public knowledge on climate change as a threat to Minnesota's wildlife and their habitats.

- a. Work with wildlife biologists to develop a resource list of iconic and unique Minnesota SGCN that are highly impacted by climate change, such as moose, boreal chickadee, migratory birds, shorebirds, wood turtle, brook trout, and monarch. Feature these species in communications products, such as stories, maps and infographics, as a way to garner public understanding of how climate change impacts wildlife and natural resources.
- b. Promote and provide trainings for educators on [AFWA's Project WILD module Climate and Wildlife](#), a curriculum guide to support integrating a wildlife lens into climate education, and Climate Generation's curriculum, [Healthy Habitats: Climate Change Action for K-2](#).
- c. As they become available, assist in communications regarding best management practices, climate-adapted management plans, and other resources that may assist habitat managers by creating materials and holding workshops.
- d. Develop a communications plan around climate change and wildlife in Minnesota. Include crafting talking points and key messages that translate scientific information, making it more easily understood. Utilize current resources from [University of Minnesota Climate Adaptation](#)

- [Partnership](#) and the [Midwest Climate Adaptation Science Center](#) among others). Couple information on climate stress with action-oriented and hopeful messages to keep people motivated (e.g. stories of success in climate adaptation and actionable solutions that individuals can take to make a difference).
- e. Develop community science projects that focus on phenology (such as through the [National Phenology Network](#)) to involve the public in documentation of seasonal changes that may be due to climate change; to help foster understanding and support for climate science.

Example SGCN: Moose

An iconic Minnesota species, [moose \(*Alces alces*\)](#) are important to the state's ecology, culture and economy. Moose are well adapted to Minnesota's traditional cold winters and the boreal forest of northern Minnesota, making them more vulnerable to changes in that climate. The moose population has experienced a 60% population decline since 2006 in northeast Minnesota, largely due to changes from the warming climate. Heat stress, increased winter ticks, and introduction of parasites carried by white-tailed deer (brainworms and liver flukes) are key impacts from a changing climate (Weiskopf et al., 2019; [Midwest Climate Adaptation Science Center](#)). Minnesota has many active organizations working to better understand climate change impacts on moose and find ways to adjust management actions to adapt to those changes. Moose offer an excellent opportunity to share local, relevant, and compelling information about climate change impacts to SGCNs and their habitats.

6. Apply existing social science research and conduct new research as needed to better understand Minnesotans' attitudes about wildlife and wildlife conservation and to help craft effective messaging.

Recent social science research, contracted by the Nongame Wildlife Program, has provided valuable insight to help guide Minnesota's public engagement efforts for SGCN (Sinkular et al., 2002; Jennings et al., 2024a; Jennings et al., 2024b) and these resources are recommended for consideration in crafting effective conservation messaging. However, it should be noted that these studies were limited to very specific groups of people, such as wildlife viewers, participatory scientists, or current supporters of the Nongame Wildlife Program. To understand the perspectives of broader groups of people within Minnesota's geographic boundaries, the Minnesota results for the America's Wildlife Values study (Dietsch et al., 2018) offers an additional helpful reference. Also valuable are The Nature of Americans (Kellert, 2017) and Conservation Words That Work (Duda, 2021), however both studies' results are provided on a national scale. Consideration should be given as to whether further state-level research is needed to better understand Minnesota's unique demographics and geographic locations (e.g. Metro, suburb, rural) and their attitudes, interests, and understanding of wildlife and wildlife conservation.

Case Study: Minnesota Bat Festival brings the fun to learning about bats

The [Minnesota Bat Festival](#) is an annual festival organized by the U.S. Fish & Wildlife Service, held at the Minnesota Valley National Wildlife Refuge, just outside of Minneapolis. It is a good example of appealing to a broader audience in a densely populated area with a fun, well-timed event about a misunderstood, yet fascinating and ecologically valuable species. The event's goal is to educate adults and youth about bat behavior, bat research techniques, and raise awareness of the many threats bats face. Up to fifteen organizations with some specialty in bat conservation and education have participated, providing live demonstrations, presentations, activities and learning opportunities appropriate for all ages. The festival takes place during International Bat Week, which is the last week of October. Also coinciding with Halloween, the event is well-timed to add additional appeal for families and adults alike seeking festive seasonal activities and draws high numbers of attendees. In 2023, 1,500 people attended. First held in 2016, many attendees have shared that the event has become a tradition in their family or that it has helped them appreciate bats more. Through this festival the Refuge has found that Minnesotans are hungry for information about bats.

Case Study: The Rare Species Guide Provides Easy Access to Information on Minnesota's Rare Animals and Plants

The DNR's [Rare Species Guide](#) (RSG) is an online guide to Minnesota's rare animal and plant species, including State-listed endangered, threatened, and special concern species. The RSG provides information on the status, distribution, ecology, conservation, and management of our rarest animals and plants. The user can query over 600 species profiles based on name, taxonomic group, status, county, Ecological Classification System (ECS) subsection, watershed, and/or habitat.

The RSG is one way the DNR addresses the mandate in the Minnesota Environmental Protection Act to "establish and maintain information systems sufficient to gauge the environmental health of the state." The RSG is a foundational resource for State-listed and SGCN species topics and for administering State-listed species rule and law. The RSG is used by professionals and the general public alike, receiving over 186,000 visits in 2024.

Environmental Education

Strategy 2: Support environmental education on SGCN and their habitats on a statewide level.

Environmental education builds knowledge of and connection with the natural world. It fosters awareness, skills, and agency, encouraging people to explore environmental issues, solve problems, and care for the environment. As described by the [Environmental Protection Agency](#), “Environmental education does not advocate a particular viewpoint or course of action. Rather, environmental education teaches individuals how to weigh various sides of an issue through critical thinking and it enhances their own problem-solving and decision-making skills.”

While environmental education techniques are referenced throughout this chapter to help adult audiences gain the knowledge, attitudes, and behaviors to support SGCNs and their habitats, Strategy 2 focuses on applying environmental education within the school age (K-12) realm. Studies show that environmental education for students has numerous benefits, including strengthening motivation to engage in environmentally friendly behavior, and equipping young people with “action knowledge” to have meaningful environmental impact (Van de Wetering et al., 2022).

To help in implementing this strategy, Minnesota has a large, established and engaged K-12 environmental education community made up of professionals and volunteers working together to increase Minnesotans’ knowledge, attitudes and behaviors that support SGCNs and their habitat. Supporting environmental education as described in the recommended actions below will provide environmental educators the resources they need to teach about SGCNs/habitats. Collaborating with and supporting K-12 educators is an effective way to meet teachers’ needs and integrate wildlife learning for youth in the short term, while building the foundation for conservationists and wildlife stewards in future generations.

Although Minnesota has a strong tradition of connecting with the outdoors and supporting conservation work, our society is still shifting to spend more time indoors (Kellert, 2017). As we do so, people are increasingly disconnected from and less knowledgeable about nature. Environmental literacy or knowledge is shown to directly foster individuals’ concern, attitudes, and level of personal responsibility towards environmental protection (Teksoz et al., 2012). K-12 environmental education is essential to building environmental literacy which is key to protecting the environment and ensuring a sustainable future for generations to come (Yildirim et al., 2025).

We recommend focusing environmental education efforts on SGCN in the following species groups: mammals, birds, native fish, reptiles, pollinating insects, amphibians, and mussels. These species groups were selected to highlight based on the Minnesota Results of the National Wildlife Viewer Survey (Sinkular et al., 2022), in which our state’s wildlife viewers selected these species groups (except for mussels) as those they were interested in observing. Pollinating insects make a good match for youth because supportive actions, such as native plantings and insect observation, are accessible to young people and provide an opportunity to make a meaningful impact on a threatened population. Knowing that the public places a high value on water quality (Duda, 2021), freshwater mussels are recommended for environmental education because they provide a strong opportunity to connect a community value to a less visible, yet highly valuable taxa.

SGCN Example: Monarch butterfly as an Ambassador Species

Equipping young people with “action knowledge” they can employ to support environmental issues they care about is one of the most profound impacts of environmental education. There are many species that stand to benefit from supportive activity driven by young people. The Monarch butterfly (*Danaus plexippus*) is a profound example of this possibility and has been identified as an excellent ‘ambassador species.’ Charismatic, colorful, and fragile, this species captures the attention and enthusiasm of young people while also facing significant threats and a declining population. Monarchs present an ideal opportunity for young people to view a declining species and take part in its conservation. Planting milkweed and other nectar-producing plants also can help them to understand the various components of habitat. Connecting with multi-national conservation efforts with concerned residents of Canada and Mexico (e.g. through Monarch Joint Venture) further enhance the conservation learning for this inspiring migratory butterfly and allows young people to understand more about wildlife conservation.

Recommended Actions for Environmental Education:

- 1. Offer free or low-cost professional development trainings to educators (both formal and non-formal) on environmental education and on SGCN and their habitats.**
 - a. Reinvigorate and support statewide coordination of reputable curriculum training programs like [Project WILD](#) and [Project Learning Tree](#), utilizing their efficient train-the-trainer approach to expand capacity for implementation. In Minnesota, we could train facilitators to train educators, naturalists, [Master Naturalists](#), and nature tourism professionals, who already engage with the public daily on how to implement the wildlife and habitat lessons from these valuable guides.
 - b. Utilize Education Minnesota’s [MEA annual conference](#) as a means for wildlife professionals to connect with educators and provide professional development sessions, and to create networking opportunities for teachers who use a given wildlife curriculum to encourage mutual support.
 - c. Promote existing professional development opportunities; e.g. [Monarch Joint Venture’s North American Monarch Institute](#), [Monarch NECTAR Hub](#) and [Wolf Ridge Environmental Learning Center’s Teacher Training Institute](#).

- 2. Promote existing SGCN/ habitat curricula that meet state standards in a variety of subject areas.**

The following curricula already designed to meet state standards are recommended: 1) Association of Fish and Wildlife Agencies’ suite of [Project WILD](#) curricula, which includes, Project WILD, Aquatic WILD, Growing Up WILD, Flying WILD, plus more specific modules; 2) Monarch Joint Venture’s [Monarchs and More](#); and 3) the DNR’s [Life on the Edge in Minnesota](#) (a supplement to Project Learning Tree’s Life on the Edge activity). It’s also recommended that organizations that have existing SGCN/habitat lessons correlate these with established K-12 courses (AP Environmental Science, AP Biology, etc.) to help encourage their implementation by educators.

- 3. Establish or identify an existing resource hub where educational resources on SGCN could be compiled into one convenient location—sorted by grade, state standards, topic, etc.**
 - a. Investigate whether the [Minnesota Association of Environmental Education’s EE Portal](#) (MAEE) could serve as this hub, with SGCN resources submitted to it.

- b. Connect organizations and schools with wildlife experts by compiling a list of organizations/ programs that can be a resource and incorporate this into the [Minnesota Association for Environmental Education's EE Portal](#).
- 4. **Provide funding and technical support to advance outdoor learning spaces at schools that would attract and support SGCN (e.g. native plant gardens for pollinators and birds) and enhance outdoor learning.**
 - a. Develop a small grants program to provide funds to schools and other local organizations doing habitat restoration projects (e.g. [Great River Greening](#)) for development of outdoor learning spaces that also provide habitat benefits.
 - b. Develop a habitat education train-the-trainer program that could promote and train professional educators and established volunteer organizations ([Master Gardeners](#), [Master Naturalists](#), etc.) to provide technical assistance to schools (See Strategy 1, Action 2b) interested in developing or enhancing outdoor learning spaces of habitat value.
 - c. Collaborate with the [DNR's School Forest Program](#) and other outdoor classroom programs to enhance habitat for SGCN through their stewardship plans and integrate other educational supports such as interpretive signs on SGCN at school forests.
- 5. **Provide and/or increase awareness of small grants for field trips and transportation to natural areas for environmental education programming.**

Case Study: Monarch Joint Venture's education program helps educators implement monarch education

Monarch Joint Venture's [Education Program](#) provides an excellent example of how an organization can integrate multiple recommended actions to support this environmental education strategy. Monarch Joint Venture offers professional development for educators ([North American Monarch Institute](#)), funding and technical support for educational gardens, curriculum ([Monarchs and More Curriculum for grades K-8](#)) aligned with the Next Generation Science Standards, educational resources for kids ([Growing with Monarchs Activity Book](#)), K-12 classroom presentations, and online courses for all ages ([Monarch NECTAR Hub](#)). The North American Monarch Institute alone has reached 126 Minnesota educators who in turn have reached 126,000 Minnesota students since 2010. These combined actions and outreach with multiple partners show how environmental education programs like Monarch Joint Venture can build support for a SGCN and its habitat.

Case Study: University of Minnesota (UMN) Bee Lab Pollinator Toolkit makes pollinator education easy by supplying resources

The UMN Bee Lab [Pollinator Education Toolkit](#) is an example of one way in which environmental education can directly support SGCN. The toolkit comes with a variety of interactive games and activities that help recipients grow in appreciation of pollinators as well as learn strategies to support these important species. Over 200 kits have been distributed to Minnesota educators and community leaders, indicating the significant interest in both pollinator education and support. Increased public awareness of pollinator needs can lead to increases in native planting in homeowners' yards, as well as broad popular support for efforts to preserve these species.

Wildlife Viewing

Strategy 3: Provide and promote wildlife viewing opportunities and programs that inspire curiosity, caring, and connection with Minnesota's SGCN.

Wildlife viewing is one of the fastest growing types of wildlife-related recreation in the U.S. and is highly popular in Minnesota, making wildlife viewers an essential audience to lead towards SGCN conservation actions. The 2022 National Survey of Hunting, Fishing, and Wildlife-Associated Recreation (National Survey of Wildlife Recreation) defines wildlife viewing as “closely observing, feeding, and photographing wildlife, visiting parks and natural areas around the home because of wildlife, and maintaining plantings and natural areas around the home for the benefit of wildlife.” (U.S. Department of the Interior, U.S. Fish and Wildlife Service, 2022) In Minnesota, 3.6 million people (ages 16 and older) watch wildlife, according to the National Survey of Wildlife Recreation. For comparison, this number far exceeds the combined number of people (ages 16 and older) who fish and hunt in Minnesota, 1.7 million and 504,000 respectively. This same survey estimated that wildlife viewers spent \$5.0 billion in 2022 on wildlife watching activities in our state.

By engaging Minnesotans in wildlife viewing, an activity that fosters connections between wildlife and people's lives, we can support the conservation of SGCN, as has been demonstrated in multiple research studies (Grooms et al., 2023; Cooper et al., 2015; Hvenegaard, 2002; McKinley et al., 2017). One of the many ways that wildlife viewing engagement could support conservation is through financial support. According to a 2025 survey from Charities Aid Foundation, the leading motivation for Americans to support a cause was feeling a personal connection to it and the leading factor for their decision to donate to a charity was because it was tied to a cause they've personally experienced or care deeply about (Vogels, 2025a; Vogels, 2025b). Additionally, by supporting wildlife viewing-based tourism in Minnesota, wildlife viewers will provide income to the conservation efforts of the sites they visit, which in turn has a beneficial ripple effect to those local economies through their tourism expenditures. By communicating these economic benefits, we can garner further support for SGCN conservation from a wider variety of audiences (i.e. state and community leaders), by demonstrating how SGCN and their habitats benefit local economies.

Wildlife viewing programs that focus on birds and mammals stand to be particularly effective. These are by far the most popular species groups to view by Minnesotans. Each group was selected by 78% of respondents in the 2022 Minnesota Results of the National Wildlife Viewer Survey as the specific types of wildlife they were interested in viewing (Sinkular et al., 2022). However, it's important to note that 87% of respondents indicated they were interested in viewing more than one type of wildlife, showing potential that this strategy could support the other species groups of interest reported, which included native fish, reptiles, insects, and amphibians (listed in order of percent of people interested from highest to lowest).

Bird watchers (or birders), in particular, have been identified as a key group that stands to make a positive impact for wildlife conservation. As defined in *Birding in the United States* (USDOI, USFWS, 2022), a birder is someone who has, “taken a trip one mile or more from home for the primary purpose of observing birds or closely observed or tried to identify birds around the home.” Participation in bird watching has been found to significantly impact pro-environmental behavior (Larson et al., 2018). One study (Cooper et al., 2015) quantified that wildlife recreationists—both hunters and birdwatchers—were each 4-5 times more likely than

non-recreationists to engage in conservation behaviors, including activities such as donating to support local conservation efforts, enhancing wildlife habitat on public lands, advocating for wildlife recreation, and participating in local environmental groups. Demonstrating this behavior in Minnesota, hundreds of bird watchers of all experience levels collected data in a field survey between 2009-2013 for the [Minnesota Breeding Bird Atlas](#), a collaborative project led by Audubon Minnesota and the University of Minnesota's Natural Resources Research Institute, that documents the distribution of every breeding bird in the state, providing a foundation for the conservation of these species.

Not only do birders demonstrate conservation behavior, but they are also a large group with significant economic might. There are 96 million birders (ages 16 and older) in the United States and birding participation rates are highest amongst those who have higher annual household incomes (USDOI, USFWS, 2022). Birders' expenditures supported \$279 billion to the economy, 1.4 million jobs, and \$90 billion in job income (at the local, state, and national level). By engaging Minnesotans in the recreation of wildlife viewing, we may enhance people's understanding and appreciation of wildlife and promote their likelihood of supporting conservation of SGCN through habitat stewardship, participatory science, and donations.

To ensure we effectively connect the people of Minnesota to the recreation of wildlife viewing, our recommended actions address the barriers and motivations for the activity, as identified by Minnesota's state-level results from the 2022 National Wildlife Viewer Survey. Our recommended actions follow the survey's recommendations to provide and promote the following: (1) wildlife viewing information, including where to go and how to do it, (2) access to wildlife viewing locations, (3) around-the-home viewing opportunities, (4) social support networks for wildlife viewers, and (5) services and information that support those who are new to the activity. To sustain and increase the number of wildlife viewers in Minnesota, our recommended actions also utilize the recreational model of recruitment, retention, and reactivation (Heberlein et al., 2008; Larson et al., 2014; Quartuch et al., 2017), most often applied to the wildlife recreational activities of hunting and angling, but also applicable to wildlife viewing (Grooms et al., 2020).

SGCN Example: Cape May Warbler

Migratory birds, such as the [cape may warbler \(*Setophaga tigrina*\)](#), face several stressors, particularly in developed areas, as they migrate between their wintering and breeding grounds each spring and fall, such as lack of suitable habitat, window collisions, light pollution, and climate change. As reported by the Minnesota Breeding Bird Atlas, cape may warblers have high mortality during migration. Loss et al (2014) found that their risk of mortality from collisions with low-rise buildings is 16.7 times more than the average risk for a bird. Cape may warblers migrate throughout Minnesota in the spring and fall to and from their breeding grounds in Northeast Minnesota and Canada. Fortunately, a high number of Minnesotans are interested in seeing birds (as described above) and birders have a particular interest in seeing warblers. By capitalizing on this existing interest in birds and especially highly attractive birds, such as cape may warbler, and engaging more folks in bird watching (a relatively easy species group to observe), an opportunity is created to increase public interest and knowledge of the birds on Minnesota's SGCN list and the stressors they face. We can leverage the relevance of migratory birds to peoples' lives and shift human behaviors towards bird-friendly conservation actions, such as those recommended in Strategy 5 on habitat stewardship.

Recommended Actions for Wildlife Viewing Efforts:

- 1. Develop a state Wildlife Viewing Plan for Minnesota, in collaboration with stakeholders and partners, that focuses on SGCN and addresses recommendations from the Minnesota Wildlife**

Viewer Survey (Sinkular et al., 2022) while tying into and promoting the state’s many existing resources and programs. The following actions provide a sample of what the plan could include.

2. Develop and promote introductory resources, programs, and social networks to support people who are new to the activity of wildlife viewing that will help them get started.

- a. To recruit people into the activity of wildlife viewing, provide basic information and instructional experiences. Create resources describing how to get started, (e.g. where to go to see birds and information on binoculars, field guides, and identification apps). And by promoting, providing, and supporting introductory activities and workshops, people can gain hands-on experience in a comfortable setting (e.g. guided instructional bird walks or workshops on how to find and identify birds). To help encourage wildlife viewing of other easy to observe taxa, such as butterflies and dragonflies, refer to the models created for introductory birding resources and programs.
- b. Provide loaner equipment (e.g. a library check-out program- backpacks with binoculars and easy-to-use field guides, etc. and tips for getting started), so that people can easily try out the activity without having to make a financial investment.
- c. Create a sense of community and social support to further enhance recruitment of Minnesotans into wildlife viewing. This can be done by promoting and supporting wildlife viewing events (e.g. birding festivals or smaller scale “meet-ups”) and connecting folks with existing social networks in Minnesota (e.g. local birding organizations). An important form of social support for wildlife viewers are their families (Sinkular et al., 2022), so providing and supporting programs and resources that engage families to observe wildlife together would be valuable. One example would be to collaborate with the DNR’s Parks and Trails Division to develop an I Can Bird! program, utilizing their successful I Can! program structure.

3. Build interest in wildlife viewing amongst broader audiences by appealing to other, related interests.

- a. Cross-promote wildlife viewing opportunities with other popular outdoor activities taking place in natural spaces (e.g. camping, trail running, hiking, paddling, disc-golf), creating awareness of SGCN in those areas
- b. Develop wildlife viewing games that get people outdoors observing wildlife who are motivated by the challenge of it (e.g. birding bingo/ scavenger hunts, a SGCN observation “bucket list,” or promoting/ hosting City Nature Challenges.)
- c. Collaborate with outdoor affinity community groups to hold introductory wildlife viewing events. Create these as an opportunity for members of the community who may be more experienced and comfortable with the activity to provide leadership and inspiration for other community members to participate.
- d. Work with the Minnesota Master Gardener organization to promote planting of native plants gardens for attracting butterflies, bumblebees, and birds to gardeners.

4. Provide and promote access to wildlife viewing opportunities that enable people to observe and learn first-hand about Minnesota’s SGCN.

- a. Create and promote online resources and printed materials that clearly identify and illustrate top locations to go wildlife viewing in Minnesota. A large-scale example of such a resource

could be creating a statewide wildlife viewing designation system, highlighting Minnesota's top wildlife viewing destinations, including site designation signage and accompanying digital guides/maps to those sites with descriptions of the locations describing what wildlife you can see there. Smaller scale examples of these resources could include a series of local level guides to birding in the state's popular travel destinations (e.g. birding guides to the Twin Cities, Duluth, North Shore, etc.). Partnering with [Explore Minnesota](#) and local tourism and historical societies will help expand the promotion of these resources.

- b. Explore developing small grant programs to provide funding for improvements to or installation of new wildlife viewing infrastructure (e.g. platforms, trails, viewing blinds, interpretive signage) that highlight and enhance SGCN wildlife viewing opportunities at parks, nature centers, and other high visitation natural areas. Sites participating in the above described statewide wildlife viewing system (Action 4a) could be prioritized.
- c. Develop live-streaming wildlife cameras of lesser known or particularly sensitive SGCN, and couple them with interpretation, to increase public interest and knowledge about species that they may otherwise have no opportunity to observe.
- d. Provide and promote guided public program opportunities that enable people to observe SGCN that are challenging for people to observe on their own. For example, the easiest entry into viewing native fish (a species group of high interest to Minnesota's wildlife viewers) is through guided experiences with professional biologists or naturalists with access to specialized equipment and identification knowledge—this could potentially include guided activities such as freshwater snorkeling, angling, microfishing, seining or electrofishing demonstrations.

5. Provide program opportunities and resources that support wildlife viewing around and near people's homes where most wildlife viewers participate.

Most of Minnesota's wildlife viewers participate in wildlife viewing around and near their homes (Sinkular et al., 2022). Creating and promoting community habitat stewardship engagement efforts (detailed in Strategy 5) will provide easily accessed opportunities to view wildlife. For instance, encouraging people to install native plant gardens in their home landscaping or at a local school (Strategy 2) creates easily accessed bird and butterfly viewing opportunities. Such activities could be further enriched by promoting participatory science projects (Strategy 6) that involve data collection of wildlife using these native plant gardens, educational curricula on the birds and butterflies (Strategy 2) observable in these spaces, and development of regional pocket identification guides to wildlife that people may observe in native plant gardens that includes SGCNs (see Action 6c).

6. Create fun, engaging opportunities for wildlife viewers to directly support SGCN conservation through their participation in the activity of wildlife viewing itself.

- a. Collaborate with partners and tourism entities to develop guided interpretive tours of charismatic Minnesota SGCNs, in which participation fees directly support SGCN conservation (e.g. supports habitat work in the area of the tour or goes into Minnesota's Nongame Wildlife Fund). These charismatic species can also work as the draw to provide interpretation on lesser known SGCN that share the same habitat. One possibility could be leveraging Blue Mounds State Park's bison buggy tours to also show and provide education about other prairie SGCN observable during the tour.

- b. Hold a birding competition that supports SGCN conservation by applying a portion of the registration fees as a donation to a conservation program (e.g., a nonprofit organization or the Nongame Wildlife Fund).
- c. Develop a series of pocket identification guides to Minnesota's wildlife that incorporate conservation actions and ethics. An example might be a guide to butterflies that also encourages planting of the host plants that attract them or a guide to birds that describe simple actions to help them (see Strategy 5).
- d. Recruit wildlife viewers to volunteer for participatory science opportunities (see Strategy 6) to engage them in deeper learning of the taxa they are interested in and taking conservation action for SGCN by collecting needed data.

Case Study: Twin Cities Bird Festival creates community amongst bird enthusiasts while promoting conservation actions

This festival, organized by the Urban Bird Conservation Partnership of the Twin Cities (a participant in the [Urban Bird Treaty Program](#)), celebrates World Migratory Bird Day (WMBD) and seeks to connect local communities with the birds around them. Hosted in 2025 by [Three Rivers Park District](#) at Mississippi Gateway Regional Park with partners providing additional bird watching experiences throughout the Twin Cities, the event provided numerous opportunities for urban residents to observe and learn about birds. The guided bird walks, presentations and exhibits were centered around the WMBD theme of creating bird-friendly cities and communities. The Twin Cities Urban Bird Festival is a great example of a partnership involving multiple conservation groups that is striving to engage folks in bird watching and connect bird watchers to conservation actions. The inaugural 2025 festival was attended by over 500 people of all ages and birding experience levels, showing that there is demand for wildlife viewing events such as this one and holds great promise for growth. This festival addresses multiple recommended actions: creating a sense of community and social support for bird watchers, including families; opportunities to observe and learn about birds firsthand on guided bird walks; supporting those who are new to birding with guided experiences and access to equipment; and an opportunity to engage urban audiences in local bird watching without having to travel far from home.

Case Study: Minnesota DNR's FalconCam engages people with a vulnerable bird

For fourteen years, the DNR's [FalconCam](#) has streamed video of peregrine falcons nesting atop a skyscraper in downtown Saint Paul. Watched by over 20,500 people in 2024 and over 12,000 people subscribed to FalconCam emails, it engages people in closely observing the breeding behaviors of this SGCN and state listed Species of Special Concern. Through an intimate look into their nesting lives of two parent birds rearing their young, the FalconCam enables its viewers to feel a connection to the species. It also creates awareness of the fascinating wildlife living in our cities and provides an opportunity to draw attention to the previous plight of peregrine falcons threatened by pesticides, the conservation success story that followed, and the mission of the DNR's donation-supported Nongame Wildlife Program.

Responsible Recreation

Strategy 4: Foster responsible recreation practices that reduce recreational impacts to SGCN and their habitats while allowing for safe and satisfying access to the outdoors.

Minnesotans benefit from abundant outdoor recreational opportunities throughout the state and outdoor recreation has been growing in popularity and importance to residents' lives (Minnesota DNR, 2017). Overall, recreational use of public lands can enhance the value of these spaces by raising awareness, increasing the importance placed on outdoor spaces by members of the public, and encouraging sustainable use and stewardship (Larson et al., 2018). Connections with nature and the outdoors also positively impact human health and wellbeing and encourage proactive conservation behaviors (Richardson et al., 2020). While outdoor enthusiasts in general want to preserve and protect the resources they value, inadvertent impacts from outdoor recreation activities can be detrimental to SGCN and their habitats through lack of awareness, misunderstanding of proper etiquette and techniques, or social tolerance of destructive behaviors.

Habitat destruction, the spread of invasive species or diseases, and pollution are all stressors that can result from outdoor recreation. These impacts are not mutually exclusive and may impact multiple SGCN within the same habitat simultaneously. For example, recreation that covers large distances such as canoe or OHV and mountain biking trails can spread invasive species, disrupt or destroy nesting habitat for turtles, salamanders, snakes, and tiger beetles, and may cause direct mortality if animals or nests/dens are run over by vehicles or bikes. Recreation on lakes can impact common loons and other waterbirds when nests and shorelines are damaged by boat wakes and pollution is introduced from lead fishing tackle. Multiple bat species can be seriously affected by recreation in caves by the spread of disease or disturbance during hibernation. Peatland habitats, such as bogs and calcareous fens, are particularly fragile ecosystems that are also sensitive to human encroachment.

Recreation can also impact SGCN by affecting their behavior. For example, some SGCN species like Canada lynx (*Lynx canadensis*) and American goshawk (*Accipiter gentilis*) are particularly sensitive to human activity, such as snowmobiling for the lynx and general human disturbance for the goshawk. These sensitive species may alter their movement patterns and activity or abandon otherwise suitable nesting territories, respectively (Squires et al., 2019; Morrison et al., 2011).

This strategy offers guidance for land managers, conservation educators, outdoor program and recreation leaders, and natural resource communicators to reduce negative impacts from outdoor recreation activities, encourage effective resource management, and foster positive behavioral changes through raising awareness and providing education focused on environmentally responsible recreation. Many of the actions outlined below focus on social marketing campaigns, which are a proven tool for informing and educating the public and affecting sustained behavioral change that contributes to natural resource conservation (McKenzie-Mohr & Tabanico, 2024). Successful messaging highlights the positive effects produced by individual actions and the cumulative effect on the environment or resource. Perceived impact of individual actions is a significant predictor of future behaviors (Lawhon et al., 2013) and people are more likely to act when they believe their actions make a difference. These tactics emphasize the importance of individual decisions, empower

individuals to feel capable of making a meaningful difference, and encourage people to promote responsible recreational behaviors as a social norm.

Stressors Addressed: Recreation (e.g. accidental take and overuse of resources), Hunting and Collecting Animals (e.g. improperly discarded gut piles causing potential exposure to lead), Fishing (e.g. use of lead tackle), Invasive Species (Problematic Non-native Species), Diseases and Pathogens, and Water-borne Pollution

SGCN Example: Northern Barrens Tiger Beetle

Tiger beetles are an intriguing group of species that feed on insects including agricultural pests, but they are increasingly threatened by anthropogenic impacts, with eight species listed as SGCN in Minnesota. Tiger Beetles are mostly found in open sandy habitats like beaches, dunes, savannas, riverbanks, and trails. They are dependent on these habitats to mate, lay eggs, construct larval burrows, and hunt for prey. Unfortunately, a substantial amount of their habitat in Minnesota has been lost to development, and much of their remaining habitat is subject to negative impacts from recreation. For example, the [northern barrens tiger beetle \(*Cicindela patruela patruela*\)](#) is native to areas in central and southern Minnesota that are also popular areas for outdoor recreation, making it particularly vulnerable. Habitat degradation stemming from off-highway vehicles (OHV) used off trails or in non-approved areas, camping in non-designated areas, or even hiking off-trail can have negative impacts. Temporary effects like soil compaction or displacement can kill adults, damage burrows, and disrupt egg-laying. Over time, repeated disturbance can render areas uninhabitable. Responsible recreational behaviors such as staying on designated trails and avoiding disturbance of other areas can lessen the negative impacts on tiger beetles, but effective communication is key. Most outdoor enthusiasts are prepared to alter their behaviors if they know why the change is needed or beneficial and knowing that specific actions will benefit at-risk animals or plants while maintaining or enhancing the recreational experience can be particularly motivating (Abrams et al., 2020). Increased attention and focus on creating specific, actionable messages are needed to create awareness around these issues and the need for behavior changes.

Recommended Actions for Responsible Recreation Efforts

1. **Collaborate with park and natural area managers to share best practices that support responsible recreation.**
 - a. Host a speaker series or symposium on topics related to creating low-impact, satisfying recreational access and its benefits for sensitive wildlife and habitat. Example topics for wildlife viewing recreation include bird blinds, wildlife viewing platforms, and wildlife cameras. Example topics addressing all types of recreation could include effective boardwalks and interpretive/instructional signage or public art installations that increase knowledge of the vulnerable SGCN and habitats on the property and how to reduce one's impact while visiting/recreating to avoid inadvertent harm.
 - b. Promote and provide information to park and natural area managers on the installation of boot brush kiosks that encourage visitors to wipe mud off their boots to prevent the spread of invasive plant seeds and earthworm egg cases, which damage habitat. See the "PlayCleanGo: Stop Invasive Species in Your Tracks" program for [example boot brush kiosk signage](#) that shares consistent messaging.
2. **Develop highly positive, fun, and memorable key messages--using social science on behavior change and social marketing--that promote and create norms around specific desired behaviors that reduce human impact on SGCN and their habitat.**

Use these key messages in campaigns, programs, materials, and products to promote the desired norm. Some ways to disseminate these messages are to collaborate with the DNR's Division of Parks and Trails and Division of Fish and Wildlife on incorporating these messages into their recreation education programs and materials. For example, the creation of a SGCN mascot (an ambassador species—see strategy 1a) that could help deliver the above messages similar to Smokey Bear would add to their appeal.

- 3. Leverage resources from national initiatives (e.g. Tread Lightly!, Leave No Trace, and PlayCleanGo) to communicate established responsible recreation practices and draw the connection to how these practices benefit SGCN by reducing inadvertent human-caused threats (e.g. not transporting potential vectors of pest and disease transmission).**
- 4. Collaborate with organizations already conducting outreach on aquatic and terrestrial invasive species to incorporate information about specific SGCNs that benefit from their recommended actions.**

The DNR has a variety of programs conducting outreach on invasive species that provide an opportunity for collaboration and inclusion of SGCN information. The DNR Division of Forestry's PlayCleanGo Program provides an opportunity for incorporating messaging relevant for local OHV clubs, mountain bike groups, and other recreationist groups. The DNR's Invasive Species Program provides behavior change messaging about aquatic invasive species to audiences like boaters, anglers, and shoreline owners. They also provide "[Prevent the Spread](#)" messaging about terrestrial invasive to species to a wide variety of outdoor recreationists.

- 5. Design and create infrastructure and materials that make it easy for people to learn and adopt responsible recreation behaviors by including specific information with "how to" style instructions.**

Placing information regarding responsible recreation behaviors on-site at recreational areas or including it in resources that the recreationists use is a way to easily raise awareness and make adoption of the behavior more likely. One example is providing simple stations at recreation access sites such as trailheads, fishing piers, or boat launches for people to take desired actions (e.g. boot cleaning, fishing line disposal, ethical canoeing practices) with accompanying signage explaining the what, why, and how of the desired actions. To reach wildlife viewers, incorporate information on ethical wildlife viewing practices within wildlife identification and visitation guides and other wildlife viewing materials/content, to ensure wildlife viewing recreationists are knowledgeable about issues and avoid harmful behavior.

- 6. Collaborate with organizations, groups, and businesses that can help develop and deliver relevant and tailored messaging to increase awareness amongst the recreationists with which they regularly interact on SGCN that are vulnerable to harm through their recreational activities and how to avoid inadvertent injury/harm to them.**

The following entities are example organizations/ businesses that could potentially help with message development and delivery: [Minnesota Outdoor Recreation Industry Partnership](#); specific recreational groups, such as OHV and paddling clubs; lake associations; and outdoor outfitters.

Case Study: Clean in, Clean Out provides a simple message to prevent the spread of Aquatic Invasive Species

The DNR's [Clean In, Clean Out](#) program is a comprehensive outreach campaign and resource to help Minnesotans avoid the spread of aquatic invasive species (AIS) throughout the state. The program is especially effective due to its positive messaging that connects people with the resources they care about, explains why AIS are an issue, empowers people to make meaningful conservation choices, and perhaps most importantly, explains exactly what actions to take based on the activity or gear involved. The main website explains that:

"Invasive plants, animals and diseases can harm lakes, rivers, streams, wetlands, water quality, and wildlife habitat. People can unintentionally spread invasive species by moving boats, gear, and equipment from one waterbody to another or releasing non-native species into the environment. No matter how you enjoy the outdoors, always take these actions to prevent the spread of aquatic invasive species (AIS)."

The website has information about a wide variety of common aquatic activities like boating, fishing, and paddling, as well as resources that apply to people engaged in potentially less common activities like water gardeners, aquarium owners, and commercial bait harvesters. There is specific information about how AIS impact different gear and activities, specifically what to do to minimize risk of spreading AIS, instructions to decontaminate gear, relevant laws and regulations, and important links to additional resources. The website is a wealth of information, and the message is simple and clear: We all play an important role in protecting Minnesota waters from aquatic invasive species.

Habitat Stewardship

Strategy 5: Engage and empower communities to improve wildlife habitats where they live, work, and recreate.

Land use change and the associated loss of habitat and habitat degradation is the number one driver of species loss and population decline for terrestrial species worldwide (Jaureguiberry et al., 2022). When lands are converted from natural landscapes to human-dominated ones, wildlife are challenged to find suitable habitat within a matrix of development and heavily modified ecological landscapes. Even though many landscape scale changes in habitat cannot be reversed through small-scale actions, anyone with access to a patch of land, whether it be small urban lots or large rural acreages, can help improve habitat in their domain. Collective efforts by individuals and communities have the ability to make a difference for some SGCN, particularly those that use developed areas (see Urban and Other Developed Lands chapter). This includes some migratory birds, pollinating insects, and bats seeking food sources; suitable nesting, egg laying and roosting sites; and migration stopover sites. Even the simple choice to include native plants in one's home landscaping, can provide critical benefits to wildlife (Tallamy, 2009; Tallamy et al., 2021; Tallamy & Shriver, 2021). For example, the use of native landscaping in suburban areas has been found to significantly support the abundance and diversity of birds, butterflies, and moths with one study finding that bird species of regional conservation concern were eight times more abundant and significantly more diverse on native landscaped properties than those landscaped with non-native plants (Burghardt et al., 2009). When individual actions are replicated throughout a community, these small-scale habitat improvements can make a powerful, large-scale change for wildlife in urban, suburban, and agricultural areas.

Our recommended actions for habitat stewardship primarily focus on practices beneficial to migratory birds, pollinating insects, and bats as there are simple conservation actions for these groups that can be easily and

practically employed by many people in most communities. Also important, these species groups hold relevancy to Minnesotans; appealing to their wildlife viewing species interests (Sinkular et al., 2002) and in the case of bats, their interests in mosquito control to better enjoy the outdoors (Minnesota DNR, 2017) or encouraging bats to roost outdoors rather than in their homes.

In addition, when engaging members of waterfront communities, we leverage Minnesotans' passion for our state bird by highlighting the benefits to the common loon while encouraging stewardship of their lake shoreline habitats.

When engaging rural communities in habitat stewardship, we highlight practices that can support grassland birds and turtles that use upland and wetland habitats like wood turtle and Blanding's turtle—all of which would stand to benefit from habitat enhancements in these settings and may hold appeal for the landowners. Our recommended actions take a community approach (to maximize habitat benefits through collective action), create relevancy for communities by highlighting taxa of interest to wildlife viewers, and address potential societal barriers, including overcoming social norms and lack of knowledge and skill.

Stressors Addressed: Development, Crop Production, Timber Harvest, Tree Plantations, Air-borne Pollution, Water-borne Pollution, Light and Noise Pollution, Invasive Species (Problematic Non-native Species), and Fire Management

SGCN Example: Rusty patched bumble bee

The [rusty patched bumble bee \(*Bombus affinis*\)](#) was once widespread and abundant across eastern North America including Minnesota but has disappeared from much of its historic range over the past 30 years, although still abundant in Minnesota. The cause of this decline is not fully understood, but disease, pesticides, and habitat loss have likely played roles. The bee was listed as federally endangered in 2017.

In 2019, the rusty patched bumble bee was named Minnesota's state bee to raise awareness of its conservation status and the importance of protecting pollinator habitat. Like many other bumble bees, rusty patched bumble bees forage on a variety of flowering plants for pollen and nectar. Bumble bees nest in cavities above or below ground and often use brush piles, leaf litter, or abandoned burrows for shelter. Rusty-patched bumble bees do not avoid developed areas, so by planting native plants that bloom throughout the growing season, reducing the use of pesticides, and leaving nesting areas alone, Minnesotans can help support our endangered state bee and other pollinator species.

Recommended Actions for Habitat Stewardship:

- 1. Educate local communities, landowners, and land managers on the habitat needs of the SGCN sharing their local area, highlighting migratory birds, pollinating insects, bats, and other species relevant to the community type/geographic area, as applicable.**
- 2. Provide coordination that connects communities to programs and resources for technical, financial, and organizational support, so that they can more easily implement habitat stewardship actions for SGCN. Engagement of the 11 federally recognized Tribes with Minnesota's borders is recommended as part of the process to ensure their access to these programs and resources and that this support meets their needs and interests. Specific examples of coordination are provided below.**

- a. Collaborate with conservation organizations and agencies to develop resource information hubs that compile the many existing resources on what individuals can do to improve habitat into one convenient place.
- b. Develop a train-the-trainer program for folks to become Master Habitat Stewards-- e.g. utilizing Soil and Water Conservation District educators, Master Gardeners, and Master Naturalists--to increase capacity for habitat education, technical assistance, and site visits.
- c. Promote existing stewardship programs to municipalities that empower them to engage their communities and parks departments in taking actions directly benefiting wildlife. Examples of programs to promote to municipalities include: National Wildlife Federation's [Mayor's Monarch Pledge](#), American Bird Conservancy's [Bird City](#), Minnesota Board of Soil and Water's [Pollinator Pathways](#) grant program, and Audubon International's [Audubon Cooperative Sanctuary Program for Golf Courses](#).
- d. Develop a plan for the creation of local-level habitat stewardship collaboratives, involving varied stakeholders (see examples below). These groups could serve as forums for building relationships and trust, holding listening sessions, crafting relevant messaging, increasing knowledge of the SGCN and habitats in their community, and maximizing capacity for on-the-ground implementation of habitat enhancement projects.
 - Example stakeholders for Urban/suburban Communities: NGOs and agencies involved in community habitat enhancement work (e.g. Great River Greening), Soil and Water Conservation Districts, local parks departments, transportation departments, local influential leaders (mayors and city councils), schools, libraries, and well-being centers.
 - Example stakeholders for Rural Communities: farm bureau, farmers union, Minnesota State Cattlemen's Association, Pheasants Forever, Ducks Unlimited, Minnesota Deer Hunters Association, Minnesota Chapter of the Wildlife Society, North Shore Forest Collaborative, Forestry for the Birds Initiative, Soil and Water Conservation Districts, Natural Resource Conservation Service's Farm Bill biologists/private lands biologists, and Minnesota Corn and Soybean Growers Associations
 - Example stakeholders for Waterfront Communities: Homeowners Associations, Watershed Organizations, Tribal governments, and researchers

3. Craft communication resources, campaigns, and media with messages that increase public understanding on how the health of habitats, which we share with SGCN, is directly tied to the individual choices Minnesotans make on their own properties and how collectively, our stewardship actions make a difference for both wildlife and humans.

- a. Promote the benefits of stewardship actions for humans (e.g. well-being, cost savings, wildlife viewing opportunities) and for wildlife, celebrating habitat as a healthy space for both people and SGCN. This action could connect with actions in Strategy 3.
- b. Promote actions that have a clear connection between larger scale benefits to wildlife and humans, resulting from our individual, smaller actions. For instance, collaborate with local watershed organizations to promote connectivity messaging on how what individuals do on their property impacts the health of our local lakes and rivers. Other messages could include promoting how planting a pocket of milkweed for monarchs builds habitat across the

landscape, how turning “lights out” supports migrating birds and reduces light pollution, and how by landscaping with native plants, we increase food and nesting opportunities for birds.

4. **Create and distribute materials and media that promote and enhance application of everyday stewardship actions and best management practices for habitats and SGCN. To be most effective these communications should be provided in easily applied formats and translate scientific, technical information into accessible, plain language and in different languages. A list of specific actions to encourage everyday conservation actions includes:**
 - a. **Simplify people’s ability to include native plants in their landscaping by providing and promoting information on what to plant for SGCN, where to buy the plants, and how to incorporate them into their gardens.** Essential resources to develop would be lists of the top beneficial native plants for Minnesota’s SGCN, organized by the state’s ecoregions. These lists should include soil and sun requirements and information on which species groups are attracted to which plants; e.g. birds, butterflies, bumble bees, etc. Helpful partners and resources in this effort could be [Minnesota’s Wild Ones Chapters](#) and [Monarch Joint Venture](#). To help people locate and obtain native plants, develop a database/list of native plant nurseries and seed sources in Minnesota and promote local native plant sales. And finally, to help people incorporate native plants into their landscaping, promote and connect people to resources with how-to information, such as Minnesota’s Board of Water and Soil Resources’ [Lawns to Legumes](#) program and [Ganawenindiwag: Working with plant relatives to heal and protect Gichigami shorelines](#). Updating and reprinting the DNR’s *Landscaping for Wildlife* and *Loonscaping* would provide additional helpful resources.
 - b. **Provide guidance on how people can stop the introduction and spread of invasive species.** Disseminate existing products on (1) how to identify and remove common invasive plants, (2) how to avoid buying invasive plants and plant native alternatives instead, and (3) how to avoid releasing invasive plants and animals into the environment. Create new products as needed.
 - c. **Promote ways people can reduce threats to wildlife around homes.** Guidance should include the following: (1) reducing the use of pesticides and other chemicals in yards, why it matters for wildlife, and information on alternatives that are less impactful to wildlife; 2) methods to prevent bird window collisions, highlighting available programs and resources (e.g. American Bird Conservancy); 3) the importance of snags for cavity-nesting birds, encouraging people to maintain them and/or supplement them with bird boxes; 4) keeping cats indoors for the benefit of birds by utilizing available resources (e.g. from [American Bird Conservancy](#) and the [Association of Fish and Wildlife Agencies](#)) and by working with veterinarians and Department of Public Health to help amplify these messages; and 5) turning off of exterior lights to reduce light pollution, explaining its impacts on migrating birds, moths, and fireflies, and by highlighting programs such as Audubon’s Lights Out Campaign and citing reports showing timed/less lights don’t impact homeowner safety.
 - d. **Promote bat boxes, along with [siting and design recommendations](#).** Little brown bats (*Myotis lucifugus*) commonly roost in artificial structures including human residences and bat boxes (Minnesota DNR, 2025). A statewide survey in Minnesota documented little brown bats as the primary species utilizing bat boxes, with colony sizes ranging from 20 to nearly 600 individuals. When little brown bats are excluded and displaced from a building or human structure, their preferred summer roost sites, bat boxes can provide alternative roosting habitat. They also provide a safe location for maternity colonies to raise their pups, which can

support the recovery of Minnesota's little brown bat population from white-nose syndrome. (DNR, 2025) Installing bat boxes is a relatively easy way for people to support little brown bats in Minnesota, however studies have shown these structures must be properly constructed, sited, and monitored through time as these long-lived bats will rely on the bat houses to raise pups for many years. Therefore, when promoting bat boxes, it is important to provide information for best practices and to promote stewardship of bat houses in order to truly benefit the species.

- e. **When working with waterfront communities, include additional information relevant to aquatic habitats:** 1) promote the importance of natural shoreline for water quality, loon habitat, and other aquatic wildlife; 2) provide plant lists of native riparian vegetation that benefit loon, other waterbirds and pollinators; and 3) encourage [behavior change to reduce the spread of aquatic invasive species](#).
- f. **When working with private landowners, provide information on best management practices (BMPs), applicable to larger acreages.** Specific recommendations include: 1) develop easily accessed products in plain-language formats to share and promote BMPs for managing SGCN habitats, such as shorelines for loons, prairie for grassland birds and pollinating insects, and forests for birds, bats, wood turtle and other forest-dependent herpetofauna; 2) promote and distribute invasive plant BMP materials for managing large landscapes; 3) provide information on how to recognize and enhance the native plants beneficial to wildlife that are already on their lands, e.g. milkweeds; and 4) promote the importance of prescribed fire, in combination with retained areas of refugia, as a necessary habitat management tool and provide information to support capacity for conducting prescribed fire in Minnesota. University of Minnesota Extension's [My Minnesota Woods](#) and [Minnesota Woodland Steward program](#) provides resources for this work.

Case Study: Lawns to Legumes enables homeowners to create pollinator habitat

The [Lawns to Legumes](#) program by Minnesota's Board of Soil and Water Resources (BWSR) exemplifies several of these recommended actions. This stewardship program helps people to install pollinator-friendly native plantings on residential lawns. They offer workshops, coaching, planting guides, and grants to individuals. Additionally, they offer Pollinator Pathways grants, which are pollinator programs run by local governments and nonprofit organizations with support from BWSR, and a public education campaign to raise awareness about creating pollinator habitat. According to their website, in their 2024/2025 fiscal year, the Minnesota Board of Water and Soil Resources (BWSR) offered over \$1.1 million in grant funding to promote and establish pollinator plantings through a residential and/or educational-based approach.

Case Study: Minnesota Loon Restoration Project engages waterfront communities in restoring shoreline for loons

The DNR's [Minnesota Loon Restoration Project](#) provides an example of how to engage waterfront communities. The project supports lake associations as they develop management plans that encourage loon-friendly activities and habitat restoration on the lakes and it educates the public about loon conservation. The Minnesota DNR has recruited lake associations to participate in the "Loon Friendly Lake Registry." Through presentations at annual lake association meetings and support in the development of lake-specific management plans, the DNR has empowered owners of lake shorelines to restore and conserve shoreline habitat. There are currently 64 lake associations enrolled in the Loon Friendly Lake Registry. In 2024 alone, the Loon Restoration Project reached approximately 2,500 people in over 40 live speaking events/booths to discuss and present on loon conservation in the state.

Participatory Science

Strategy 6: Offer participatory science programming for Minnesotans to take part in conservation volunteerism that addresses SGCN data needs.

Participatory science (also known as citizen science, community science, or crowd-sourced science) engages the public in scientific projects that produce reliable data and information, often through public data collection, that may be used by scientists, decisionmakers, or the public (McKinley et al., 2017). Involving non-professional participants in wildlife research and data collection increases program capacity and fosters ongoing public involvement in conservation. One of the most significant advantages of participatory science collaborations is the ability to collect data at a scale well beyond the normal reach for many agencies and organizations (Dickinson et al., 2010), including the DNR. Participatory scientists help fill the gap by collecting data across wide geographic areas and over long periods of time. These datasets are invaluable to conservation at local, regional, and international scales (McKinley et al., 2017). A few longstanding examples are the [Breeding Bird Survey](#), [eBird](#), [Minnesota Loon Monitoring Program](#), and [Minnesota Frog & Toad Calling Survey](#). Ongoing public participation fosters appreciation for Minnesota's unique natural resources, encouraging lifelong learning, engagement and conservation action.

Minnesota has a strong history of participatory science and conservation. Universities, state agencies, local governments and parks, nongame wildlife interest groups, and others have all contributed to wildlife conservation for many decades. Participatory science has emerged as way to empower communities to learn and invest in their local environments.

Recommended species groups for participatory science projects in Minnesota include, frogs and toads, loons and other birds, bats, butterflies, bees, and mussels. All these groups represent SGCN with data needs, while also being appropriate for volunteer researchers, thus ensuring that participatory science efforts successfully obtain useable data. These species groups are relatively easy for non-experts to identify with minimal training and simple resources and observing them requires relatively little or easily operated equipment. They are also species that can be readily identified with photographs, recordings, or other simple technology. Another important consideration is that all of these recommended species groups are those that Minnesotans are interested in observing (Sinkular et al., 2022) or in the case of mussels, speak to other human conservation interests, like water quality (Duda, 2021). Participatory science projects are reliant upon volunteers, so they

must hold relevancy to people’s interests. The 2025-2035 Minnesota State Wildlife Action Plan seeks to sustain and further develop the collective power of participatory science in Minnesota by supporting existing initiatives, collaborating with partners, and the developing new projects and programs. The recommended actions in this section draw upon social science research of Minnesota’s participatory scientists (Jennings et al., 2024a). They are designed to contribute to a broad framework of opportunities for Minnesotans statewide to engage in participatory science that benefits SGCN and habitat and is meaningful to the interests of our residents.

Stressors Addressed: Participatory science does not necessarily address a specific stressor; rather, it is a solution to address data gaps/ lack of information.

SGCN Example: Blanchard’s cricket frog

The [Blanchard’s cricket frog \(*Acris crepitans*\)](#) is Minnesota’s only endangered amphibian. Historically, they had a patchy distribution in Minnesota, with populations reported in the extreme southwestern and southeastern corners of the state. In the 1960s, their populations declined drastically for unknown reasons, and they were thought to be extirpated from the state.

In the 1990s, a group of participatory scientists found two very small populations of Blanchard’s cricket frogs. For the next 20 years, these were the only known locations of Blanchard’s cricket frogs in Minnesota.

In the mid-2010s, DNR biologists and partners began discovering new, small populations of Blanchard’s cricket frogs throughout their historic range in Minnesota. Hope for the future of the Blanchard’s cricket frog hopped back onto the scene and the Nongame Wildlife Program partnered with HerpMapper and the Amphibian and Reptile Survey of Minnesota to create a Blanchard’s Cricket Frog Survey, using participatory scientists to gather more information about this rare amphibian.

Recommended Actions for Participatory Science:

- 1. Promote existing participatory science initiatives in Minnesota to help recruit volunteers to those projects.**

While several participatory science projects already exist in Minnesota, people need an easy way to learn about these opportunities and increased promotional efforts would help (Jennings et al., 2024b). To help boost promotion, it is recommended to create a Minnesota-based resource hub for wildlife participatory science opportunities. This could be modeled after [The Wisconsin Aquatic and Terrestrial Resources Inventory](#). Some examples of projects that would benefit from additional promotion and that could be part of the resource hub include: University of Minnesota’s Bumble Bee Atlas; [University of Minnesota Extension participatory science projects](#); Monarch Joint Venture’s Integrated Monarch Monitoring Program; and the DNR’s Minnesota Frog and Toad Calling Survey, Minnesota Cricket Frog Survey, Loon Watcher Survey, Minnesota Loon Monitoring Program, and Chimney Swift Survey.

- 2. Collaborate with wildlife biologists and participatory science professionals to identify gaps in existing participatory science programs, identify project priorities, and to understand what specific data would be most useful and appropriate and relevant for volunteers to collect.**

Although Minnesota has several participatory science projects, there are many areas of opportunity with other species groups, including many bird species, bats, butterflies, and other easily identified insects, and mussels – all represent SGCN with data needs and taxa relevant to Minnesotans. These should be explored further with wildlife biologists to determine specific project needs and design.

To enhance the success of newly developed projects, partnerships and collaborations with the following organizations are recommended: 1) museums, science centers, nature centers, and zoos to identify projects of mutual interest and maximize resources for implementation; 2) The University of Minnesota Extension's Minnesota Master Naturalist Program for assistance in designing projects that are appealing and volunteer-friendly, volunteer recruitment, and engagement in train-the-trainer opportunities to maximize capacity; 3) existing conservation projects/ programs to add a community science component (e.g. Lawns to Legumes to add pollinator monitoring opportunity for their grant recipients); 4) conservation agencies and organizations that have community connections (e.g. USFWS National Wildlife Refuges, Local Parks Departments, and community-based non-profit organizations); and 5) entities that may have monitoring needs (e.g. the DNR's Minnesota Biological Survey, Scientific Natural Areas, and Wildlife Management Areas; Pheasants Forever; Ducks Unlimited; and The Prairie Enthusiasts).

3. Develop community science opportunities to collect data on SGCN occupying less developed and rural habitats through unique partnerships.

SGCN inhabiting less populated areas (e.g. burrowing owls and Richardson's ground squirrels), may pose challenges for volunteer recruitment, so unique partnerships should be explored.

Recommendations include: 1) develop a project in partnership with rural postal offices to train mail carriers to identify SGCN of interest and 2) build relationships with educators and collaborate on project ideas, in which students and their families could participate.

4. Engage broader groups of volunteers by providing training, meeting people where they are, and suiting their needs and interests.

- a. Eliminate potential barriers and help folks get started in participatory science by: 1) designing projects for variable skill levels including entry level, offering multiple trainings in varying formats, and creating opportunities requiring different levels of commitment; 2) ensuring easy to use species identification resources are available (this also relates to Strategy 1); and 3) promoting entry-level opportunities (e.g. eBird and iNaturalist), coupled with supportive materials or trainings, to anyone new to participatory science and to educators to encourage youth involvement.
- b. Create locally and personally relevant opportunities for participatory science by 1) encouraging groups to adopt-a-site (e.g. their local park) for on-going and varied community science projects, with recommendations for projects that would be a good fit for the site; 2) designing projects around community interests and needs and build the conservation effort off of that, so that it benefits the participating community as well as the organizations involved (e.g. working with a community to restore bluff habitats to help keep timber rattlesnakes out of their yards); 3) developing and promoting existing participatory science opportunities for collection of data around people's homes (e.g. Project FeederWatch, Monarch Monitoring,

Bumble Bee Watch), which also supports Strategy 3; and 4) promoting participatory science opportunities that can be done without leaving one's home (e.g. Zooniverse and iNaturalist).

5. **Ensure that data from participatory science projects may be accessed for research conducted by conservation organizations and academic institutions. Conduct outreach to promote the use and application of the data.**

Case Study: Minnesota Bumble Bee Atlas volunteers contribute to long-term, regional dataset

The [Minnesota Bumble Bee Atlas](#) is a statewide program run by the University of Minnesota Extension (with support from USFWS and DNR) in which participatory scientists gather population data to track Minnesota's native bumble bees. Volunteers receive training on how to identify and survey bees. Participants select an area within the state in which to survey and run catch-and-release surveys from June to September, submitting all of their data to the Atlas. Each bee that is captured is well photographed so that its identification may be verified by experts after the photographs are submitted. In 2024, more than 240 volunteers conducted almost 200 surveys. Almost 2,400 individual bees were recorded, resulting in valuable data that adds to a region wide, long-term dataset. The scientific power of this program is multiplied because it is part of the Midwest Bumble Bee Atlas, in which a number of other states and lead partner, the Xerces Society, are all using the same approach and protocols to create a seamless dataset across the region.

Case Study: Volunteer Loon Watchers Collect Data to Guide Loon Conservation

Minnesota [DNR's Volunteer Loon Watcher Survey](#) is an excellent example of crafting a participatory science project that is relevant to Minnesotans (in this case focusing on the state bird) and fulfills a SGCN data need. For over 40 years, approximately 200 volunteer loon watchers have recorded their loon observations on lakes and provided great insight into loons. Volunteer observations have provided the Minnesota DNR's Nongame Wildlife Program with data on nesting success, number of loons observed, interesting occurrences and problems that may negatively affect the loons. These data have deepened our knowledge about loons in Minnesota and guided conservation of our state bird.

Financial Support

Strategy 7: Cultivate financial support for the conservation of SGCN through the development and promotion of public funding opportunities and public-private partnerships.

Minnesotans care about the outdoors and the state's natural heritage and connecting to that care is imperative for supporting SGCN conservation and research. By building a stronger relationship between the DNR's Nongame Wildlife Program and Minnesotans through targeted partnerships and marketing, we can leverage their support to help SGCN.

The DNR's Nongame Wildlife Program (NWP) leads the conservation of SGCN, with funding from USFWS State Wildlife Grants (SWG), donations to the Nongame Wildlife Fund, Reinvest In Minnesota (RIM) funds that are generated from purchases of Minnesota's critical habitat license plates (see Case Study below) and the Environment and Natural Resources Trust Fund (Minnesota lottery funds made available through competitive grants). The NWP does not receive any 'base funding' from the state and is reliant on this unique blend of private, state, and federal funds, none of which are guaranteed. Donations to the Nongame Wildlife Fund provide essential matching dollars required to receive both SWG and RIM funds. Donations are obtained through direct giving as well as through a 'check-off' option on Minnesota tax forms, in which tax payers can donate a portion of their tax refund.

While the amount of funds from donations has been relatively stable, the number of individual donations to the NWP have been on the decline (see Case Study below). A survey of NWP supporters (Jennings et al., 2024b) found that lack of awareness was the largest barrier for engaging with the program. They found that 52% of respondents had little to no familiarity with the NWP and 75% listed being "unaware of opportunities" as the main barrier to volunteering for participatory science programs and donating to the program. However, after receiving an introduction to the program's work, 60% of respondents were interested in volunteering and 70% were interested in donating.

In addition, wildlife viewers indicated that they were most likely to increase their contributions to the DNR if they knew their funds would be used for the conservation of rare or vulnerable wildlife species or if used to support conservation of the types of wildlife they like to view (Sinkular et al., 2022). This tracks with another finding that having a conservation impact is a primary driver of contributions for the NWP's current and potential donors (Jennings et al., 2025). Therefore, another approach the NWP can take to motivate public support will be highlighting their conservation project work and especially their success stories, instilling confidence and trust around how donations to the Nongame Wildlife Fund benefit SGCN, like the recovery of bald eagles or trumpeter swans or the improved status of peregrine falcons. It's important to highlight species of relevance to Minnesotans. Species groups to especially highlight include mammals and birds, which most wildlife viewers reported an interest in observing (Sinkular et al., 2022). Another species group to consider highlighting would be mussels, given their connection to water quality, another strong human conservation value (Duda, 2022).

Case Study: Trumpeter Swans Success Story

[Trumpeter swans \(*Cygnus buccinator*\)](#) originally inhabited wetlands across a broad region of North America, including Minnesota. However, hunting and loss of habitat greatly reduced their populations and by the 1930s, only 69 trumpeter swans remained in the lower 48 states. In the 1960s, Three Rivers Park District (then known as Hennepin Parks) began the nation's first major trumpeter swan restoration project. When the Nongame Wildlife Program was founded in 1977, they joined these conservation efforts. However, it became apparent that more funding was needed to bring this iconic wildlife species back to Minnesota. In 1980, the Minnesota legislature passed the Nongame Wildlife Checkoff Law, establishing a tax checkoff on Minnesota state tax forms that allowed Minnesotans to donate a portion of their tax return to the Nongame Wildlife Program. In 1981, Minnesotans donated over \$500,000 to support nongame wildlife research and conservation. Over the next two decades, funds from the Nongame Wildlife Fund supported the trumpeter swan restoration project. Hundreds of swans were released back into Minnesota and today their numbers range in the thousands. This conservation success story demonstrates that we can make a difference for rare wildlife and has the potential to inspire conservation action.

Recommended Actions for Financial Support

1. **Increase public interest in donating to the Nongame Wildlife Fund with strategic marketing and promotions that expand Minnesotans' familiarity with and connection to the Nongame Wildlife Program's mission to conserve SGCN and their habitats.**

Develop a marketing plan for the Nongame Wildlife Program that helps communicate the program's goals in a manner that creates a sense of pride and relevancy and clearly shares how to contribute. Include the following elements within the plan: 1) materials and promotional activities that share the NWP mission, build relationships with future donors, and connect the needs of vulnerable wildlife to Minnesotan's pride in the outdoors and the state's natural heritage; 2) methods to nurture relations with current donors through regular acknowledgment and communications about the NWP; and 3) a survey of Nongame Wildlife Program donors every five years to better understand who they are and evaluate and improve our outreach and engagement efforts.

2. **Reinvigorate critical habitat license plate promotions.**

Promote how the Reinvest in Minnesota Critical Habitat License plate fund benefits the Nongame Wildlife Program and SGCN conservation. Develop new Critical Habitat License plate outreach materials and plates, led by the DNR. Collaborate on critical habitat license plate promotions between DNR programs/divisions and other organizations. Increase Minnesotans' understanding about how these plates provide matching dollars to donations for the Nongame Wildlife Fund, Minnesota Biological Survey, and Minnesota's Scientific and Natural Areas, all of which support SGCN and their habitats.

3. **Promote the Nongame Wildlife Fund Tax Check Off and highlight how donating to the Nongame Wildlife Fund benefits SGCN conservation.**

Develop information products and communication campaigns encouraging Minnesotans to donate to the Nongame Wildlife Fund during the tax season. Work with Certified Public Accountants (CPAs) and other tax professionals to promote the Nongame Wildlife Fund to their clients. Promote the Nongame tax checkoff on corporate tax forms to corporations.

4. **Develop relationships with philanthropic organizations and individuals who have an interest in wildlife conservation.**

Strengthen support for the Nongame Wildlife Program by corporate donors and foundations through researching conservation-inclined philanthropic organizations, relationship building, and the promotion of corporate giving. Leverage existing partnerships and connect with local businesses interested in donating portions of their sales to the Nongame Wildlife Program. Promote the Nongame Wildlife Fund to estate attorneys as an option for planned giving.

5. **Explore new funding mechanisms that could create a new revenue stream(s) for the Nongame Wildlife Fund.**

Collaboration between the Nongame Wildlife Program, local businesses/corporations, and/or the DNR's legislative liaisons to develop new funding mechanisms could generate additional revenue for the Nongame Wildlife Fund.

Case Study: DNR EagleCam fosters connection to the Nongame Wildlife Program

The [DNR's EagleCam](#) was established in December 2012 to raise awareness of the Nongame Wildlife Program and the conservation success story of the bald eagle recovery effort in Minnesota. Over the last thirteen years, the EagleCam has soared in popularity, streaming in over 160 countries across the world. Between 2020-2024, the EagleCam webpage has been visited over 4.1 million times with over 795,000 users. In 2022 alone, the webpage had over 2.2 million views with over 350,000 users. Many people attribute their familiarity with the Nongame Wildlife Program to the EagleCam, and numerous donors to the Nongame Wildlife Fund indicate that the EagleCam motivated their donation.

Case Study: Critical habitat license plates provide double match for donations to the Nongame Wildlife Program

The Minnesota [critical habitat license plate](#) was initiated in 1995 to provide an opportunity for citizens to contribute to conservation of Minnesota's natural resources and show their commitment and individuality with a special vehicle license plate. As of 2025, there are 9 specialty plates featuring both game and nongame wildlife and plant species. Minnesotans can obtain a critical habitat plate with a minimum \$30 annual contribution added to their tab renewal. This contribution goes into the Reinvest in Minnesota (RIM) Critical Habitat Program, which provides a 2-to-1 match for each individual donation made to the Nongame Wildlife Program. Half of the matching funds go to the Nongame Wildlife Fund and the other half go to support the Minnesota Biological Survey and Scientific and Natural Areas, all of which benefit Species in Greatest Conservation Need. In 2024, these plate sales provided 2.3 million for the conservation of SGCN and their habitats.

Case Study: Updated advertising campaign for the Nongame Wildlife Fund boosts Tax Check-off donations in 2025

The Nongame Wildlife Fund has been present on Minnesota state tax forms since 1980. Once a strong source of funding for the Nongame Wildlife Program, donations to the tax checkoff have waned over the last decades. In 2025, the program developed new outreach initiatives for promoting the Nongame Wildlife Fund, such as advertising on Meta (Facebook and Instagram), revamping their donation webpage, and advertising through radio spots on Minnesota Public Radio. Across all advertising platforms, messaging was overhauled to reflect Minnesotan wildlife viewers interest in supporting rare or vulnerable wildlife (Sinkular et al., 2022), coupled with images of wildlife belonging to species groups that they are interested in viewing as found in that same study. The updated and enhanced campaign was a success, resulting in an almost 100% increase in new visitors to the Nongame Wildlife Program’s donation webpage and over 500 more individual donations amounting to a \$30,000 increase in donations compared to the 2024 campaign period.

References for Public Engagement

- Abrams, K. M., Molder, A. L., Nankey, P., & Leong, K. (2023). Encouraging Respectful Wildlife Viewing Among Tourists: Roles for Social Marketing, Regulatory Information, Symbolic Barriers, and Enforcement. *Social Marketing Quarterly*, 29(1), 67–86. <https://doi.org/10.1177/15245004231153085>
- Ardoin, N. M., Bowers, A. W., & Gaillard, E. (2020). Environmental education outcomes for conservation: A systematic review. *Biological Conservation*, 241, 108224. <https://doi.org/10.1016/j.biocon.2019.108224>
- Bergman, J.N., Buxton R.T., Lin H-Y., Lenda M., Attinello K., Hajdasz, A.C., Rivest, S.A., Tran Nguyen, T., Cooke S.J., & Bennett, J.R. (2022). Evaluating the benefits and risks of social media for wildlife conservation. *FACETS*, 7(1), 360–397. <https://doi.org/10.1139/facets-2021-0112>
- Burghardt, K. T., Tallamy, D. W., & Gregory Shriver, W. (2009). Impact of Native Plants on Bird and Butterfly Biodiversity in Suburban Landscapes. *Conservation Biology*, 23(1), 219–224. <https://doi.org/10.1111/j.1523-1739.2008.01076.x>
- Cooper, C., Larson, L., Dayer, A., Stedman, R., & Decker, D. (2015). Are wildlife recreationists conservationists? Linking hunting, birdwatching, and pro-environmental behavior. *The Journal of Wildlife Management*, 79(3), 446–457. <https://doi.org/10.1002/jwmg.855>
- Dickinson, J. L., Zuckerberg, B., & Bonter, D. N. (2010). Citizen Science as an Ecological Research Tool: Challenges and Benefits. *Annual Review of Ecology, Evolution, and Systematics*, 41(1), 149–172. <https://doi.org/10.1146/annurev-ecolsys-102209-144636>
- Dietsch, A.M., Don Carlos, A.W., Manfredo, M.J., Teel, T.L., & Sullivan, L. (2018). State report for Minnesota from the research project entitled “America’s Wildlife Values.” Fort Collins, CO: Colorado State University, Department of Human Dimensions of Natural Resources. <https://content.warnercnr.colostate.edu/AWV/MN-WildlifeValuesReport.pdf>

- Duda, M. D. (2021). *Conservation Words That Work: Determining How to Engage the American Public Through the Language of Conservation*. Gardners, PA: Wildlife Management Institute.
<https://wildlifemanagement.institute/sites/default/files/WMI-Words-Matter-Report-2021.pdf>
- Grooms, B., Dayer, A., Barnes, J., Peele, A., Rutter, J., & Cole, N. (2023). Exploring the relevance of the multidimensionality of wildlife recreationists to conservation behaviors: A case study in Virginia. *Conservation Science and Practice*, 5(6), e12915. <https://doi.org/10.1111/csp2.12915>
- Grooms, B., Rutter, J. D., Barnes, J. C., Peele, A., & Dayer, A. A. (2020). *Supporting Wildlife Recreationists in Virginia: Survey report to inform the Virginia Department of Wildlife Resources' Wildlife Viewing Plan*.
<http://hdl.handle.net/10919/101046>
- Heberlein, T. A., Serup, B., & Ericsson, G. (2008). Female Hunting Participation in North America and Europe. *Human Dimensions of Wildlife*, 13(6), 443–458. <https://doi.org/10.1080/10871200802294265>
- Hvenegaard, G.T. (2002). Birder Specialization Differences in Conservation Involvement, Demographics, and Motivations. *Human Dimensions of Wildlife*, 7(1), 21-36. <https://doi.org/10.1080/108712002753574765>
- Jaureguiberry, P., Titeux, N., Wiemers, M., Bowler, D. E., Coscieme, L., Golden, A. S., Guerra, C. A., Jacob, U., Takahashi, Y., Settele, J., Díaz, S., Molnár, Z., & Purvis, A. (2022). The direct drivers of recent global anthropogenic biodiversity loss. *Science Advances*, 8(45), eabm9982. <https://doi.org/10.1126/sciadv.abm9982>
- Jennings, K. K., Chaves, W. A., & Dayer, A. A. (2024a). *Minnesota Report of Black, Indigenous, and Community Scientists of Color*. Blacksburg, Virginia: Virginia Tech. <https://hdl.handle.net/10919/119014>
- Jennings, K. K., Dayer, A. A., & Chaves, W. A. (2024b). *Insights into Supporters of the MN Department of Natural Resources Nongame Wildlife Program: A Report*. Blacksburg, Virginia: Virginia Tech.
<https://hdl.handle.net/10919/120596>
- Jennings, K. K., Dayer, A. A., Chaves, W. A., & Osmundson, C. L. (2025). How expectations of reciprocity impact donation and volunteer behavior to Nongame Wildlife Programs: a case study in Minnesota. *Human Dimensions of Wildlife*, 0(0), 1–17. <https://doi.org/10.1080/10871209.2024.2447848>
- Kellert, S.R. (2017). *The Nature of Americans: Disconnection and Recommendations for Reconnection*, National Report. Mishawaka, IN: DJ Case & Associates. <https://natureofamericans.org/>
- Larson, L. R., Cooper, C. B., Stedman, R. C., Decker, D. J., & Gagnon, R. J. (2018). Place-Based Pathways to Proenvironmental Behavior: Empirical Evidence for a Conservation–Recreation Model. *Society & Natural Resources*, 31(8), 871–891. <https://doi.org/10.1080/08941920.2018.1447714>
- Larson, L. R., Stedman, R. C., Decker, D. J., Siemer, W. F., & Baumer, M. S. (2014). Exploring the Social Habitat for Hunting: Toward a Comprehensive Framework for Understanding Hunter Recruitment and Retention. *Human Dimensions of Wildlife*, 19(2), 105–122. <https://doi.org/10.1080/10871209.2014.850126>

- Lawhon, B., Newman, P., Taff, D., Vaske, J., Vagias, W., Lawson, S., & Monz, C. (2013). Factors Influencing Behavioral Intentions for Leave No Trace Behavior in National Parks. *Journal of Interpretation Research*, 18(1), 23–38. <https://doi.org/10.1177/109258721301800103>
- McFarlane, B. L., & Boxall, P. C. (1996). Participation in wildlife conservation by birdwatchers. *Human Dimensions of Wildlife*, 1(3), 1–14. <https://doi.org/10.1080/10871209609359066>
- McKinley, D. C., Miller-Rushing, A. J., Ballard, H. L., Bonney, R., Brown, H., Cook-Patton, S. C., Evans, D. M., French, R. A., Parrish, J. K., Phillips, T. B., Ryan, S. F., Shanley, L. A., Shirk, J. L., Stepenuck, K. F., Weltzin, J. F., Wiggins, A., Boyle, O. D., Briggs, R. D., Chapin, S. F., ... Soukup, M. A. (2017). Citizen science can improve conservation science, natural resource management, and environmental protection. *Biological Conservation*, 208, 15–28. <https://doi.org/10.1016/j.biocon.2016.05.015>
- Metropolitan Council. (2024). *Metro population growth slows, construction strong*. Retrieved July 24, 2025, from <https://metrocouncil.org/News-Events/Council-News/Newsletters/Metro-population-growth-2024.aspx>
- McKenzie-Mohr, D. & Tabanico, J. (2024). *Fostering sustainable behavior: An Introduction to community-based social marketing* (4th ed). Wren's Gate Books.
- Minnesota Department of Natural Resources. (2005). [*Guide to artificial roosts for bats*](#). Saint Paul, MN. <https://www.dnr.state.mn.us/mammals/bats.html>
- Minnesota Department of Natural Resources. (2017). [*2017 Minnesota outdoor activities survey: Report on findings*](#). https://files.dnr.state.mn.us/aboutdnr/reports/recreation/2017_minnesota_outdoor_activities_survey_findings_report.pdf
- Minnesota Department of Transportation. (2022a). *Minnesota GO Planning Minnesota's Transportation Future: Urban and Rural Population Trend Analysis*. https://www.minnesotago.org/application/files/6416/5270/8798/MNDOT_Urban_Rural_Trend_FINAL.pdf
- Minnesota Department of Transportation. (2022b). *Minnesota GO Planning Minnesota's Transportation Future: Demographic Trend Analysis*. https://www.minnesotago.org/application/files/5016/2437/5306/Demographics_Trend_Analysis_FINAL.pdf
- Quartuch, M. R., Stedman, R. C., Decker, D. J., Larson, L. R., Siemer, W. F., & Baumer, M. S. (2017). Exploring Nontraditional Pathways into Hunting in New York State: Implications for Recruitment and Retention. *Human Dimensions of Wildlife*, 22(5), 391–405. <https://doi.org/10.1080/10871209.2017.1334247>
- Restoring Forest Raptors: Influence of Human Disturbance and Forest Condition on Northern Goshawks | Request PDF. (2009). *ResearchGate*. https://www.researchgate.net/publication/249432552_Restoring_Forest_Raptors_Influence_of_Human_Disturbance_and_Forest_Condition_on_Northern_Goshawks

- Richardson, M., Passmore, H.-A., Barbett, L., Lumber, R., Thomas, R., & Hunt, A. (2020). The green care code: How nature connectedness and simple activities help explain pro-nature conservation behaviors. *People and Nature*, 2(3), 821–839. <https://doi.org/10.1002/pan3.10117>
- Salazar, G., Mills, M., & Verissimo, D. (2018). Qualitative impact evaluation of a social marketing campaign for conservation. *Conservation Biology*, 33(3), 634–644. <https://doi.org/10.1111/cobi.13218>
- Sinkular, E. N., Jennings, K. K., Karns, M. J., Pototsky, P. C., & Dayer, A. A. (2022). *Minnesota Results of the Wildlife Viewer Survey: Enhancing relevancy and engaging support from a broader constituency*. <http://hdl.handle.net/10919/112191>
- Squires, J. R., Olson, L. E., Roberts, E. K., Ivan, J. S., & Hebblewhite, M. (2019). Winter recreation and Canada lynx: reducing conflict through niche partitioning. *Ecosphere*, 10(10), e02876. <https://doi.org/10.1002/ecs2.2876>
- Tallamy, D. (2009). *Bringing Nature Home: How You Can Sustain Wildlife with Native Plants* (2nd ed). Portland, OR: Timber Press.
- Tallamy, D. W., Narango, D. L., & Mitchell, A. B. (2021). Do non-native plants contribute to insect declines? *Ecological Entomology*, 46(4), 729–742. <https://doi.org/10.1111/een.12973>
- Tallamy, D. W., & Shriver, W. G. (2021). Are declines in insects and insectivorous birds related? *Ornithological Applications*, 123(1), duaa059. <https://doi.org/10.1093/ornithapp/duaa059>
- Teksoz, G., Sahin, E., & Tekkaya-Oztekin, C. (2012). Modeling Environmental Literacy of University Students. *Journal of Science Education and Technology*, 21(1), 157–166. <https://doi.org/10.1007/s10956-011-9294-3>
- U.S. Department of the Interior, U.S. Fish and Wildlife Service. (2024). *2022 Birding in the United States: A Demographic and Economic Analysis, Addendum to the 2022 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation*. <https://www.fws.gov/media/2022-birding-united-states-demographic-and-economic-analysis>
- U.S. Department of the Interior, U.S. Fish and Wildlife Service. (2023) *2022 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation*. <https://www.fws.gov/media/2022-national-survey-fishing-hunting-and-wildlife-associated-recreation>
- Van De Wetering, J., Leijten, P., Spitzer, J., & Thomaes, S. (2022). Does environmental education benefit environmental outcomes in children and adolescents? A meta-analysis. *Journal of Environmental Psychology*, 81, 101782. <https://doi.org/10.1016/j.jenvp.2022.101782>
- Vogels, E. A. V. (2025a, January 13). From Inspiration to Action: How Donors Find Their Causes. *CAF America*. <https://cafamerica.org/blog/from-inspiration-to-action-how-donors-find-their-causes/>
- Vogels, E. A. V. (2025b, May 1). What Motivates U.S. and U.K. Donors to Give. *CAF America*. <https://cafamerica.org/draft/what-motivates-u-s-and-u-k-donors-to-give/>

Weiskopf, S. R., Ledee, O. E., & Thompson, L. M. (2019). Climate change effects on deer and moose in the Midwest. *The Journal of Wildlife Management*, 83(4), 769–781. <https://doi.org/10.1002/jwmg.21649>

Winter, T. J. (2024). Minnesota’s Native Fish Bill: a case study in shifting the “rough fish paradigm.” *Environmental Biology of Fishes*, 107(12), 1327–1338. <https://doi.org/10.1007/s10641-024-01644-w>

Yildirim, M. S., Elkoca, A., Gökçay, G., Yilmaz, D. A., & Yıldız, M. (2025). The relationship between environmental literacy, ecological footprint awareness, and environmental behavior in adults. *BMC Public Health*, 25(1), 551. <https://doi.org/10.1186/s12889-025-21340-3>

Chapter 5. Monitoring and Research

Overview

The importance of monitoring wildlife populations has long been recognized as a critical component of wildlife conservation. This is not only true for species that are actively managed using harvest regulations, such as waterfowl, but also of rare species that may be at risk of extirpation or extinction if we are unaware of their declining status. Today, monitoring elements of biodiversity, from bees and salamanders to orchids and oak savannas, is recognized as one of the most important responsibilities of natural resource agencies.

Building on the two Prior Plans

Monitoring rare and uncommon wildlife species began in the late 1970s with the initiation of the Nongame Wildlife Program. Initial efforts focused on common loons, bald eagles, and colonial waterbird nesting sites. Soon after, the launch of the Minnesota Natural Heritage Program placed an equal emphasis on native plants and plant communities with monitoring efforts focused on rare species such as Prairie Bush Clover (*Lespedeza leptostachya*) and Western Prairie Fringed Orchid (*Plantanthera praeclara*). The state was well-primed to address the importance of monitoring when the state wildlife action planning process began in the early 2000s.

2005-2015 Tomorrow's Habitat for the Wild and Rare

One of the three goals established by the 2005-2015 SWAP was to “Improve knowledge about Species in Greatest Conservation Need (SGCN)”. A key strategy for implementing that goal was to “Monitor long-term changes in SGCN populations and habitats”. An important statewide project funded with State Wildlife Grant dollars was the completion of a statewide survey of freshwater mussels that began in 1999 and the subsequent launch of a long-term monitoring program. Other projects included continued monitoring of American goshawks, double-crested cormorant and American white pelican nesting colonies, and a small but critically important Henslow's sparrow population at Great River Bluffs State Park. Monitoring habitats that rare species depend upon for their survival is equally important. Once covering nearly one-third of Minnesota, native prairie now only covers approximately 235,000 acres. The species dependent on this rare resource are among the most endangered and threatened species in the state. It was only natural, therefore, to design and launch a long-term monitoring program (SPICE; Sustaining Prairies in a Changing Environment) to track the status and trends of this critical habitat.

Minnesota's Wildlife Action Plan 2015-2025

In the second update of the SWAP, monitoring SGCN and their habitats, and applying the knowledge gained to design new management approaches or to adapt existing ones, was a major theme. The plan recognized the need to apply monitoring strategies at three levels: the Wildlife Action Network, in Conservation Focus Areas, and Species Monitoring. Numerous survey, monitoring, and research projects were completed under this Plan.

For the third iteration of the Plan, this chapter focuses on the role of monitoring and research activities within Minnesota's SWAP. We provide an overview of our vision for what we want to achieve in the context of surveys, monitoring, and research for animals, plants, and habitats in Minnesota. We discuss the relevant questions we need answered and structure them into three pathways: baseline diagnostic information needs, determining positive actions and management activities that promote effective conservation, and estimating impacts from threats on species and habitats. We then follow up with a section on best practices for survey,

monitoring, and research design, which is subdivided into handling existing data versus collecting new data. Before diving into this discussion, we define relevant terms pertinent to the presentation of the material.

Important Definitions

Surveys

Wildlife and plant surveys are the primary tool used by the Minnesota Department of Natural Resources (DNR) and its partners to gather scientific data at specific locations and points in time. Surveys can be used to collect vital information on a variety of ecological metrics, including (1) population abundance or relative abundance, (2) population demography (e.g., survival, fecundity, dispersal), (3) habitat conditions and species-habitat relationships, (4) threats or stressors affecting species or habitats, and (5) the geographic distribution of species across Minnesota. Even single surveys can be valuable especially when it leads to the discovery of rare species in new locations and informs conservation efforts.

Monitoring

Monitoring is generally defined as a structured activity in a set of sites, repeated through time. These measurements can include surveys of some species (a type of structured counting effort), evaluation of habitat features (e.g., density of trees), or assessments of ecosystem health (e.g., water quality). Monitoring is distinguished from surveys by employing data collection methods and efforts that are repeated and standardized at some set of sites to detect changes (such as trends in abundance, species richness, or occupancy) through time or across space. The temporal spacing of repeated sampling (e.g., daily, seasonally, or annually) and the spatial arrangement of sampling units such as the distance between and distribution of sampling locations are determined by the focal species or system and the specific questions being addressed. Key features of appropriately designed monitoring efforts are sampling designs built around a focal question(s) or objective(s), standardized efforts, repeatable procedures, well-curated data, and recording of negative data (the absence of the target). The presence and absence of target species recorded through time permit the estimation of detection probability, either explicitly or implicitly, depending on the timing of recurrent surveys and assumptions about survival and movement in the study area. Accounting for detectability helps remove bias in metrics of interest (e.g., abundance, survival, etc.) and can improve trend estimates (Kéry and Schmidt 2008). **Surveillance monitoring** involves tracking trends in species, communities, or habitat metrics over time to understand their status and dynamics. In contrast, **effectiveness monitoring** is structured around specific questions regarding the success of management or conservation actions intended to produce beneficial outcomes, such as habitat creation for threatened or endangered species, invasive species control, or the restoration and reconstruction of native habitats or ecosystems.

Research

Research is the process of obtaining information to answer a specific question(s), involving the collection and analysis of data to test predictions or hypotheses. While the term is frequently associated with formal experimentation, well-designed surveys and monitoring efforts can also be structured to address research questions. Research approaches vary in purpose, design, and methodology, yet they are not mutually exclusive and can overlap, combining elements from multiple types. The strength and utility of research increases when questions and objectives are clearly articulated and quantitatively linked to biologically meaningful outcomes. The DNR primarily engages in the following research approaches:

- *Basic or Observational Research*, which serves to expand general knowledge and understanding of natural systems and the plants and animals that are found within them. This type of research is meant to gather information by documenting characteristics, patterns, trends, or species within ecological systems and may be used to inform conservation actions. An example would be a survey to detect the presence of the plains pocket mouse (*Perognathus flavescens*) on the property of the Twin Cities Army Ammunition Plant in Ramsey County, Minnesota.
- *Applied Research* is defined by its purpose for solving practical problems or informing decision-making in natural resources. While not all applied research is experimental, many applied studies employ formal experimental designs. For example, testing whether prescribed burning or goat-grazing increases native plant and pollinator species richness in grassland habitats and measuring the magnitude of those changes compared to untreated controls constitutes applied research.
- *Experimental Research* involves intentional manipulation of variables through controlled interventions, often including randomization, replication, and pre- and post-treatment measurements. While experimental research can also fit under the umbrella of being applied, some is purely theoretical. An example would be the use of a Before-After-Control-Impact (BACI) design to assess the impact of cattle grazing on composition of the native plant community or on reproductive success in prairie obligate birds.
- *Mechanistic Research* seeks to answer why patterns in nature occur, focusing on identifying underlying biological or ecological processes driving patterns that we see in nature. An example might be a study to determine the relationship between coyote abundance and snowshoe hare survival to understand how predation pressure influences their population and recruitment.

Surveys, monitoring, and research are interrelated tools essential for understanding and managing ecological systems. Each plays a distinct yet complementary role in collecting and interpreting ecological data. Surveys are the primary method for gathering measurement data at specific locations and times; often used to detect the presence or absence of individual species or species groups and help define species distributions and range limits.

Monitoring builds on surveys by structuring data collection in a consistent, standardized, and repeatable way across time and space. This structured approach enables the detection of trends in abundance, occupancy, species richness, and habitat conditions. A critical component of monitoring and survey efforts is accounting for species detectability, which improves the accuracy of ecological assessments and trend estimates.

Research addresses specific scientific questions or objectives, often through hypothesis testing or predictive modeling, and may use data from surveys and monitoring when aligned with its objectives. Research can take various forms including observational, applied, experimental, or mechanistic with each distinguished by the level of intervention and the nature of the questions asked. Together, these tools provide baseline data, reveal ecological trends, and deepen system understanding to inform conservation and resource management.

Minnesota supports a wide range of terrestrial and aquatic ecosystems, which in turn provide habitat for diverse native plants and animals. These ecosystems deliver critical services, including water filtration, flood protection, carbon storage, timber production, and recreational opportunities. However, they are increasingly threatened by direct and indirect stressors such as land and water use pressures, pollution, invasive species, pathogens, and changes in climate and land management.

In response to these challenges, surveys, monitoring, and research form the scientific foundation for biodiversity conservation and natural resource stewardship in Minnesota. Rigorous scientific approaches are critical for guiding conservation strategies, informing adaptive management, supporting technical assessments, and conducting environmental reviews. Later sections of this chapter provide guidance on utilizing existing ecological data and outline recommendations for future data collection efforts.

Baseline Information Needs

The SWAP provides a framework to guide conservation efforts, emphasizing the need for data-driven decisions to protect declining and vulnerable species. The DNR has set clear goals to sustain biodiversity, identify conservation challenges, and support species viability and recovery now and for future generations of Minnesotans. By systematically tracking the distribution and abundance of wildlife, particularly species with low and declining populations, the DNR can develop targeted strategies to mitigate threats to species, populations, and habitats.

In order to have a starting point for assessing species and habitat status in Minnesota, we need baseline diagnostic information about presence/absence, spatial distribution, relative abundance, and population trends. The process of assigning whether a species should be considered a Species in Greatest Conservation Need (SGCN) starts with understanding the rarity, trends, and threats of an animal or plant (see Chapter 2). It is not enough to have survey information at one location and time because species and habitats are dynamic and their status is influenced by a multitude of factors, which can be positive or negative.

Species

For the SWAP to be successful, a strong foundation of baseline information is essential. This includes comprehensive surveys to estimate the distributions of animal and plant species across Minnesota's diverse landscapes, and monitoring efforts designed to detect trends in species occurrence, abundance, or relative abundance over time.

Habitats

Equally critical is the collection of habitat-use data for SGCN, especially given Minnesota's ecological diversity, which spans multiple biomes and supports a wide range of species and community types. Understanding how, when, and where species interact with their habitats is inherently challenging, even for common species with relatively abundant data. For rare species, these challenges are magnified due to limited detections, sparse distribution, and often cryptic behaviors, making it difficult to draw reliable conclusions about habitat associations and requirements. Nevertheless, this understanding is vital for identifying the environmental conditions necessary for the survival, reproduction, and persistence of SGCN. Moreover, a focus on SGCN can yield broader conservation benefits, as their habitat needs often overlap with those of other vulnerable, under-studied, and even common species, thereby supporting more inclusive, ecosystem-based conservation strategies.

Life History and Population Traits

These patterns of habitat use are closely linked to species-specific life history traits, such as reproductive strategies, dispersal capacities, and habitat specialization. Such traits determine how resilient populations are

to natural events (e.g., drought, fire regimes) and anthropogenic pressures (e.g., habitat fragmentation, pollution). This information helps predict population trajectories, assess extinction risks, and develop science-based conservation strategies.

Additionally, data on population demographic status including survival rates, dispersal, migration, and population viability are needed to assess the health and resilience of populations.

Threats and Actions

Coupled with detailed knowledge of threats facing SGCN and many other species, along with management actions available to mitigate those threats, this comprehensive baseline data enables the DNR and its partners to identify vulnerable life stages, critical habitats, and priority areas for conservation, protection, or restoration.

Surveillance Monitoring

Incorporating robust monitoring into Minnesota's SWAP is critical for sustaining wildlife and plant populations, preserving biodiversity, and conserving the habitats that serve as homes for terrestrial and aquatic taxa. Surveillance monitoring establishes baseline conditions and can detect long-term trends, while effectiveness monitoring evaluates the value of conservation actions. Together, they provide the information needed to sustain biodiversity, manage habitats effectively, and respond to emerging conservation challenges. In Minnesota, these monitoring approaches support the SWAP by ensuring that conservation efforts are grounded in scientific evidence and tailored to the evolving needs of wildlife and their habitats.

Surveillance monitoring focuses on understanding species, population, and habitat distributions and tracking trends in various metrics through time. It provides a long-term view of ecosystem health by detecting changes in species abundance, distribution, and community composition. This type of monitoring can be used to establish actionable thresholds and triggers, signaling potential ecological problems before they become critical or conditions for targeted management prescriptions (Nichols & Williams, 2006; Lindenmayer & Likens, 2010). For example, sustained declines in a SGCN could indicate habitat degradation, climate change effects, or emerging threats such as disease. Surveillance monitoring also provides valuable data on habitat trends, helping natural resource managers anticipate shifts in wildlife distributions and potential barriers to conservation efforts. However, while it is useful for identifying patterns and broad-scale concerns, it does not pinpoint causal mechanisms behind observed changes in trends. In the context of trend estimation, additional limitations may include the absence of long-term, standardized data collection methods, making it crucial to establish strong roots with defensible sampling design principles incorporating sampling unit independence, replication, and stratification among Minnesota's diverse biomes and habitats.

Species-level Monitoring

Surveillance monitoring may occur at the species-level across a variety of terrestrial and aquatic animal and plant taxa, ranging from common species to those that are declining, rare, or in need of further research (i.e., SNI - Species Needing Information). Of particular importance to the SWAP are SGCN, which are native animals and plants whose populations are rare, declining, or at risk of decline. Also of primary focus are stewardship species, which have a stable population in MN but are declining within the region. The DNR also recognizes the ecological importance of keystone species and ecosystem engineers—organisms whose roles in structuring

habitats and ecosystem function make them critical for enhancing biodiversity. It is not possible for DNR to track population status and trends of all species of animals and plants in Minnesota, so over the next 10-year cycle of the SWAP, DNR and partners will be exploring alternative prioritization strategies for survey and monitoring efforts. Solutions may incorporate:

- Use of structured decision-making tools to guide the prioritization process (Box 5.1)
- Stratification of species and habitats according to their conservation status (for example, as recently assessed in the s-rank process see chapter 2).
- Incorporating data emerging from parallel efforts, such as aquatic community data from the Minnesota Pollution Control Agency or participatory science programs such as the nationally known eBird (Box 5.2).
- Use of focal monitoring sites to sample a subset of habitat locations for a variety of animals through time, such as the Ecological Monitoring Network (Box 5.3).
- Development and testing of species distribution models (Yackulic et al. 2012, Hefley and Hooten 2016) to efficiently target potential habitat for threatened, endangered, and species defined as SGCN.
- Adoption and transition from finer-scale relative abundance surveys to broader-scale occupancy designs (Mackenzie et al. 2006) that have the potential to make more efficient use of limited agency resources.
- Landscape stratification with revolving spatial and temporal panels organized through time to rotate sampling and resampling efforts across all regions of Minnesota.
- Direct participatory science volunteers to monitor readily identifiable species to maximize information gain and geographic coverage given limited agency resources. It can be applied efficiently to gather broad distributional information or for targeted rare-species surveys.

Box 5.1. What is Structured Decision Making?

Structured Decision Making arises from the formal field of decision analysis, and its intent is to decompose natural resource problems into a structured sequence, while accounting for underlying values of all stakeholders (Runge and Bean 2020). A common form utilizes the ProACT cycle to lay out the decision framework, which structures the issue into problem framing, objectives, alternative actions, consequences, and evaluation of trade-offs among alternatives that will lead to choosing a preferred alternative action (Figure 5.1). Adaptive management is derived as a sub-discipline, where a cycle of iterative decision-making proceeds about a management objective through time.




Figure 5.1. Diagram of the standard iterative steps involved with structured decision making, of which adaptive management is a sub-discipline. The figure presents a generalized form of the ProACT cycle; Problem, Objectives, Alternatives, Consequences and Tradeoffs.

Box 5.2. Summary of Participatory Science Programs

Participatory science, community science, or citizen science, refers to programs that engage members of the public in data collection and research activities—often in collaboration with or under the guidance of professional scientists. In the field of ecology, participatory science programs have become increasingly valuable for monitoring biodiversity, tracking species distributions, assessing ecosystem health, and providing outreach opportunities for various user groups. These programs harness the power of widespread public engagement to gather large quantities of ecological data across broad spatial and temporal scales that would be otherwise difficult or prohibitively expensive for agencies to collect alone. Minnesota DNR sponsored examples include MN PlantWatch, the Minnesota Loon Monitoring Program, the Frog and Toad Calling Survey, and the Chimney Swift Survey. In addition to generating scientifically useful data, participatory science fosters environmental stewardship, increases scientific literacy, and strengthens community involvement in conservation efforts. When well-designed, these programs include standardized protocols, training for participants, and rigorous data validation methods, ensuring that the resulting data can inform ecological research and natural resource management decisions.

Box 5.3. Highlights of the DNR-led Ecological Monitoring Network.

The DNR established the Ecological Monitoring Network (EMN) in 2017 with approximately 550 sites across Minnesota to track ecological change through time. Staff for the program provide data on the status of SGCN habitats, integrating a metric for ecological integrity, and how those habitats are changing by natural and human-derived mechanisms. This effort is being led by the Minnesota Biological Survey (MBS), in collaboration with other DNR divisions and partners. Before EMN, there was no comprehensive statewide monitoring network that consistently measured and evaluated status and trends in native grassland, wetland, and forest habitats. Without such information, it is very difficult for Ecologists and DNR leadership to make informed land management, land use, and conservation decisions.

EMN employs a nested plot design to sample all major habitat types across Minnesota, ensuring a statewide distribution that is independent of habitat quality or land ownership. Data collection focuses on ground layer vegetation and trees, alongside assessments of ecological factors such as deer browse, earthworms, water chemistry, prairie structure, and coarse woody debris. Early findings have provided valuable insights into plant community variation across the state (see [Ecological Monitoring Network | Minnesota DNR](#)). As we begin our first resampling cycle, we aim to assess how SGCN habitats have changed over the past decade. There is potential to expand this framework to include monitoring of animal elements, bridging the gap between flora and fauna conservation.

Regardless of the approach chosen, monitoring SGCN and SNI species provides baseline data to detect population changes through time, identifying organisms that require conservation attention or those that may become SGCN in the future. This process can inform near-term conservation priorities and guide targeted management actions. The DNR and its partners are actively engaged in current and planned efforts focused on species-level surveillance in Minnesota including:

- [Minnesota Frog and Toad Calling Survey](#) (DNR)
- [North American Breeding Bird Survey](#) (U.S. Geological Survey)
- [HerpMapper](#) (a variety of biologists, University personnel, and NGOs)
- [Minnesota National Forest Breeding Bird Monitoring Program](#) (University of Minnesota Duluth Natural Resources Research Institute)
- [Minnesota Statewide Mussel Survey](#) (DNR)
- American Goshawk Nest Monitoring (DNR)
- [Spongy Moth Survey](#) (Minnesota Department of Agriculture)
- Blanding's Turtle Population Monitoring (DNR)
- Timber Rattlesnake Population Assessment (DNR)
- [Early Detection and Distribution Mapping System](#) (EDDMapS; Center for Invasive Species and Ecosystem Health, University of Georgia)
- Dakota Skipper Monitoring (DNR)
- Bat Acoustic Monitoring (DNR, USFS and USFWS)
- [Monarch Larva Monitoring Project](#) (Monarch Joint Venture, University of Wisconsin-Madison Arboretum)
- [Journey North](#) (University of Wisconsin-Madison Arboretum)
- [Integrated Monarch Monitoring Program](#) (Monarch Joint Venture)
- [Minnesota Loon Monitoring Program](#) (DNR)
- [MN PlantWatch](#) (DNR; see Box 5.4)
- Federally Listed Plant Species Monitoring (DNR; see Box 5.5)
- [Minnesota Bumble Bee Atlas](#) (University of Minnesota, Xerces Society, DNR)
- Minnesota Colonial Waterbird Survey (DNR and NRRI)
- [Minnesota Breeding Trumpeter Swan Survey](#) (DNR)
- [Minnesota Waterfowl Breeding Population Survey](#) (DNR and USFWS)
- [Minnesota Amphibian & Reptile Survey](#) (HerpMapper.org, Bell Museum of Natural History, and Minnesota Herpetological Society)
- [Minnesota Cricket Frog Survey](#) (DNR)
- [Terrestrial Invasive Species Participatory Science](#) (TIPS) projects (University of Minnesota Extension)

Box 5.4. Highlights of the DNR-led program, MN PlantWatch

Rare plants are a strong indicator of overall ecological health and areas with a higher diversity of native plants can support more birds, insects, and other wildlife species. Healthy ecological communities are more resilient to threats such as climate change, invasive species, and disease. Established in 2023, MN PlantWatch is a community science-driven initiative that supports and informs the conservation of Minnesota's rare native plant species. Minnesota is home to more than 1700 vascular plant species, of which over 300 are listed as Endangered, Threatened or of Special Concern. For many of these rare plants, data on their distribution, population status, threats and potential management needs are incomplete. Current information is needed to inform the best methods for protecting and managing these unique species. Volunteers are recruited for the program and trained to conduct search and survey efforts for rare plants in their natural habitats, documenting their location, collecting seeds, and providing information about population health and habitat threats. This program is a partnership between the DNR and the [University of Minnesota Landscape Arboretum](#) (UMLA), in collaboration with the [Bell Museum Herbarium](#), [Minnesota Master Naturalist Program](#) and the [Minnesota Native Plant Society](#).

Box 5.5. Highlights of the DNR-led Federally Listed Plant Species Monitoring Program.

The DNR has been monitoring federally listed plant species since the mid-1980s. The level and intensity of monitoring has varied for each species including the western prairie fringed orchid (*Platanthera praeclara*), prairie bush clover (*Lespedeza leptostachya*), Dwarf Trout Lily (*Erythronium propullans*), and Leedy's Roseroot (*Rhodiola integrifolia* subsp. *Leedyi*) and is dependent on their life history and informational needs. Basic monitoring has been done at two levels that has consisted of updating the NHIS database with more accurate locational information using GPS technology and a census count of the individuals found at a given location each year. This information has been critical in guiding conservation actions of the species, be it habitat management or environmental review projects.

Demographic monitoring has been directed toward the western prairie fringed orchid and prairie bush clover. This monitoring is more substantive and involves returning to marked plots on the landscape and tracking individual plants from year to year as well as any new recruits that establish within the transect.

In the case of western prairie fringed orchids, we have pursued an even more intensive phenological monitoring protocol. Under this monitoring, individuals are marked multiple times within the same year to document their emergence, growth, capsule development and seed dehiscence. This combination of nested monitoring has led to the publication of several peer-reviewed papers (Biederman et al. 2018) as well as habitat management guidelines that assist land managers in making decisions that further enhance the conservation of these species.

The western prairie fringed orchid is found throughout the prairie region of Minnesota. Despite this wide geographic distribution, the DNR revisits and completes a flowering census of approximately 80% of the state's populations each season. This effort is only possible because of a collaborative effort between the DNR, The Nature Conservancy, The US Fish and Wildlife Service and the National Park Service. This collaborative partnership also means that information about the species life history and management needs are shared across the entities responsible for the conservation of this rare prairie orchid.

Habitat and Community Monitoring

In addition to species-level surveillance monitoring, the DNR and its partners are actively engaged in ongoing and planned efforts focused on habitats in Minnesota. Habitats consist of the natural environment in which a species lives, grows, and reproduces and provides essential resources like food, shelter, and breeding sites. It

includes all the biotic (living) and abiotic (non-living) factors that influence a species' survival, such as soil type, climate, vegetation, and water availability. These factors come together to support the ecological niche for each species, which range from general to very specific. Collections of interacting species that live in the same habitat at the same time are considered ecological communities (e.g., calcareous fens and vernal pools). Not only is it important to understand status and trends of species and their habitats, but also the dynamics of how species interact as communities within habitats through processes such as predation, competition, parasitism, and mutualism. The form and structuring of these kinds of interactions vary in spatial scale and shape the broader structure of ecosystems. The exact composition and diversity of a community depends on many factors such as landscape geography, resource availability, disturbance, species interactions, and climate. Examples of habitat-level surveillance monitoring efforts in Minnesota led by the DNR and its partners include:

- Prairie Status and Trend Monitoring Project (SPICE: Sustaining Prairies in a Changing Environment) (DNR; see Box 5.6)
- Ecological Monitoring Network (DNR; see Box 5.3)
- Forest Health Index (DNR)
- Wetland Status and Trend Monitoring Program (DNR)
- Minnesota Wetland Hydrology Network (WHyN; DNR)
- Stream Index of Biological Integrity (IBI) Monitoring Program (DNR)
- Lake IBI Monitoring Program (DNR)
- Minnesota Native Plant Community monitoring by relevé sampling (DNR)
- High Conservation Value Forest Monitoring (DNR)
- Watershed Health Assessment Framework (DNR)
- Prairie Reconstruction Initiative (USFWS, USGS, DNR, and TNC)

Box 5.6. Prairie Status and Trend Monitoring Project

The Sustaining Prairies in a Changing Environment (SPICE) project, led by the Minnesota Department of Natural Resources (DNR), monitors grassland structure and composition, bird species, and—soon—pollinators across 40 high-quality prairies in western Minnesota (Figure 5.6). A primary goal of the project is to evaluate trends in grassland bird populations and assess the conservation value of these remnant prairies relative to more marginal sites monitored through the Breeding Bird Survey (BBS). Since 2008, this well-designed, standardized effort has produced temporally and spatially replicated data. Analyses through 2023 indicate little change in vegetation structure, but reveal notable declines in certain grassland bird species. As a long-term monitoring initiative, SPICE also aims to understand how high-quality prairie remnants in Minnesota respond to habitat fragmentation and climate change. It exemplifies the importance of long-term, multi-objective monitoring efforts in guiding effective conservation.




Figure 5.6. Distribution of 40 monitoring sites of high-quality remnant prairie scattered across western Minnesota that make up the Sustaining Prairies in a Changing Environment (SPICE) project led by the DNR.

Conservation Effectiveness

To evaluate whether conservation efforts are achieving their goals, effectiveness monitoring serves as a critical tool. It is a form of applied or experimental research, focused on evaluating the impact of targeted conservation actions on ecosystem attributes. This type of monitoring is essential for determining the effectiveness of management interventions such as habitat restoration, invasive species control, or policy implementation, in producing desired ecological responses. Grounded in the principles of adaptive management, effectiveness monitoring enables agencies to test specific management hypotheses or learning objectives, which can then be refined based on real-world data (Box 5.7). By systematically measuring the success of interventions, it helps identify which actions are yielding positive results and which need modification to improve outcomes for species and habitats. For example, if a restoration project intended to improve nesting success for a bird species fails to meet its objectives, effectiveness monitoring can guide refinements, such as altering vegetation structure or implementing predator control. Ultimately, this approach allows natural resource managers to respond adaptively to dynamic conditions, including climate change and human-induced habitat changes, ensuring that conservation strategies remain effective and evidence-based.

Resources for conservation are chronically limited, and delays in action can push species and ecosystems toward irreversible decline. While there is debate about return-on-investment (ROI) frameworks which prioritize actions with the greatest benefit per cost (Joseph et al. 2009; Carwardine et al. 2012; Vucetich et al. 2017), there is a very strong argument made for proactive conservation (Walls 2018). This idea emphasizes conservation efforts for species and habitats before critical thresholds are met, thereby reducing extinction (or extirpation) risk and high restoration costs. Going further, Walls (2018) suggests taking actions that leverage and utilize stakeholder collaboration like participatory science initiatives and complementing time- and labor-intensive methods for long-term monitoring with alternative methodological approaches for detecting species population declines and estimating extinction risk. While there is much peer-reviewed literature on successfully using formal decision theory and ROI approaches to prioritize conservation efforts known as “conservation triage”, collectively we must push forward with refining existing methods and applying novel prioritization strategies.

The DNR and its partners are actively engaged in current and planned efforts focused on conservation effectiveness monitoring with projects including:

- Response of four-toed salamanders (*Hemidactylium scutatum*) to timber management (DNR; Box 5.8)
- Evaluating techniques for controlling Smooth Brome in native Prairie (DNR)
- Response of red-shouldered hawks (*Buteo lineatus*) to timber management (DNR)
- Evaluation of management alternatives on bluff prairie habitat (TNC)
- Targeted control of the non-native invasive common reed (*Phragmites australis subsp. australis*) (DNR)
- Patch-burn grazing projects at Chippewa Prairie & Hole in Mountain Wildlife Management Area (DNR)
- Prairie reconstructions in the Prairie Coteau Conservation Opportunity Area and monitoring of birds, butterflies, and bumble bees (DNR)
- Conservation grazing at Hole-in-Mountain Wildlife Management Area (DNR, The Nature Conservancy, Sustainable Farming Association, private cattle producer; Box 5.9)
- Prairie reconstruction supplementation with early season forbs for pollinator communities (DNR)
- Bat artificial roost project (DNR; see [Guide to Artificial Roosts for Bats](#))

Box 5.7. What is Adaptive Management?

Adaptive management is a sub-discipline of structured decision making that offers a science-based, iterative approach to decision-making under uncertainty. It emphasizes the use of multiple working hypotheses, coupled with the collection of new data, to update the likelihood of outcomes predicted by those hypotheses (Williams 2011a). Through repeated cycles of data collection and analysis, the approach seeks to identify the model, represented in each hypothesis, that best describes system dynamics (Figure 5.7).

The process, like structured decision making, involves two fluid and iterative stages: a deliberative phase and an iterative phase. In the deliberative phase, stakeholders collaboratively frame the problem, articulate objectives, identify potential management actions, predict the consequences of those actions, and design appropriate standardized monitoring protocols (Williams and Brown 2014). In the iterative phase, monitoring data are collected and used to evaluate management effectiveness and inform the relative support for competing hypotheses. This process of confronting hypotheses with data enables technical learning about ecological systems.

Importantly, adaptive management also facilitates institutional or “double-loop” learning, wherein stakeholders can revisit and revise objectives, actions, or assumptions based on new insights (Argyris and Schön 1978). Adaptive management can be conducted in either an active or passive form. Active adaptive management emphasizes learning about system dynamics as a primary goal, while passive adaptive management focuses on achieving specific management objectives, with learning considered a secondary benefit (Williams 2011b). Regardless of the form, effective monitoring and iterative data collection compared against working hypotheses of system dynamics are central to the adaptive management cycle, allowing project refinement and evolving understanding.

A prime example of adaptive management in action is the Grassland Monitoring Team (GMT; [Grassland Monitoring Team](#)), a collaboration among the DNR, the U.S. Fish & Wildlife Service, and The Nature Conservancy. Formed in 2007, the GMT aims to improve management of native prairies in Minnesota using standardized monitoring protocols embedded within an adaptive management framework and represents a model for collaborative, evidence-based conservation across the state.




Figure 5.7. Diagram of the standard iterative steps involved with adaptive management, a sub-discipline of structured decision making. The figure presents the generalized form along with opportunities for technical and institutional learning, adapted from (Williams and Brown 2014).

Box 5.8. Response of four-toed salamanders to forest management

The four-toed salamander was first found in Minnesota in 1994, has been documented in seven counties, and is listed a species of special concern (Figure 5.8). It breeds in ephemeral wetlands devoid of fish with a sphagnum component among mature upland deciduous or mixed deciduous-conifer forests. Little is known beyond their breeding ecology, with much to be learned about their migratory patterns and upland forest usage.

Since they are found in small, isolated breeding colonies in Minnesota, they are particularly vulnerable to habitat alterations. Of concern is the impact of forest harvest on ephemeral wetlands they breed in and the upland forest surrounding these sites that they migrate to. In response, DNR is leading a study to better understand if there are negative impacts on four-toed salamander breeding colonies due to forest management practices.

This project falls under the category of quasi-experimental research because there are a number of confirmed locations with this salamander that are both unmanaged and planned for future forest management. This permits the implementation of a BACI design where control sites and managed sites can be monitored through time to estimate impacts, if present, of forest management on salamander occupancy and relative abundance.

Box 5.9. Conservation grazing at Hole-in-Mountain Wildlife Management Area

In 2018, the DNR in partnership with The Nature Conservancy, U.S. Fish and Wildlife Service, The Sustainable Farming Association and a local producer launched a long-term monitoring project on a 566-acre parcel of remnant prairie at the Hole-In-Mountain Wildlife Management Area (WMA) to study how vegetation structure and species composition respond to a patch-burn cattle grazing approach. The project was expanded in 2021 to include evaluating a response to adaptive management grazing. The objective was to evaluate whether innovative grazing techniques can increase structural heterogeneity and species diversity in a prairie invaded by smooth brome, an invasive grass. A central aim is to improve wildlife habitat through cooperation with a local cattle operation. Lessons from this project could help enhance habitat on private working lands, which is critical for slowing or reversing wildlife population declines in prairie region ecosystems. Vegetation data show that grazing is increasing overall plant species richness, including both native and non-native species.

Because prairie ecosystems include more than plants, this project has evolved into an interdisciplinary monitoring approach. The DNR Nongame Wildlife Program and Minnesota Biological Survey are also monitoring responses of grassland-obligate birds, bumble bees, and butterflies including several Species in Greatest Conservation Need (SGCN). Most SGCN birds show no negative effects from grazing, except Bobolinks, which appear sensitive to the shorter grass structure. Some species, like the western meadowlark, are declining at both the treatment and nearby control sites—mirroring broader regional trends and suggesting no evidence of a positive grazing effect. DNR initiated monitoring for bumble bees and butterflies on the WMA, along with an adjacent TNC preserve in 2023, to assess effects of patch-burn grazing and will resume efforts in 2025.

We anticipate future data will offer clearer insights into how SGCN and prairie specialist species respond to these grazing techniques. Past low management grazing and subsequent lack of disturbance at the site has led to smooth brome invasion and altered soil conditions, which will take time to address. This project underscores the importance of managing for resilience amid increasing variability in precipitation. Diversity and connectivity across space and time is the foundation of that resilience, and these grazing techniques may offer a viable and economically sound approach to restore prairie ecosystems.

Estimating Impacts from Threats

Gaining a better understanding of conservation threats and their impacts on species, habitats, and communities across Minnesota is an important topic for the next 10-year cycle of the SWAP. When updating our SGCN list by following the NatureServe species conservation status assessment (S-rank process; see Chapter 2), we found that the threats assessment component was the most challenging (Salafsky et al. 2008, Master et al. 2012, Keith et al. 2013, IUCN 2023). The conservation assessment process scores threats according to scope and severity, both measured on an ordinal scale. Scope measures how widespread a threat is in space, while severity describes how damaging a threat is to the affected portion of the population. Geographic Information System (GIS) layers representing area of species occupancy (via distribution maps, known occurrences, and modelled habitat) intersected with spatial information about where in space a given threat occurs can be used to represent scope. However, severity relates to how much a SGCN population is expected to decrease over some time frame (e.g., 10 years) as a result of exposure to a particular level of a specific threat. Data to inform these estimates are hard to find for many types of threats, and in these cases, expert elicitation and ecological analogs (impact data from similar species in similar systems) are integrated into the assessment. We emphasize the need to better understand the interactions between threats and species impacts in the next 10-year SWAP cycle, not only to improve conservation status assessments, but also to help identify actions that will have the greatest impact. The area of need described here falls under the umbrella of experimental or mechanistic research, which requires well-designed studies around clear and focused questions.

Best Practices for Survey, Monitoring, and Research Design

Principles for Effective Use of Existing Data

Resource management agencies, such as the DNR, manage two primary categories of ecological data: (1) data that has already been collected, and (2) data that will be collected in the future. While future data collection can be strategically planned and aligned with current decision-making needs, existing data requires a different perspective. Existing data collected from past survey, monitoring, and research efforts offers a potential goldmine of opportunity for conservation because of the many uses of this information for targeted habitat protection and protected area designation (Guisan et al. 2013, Bennett et al. 2018), improved cross-scale data integration for invasive species management (Xie et al. 2022), integrated data models for improved abundance, survival, and occupancy estimation (Sun et al. 2019, Doser et al. 2022), redefining key metrics like extent of occurrence, area of occupancy, and population trends (Bosso et. al. 2024, Gallagher et al. 2025), and adaptive management (Moore et al. 2011). These are but a small sample of potential uses and benefits of existing datasets.

Collecting new data, particularly in the context of ecological monitoring, demands significant time and resources which can delay needed conservation interventions. For example, the Christmas Island Pipistrelle (*Pipistrellus murrayi*), an island bat in the Indian Ocean was monitored for decades until it went extinct after the Australian government failed to take conservation actions to reverse documented negative trends (Martin et al. 2012). Long-term monitoring programs may take years to uncover meaningful trends and may absorb large portions of project budgets, leaving little left for immediate action (Caughlan & Oakley, 2001). While waiting for more complete datasets, critical decline signals may go unnoticed or become too late to act upon, undermining potential conservation outcomes.

To avoid these delays, it is essential to make efficient and effective use of existing ecological data whenever possible. Open-source data, previously collected datasets, and living data repositories like the [Minnesota Natural Heritage Information System](#) (NHIS) can provide rapid insights and help identify priority conservation needs. For example, the NHIS provides comprehensive information on Minnesota’s rare plants, animals, native plant communities, and other significant natural features. Continuously updated and standardized as new data become available, it serves as the most complete and authoritative source on the state’s rare and noteworthy biodiversity. Its primary purpose is to promote a deeper understanding of these features and to support their effective conservation.

Decision science tools like Value of Information (VOI) analysis can help assess whether new data collection is necessary or if existing data suffice to guide action (Bennett et al., 2018). When gaps are identified, targeted monitoring focused on key areas or species can supplement existing datasets, making overall efforts more strategic and cost-effective. Although integrating existing data from multiple sources comes with challenges such as varying protocols, sampling biases, or data scale differences, careful analytical methods can address these issues (Pacifi et al., 2017; Zipkin et al., 2021). Combining datasets and integrating previously collected data with real-time monitoring can reduce bias associated with limited single-source data and strengthen inference when used judiciously. Where appropriate, existing data can also be used to test hypotheses and inform adaptive decision-making, ensuring that conservation actions proceed on a solid, evidence-based foundation (Buxton et al., 2021; Binley et al., 2024).

To maximize the value of existing ecological data, several principles and best practices should guide data curation, analysis, and application. First, it cannot be overstated how important ecological data husbandry—including sound data management, relational database design, and modern data engineering—is to the integrity and utility of ecological information. As conservation professionals collect increasingly complex and voluminous datasets from field surveys, monitoring efforts, geospatial data, and automated monitoring technologies, the need for structured, reliable, and scalable data repositories becomes essential.

Relational databases such as PostgreSQL, Oracle, and ESRI’s Spatial Database Engine (SDE) are central to this effort. PostgreSQL is an open-source, highly extensible relational database system widely used in ecological and geospatial contexts. It allows for powerful spatial data operations, making it well-suited for managing species observations, habitat maps, and environmental variables. Oracle, by contrast, is a robust, enterprise-grade relational database system commonly used in larger institutions that require advanced performance, security, and scalability features, though it comes with significant licensing costs. SDE is a spatial database technology integrated with ESRI’s ArcGIS software and is designed specifically for managing spatial datasets within a GIS environment. While not a standalone database itself, SDE acts as a middleware layer that enables spatial data to be stored and queried efficiently within relational databases like SQL Server or Oracle.

For smaller-scale ecological projects, spreadsheets (e.g., Microsoft Excel or Google Sheets) and text files (e.g., CSV or TXT) offer practical and accessible tools for storing and managing data. These formats are widely used due to their simplicity, low barrier to entry, and compatibility with most data analysis and visualization software. For Biologists and researchers with limited resources or technical support, spreadsheets provide an intuitive interface for entering, organizing, and performing basic analyses on ecological data.

Text-based formats like CSV files are especially valuable for their flexibility and interoperability. They can be easily imported into statistical software (e.g., R or Python), databases (e.g., PostgreSQL), and GIS tools (e.g.,

QGIS or ArcGIS), making them an ideal intermediate format for data exchange. These formats are also readable and facilitate quick inspection or troubleshooting.

However, while spreadsheets and text files are suitable for simpler datasets and single studies, they come with limitations. Spreadsheets lack the relational structure and data integrity controls of formal databases, making them more prone to data entry errors, version confusion, and difficulty managing large or hierarchical datasets. Best practices when using spreadsheets or text files include maintaining consistent column names, avoiding merged cells or embedded formulas, documenting metadata clearly, and backing up files regularly. For projects that may eventually grow in scale or require collaboration across teams, planning for future migration into a more robust data management system is advisable.

Documenting and archiving metadata in particular are essential. This involves capturing key contextual information about why data were collected, what was measured, how it was collected, where and when was it collected, and who conducted the work. When documentation is lacking, institutional knowledge becomes vital. Interviews with long-serving staff members can help fill in the gaps recovering lost metadata and providing critical context. Next, standardization and integration ensure that data from different sources can be aligned to a common format, taxonomy, measurement unit, and coordinate system. Combining related datasets—for example, surveys for Dakota Skippers (*Hesperia dacotae*) from different years and parts of Minnesota—can greatly enhance analytical utility.

Identifying and addressing biases is another core element of good data stewardship. Datasets should be evaluated for representativeness, and known spatial, temporal, or methodological biases should be documented. Transparency about these limitations helps avoid misinterpretation and supports proper use and integration of the data. Additionally, reproducible curation workflows should be implemented using tools such as R, Python, or Jupyter Notebooks. These platforms facilitate transparent data cleaning and transformation processes and support version control to track all changes made to a dataset. Lastly, effective data governance ensures proper management and accessibility. This includes defining data ownership, establishing sharing and access policies, and enhancing discoverability through searchable metadata catalogs.

Curated existing data, when responsibly integrated into decision-making, can fill gaps in current monitoring programs, offer historical baselines for understanding ecological change, support predictive modeling efforts such as habitat suitability or species distribution, and validate new data collection methods. It also provides a foundation for stakeholder engagement and greater transparency. However, care must be taken to ensure that analyses based on existing data are transparent about their limitations, grounded in the data's original context, and supplemented with new data where appropriate.

Effectively curating and utilizing existing ecological data is both a challenge and an opportunity. While some datasets may initially appear difficult to use, the information they contain can significantly enhance the DNR's capacity for data-driven, adaptive management. By investing in thoughtful data stewardship and integration, resource agencies can unlock the full value of their ecological data assets, supporting more robust and informed management and policy decisions for the future.

Principles for Effective Study Design and Collection of New Data

The following sections provide an overview of key statistical and experimental design principles to consider when developing a new project aligned within the scope of the SWAP, prior to collecting data. Whether for

surveys, monitoring, or research, designing an ecological study requires careful consideration of the specific goals, objectives, spatial and temporal scales, methodological approaches, and logistical constraints of a given project, ensuring that the data collected are both valid and informative for the question(s) at hand. The subject is explored in depth across numerous textbooks and peer-reviewed publications, reflecting the nuanced challenges of capturing and understanding ecological patterns and processes reliably. While not exhaustive, we provide citations below to excellent resources that expand upon the ideas presented here.

Defining Objectives

The overarching goal of SWAP-related projects is to generate data that will provide useful information about ecological system states, population trends, effectiveness of conservation actions, or negative impacts from stressors. To reach any of these goals requires clearly defining the objectives and question(s) and articulating precisely what metrics are to be measured or tested and why. In the context of the SWAP, the study's objectives must align with broader management or conservation goals. To be effective, objectives should be *specific*, defining the purpose, scope, and responsible parties; *measurable*, by including clear metrics for evaluating ecological responses; *achievable*, considering time, personnel, and logistical constraints; *relevant*, in that they address current conservation needs or priorities; and *timely*, with explicit benchmarks and timelines to guide implementation and assessment (Doran 1981, but see Bjerke and Renger 2017).

Defining the Target Population

An essential element of a good study design is the definition of a relevant target population and a corresponding sampling frame. The target population refers to the ecological entities or areas to which inferences will be made—such as a species across its known range, or specific ecosystems like Bluff prairies in the Driftless region of Minnesota. The sampling frame, in turn, is the set of all possible units (e.g., geographic locations with a species' range, or individuals within a larger animal population) that could potentially be sampled. It is crucial that the sampling frame is explicitly defined, comprehensive, and accessible, as it directly determines the validity of any statistical inference. A poorly defined frame increases the risk of producing biased or unrepresentative results, thereby reducing their value for informing decision-making. Sound experimental design, rooted in these principles, ensures that data collected in the field are robust, reproducible, and directly informative for science-based resource management.

Probability-based Sampling

Another critical principle is to use a probability-based sampling design for collecting data in a way that supports unbiased statistical inference by selecting sampling units that represent larger populations (Thompson 1992). Simple random sampling is the most basic approach, where every sampling unit in the study area has an equal probability of being selected. It is easy to implement, but can lead to uneven spatial coverage, sometimes resulting in clustered sampling locations with gaps in landscape representation. This is more likely to occur in ecological studies examining patterns across space.

To account for this reality, stratified random sampling improves upon the design by dividing the study area into distinct groups or strata based on characteristics like habitat type, geographic zone, or elevation. Random samples are taken within each stratum helping to ensure that important groups are represented in the sample.

This increases the precision of estimates, particularly when strata are distinct, but it does not guarantee that sampling units will be evenly spaced across the landscape.

Generalized Random Tessellation Stratified (GRTS) sampling goes further by combining the value of random sampling with a spatial balancing mechanism (Stevens and Olson 2004). It is a form of probability-based sampling within the broader framework of sampling designs where each spatial unit has a known non-zero probability of being selected. It can accommodate unequal inclusion probabilities, stratification, and hierarchical designs that reduce clustering of samples by spreading them evenly across the spatial scope of inference. It is particularly useful for broad-scale environmental monitoring, where consistent coverage of diverse landscapes is important. GRTS supports flexible designs including rotating panels for long-term trend detection. The spatial balance and design flexibility make this design particularly useful for ecological applications.

In some situations, randomization may not be feasible due to logistical constraints and study areas covering large geographic expanses; in these cases, systematic sampling—where samples are placed at regular intervals—can be a suitable alternative. However, care must be taken to confirm that there are no hidden spatial or temporal patterns in the sampling frame that could introduce bias into the results.

Sampling Unit Replication

Replication is another cornerstone of effective study design. Including multiple independent sampling units, such as plots or sites, allows researchers to accurately estimate variability and capture the natural heterogeneity of a target population. Replication also strengthens the statistical power of a study, increasing confidence in detecting real patterns or treatment effects.

Randomization

Randomization is another key design principle that strengthens the validity of field studies. By randomly assigning management treatments and controls among experimental units—such as grazed versus ungrazed plots—one can avoid confounding effects and help ensure that observed differences in outcomes are attributable to the treatment itself rather than other uncontrolled variables. This approach enhances the internal validity of a study and supports stronger causal inference. Even in basic research applications that do not warrant control plots, randomization of sampling plots among strata within the sampling frame is essential for unbiased estimation of occupancy, abundance, or relative abundance.

Statistical Power

Prior to collection of any new data, it is important to ensure that studies are designed with sufficient sample sizes and statistical power to detect biologically meaningful effects. Also, without enough replication (and beware of pseudo-replication; Hurlbert 1984), even well-designed studies may fail to identify important patterns or changes due to high variability or low signal-to-noise ratio. Conducting pilot studies or using simulations can help inform the number of samples needed to achieve acceptable power for detecting ecologically relevant effect sizes; prospective power analysis is key (Steidl et. al. 1997). This planning step is particularly important when resources are limited, as it helps prioritize efforts and ensure that results will be informative for decision-making.

Incorporating Controls

Equally important is the inclusion of controls units—areas or plots where no treatment or management action has occurred—which serve as a baseline for comparison. This is especially critical when evaluating the outcomes of conservation interventions or disturbances. Designs such as Before-After-Control-Impact (BACI) are particularly powerful in assessing the effects of management actions, as they compare changes over time at both treatment and control sites, helping to isolate the true impact of the intervention from natural background variability (Green 1979).

Temporal and Spatial Designs

A thoughtful temporal and spatial design are also important, especially in ecological studies where environmental conditions in the field can fluctuate substantially over time and space. Incorporating repeated observations through time—whether seasonal, annual, or over multiple years—allows researchers to capture trends, identify cyclical patterns, and reduce the impact of short-term variability that may obscure true ecological responses. In the analysis phase, it is ideal to account for temporal autocorrelation, where observations collected closer together in time are more similar than those further apart. Ignoring this structure can lead to underestimated uncertainty and overconfidence in conclusions.

Similarly, the spatial design must be carefully planned to ensure that landscape heterogeneity is properly represented. The spatial distribution of sampling locations should reflect the spatial scale at which the focal species, ecological system, or key processes operate, such as home range, habitat extent, or the geographic scope of ecological dynamics. For instance, heterogeneous habitats require spatial sampling that captures variability within and across patches. Stratified random sampling across different habitat types, elevation zones, or land-use categories can improve inference and help ensure that results are generalized beyond the sampled sites. Moreover, spatial autocorrelation—where nearby sites tend to have more similar conditions or species assemblages than distant ones—should be considered both in study design and in statistical modeling. Failing to do so may inflate the apparent sample size and lead to spurious findings. Together, thoughtful temporal and spatial designs are essential for robust ecological inference. They increase the reliability of estimates, improve the ability to detect meaningful changes or relationships, and provide critical context for understanding the dynamics of complex natural systems.

Counting Errors

Counts of animals and plants in nature rarely reflect true population dynamics because of detection and measurement error, which can significantly bias results if unaccounted for. Many ecological field studies face challenges with imperfect detection—failing to observe a species or event when it is present. To mitigate this, personnel should design sampling protocols that incorporate methods to estimate and correct for detection probability (Otis et al. 1978). These may include repeated surveys at the same site to estimate occupancy, distance sampling to model detectability as a function of distance from the observer, time-to-detection methods, double-observer approaches, or mark-recapture techniques (Williams et al. 2002). When it is not feasible to directly estimate detection probability, models should include covariates that influence detection (such as time of day, weather, or observer experience) to control for this variability implicitly. By integrating these strategies, field studies can produce more accurate and reliable inferences about the true ecological processes that are driving observed patterns.

Emerging technologies are also transforming wildlife survey methods by offering powerful tools to augment traditional survey and monitoring approaches, helping biologists and researchers detect animals more efficiently, across broader spatial and temporal scales, and with reduced bias. Tools such as environmental DNA (eDNA), Adapted-Hunt Drift Fence Technique (AHDriFT), and automated acoustic monitoring are increasingly being adopted or studied for their potential to improve species detection and monitoring outcomes, particularly for elusive, rare, or low-density species (Lugg et al., 2018).

Technologies like eDNA allows researchers to detect species by collecting and analyzing DNA that organisms shed into the environment (e.g., water, soil, snow). This method can be highly sensitive and is particularly useful for detecting aquatic or cryptic terrestrial species without the need for direct observation or capture (Goldberg et al., 2015; Rees et al., 2014). Likewise, AHDriFT combines low-profile drift fences with motion-triggered cameras to document small-bodied amphibians, reptiles, and mammals, offering a passive and non-invasive approach to detecting ground-dwelling species (Martin et al. 2017; Amber et al. 2021). Of particular utility are automated acoustic monitoring systems that use programmable audio recorders to capture vocalizations such as bird songs, frog calls, or bat echolocation. These recordings can be analyzed using software or expert review and have proven useful for long-term, broad-scale surveys (Sugai et al., 2019). When used in concert with conventional methods, these technologies enhance detection probability, reduce effort and observer bias, and help fill key ecological knowledge gaps needed for informed conservation decisions.

Standardized Protocols

Establishing clear and consistent data collection protocols is fundamental to ensuring data quality and enhancing the value of field studies. Standardized protocols—especially those aligned with nationally or regionally recognized methods—promote consistency in how data is collected, making it easier to compare results across studies, contribute to broader ecological databases, and support meta-analyses. Calibration of equipment, consistent training for field staff, and detailed documentation of procedures all contribute to the reliability and reproducibility of findings. When protocols are widely adopted, they also support data integration efforts that can increase statistical power and expand the scope of ecological inference.

Estimating and Communicating Uncertainty

Inferential uncertainty can arise from various sources in ecological modeling and parameter estimation, including natural variability in the underlying process(es) at work, data limitations, and structural assumptions in models. Explaining the nature, sources, and implications of uncertainty can be especially difficult when communicating with decision-makers, landowners, or the general public. Point estimates alone can be misleading if not accompanied by measures that convey the range of plausible values, such as confidence intervals (in frequentist approaches) or credible intervals (in Bayesian frameworks). It is also essential to acknowledge and describe the sources of uncertainty in a project or study, which may include sampling error, measurement error, and model-based uncertainty.

Communicating scientific and model-based uncertainty is challenging, particularly when engaging with audiences unfamiliar with the concept. Many stakeholders are accustomed to receiving definitive answers rather than probabilistic information. For example, people frequently encounter weather forecasts that express uncertainty, such as a 40% chance of rain, but may interpret such predictions in binary terms (either rain will occur or will not), highlighting the cognitive challenge of understanding probabilistic information

(Gigerenzer et al., 2005). This tendency can be amplified in policy and natural resource decision-making contexts, where stakeholders may expect clear guidance or action, not ranges or degrees of confidence. Misinterpretations of uncertainty may erode trust in scientific findings or be used to justify inaction (Fischhoff & Davis, 2014). Furthermore, the language used by scientists with terms like “confidence intervals,” “model error,” or “credible intervals” can be opaque or intimidating to non-technical audiences.

Effective communication strategies include using visual aids (e.g., shaded confidence bands or probability distributions), focusing on the implications of uncertainty rather than the technical details, and framing uncertainty as a normal and informative part of the scientific process. Trust is critical; when science communicators are perceived as transparent and credible, stakeholders are more likely to accept and act on uncertain information (McInerney et al., 2014). Transparent communication of uncertainty not only strengthens scientific integrity but also supports sound decision-making by making the limits of inference clear to policymakers, managers, and the public.

Use of Statistical Models

A cornerstone of effective field study design is the use of appropriate statistical models to address questions of interest. Depending on the study question, the type of statistical model will differ. It is suggested that models can fit into one of three applications: exploration, inference, or prediction (Tredennick et al 2021). Arguably the most prevalent use of statistical models is for exploration, where one seeks to describe patterns in the data and generate hypotheses about what drives ecological processes. On the other hand, modeling for inference formalizes sets of alternative hypotheses describing how a system functions or responds, which in turn are evaluated against empirical data. This effort comprises evaluation of model coefficients associated with hypothesized covariates of importance, which can be positive or negative, and have lesser or greater influence relative to other covariates. Modeling for prediction is likely the least utilized function by state natural resource agencies. This type of modeling simply seeks to determine which of some set of models will best predict response values for new observations of essential covariates. The use of models should align with what the work is being conducted for in the first place. Regardless of the purpose for modeling, be mindful to avoid the trap of relying on p-values within a significance testing framework to decide whether and which factors are more important for explaining variation attributable to some factor(s) (Johnson 1999). Rather, it is good practice to consider effect sizes and biological significance of factors used in models (Johnson 1999). The forms of these models, the paradigm of statistical thinking, and model selection uncertainty are beyond the scope of this brief narrative, but suffice it to say that there are many volumes dedicated to this diverse subject (McCullagh & Nelder, 1983; Gelman et al., 1995; Burnham & Anderson, 1998; Hastie et al., 2001).

Adaptive Designs

Finally, an effective ecological study design should be adaptive and iterative, allowing for refinement as new insights emerge. Adaptive monitoring acknowledges that ecological systems are dynamic and that initial assumptions or methods may need adjustment over time. By regularly evaluating data as it is collected, researchers can modify sampling designs, protocols, or analytical approaches to improve efficiency, address emerging questions, or respond to unexpected challenges. Embedding monitoring within a broader decision framework—such as structured decision making or adaptive management—further enhances its value (Boxes 5.1 and 5.7). These frameworks emphasize learning by doing, using monitoring results to inform and update management actions in a continuous feedback loop. This iterative process not only increases the relevance and

responsiveness of ecological studies but also ensures that monitoring contributes directly to evidence-based conservation and resource management. Together, all these principles form the foundation of statistically sound, reproducible, and management-relevant ecological field studies.

References for Monitoring and Research

Amber, E. D., Lipps, G. J., & Peterman, W. E. (2021). Evaluation of the AHDriFT Camera Trap System to Survey for Small Mammals and Herpetofauna. *Journal of Fish and Wildlife Management*, 12(1), 197–207.

<https://doi.org/10.3996/JFWM-20-016>

Argyris, C., & Schön, D. A. (1978). *Organizational learning: A theory of action perspective*. Addison-Wesley Pub. Co.

Bennett, J. R., Maxwell, S. L., Martin, A. E., Chadès, I., Fahrig, L., & Gilbert, B. (2018). When to monitor and when to act: Value of information theory for multiple management units and limited budgets. *Journal of Applied Ecology*, 55(5), 2102–2113. <https://doi.org/10.1111/1365-2664.13132>

Biederman, L. A., Derek, A., Nancy, S., John, P., Judy, B., & Jeanne, P. (2018). Using phenological monitoring in situ and historical records to determine environmental triggers for emergence and anthesis in the rare orchid *Platanthera praeclara* Sheviak & Bowles. *Global Ecology and Conservation*, 16, e00461. <https://doi.org/10.1016/j.gecco.2018.e00461>

Binley, A. D., Vincent, J. G., Rytwinski, T., Soroye, P., & Bennett, J. R. (2024). Making the most of existing data in conservation research. *Perspectives in Ecology and Conservation*, 22(2), 122–128. <https://doi.org/10.1016/j.pecon.2023.11.004>

Bjerke, M. B., & Renger, R. (2017). Being smart about writing SMART objectives. *Evaluation and Program Planning*, 61, 125–127. <https://doi.org/10.1016/j.evalprogplan.2016.12.009>

Bosso, L., Panzuto, R., Balestrieri, R., Smeraldo, S., Chiusano, M. L., Raffini, F., Canestrelli, D., Musco, L., & Gili, C. (2024). Integrating citizen science and spatial ecology to inform management and conservation of the Italian seahorses. *Ecological Informatics*, 79, 102402. <https://doi.org/10.1016/j.ecoinf.2023.102402>

Burnham, K. P., & Anderson, D. R. (1998). *Model selection and inference: a practical information-theoretic approach* (corr. 2. printing). Springer.

Buxton, R. T., Bennett, J. R., Reid, A. J., Shulman, C., Cooke, S. J., Francis, C. M., Nyboer, E. A., Pritchard, G., Binley, A. D., Avery-Gomm, S., Ban, N. C., Beazley, K. F., Bennett, E., Blight, L. K., Bortolotti, L. E., Camfield, A. F., Gadallah, F., Jacob, A. L., Naujokaitis-Lewis, I., ... Smith, P. A. (2021). Key information needs to move from knowledge to action for biodiversity conservation in Canada. *Biological Conservation*, 256, 108983. <https://doi.org/10.1016/j.biocon.2021.108983>

Carwardine, J., O'Connor, T., Legge, S., Mackey, B., Possingham, H. P., & Martin, T. G. (2012). Prioritizing threat management for biodiversity conservation. *Conservation Letters*, 5(3), 196–204. <https://doi.org/10.1111/j.1755-263X.2012.00228.x>

- Caughlan, L., & Oakley, K. L. (2001). Cost considerations for long-term ecological monitoring. *Ecological Indicators*, 1(2), 123–134. [https://doi.org/10.1016/S1470-160X\(01\)00015-2](https://doi.org/10.1016/S1470-160X(01)00015-2)
- Doran, G. T. (1981). There's a S.M.A.R.T. way to write management's goals and objectives. *Management Review*, 71(11), 35–36.
- Doser, J. W., Leuenberger, W., Sillett, T. S., Hallworth, M. T., & Zipkin, E. F. (2022). Integrated community occupancy models: A framework to assess occurrence and biodiversity dynamics using multiple data sources. *Methods in Ecology and Evolution*, 13(4), 919–932. <https://doi.org/10.1111/2041-210X.13811>
- Fischhoff, B., & Davis, A. L. (2014). Communicating scientific uncertainty. *Proceedings of the National Academy of Sciences*, 111(supplement_4), 13664–13671. <https://doi.org/10.1073/pnas.1317504111>
- Gallagher, R., Roger, E., Packer, J., Slatyer, C., Rowley, J., Cornwell, W., Ens, E., Legge, S., Simpfendorfer, C., Stephens, R., & Mesaglio, T. (2025). Incorporating citizen science into IUCN Red List assessments. *Conservation Biology*, 39(2), e14329. <https://doi.org/10.1111/cobi.14329>
- Gelman, A., Carlin, J., Stern, H., & Rubin, D. (1995). *Bayesian data analysis* (1st Edition). Chapman and Hall/CRC.
- Gigerenzer, G., Hertwig, R., Van Den Broek, E., Fasolo, B., & Katsikopoulos, K. V. (2005). A 30% Chance of Rain Tomorrow: How Does the Public Understand Probabilistic Weather Forecasts? *Risk Analysis*, 25(3), 623–629. <https://doi.org/10.1111/j.1539-6924.2005.00608.x>
- Goldberg, C. S., Strickler, K. M., & Pilliod, D. S. (2015). Moving environmental DNA methods from concept to practice for monitoring aquatic macroorganisms. *Biological Conservation*, 183, 1–3. <https://doi.org/10.1016/j.biocon.2014.11.040>
- Green, R. H. (1979). *Sampling design and statistical methods for environmental biologists*. Wiley.
- Guisan, A., Tingley, R., Baumgartner, J. B., Naujokaitis-Lewis, I., Sutcliffe, P. R., Tulloch, A. I. T., Regan, T. J., Brotons, L., McDonald-Madden, E., Mantyka-Pringle, C., Martin, T. G., Rhodes, J. R., Maggini, R., Setterfield, S. A., Elith, J., Schwartz, M. W., Wintle, B. A., Broennimann, O., Austin, M., ... Buckley, Y. M. (2013). Predicting species distributions for conservation decisions. *Ecology Letters*, 16(12), 1424–1435. <https://doi.org/10.1111/ele.12189>
- Hastie, T., Friedman, J. H., & Tibshirani, R. (2001). *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*. Springer.
- Hefley, T. J., & Hooten, M. B. (2016). Hierarchical Species Distribution Models. *Current Landscape Ecology Reports*, 1(2), 87–97. <https://doi.org/10.1007/s40823-016-0008-7>
- Hurlbert, S. H. (1984). Pseudoreplication and the Design of Ecological Field Experiments. *Ecological Monographs*, 54(2), 187–211. <https://doi.org/10.2307/1942661>
- IUCN. (2023). *Threats Classification Scheme (Version 3.2)*. International Union for Conservation of Nature. <https://www.iucnredlist.org/resources/threat-classification-scheme>

- Johnson, D. H. (1999). The Insignificance of Statistical Significance Testing. *The Journal of Wildlife Management*, 63(3), 763. <https://doi.org/10.2307/3802789>
- Joseph, L. N., Maloney, R. F., & Possingham, H. P. (2009). Optimal Allocation of Resources among Threatened Species: a Project Prioritization Protocol. *Conservation Biology*, 23(2), 328–338. <https://doi.org/10.1111/j.1523-1739.2008.01124.x>
- Keith, D. A., Rodríguez, J. P., Rodríguez-Clark, K. M., Nicholson, E., Aapala, K., Alonso, A., Asmussen, M., Bachman, S., Basset, A., Barrow, E. G., Benson, J. S., Bishop, M. J., Bonifacio, R., Brooks, T. M., Burgman, M. A., Comer, P., Comín, F. A., Essl, F., Faber-Langendoen, D., ... Zambrano-Martínez, S. (2013). Scientific Foundations for an IUCN Red List of Ecosystems. *PLoS ONE*, 8(5), e62111. <https://doi.org/10.1371/journal.pone.0062111>
- Kéry, M., & Schmidt, B. (2008). Imperfect detection and its consequences for monitoring for conservation. *Community Ecology*, 9(2), 207–216. <https://doi.org/10.1556/ComEc.9.2008.2.10>
- Lindenmayer, D. B., & Likens, G. E. (2010). The science and application of ecological monitoring. *Biological Conservation*, 143(6), 1317–1328. <https://doi.org/10.1016/j.biocon.2010.02.013>
- Lugg, W. H., Griffiths, J., Van Rooyen, A. R., Weeks, A. R., & Tingley, R. (2018). Optimal survey designs for environmental DNA sampling. *Methods in Ecology and Evolution*, 9(4), 1049–1059. <https://doi.org/10.1111/2041-210X.12951>
- MacKenzie, D. I. (Ed.). (2006). *Occupancy estimation and modeling: inferring patterns and dynamics of species*. Elsevier.
- Martin, S. A., Rautsaw, R. M., Robb, F., Bolt, M. R., Parkinson, C. L., & Seigel, R. A. (2017). Set AHDriFT: Applying game cameras to drift fences for surveying herpetofauna and small mammals. *Wildlife Society Bulletin*, 41(4), 804–809. <https://doi.org/10.1002/wsb.805>
- Martin, T. G., Nally, S., Burbidge, A. A., Arnall, S., Garnett, S. T., Hayward, M. W., Lumsden, L. F., Menkhorst, P., McDonald-Madden, E., & Possingham, H. P. (2012). Acting fast helps avoid extinction. *Conservation Letters*, 5(4), 274–280. <https://doi.org/10.1111/j.1755-263X.2012.00239.x>
- Master, L. L., Faber-Langendoen, D., Bittman, R., Hammerson, G. A., Heidel, B., Ramsay, L., Snow, K., Teucher, A., & Tomaino, A. (2012). *NatureServe Conservation Status Assessments: Factors for Evaluating Species and Ecosystem Risk*. NatureServe, Arlington, VA.
- McCullagh, P., & Nelder, J. A. (1983). *Generalized linear models*. Chapman and Hall.
- McInerney, G. J., Chen, M., Freeman, R., Gavaghan, D., Meyer, M., Rowland, F., Spiegelhalter, D. J., Stefaner, M., Tessarolo, G., & Hortal, J. (2014). Information visualisation for science and policy: engaging users and avoiding bias. *Trends in Ecology & Evolution*, 29(3), 148–157. <https://doi.org/10.1016/j.tree.2014.01.003>
- Moore, C. T., Lonsdorf, E. V., Knutson, M. G., Laskowski, H. P., & Lor, S. K. (2011). Adaptive management in the U.S. National Wildlife Refuge System: Science-management partnerships for conservation delivery. *Journal of Environmental Management*, 92(5), 1395–1402. <https://doi.org/10.1016/j.jenvman.2010.10.065>

- Nichols, J., & Williams, B. (2006). Monitoring for conservation. *Trends in Ecology & Evolution*, 21(12), 668–673. <https://doi.org/10.1016/j.tree.2006.08.007>
- Otis, D. L., Burnham, K. P., White, G. C., & Anderson, D. R. (1978). Statistical Inference from Capture Data on Closed Animal Populations. *Wildlife Monographs*, 62, 3–135. <https://www.jstor.org/stable/3830650>
- Pacifici, K., Reich, B. J., Miller, D. A. W., Gardner, B., Stauffer, G., Singh, S., McKerrow, A., & Collazo, J. A. (2017). Integrating multiple data sources in species distribution modeling: a framework for data fusion. *Ecology*, 98(3), 840–850. <https://doi.org/10.1002/ecy.1710>
- Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R. M., & Gough, K. C. (2014). REVIEW: The detection of aquatic animal species using environmental DNA – a review of eDNA as a survey tool in ecology. *Journal of Applied Ecology*, 51(5), 1450–1459. <https://doi.org/10.1111/1365-2664.12306>
- Runge, M. C., & Bean, E. A. (2020). Decision analysis for managing public natural resources. In M. C. Runge, S. J. Converse, J. E. Lyons, & D. R. Smith (Eds.), *Structured decision making: Case studies in natural resource management* (pp. 1–11). John Hopkins University Press.
- Salafsky, N., Salzer, D., Stattersfield, A. J., Hilton-Taylor, C., Neugarten, R., Butchart, S. H. M., Collen, B., Cox, N., Master, L. L., O'Connor, S., & Wilkie, D. (2008). A Standard Lexicon for Biodiversity Conservation: Unified Classifications of Threats and Actions. *Conservation Biology*, 22(4), 897–911. <https://doi.org/10.1111/j.1523-1739.2008.00937.x>
- Steidl, R. J., Hayes, J. P., & Schaubert, E. (1997). Statistical Power Analysis in Wildlife Research. *The Journal of Wildlife Management*, 61(2), 270. <https://doi.org/10.2307/3802582>
- Stevens, D. L., & Olsen, A. R. (2004). Spatially Balanced Sampling of Natural Resources. *Journal of the American Statistical Association*, 99(465), 262–278. <https://doi.org/10.1198/016214504000000250>
- Sugai, L. S. M., Desjonquères, C., Silva, T. S. F., & Llusia, D. (2020). A roadmap for survey designs in terrestrial acoustic monitoring. *Remote Sensing in Ecology and Conservation*, 6(3), 220–235. <https://doi.org/10.1002/rse2.131>
- Sun, C. C., Royle, J. A., & Fuller, A. K. (2019). Incorporating citizen science data in spatially explicit integrated population models. *Ecology*, 100(9), e02777. <https://doi.org/10.1002/ecy.2777>
- Thompson, S. K. (1992). *Sampling*. Wiley.
- Tredennick, A. T., Hooker, G., Ellner, S. P., & Adler, P. B. (2021). A practical guide to selecting models for exploration, inference, and prediction in ecology. *Ecology*, 102(6), e03336. <https://doi.org/10.1002/ecy.3336>
- Vucetich, J. A., Nelson, M. P., & Bruskotter, J. T. (2017). Conservation Triage Falls Short Because Conservation Is Not Like Emergency Medicine. *Frontiers in Ecology and Evolution*, 5, 45. <https://doi.org/10.3389/fevo.2017.00045>
- Walls, S. C. (2018). Coping With Constraints: Achieving Effective Conservation With Limited Resources. *Frontiers in Ecology and Evolution*, 6, 24. <https://doi.org/10.3389/fevo.2018.00024>

- Williams, B. K. (2011a). Adaptive management of natural resources—framework and issues. *Journal of Environmental Management*, 92(5), 1346–1353. <https://doi.org/10.1016/j.jenvman.2010.10.041>
- Williams, B. K. (2011b). Passive and active adaptive management: Approaches and an example. *Journal of Environmental Management*, 92(5), 1371–1378. <https://doi.org/10.1016/j.jenvman.2010.10.039>
- Williams, B. K., & Brown, E. D. (2014). Adaptive Management: From More Talk to Real Action. *Environmental Management*, 53(2), 465–479. <https://doi.org/10.1007/s00267-013-0205-7>
- Williams, B. K., Nichols, J. D., & Conroy, M. J. (2002). *Analysis and management of animal populations: modeling, estimation, and decision making*. Academic Press.
- Xie, D.-J., Zhang, F.-X., Wang, C.-J., & Wan, J.-Z. (2022). Integrating Different Scales into Species Distribution Models: A Case for Evaluating the Risk of Plant Invasion in Chinese Protected Areas under Climate Change. *Applied Sciences*, 12(21), 11108. <https://doi.org/10.3390/app122111108>
- Yackulic, C. B., Chandler, R., Zipkin, E. F., Royle, J. A., Nichols, J. D., Campbell Grant, E. H., & Veran, S. (2013). Presence-only modelling using MAXENT: when can we trust the inferences? *Methods in Ecology and Evolution*, 4(3), 236–243. <https://doi.org/10.1111/2041-210x.12004>
- Zipkin, E. F., Zylstra, E. R., Wright, A. D., Saunders, S. P., Finley, A. O., Dietze, M. C., Itter, M. S., & Tingley, M. W. (2021). Addressing data integration challenges to link ecological processes across scales. *Frontiers in Ecology and the Environment*, 19(1), 30–38. <https://doi.org/10.1002/fee.2290>

Chapter 6. Implementation

This Chapter contains three sub-chapters, each of which are applied to guide our conservation work under this Plan. The first, the Conservation Action Network, represents biodiversity and high-quality habitat in our state spatially as a resource to prioritize conservation actions geographically. Secondly, we describe Conservation Opportunity Areas, which identify locations for partners to focus on-the-ground conservation efforts that build and improve areas in and around the Conservation Action Network. Third, we provide foundational information regarding climate trends in Minnesota and strategies for climate adaptation that are broadly applicable throughout many types of conservation actions.

Conservation Action Network

Of the more than 7,000 known plant and animal species in Minnesota, 1,155 have been identified as Species in Greatest Conservation Need (SGCN) following a rigorous conservation status assessment described in Chapter 2. Given the sheer number of SGCN, one of the main goals of the Plan is to prioritize and focus on conservation actions for habitats needed by SGCN (see Chapter 3). This habitat approach can meet the needs of multiple species and emphasizes sustaining and enhancing terrestrial and aquatic habitats for SGCN in the context of both the local and larger landscape scale (including watersheds). Habitat loss and fragmentation impact biodiversity, therefore habitat conservation and restoration have become central objectives in conservation planning. Evaluating landscape functional connectivity and mapping habitat networks are key steps (and challenges) for implementing effective actions (Duflot et.al., 2018)

The goals for managing local habitat should include sustaining or enhancing landscape-scale biological diversity, improving the functions of conservation lands and waters, and supporting ecosystem resilience. Resilience, as it applies to Minnesota's Wildlife Action Plan, is the capacity of an ecological system to absorb some level of disturbance and reorganize while still retaining essential functions, structures, and feedbacks. The focus of the habitat approach is to implement conservation actions that will maintain or enhance the conditions that increase a system's resilience. Examples of such actions include:

- Protecting large habitat areas from fragmentation.
- Restoring natural levels of connectivity while maintaining natural barriers.
- Reducing invasive species.
- Prioritizing habitats for biological and functional diversity (vs. single-species needs).
- Minimizing pollution and impervious surfaces.
- Restoring watershed hydrology.
- Reintroducing disturbance when appropriate.

Measuring ecosystem resilience is an emerging concept which includes developing ways to assess functional groups (e.g., decomposers, producers, predators) and understanding the niche or redundancy of functional groups at multiple scales, while also identifying structural diversity, ecosystem services, and human social/ecological connections.

Building on the two Prior Plans

2005-2015 Tomorrow's Habitat for the Wild and Rare

Minnesota's original State Wildlife Action Plan, *Tomorrow's Habitat for the Wild and Rare*, identified significant loss and degradation of habitat as the principal management challenge for achieving its goal to stabilize and increase SGCN populations (DNR 2005). The principal strategy for addressing this management challenge was the need to identify key SGCN habitats and focus management efforts on them. Habitat descriptions were subsequently provided for 16 key habitats in profiles of all 25 ecological subsections in Minnesota ([Ecological Classification System](#)). Each subsection profile included a matrix of SGCN use for all habitats that occurred in the subsection along with an assessment of which habitats were the most important to the greatest number of SGCN. Conservation actions to maintain and enhance the key habitats in each subsection were delineated.

Minnesota's Wildlife Action Plan 2015-2025

Minnesota's Wildlife Action Plan 2015-2025 identified a three-pronged approach to ensuring the long-term health and viability of Minnesota's wildlife, the first and most comprehensive being the habitat approach (DNR 2015). In the 2015 Plan, the Wildlife Action Network (WAN) was developed as a tool for the implementation of a habitat-based conservation approach, resulting in a map illustrating the distribution of high-quality habitats for both terrestrial and aquatic SGCN and identifying potential movement corridors among the prioritized habitat cores. To prioritize areas for on-the-ground conservation efforts, sites within the WAN were assigned priority rankings based on five metrics, including SGCN population viability and species richness. These rankings were used in combination with a set of Conservation Focus Areas where resources, such as partners and/or funding, were available to address conservation needs through on the ground habitat restoration and management.

The 2025-2035 Plan maintains the same concept of the WAN, applying updated and new information to better identify and prioritize areas of high conservation value. With the addition of plants as SGCN and the availability of additional current landcover GIS layers reflecting habitat quality, the resulting WAN revision is more comprehensive and therefore called the Conservation Action Network (CAN).

The Conservation Action Network serves as the foundational prioritization tool to help accomplish the five habitat strategies laid out in Chapter 3, which are to:

1. **Protect**, buffer, and connect high quality habitats to optimize biodiversity, SGCN, and landscape benefits.
2. **Restore**, enhance, and maintain lands and waters to benefit SGCN, biodiversity, and ecosystem resilience
3. **Collaborate** with conservation partners and landowners to enhance conservation delivery
4. **Monitor** SGCN, native plant communities, habitats, and ecosystems for changes through time including responses to stressors, natural disturbances, conservation actions, and climatic conditions.
5. **Connect** to develop, innovate, incentivize, and disseminate evidence-based habitat management practices to benefit SGCN

In practice the CAN is well suited to assist in large scale and multi-faceted conservation planning efforts. As climate change becomes more pronounced, its effects will be both direct and indirect. Impacts from climate change in addition to those from other large-scale drivers will be synergistic; that is, multiple stressors will have an effect that is greater than what may be considered simply additive. Species are already moving in response to climate change, and it is expected that the ranges of plant and animal species will continue to

shift, and habitats will change. The Conservation Action Network will facilitate planning for adaptation to these changes by, identifying core areas large enough to contain a diversity of ecotones and habitats to allow for local shifts (e.g., dry to mesic prairie), and highlighting ecological corridors to allow for species movements and the flow of energy and materials. These conditions will support the biological diversity already present in the Conservation Action Network and make it more likely that ecosystem resilience can be maintained in response to climate change and other stressors, such as invasive species and other forms of habitat degradation. The CAN approach in helping prioritize areas in which to maintain and increase native biological diversity has also been shown to generally decrease the ability of invasive plants to become established (Yan et al. 2015; Levine and D'Antonio 1999; Hooper et al. 2005; Fridley et al. 2007).

While corridors are necessary for connectivity, considerations of potential negative impacts and local knowledge are also important to recognize and prevent against unintended consequences, such as the novel transmission of pathogens and invasive species. However, one review found little strong evidence for negative consequences of landscape corridors (Haddad et al. 2014), and others have found strong evidence of positive effects, such as increased biological diversity (Gilbert-Norton et al. 2010). But knowing how dynamic a system is and what the potential impacts of connectivity within that system could be is important. Corridors between naturally unconnected aquatic or wetland systems might be inappropriate, for instance.

Ultimately the CAN was developed with the aim of increasing the effectiveness and efficiency of conservation actions when resources and capacity are limited. A network focus will increase efficiency by facilitating coordination of management, prioritizing outreach efforts, and targeting technical assistance. Coordination of management will also increase its effectiveness. For example, coordinating activities across multiple ownerships within the network can benefit species that require large areas of habitat in particular successional stages (i.e., time since disturbance).

In essence, the Conservation Action Network is an effort to identify the best places for SGCN and Minnesota's diverse plants, animals and many other organisms to persist and adapt to a changing climate as well as physical changes on the land. The CAN identifies and scores core areas of high biodiversity and ecosystem function as well as areas of potential connectivity between the cores. This prioritization is intended to optimize conservation efforts given limited resources. In addition, the CAN identifies a network of lands with connections to facilitate species movements given a changing climate. The CAN intends to guide conservation partners to maximize collective outcomes and will continue to build on the utility of the 2015 WAN (see Case Study about the Wildlife Action Network in action).

Case Study: Minnesota Land Trust's Use of the 2015 Wildlife Action Network (WAN)

The Minnesota Land Trust (MLT) is a conservation non-profit organization that works with private landowners to conserve land through voluntary land conservation agreements. The Land Trust's mission is to protect and restore Minnesota's most vital natural lands to provide wildlife habitat, clean water, outdoor experiences, and scenic beauty for generations to come. Since 1991, the Land Trust has partnered with hundreds of Minnesota landowners to permanently protect over 80,000 acres of natural land and 540 miles of fragile shoreline throughout the state. In addition, they have restored over 9,000 acres of critical habitat.

The MLT's conservation easement acquisition program works throughout Minnesota in a range of different "program areas" such as the Blufflands and the North Shore that largely corresponds with state grants they receive from the Outdoor Heritage Council. A main component of this work is partnering with private landowners to place permanent conservation easements on their property. MLT's focus is typically areas of high-quality natural habitat. MLT also has a restoration program to provide restoration assistance to landowners once the conservation easement is complete. All of our program areas have slightly different goals and objectives, but use of the Wildlife Action Plan is consistent across all of them.

MLT uses the Wildlife Action Network to help prioritize where our work will have the maximum conservation impact. To prioritize among a large volume of landowner applications across a given landscape, the Wildlife Action Network is a key data point in sorting through these potential parcels and identifying priorities for protection. In southeastern Minnesota, for example, MLT uses a ranking system that weighs the parcel's size, the quality of its natural resources/habitat, and its landscape context to come up with a composite ranking for prioritization. The Wildlife Action Network is an outstanding metric for use in the "natural resources/habitat" quality score because it synthesizes so many other data sets into one, easy-to-use metric.

Developing the Conservation Action Network

The first step to develop the CAN was to select aquatic and terrestrial core areas with high concentrations of SGCN species, high-quality or rare habitats, or healthy ecosystems. Multiple Geographic Information System (GIS) layers, based on aerial and ground-truthed data, were carefully reviewed before acceptance as a core (Table 6.1). This process was advised by a broad team of conservation professionals in 3 separate meetings. These layers were then combined to create a base core map (Figure 6.1).

To address connectivity, we utilized separate prairie and forest connectivity maps recently created by ecologists in The Nature Conservancy's North America Science team, using a wall-to-wall Circuitscape model (Pelletier et al. 2014). The prairie connectivity layer includes other open habitats such as pasture, peatlands, and wetlands. The forest connectivity map includes all areas classified as forest with woodlands and shrublands classified as a higher resistance. These maps indicate land likely to facilitate the movement of plants and animals in response to a changing climate. Climate flow refers to the gradual movement of populations in response to changes in the climate. Over time, climate flow results in range shifts and the formation of novel communities. Diffuse flow represents areas that are extremely intact and consequently facilitate high levels of dispersed flow that spreads out to follow many different and alternative pathways. Concentrated flow represents areas where large quantities of flow are concentrated through a narrow area. For both forest and prairie connectivity, areas of high diffuse or high concentrated flow were included in the CAN (Figure 6.2).

The final step was to score the core and connectivity layers using five layers that represent aquatic habitats (lakes and streams), SGCN diversity (All SGCN richness grids; End, Thr, S1, S2 SGCN Richness Grids), and plant communities/ecosystems (Sites of Biodiversity Significance, Table 6.2 and Figures 6.3-6.7). The SGCN richness grids used Jenks Natural breaks to classify the number of SGCN species into five categories. The Jenks natural breaks classification method is a data clustering method designed to determine the best arrangement of values into different classes. This is done by seeking to minimize each class's average deviation from the class mean, while maximizing each class's deviation from the means of the other classes. In other words, the method seeks to reduce the variance within classes and maximize the variance between classes (Jenks 1967). These 5 scoring layers were combined into a single scoring layer where the individual layer scores were first summed together and then normalized so the overall score ranged between 0 and 1. This combined scoring layer was then used to separately score the terrestrial habitats and wetlands, lakes and streams, and prairie and forest connectivity layers. These scored layers were overlayed to create the final scored Conservation Action Network (Figure 6.8).

Identification of the CAN is based on a rich set of species occurrence data and biodiversity rankings from several decades of intensive survey efforts by Minnesota Biological Survey staff; surveys and public reports collected by Nongame Wildlife Program, Scientific and Natural Areas, and Parks and Trails staff; intensive sampling efforts by the Minnesota Pollution Control Agency for stream Indices of Biological Integrity; and other sources of information. The CAN does not capture potentially important areas lacking survey information. Over the next 10 years, the network should be further refined to include new information on additional rare species occurrences.

Table 1. Geographical Information System (GIS) layers used to map habitat cores in the Conservation Action Network (CAN).

GIS layer	Description
Species in Greatest Conservation Need (SGCN) Richness Hotspots	Number of SGCN species summed in 2 km square hexagons. Hotspots were selected using the following rules: 1. single hexagon of ten or more species 2. cluster of at least 4 contiguous hexagons each comprising 4 or more species 3. cluster of at least 6 contiguous hexagons each comprising 2 or more species; cluster must contain hotspot already defined in 1 or 2 above Once hexagon clusters were identified, we selected all “natural” habitat within the clusters from a composite landcover map.
A to C ranked Element Occurrences of Endangered and Threatened Plants	From Biotics Rare Natural Features maintained in The Natural Heritage Information System, Minnesota DNR - Division of Ecological Resources (NatureServe , DNR Natural Heritage Information System).
2015 Mapped Endangered and Threatened Animal Populations and 2025 Federal Critical Habitat designations	Populations of Endangered and Threatened Animals mapped in 2015 for the previous Wildlife Action Plan. Added to this layer were current Federal Critical Habitat designations for Dakota Skipper, Piping Plover, Poweshiek Skipperling, and Topeka Shiner.
Forest cover within Bat Habitat Conservation Plan (HCP) restriction buffers	Selected all forest classes within buffers around known hibernacula entrances and those around known occupied maternity roost trees that are used by northern long-eared bat (<i>Myotis septentrionalis</i>), little brown bat (<i>Myotis lucifugus</i>), and tricolored bat (<i>Perimyotis subflavus</i>) (https://www.dnr.state.mn.us/bathcp/index.html).
Outstanding and high Sites of Biodiversity Significance	A biodiversity significance rank is assigned by staff within the Minnesota Biological Survey (Minnesota DNR) on the basis of the number of rare species, the quality of the native plant communities, size of the site, and context within the landscape (MBS Site Biodiversity Significance Ranks).
S1 and S2 Native Plant Communities	Native Plant communities mapped by staff within the Minnesota Biological Survey (Minnesota DNR) that have a state conservation rank of S1 or S2 (Minnesota’s Native Plant Communities).
Minnesota Prairie Conservation Plan Core Areas	Represents areas with concentrations of native prairie and grasslands as identified using DNR Biological Survey, USFWS, and land cover data. Where appropriate, these boundaries were adjusted to reflect watershed divides and ownership/field boundaries (Minnesota Prairie Conservation Plan).
Wetland Native Plant Communities ranked A to C	Wetland Native Plant communities mapped by staff within the Minnesota Biological Survey (Minnesota DNR) that have a condition rank of A, B, or C (Minnesota’s Native Plant Communities).

GIS layer	Description
Forest Management Opportunity Areas	State forest land with management emphasis to improve forest habitat for wildlife species. The following types were included: Landscape (LAND), Old Forest Management Complexes (OFMC), White Pine/Conifer Management Area (WPMA), Owl Management Area (OWMA), Large Block/Moose, Management Area (MMA), Open Landscape Management Area (OLMA), Forest Patch (PATCH), Deer Management Area (DMA), INT, Pilot INT MOA, and Pilot UPLD MOA (Forest Resource Management Planning)
DNR Forest Inventory Old Growth and old growth Special Management Zones (SMZ)	Currently designated or candidate old growth forest stands on lands administered by the MN Department of Natural Resources (DNR). Also included is the old growth Special Management Zone (SMZ), a 330-foot buffer around the Old Growth layer (Old growth forests).
Forest Representative Sample Areas (RSAs)	Ecologically viable representative samples of forest types on lands administered by the MN Department of Natural Resources (DNR) designated to serve one or more of three purposes: 1) To establish and/or maintain an ecological reference condition; or 2) To create or maintain an under-represented ecological condition; or 3) To serve as a set of protected areas or refugia for species, communities and community types not captured in other Criteria of this Standard.
High Conservation Value Forests (HCVF)	Forest areas administered by the MN Department of Natural Resources (DNR) selected by Interdisciplinary Regional Teams from a subset of sites of Biodiversity Significance identified by the Minnesota Biological Survey (MBS) that was recommended for consideration by Division of Ecological and Water Resources biologists (High conservation value forests).
Lakes with high Watershed Health Assessment Framework (WHAF) Lakes Health Cores scores and Lakes with high bird rankings identified in the Lakes of Biological Significance layer.	Lakes with an overall WHAF health score of 75 or greater and lakes with a lower overall health score but had a biology component score of 70 or greater. Also added lakes ranked high or outstanding from the Lakes of Biological Significance layer with a bird rank of 1 (Outstanding) or 2 (High). (Watershed Health Assessment Framework: Lakes , Lakes of Biological Significance)
Stream reaches meeting Exceptional Use criteria for Fish or Invertebrates	This layer includes stream reaches (WIDS) that met Exceptional Use criteria for Fish and/or Invertebrates based on Indices of biological integrity surveyed and developed by the Minnesota Pollution Control Agency (MPCA). To depict continuity of streams on the map, adjacent reaches that met General Use criteria were also added as well as all streams with an order size of 5 to 9. While this approach over-represents quality stream areas, the scoring step identifies those areas (MPCA Index of biological integrity).

Table 2 GIS layers used to score habitat cores and connectivity in the Conservation Action Network (CAN)

Layer	Scoring Parameters
Sites of Biodiversity Significance	Outstanding = .8, High = 0.6, Medium = 0.4
Species in Greatest Conservation Need (SGCN) Richness Grids	Number of SGCN species using Jenks natural breaks: 0=0, 1-2 =.1, 3-7 = .3, 8-14 = .6, 15-24 = .8, 25-48 = 1
Endangered, Threatened, S1, S2 SGCN Richness Grids	Number of species using Jenks natural breaks: 0=0, 1-2 =.1, 3-5 =.3, 6-10 =.6, 11-16 =.8, 17-21 =1
Stream Indices of Biological Integrity (IBI)	[Max WID Fish IBI or Invert IBI score]/100
Lakes a. WHAF Lake Health Score (LHS) b. Lakes of biological significance	Combine a & b below, using b where a was not available. a. Lake Health Score/100 (minimum value 0.45); b. Lakes of biological significance where Outstanding = 0.8, High = 0.6




Figure 6.1. Core areas of the CAN depicting streams and lakes (blue) and terrestrial habitat and wetlands (Green).




Figure 6.2. Prairie or open habitat (orange/brown) and forest (magenta) areas of connectivity.




Figure 6.3. Sites of Biodiversity Significance scoring layer, with Below (gray) = 0, Medium (Light mint green) = 0.4, High (green) = 0.6, and Outstanding (dark green) = 0.8.




Figure 6.4. Number of all SGCN records summed within 2 km² hexagons and scored using Jenks natural breaks.




Figure 6.5. Number of MN Endangered, Threatened, S1, and S2 SGCN records summed within 2 km² hexagons, and scored using Jenks natural breaks.




Figure 6.6. Stream score from the maximum fish or invertebrate IBI value by stream segment (WID) normalized to a range between 0 and 1.




Figure 6.7. Lake scores based on Watershed Health Assessment Framework (WHAF) Lake Health Scores and Lakes of Biological Significance rankings where Lake Health Scores were not available.




Figure 6.8. The Conservation Action Network, including core areas, forest connectivity, and prairie connectivity scored.

References for the Conservation Action Network

- Duflot, R., Avon, C., Roche, P., & Bergès, L. (2018). Combining habitat suitability models and spatial graphs for more effective landscape conservation planning: An applied methodological framework and a species case study. *Journal for Nature Conservation*, 46, 38–47. <https://doi.org/10.1016/j.jnc.2018.08.005>
- Fridley, J. D., J. J. Stachowicz, S. Naeem, D. F. Sax, E. W. Seabloom, M. D. Smith, T. J. Stohlgren, D. Tilman, and B. Von Holle. 2007. The invasion paradox: Reconciling pattern and process in species invasions. *Ecology* 88, no. 1: 3–17.
- Gilbert-Norton, L., R. Wilson, J. R. Stevens, and K. H. Beard. 2010. A meta-analytic review of corridor effectiveness. *Conservation Biology* 24, no. 3: 660–68.
- Haddad, N. M., L. A. Brudvig, E. I. Damschen, D. M. Evans, B. L. Johnson, D. J. Levey, J. L. Orrock, J. Resasco, L. L. Sullivan, J. J. Tewksbury, S. A. Wagner, and A. J. Weldon. 2014. Potential negative ecological effects of corridors. *Conservation Biology* 28, no. 5: 1178–87.
- Hooper, D. U., F. S. Chapin III, J. J. Ewel, A. Hector, P. Inchausti, S. Lavorel, J. H. Lawton, D. M. Lodge, M. Loreau, S. Naeem, B. Schmid, H. Setälä, A. J. Symstad, J. Vandermeer, and D. A. Wardle. 2005. Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs* 75, no. 1: 3–35.
- Jenks, George F. 1967. "The Data Model Concept in Statistical Mapping", *International Yearbook of Cartography* 7: 186–190.
- Levine, J. M., and C. M. D'Antonio. 1999. Elton revisited: A review of evidence linking diversity and invasibility. *Oikos* 87: 15–26.
- Pelletier D, Clark M, Anderson MG, Rayfield B, Wulder MA, Cardille JA (2014) Applying Circuit Theory for Corridor Expansion and Management at Regional Scales: Tiling, Pinch Points, and Omnidirectional Connectivity. *PLoS ONE* 9(1): e84135. <https://doi.org/10.1371/journal.pone.0084135>
- Yan, Sun, H. Müller-Schärer, J. L. Maron, and U. Schaffner. 2015. Origin matters: Diversity affects the performance of alien invasive species but not of native species. *American Naturalist* 185, no. 6: 725–36.

Conservation Opportunity Areas

Conservation Opportunity Area: What are They?

Conservation Opportunity Areas (COAs) are designated geographic areas where high biodiversity and active partnerships overlap and where on-the-ground actions are focused to directly benefit Species in Greatest Conservation Need (SGCN) and their associated habitats. This network of partnerships provides a non-regulatory approach to help focus and prioritize statewide conservation actions to prevent future Federal or state listing of species and to promote the recovery of species that have already been listed. Partners may include private landowners and non-government organizations as well as federal, state, and local government agencies collaborating to identify, design, and implement coordinated on-the-ground conservation actions that not only benefit SGCNs but help partners reach their specific conservation goals.

Conservation Opportunity Area Delineation

Locations of COAs are based on core areas of high biodiversity, SGCN, high-quality habitats, and buffers and corridors identified in the Conservation Action Network (CAN). However, not all important areas identified in the CAN have an associated COA. COAs must also include resources such as organized and willing partners, and/or funding available to address conservation needs. As a result, many of the COAs expand beyond those priority areas identified in the CAN to incorporate areas where established partnerships are actively collaborating to deliver habitat management projects that have the potential to benefit SGCN. These partnership efforts present unique opportunities to capitalize on established working relationships and shared conservation goals to achieve the goals of Minnesota's Wildlife Action Plan (MNWAP). The 2025 version of the MNWAP designates 39 COAs (see Fig 6.9).

Focusing Habitat Management Efforts: the Conservation Action Network and Conservation Opportunity Areas

The footprint of many of the COAs have been expanded since 2015 to incorporate areas where established partnerships are actively collaborating. To focus habitat management efforts within critical areas designated by the CAN, the COAs will prioritize State Wildlife Grant project funding based on the location of the project in relation to the priority areas designated by the CAN. Focusing habitat management, restoration, and protection for SGCN within the CAN will enhance core areas of habitat and connections between habitats. This approach will also allow land managers/owners to use existing partnerships while focusing their efforts on priority areas.

Building Relationships, Capitalizing on Opportunities, and Working in Partnership

Since the first SWAP was established in 2005, broad-based collaborative partnerships have proven to be a very successful way to implement natural resource-based projects nationwide. By sharing resources and working collaboratively to overcome roadblocks, partnerships between federal, state, and local government entities, private landowners, non-government organizations and local stakeholder groups have been able to implement




Figure 6.9. This map of Conservation Opportunity Areas (COAs) for 2025-2035 was developed with input from internal and external partners. The total number of COA's grew from 37 to 39. Two additional COAs proposed in north central Minnesota require further consultation with Tribes for delineation.

large-scale watershed level projects that cross jurisdictional boundaries as well as smaller site-based projects. Since the COA coordinator position was created 7 years ago, the partnership approach and the relationships built through the local COAs have increased the capacity of the DNR Nongame Wildlife Program to implement effective habitat management, restoration, and improvement projects by facilitating the coordination of management, prioritizing outreach efforts, and targeting technical assistance.

What Makes a COA Effective?

Effective COAs typically hold partnership meetings at least twice a year. These meetings generally occur in the late winter to plan for the upcoming field season and in fall to evaluate and discuss how much progress was made on various efforts. Besides being an opportunity to conservation projects, the meetings often accomplish the following:

- Develop/update dynamic planning documents to guide partnership efforts.
- Provide opportunities to coordinate the implementation of on-the-ground habitat conservation projects to meet the conservation goals of partner organizations while addressing Wildlife Action Plan objectives.
- Identify and prioritize potential future projects to be implemented. This could include assessing available resources needed to implement projects, reviewing new information on species and habitats that has become available, defining project effectiveness, and implementing monitoring to evaluate and report on effectiveness.
- Address opportunities for watershed/landscape project planning within the COA and the Conservation Action Network (CAN). This can include considering how management objectives and actions at the site level can be coordinated across sites to sustain or enhance landscape-scale biological diversity and improve the ecological functionality of the habitat types involved.

Case Study: Build Biodiversity or Bus(t): Grazing, Wildlife, and Cover Crops

A field tour was an identified action item by partners of the Prairie Coteau Conservation Opportunity Area in southwest Minnesota. Many of the SGCN are imperiled due to conservation issues related to prairie/grassland loss and fragmentation, stream sedimentation and contamination, channelization, and impoundments which this tour hoped to address. The goal of this field tour was to showcase good examples of management, especially for grazing and cover crops, and to emphasize how these practices can directly or indirectly improve wildlife habitat. The target audience was private landowners and agency personnel. Around forty people attended the tour which started at Southwest Minnesota State University in Marshall, Minnesota and travelled to a local farm that showcased regenerative grazing and cover crops. Breakout sessions at this farm included Grazing for Soil Health, Small Ruminant Economics, Water and Stream Quality, and Bird Use of Cover Crops. The bus then travelled to Hole in the Mountain Wildlife Management Area to showcase a patch-burn grazing operation in partnership with a local agricultural producer. Breakout groups included Effects of Grazing on Nongame Species, and a discussion on forage differences between warm and cool season grasses and goal-oriented grazing.

Funding for this field tour was provided by a USFWS State Wildlife Grant. Partners who were involved in the planning, advertisement, and implementation of the field day included the Minnesota Grazing Lands Conservation Association, the DNR Nongame Wildlife Program, Sustainable Farming Association, Pheasants Forever, Natural Resources Conservation Service, Pipestone Soil and Water Conservation District, and the Minnesota Pollution Control Agency.

Improving Habitat for SGCN While Meeting Shared Objectives

Minnesota has an abundance of habitat management and restoration work being accomplished by state, federal, and local government entities as well as non-government organizations and private landowners.

Conservation Opportunity Areas often contain multiple partnership groups that are actively collaborating on project implementation. Each of these groups may have a slightly different approach to their efforts and/or a slightly different set of habitat types in which they work. While there may be different goals associated with each partner's involvement, each partner is able to bring resources to the table to address those goals while enhancing the overall effectiveness of a project. In most cases, managing habitat for the benefit of SGCN incorporates many of the same land management practices as managing for non-SGCN wildlife. Often just changing the timing of implementation, leaving patches of a specific habitat type untouched, or the implementation of follow-up practices such as reseeding, grazing, burning, or mowing can address the needs of SGCN within a project site. Working with partners to coordinate project planning can incorporate these types of practices within the existing framework of the project. In this way, all the partners involved can achieve their intended goals while benefitting a wider range of wildlife species.

Case Study: “Driftless Habitat Revival: Partnering with Private Landowners”

Many projects are completed on public properties within COAs; however, many agencies and NGO's work with private landowners to implement conservation work on private properties. One private land steward started their conservation journey in 1987 when they bought their first parcel of property in Houston County. The property consisted of agricultural fields, old pastures, and degraded native plant communities. Conservation work began when the land steward received a copy of the Minnesota Conservation Volunteer magazine. In it, they read an article about reforestation and decided they would like to convert one of the old fields to forest, so they partnered with a DNR forester and planted a combination of red and white pine. This was a start, but conversion of the agricultural fields didn't stop there. Many more were enrolled in the Conservation Reserve Program and were planted to a diverse mixture of native grasses and forbs. With many of the fields retired, the land steward began to make a concerted effort to restore the native plant communities on the property. These consisted of oak-hickory forest and woodlands, savanna, and remnant bluff prairie, which were all suffering from non-native species pressure from plants such as honeysuckle and common buckthorn as well as native eastern red cedar on the savanna and bluff prairie. The land steward partnered with several agencies including the NRCS and their Environmental Quality Incentive Program as well as the DNR's Nongame Wildlife Program. Together, efforts were made to first remove the eastern red cedar on the south facing slope of the bluff, and since then, a variety of follow-up treatments including forestry mowing, drone seeding, prescribed fire, and goat grazing have been utilized in an effort to control the woody invasives. These habitats are important for a variety of species near the property including timber rattlesnakes and Dekay's brown snake. Partnering with private landowners will be crucial for the survival of these species in a landscape where most land is privately owned.

Conservation Opportunity Areas in Your Area are a Great Starting Place to Direct Conservation Efforts

5. Local governments can work together on referencing COAs in land use planning and capitalize on opportunities for conservation within those areas
6. Work with property owners whose lands host significant natural features and share and discuss voluntary conservation options ranging from habitat management actions to permanent land conservation
7. Become familiar with the rare plants, animals, and habitats that occur within the COA and learn how to better manage their habitats for long-term conservation
8. Educate landowners and the public about significant natural areas within the COA and create support for local conservation initiatives to conserve them
9. Collaborate with state agencies for funding support

Climate Adaptation

Overview

Climate change is a pervasive stressor that influences every habitat and species in Minnesota. As such, integrating climate adaptation strategies into conservation planning is not optional—it is necessary. This plan addresses climate change throughout, with climate-related stressors and actions incorporated into the chapters on Species of Greatest Conservation Need (SGCN), various habitat sub-chapters, and identified as a priority within the public engagement chapter.

This section provides a concise overview of current and projected climate patterns in Minnesota, introduces climate-smart strategies, and includes references to additional resources. Scientific data, concepts, and strategies are presented to support conservation practitioners in adapting their work in a changing climate. Climate was prominently featured in the 2015 Plan, and the trends identified then, such as rising temperature and shifting precipitation patterns, have continued and are clearly depicted in updated charts and statistics.

Increasing temperatures, altered precipitation patterns, more frequent flooding, and changes in winter duration and growing seasons are driving stress across all ecosystems and are addressed in the habitat sub-chapters. These climate-induced changes are further compounded by other stressors, including invasive non-native species, disease, and challenges related to fire management. These interactions amplify impacts on both wildlife and their habitats.

Minnesota's wildlife already faces significant threats from habitat loss, invasive species, fragmentation, and degradation, as well as exposure to pollutants and pathogens. Climate change exacerbates these threats by altering habitat suitability, increasing prevalence of disease and invasive species, and introducing new physiological challenges to SGCN, such as heat stress or phenological mismatches. For example, shifts in seasonal timing can cause key life stages of one species to fall out of sync with those of its food sources or mutualistic partners.

The following content is intended to equip conservation professionals with foundational knowledge, practical strategies, and illustrative examples to help them plan and act effectively in the face of climate change.

Changes in Minnesota's climate are easily detected through historical records and are projected to continue into the future. Climate remains a topic of great interest, with a growing body of scientific literature exploring its impacts. In this Plan, we present climate information from global trends to localized profiles for Minnesota's nine sub-regions. We also include references and links to current research efforts aimed to understanding how climate change may affect the state's vulnerable wildlife and habitats. To support the implementation of conservation actions outlined in this SWAP, we will provide climate data as part of our online spatial layers. These will depict seasonal temperature and precipitation patterns across the state, along with deviations from both the 30-year baseline and the full period of record.

Building on the two Prior Plans

2005-2015 Tomorrow's Habitat for the Wild and Rare

The first SWAP focused generally on habitat loss and degradation but did not focus specifically on climate change.

Minnesota's Wildlife Action Plan 2015-2025

Climate was identified as a stressor in *Minnesota's Wildlife Action Plan 2015-2025*, which outlined a range of projected impacts on the state's wildlife and their habitats. To inform the Plan, staff reviewed scientific literature and technical reports, and the DNR conducted a climate change vulnerability assessment with habitat experts. This assessment helped identify the habitats most at risk from climate change. The Plan incorporated goals, objectives, and conservation actions aimed at: (1) reducing the impacts of current stressors on habitats and species, (2) increasing the resilience of species and habitats, and (3) addressing species-specific vulnerabilities related to life-history traits that heighten sensitivity to stressors.

In our Strategies section, we present strategies that can assist in making our conservation work more 'climate-smart' following seven principles from the document, *'Voluntary Guidance for States to Incorporate Climate Adaptation into State Wildlife Actions Plans and other Management Plans'* (Association of Fish and Wildlife Agencies 2022). These principles include:

10. Fully integrate climate change into SWAPs—ensuring climate adaptation is embedded throughout planning, rather than treated as a standalone section.
11. Adopt forward-looking goals—setting conservation objectives based on projected future conditions instead of only historic baselines.
12. Explicitly link actions to climate vulnerabilities—directly tying conservation actions to identified climate risks and vulnerabilities in species and habitats.
13. Manage for change, not just persistence—planning for ecological transitions and adaptive responses rather than aiming to maintain current conditions forever.
14. Consider broader landscapes and longer timeframes—expanding our planning beyond immediate areas with short time horizons to account for greater-scale shifts.
15. Address uncertainty by considering future scenarios and use of adaptive management—incorporating scenario planning and iterative monitoring to navigate uncertainty in climate projections.
16. Engage diverse partners with climate experience and expertise—collaborating with individuals and organizations that bring climate-specific knowledge and capabilities.

Global and National Context

Globally, average temperature has risen approximately 2 degrees Fahrenheit (between 1.7 and 2.2 degrees F°; 0.95 and 1.20 degrees Celsius) since 1850-1900; these changes are related to the production of greenhouse gasses by human activities that warm the earth's surface and atmosphere (Intergovernmental Panel on Climate Change (IPCC) 2023; Marvel et al. 2023). The sources of those gasses are primarily the burning of fossil fuels for transportation and energy generation, followed by industrial processes, deforestation, and agricultural practices (Canadell et al. 2021). These higher temperatures have already driven sea level rise and

caused more frequent natural disasters (IPCC 2023). Currently, up to 3.6 billion people globally live in highly vulnerable climate contexts (IPCC 2023).

In the United States, temperatures have increased more than the global average, as higher latitudes are warming more rapidly than lower ones. Since 1970, average temperatures in the contiguous U.S. have risen by 2.5°F, compared to a global increase of approximately 1.7°F during the same period (Marvel et al. 2023). Winter temperatures in many northern states are rising nearly twice as fast as those in summer (Marvel et al. 2023). Precipitation has also increased, with annual totals across much of the Midwest rising by 5%–15% from 1992 to 2021, relative to the 1901–1960 average (Wilson et al. 2023).

These climatic shifts have contributed to more frequent and intense extreme events, including hurricanes, floods, droughts, and catastrophic wildfires. Such conditions increasingly affect human health and safety, with extreme heat alone linked to over 700 deaths per year nationwide between 2004 and 2018 (Vaidyanathan et al. 2020). Additionally, major disasters are occurring more frequently and causing greater economic and infrastructural damage (National Centers for Environmental Information 2022).

Disaster risk is not evenly distributed and geographic, socioeconomic, and demographic factors all influence vulnerability to extreme events. Certain populations face significantly higher risks; for example, older adults and Black Americans are more than twice as likely to die from heat-related illnesses compared to younger individuals or the general population, respectively (Environmental Protection Agency 2022). Discriminatory urban zoning policies, have contributed to environmental inequities, with some low-income neighborhoods experiencing temperatures up to 12°F higher during heatwaves than wealthier areas within the same city (Hoffman et al. 2020). These communities also face a substantially greater risk of flooding (Wing et al. 2022).

Rising temperatures, along with corresponding shifts in precipitation patterns, place increasing stress on both natural and built environments affecting people, wildlife, and plant communities. The magnitude of future climate change will depend largely on global greenhouse gas emissions, which could range from aggressive reductions to more likely scenarios based on current policies. These projections estimate an increase in global temperatures of approximately 3.6 to 6.7 ° F (2 to 3.7 ° C) by 2100 (Marvel et al. 2023). Even if emissions were reduced to net zero, residual climate impacts would persist due to changes already set in motion, such as elevated ocean surface temperatures, rising sea levels, and diminishing sea ice extent (IPCC 2023).

Climate Context for Minnesota

Between 1895 and 2020, average temperatures in Minnesota increased by 3.0 °F, while annual precipitation rose by an average of 3.4 inches ([Climate Trends](#)). The most significant changes have occurred in recent decades. Compared to 20th-century average, nearly every year since 1970 has been warmer, wetter, or both, and all the top ten warmest and wettest years on record occurred between 1998 and 2020 ([Climate Trends](#)).

Winter warming in Minnesota is especially pronounced (see Figure 6.10). In fact, Minnesota has experienced the most significant winter temperature increases of any contiguous U.S. state ([Climate Change in Minnesota](#) citing Liess et al. 2022). Since 1895, average temperatures during winter (December through February) have risen two to three times faster than those in summer (June through August) ([Climate Trends](#)). The pace of winter warming has accelerated in recent decades. Between 1970 and 2021, average daily winter low temperatures increased more than 15 times faster than average daily summer high temperatures. Since 1985, Minnesota’s average winter temperatures have risen by 5.4°F, with average winter low temperatures

increasing by 6.8°F ([Climate Change in Minnesota](#)). These changes have led to a shorter season of snow cover across the region (Contosta et al. 2020), And Minnesota has lost approximately 10-14 days of lake ice over the past 50 years (MPCA and MDOC 2025).

Heavy rainfall events in Minnesota have become more common and intense than at any point on record. Since 2000, the state has experienced a significant increase in large-scale, extreme rainstorms. Rainfall events that once fell in the 98th percentile (historically among the largest 2% annually) are now occurring more regularly ([Climate Trends](#)). The number of very heavy rain events, defined as six inches or more in a single day, has been two to three times greater since 2000 compared to 20th century (Runkle et al. 2022; [NOAA National Centers for Environmental Information State Climate Summaries 2022: Minnesota](#)). In addition, a range of extreme weather events including extreme droughts, summer heat waves, severe storms, flooding, and tornadoes caused more than \$4 billion in property damage between 2000 and 2012 (Runkle et al. 2022).

DNR recently developed a set of geospatial layers illustrating climatic conditions across the state over the past 30 years (1991 – 2020), known as [Minnesota Climate Normal, 1991-2020](#). These layers are publicly available through the [Minnesota Geospatial Commons](#), a website that provides free and open access to Minnesota spatial data. The layers include annual and seasonal climate summaries, as well as comparisons between the most recent 30-year period and the historical record (1895 – 2020). Figures 6.11 and 6.12 present maps of average annual temperature and precipitation during 1991-2020, in comparison to the long-term historical averages, based on this dataset.

Information about Minnesota’s goals and trends regarding greenhouse gas emissions are reported biannually (Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC) 2025). Data visualization for the information in that report is provided in a set of graphics online in the [Minnesota Greenhouse Gas Inventory](#).




Figure 6.10. Chart depicting Minnesota Average Winter Daily Minimum Temperatures (December through February 1896 – 2021) by the DNRState Climatology Office based on data from the Minnesota Climate Explorer.




Figure 6.11. Annual average temperatures (1991-2020) deviation when compared to historical average (1895-2020) (DNR Geospatial Commons 2024).




Figure 6.12. Annual average precipitation (1991-2020) deviation when compared to historical average (1895-2020) (DNR Geospatial Commons 2024).

Future Climate Projections for Minnesota

Trends for warmer and wetter conditions are projected to continue into the future (see Figure 6.13). By mid-century (2040-2059), average annual temperatures in Minnesota are projected to increase by 3.8 - 4.5 °F (depending on future emissions scenarios) and average annual precipitation is projected to increase by up to 1.2 inches, depending on the emissions scenario (Coffman et al. 2024; [Climate Change in Minnesota](#)). When looking toward the end of century, both temperature and precipitation are projected to continue increasing.




Figure 6.13. Observed and projected temperature changes in Minnesota under “lower” (teal) and “higher” (red) emissions scenarios out to 2100 compared to historical temperature observations (orange). Figure from: Runkle, J., K.E. Kunkel, R. Frankson, D.R. Easterling, S.M. Champion, 2022: Minnesota State Climate Summary 2022. NOAA Technical Report NESDIS 150-MN. NOAA/NESDIS, Silver Spring, MD, 4 pp.

Another valuable source of projected rainfall and temperature data for Minnesota comes from reports developed by the U.S. Geological Survey (USGS) Midwest Climate Adaptation Science Center, prepared in support of [State Wildlife Action Planning in the Midwest](#). These reports include tables that allow users to explore climate projections under different models and greenhouse gas emissions scenarios. They offer comparisons to historical baseline conditions (1971-2000) for both mid-century (2040 -2069) and late-century (2070-2089) timeframes. For example, the table below presents projected annual temperature changes for the Mixed Wood Plains Ecoregion, which extends through central Minnesota, Wisconsin, and into Michigan (Ratcliffe et al. 2025).

Table 30 Excerpted from Ratcliffe et al. 2025: Projected changes in annual temperature in the Mixed Wood Plains. Projections use data obtained via the Climate Toolbox Climate Mapper (Hegewisch and Abatzoglou, 2024) based on 20 CMIP5 climate models downscaled using MACAv2-METDATA for moderate- and high-emissions (RCP 4.5 and 8.5) scenarios (Abatzoglou, 2013; Abatzoglou and Brown, 2012). Values show average historical (1971-2000) and projected annual mean temperature (°F) by mid- (2040-2069) and end of the century (2070-2099), as well as the percent change relative to the historical value (in parentheses), for each emissions scenario. To capture variation across climate models, models with the lowest (minimum model) and highest (maximum model) projected temperature under RCP 8.5 are presented in addition to the mean of all models.

Type of model	1971-2000 Historical	2049-2069 RCP 4.5	2049-2069 RCP 8.5	2070-2099 RCP 4.5	2070-2099 RCP 8.5
Minimum model	43.7 °F	46.4 °F (6%)	47.4 °F (8%)	47.3 °F (8%)	50.1 °F (15%)
Mean of models	43.7 °F	48.9 °F (12%)	50.5 °F (16%)	50 °F (14%)	54.6 °F (25%)
Maximum model	43.9 °F	51.9 °F (18%)	53.4°F (22%)	53.6 °F (22%)	59.1 °F (35%)

The timing and intensity of precipitation is also expected to shift, with wetter springs, drier summers, shorter snow seasons, heavier rainfall events, and longer dry periods without measurable precipitation ([Climate Change in Minnesota](#)). While weather and climate conditions vary from year to year, long-term trends indicate that overall precipitation increases are expected to continue throughout this century (Climate Trends). An increase in the frequency of severe storms poses challenges for the people of Minnesota, their communities, economies and ecosystems. These impacts extend across a wide range of sectors, including water and soil resources, agriculture, stormwater and wastewater infrastructure, public health and safety, waste management, and emergency response systems.

Despite these overall increases in precipitation, it may seem counter-intuitive, but future droughts are likely to become more frequent due to rising temperatures, which accelerate soil moisture loss (Runkle et al. 2022).

For more detailed projections of various climate factors such as temperature, precipitation, humidity, and snow under different future climate scenarios, please refer to the [Minnesota CliMAT \(Climate Mapping and Analysis Tool\)](#). This interactive tool displays modeled climate data across a statewide map and allows users to explore changes over multiple time horizons, helping to visualize how conditions may vary both spatially and temporally.

Climate Projections by Region within Minnesota

In 2024, the Minnesota Climate Adaptation Partnership released a set of [regional climate summaries](#) for each of the state’s nine regions. These detailed accounts explain observed changes in summer and winter temperatures and precipitation patterns over the past century (since 1895), along with projected changes for coming decades. Each summary also highlights key climate-related impacts on water resources, human health, tribal lifeways, agriculture, and tourism and recreation. Use the links below to access the climate profile for your region of the state.

17. [Northwest](#) - Kittson, Roseau, Marshall, Polk, Pennington, Clearwater, Red Lake, Norman, Mahnomen, Clay, and Becker counties
18. [North Central](#) - Lake of the Woods, Beltrami, Koochiching, Itasca, Cass, and Hubbard counties
19. [Northeast](#) - St. Louis, Lake, and Cook counties
20. [West Central](#) - Otter Tail, Wilkin, Grant, Douglas, Traverse, Stevens, Pope, Big Stone, Swift, Lac qui Parle, Chippewa, and Yellow Medicine counties
21. [Central](#) - Wadena, Todd, Morrison, Benton, Stearns, Sherburne, Kandiyohi, Wright, Meeker, McLeod, Carver, Renville, Scott and Sibley counties
22. [East Central](#) - Aitkin, Crow Wing, Carlton, Pine, Mille Lacs, Kanabec, Isanti, Chisago, Anoka, Washington, Hennepin, and Ramsey counties
23. [Southwest](#) - Redwood, Lincoln, Lyon, Pipestone, Murray, Cottonwood, Rock, Nobles, and Jackson counties
24. [South Central](#) - Le Sueur, Rice, Brown, Nicollet, Blue Earth, Waseca, Steele, Watonwan, Martin, Faribault, Freeborn counties
25. [Southeast](#) - Dakota, Goodhue, Wabasha, Dodge, Olmsted, Winona, Mower, Fillmore and Houston counties

Climate and Species in Greatest Conservation Need

A number of reports have identified characteristics that can increase the sensitivities of species or populations to climate change (e.g., *National Fish, Wildlife, and Plants Climate Adaptation Partnership* [2012]; *Integrating Climate Change into Northeast and Midwest State Wildlife Action Plans* [Staudinger et al. 2015]; *Persist in place or shift in space? Evaluating the adaptive capacity of species to climate change* [Thurman 2022]; and *Wisconsin Initiative on Climate Change Impacts* [Wisconsin's Changing Climate: Impacts and Solutions for a Warming Climate 2021]).

Such characteristics include the following:

- Highly specialized habitat requirements.
- Dependencies on interspecific interactions (host plants or animals).
- Temperature limits or having narrow environmental tolerances.
- Isolated, rare, or declining populations with poor dispersal abilities.
- Long generation times, low fecundity, or reproductive potential.
- Narrow or restricted distribution.
- Special sensitivity to pathogens.
- Sensitivity to human disturbance.

An assessment of wildlife vulnerability to climate change in Wisconsin identified several key patterns (LeDee and Ribic 2015). Amphibians and reptiles were found to be most vulnerable, primarily due to their dependence on specialized habitats microhabitats, and environmental cues. Among birds, approximately two-thirds of the 236 species assessed were considered climate-sensitive, as many have narrow environmental tolerances that are likely to be exceeded under future conditions. Mammals were projected to be affected largely due to increased risks from disease and parasitism (LeDee and Ribic 2015).

A synthesis of four climate change vulnerability assessments focusing on resident species in the Midwest found that those associated with freshwater ecosystems, particularly species affiliated with ephemeral wetlands, cold- or cool-water environments, and shallow streams, were among the most vulnerable to climate change (Small-Lorenz et al. 2013). Species and their associated vulnerabilities include the following:

- Mollusks
 - o Anthropogenic and natural dispersal barriers.
 - o Hydrologic changes: flood regimes, sediment and streambed degradation.
 - o Climate change mitigation impacts (increased ditching, tiling, irrigation).
 - o Interspecific interactions: host fish required.
- Amphibians and reptiles
 - o Anthropogenic barriers to movements (roads).
 - o Hydrologic changes to damp and aquatic habitats; drought.
 - o Climate change mitigation impacts (increased ditching, tiling, irrigation).
 - o Restricted habitat/geology.
- Fish
 - o Temperature changes: cool- and cold-water species.
 - o Precipitation changes: shallow water spawners, wetland species, species intolerant of turbidity.
 - o Water chemistry: sensitive to low dissolved oxygen.
 - o Barriers to movement: anthropogenic, large rivers.

Birds

Studies have documented that some bird species are arriving earlier at their breeding grounds across the northern U.S., a trend linked to climate change (Butler 2003; Wilson 2013). For migratory species, phenological mismatches—when the availability of resources such as food are no longer aligned with species' migratory or breeding cues—are a growing concern. The scarlet tanager (*Piranga olivacea*), for example, has been identified as vulnerable to shifting seasonal patterns and the mistiming of spring conditions (Zumeta & Holmes 1978).

While some species are negatively impacted, others may benefit from a warming climate. In New Hampshire, black-throated blue warblers (*Setophaga caerulescens*) were observed initiating breeding earlier during warmer springs. Early breeders were more likely to produce a second brood, resulting in higher overall reproductive success (Townsend et al. 2013).

Conversely, species such as the Canada warbler (*Cardellina canadensis*) are projected to shift their range northward in tandem with the retreat of the boreal and northern hardwood forests they depend on. Under the most severe climate projections, this species could face complete extirpation from the northeastern U.S. (Rodenhouse *et al.* 2008). Similar declines are anticipated for other bird species that inhabit montane spruce-fir forests at the southern edge of their range in the Midwest and Northeast. These include spruce grouse (*Alcipennis canadensis*), three-toed woodpecker (*Picoides tridactylus*), black-backed woodpecker (*P. arcticus*), yellow-bellied flycatcher (*Empidonax flaviventris*), gray jay (*Perisoreus canadensis*), boreal chickadee (*Poecile hudsonica*), ruby-crowned kinglet (*Corthylio calendula*), blackpoll warbler (*Setophaga striata*), and white-winged crossbill (*Loxia leucoptera*) (Rodenhouse *et al.* 2008).

A study by Culp et al. (2017) completed a detailed climate change vulnerability assessment for 46 migratory birds of the Great Lakes and Upper Midwest region, including a variety of types from waterbirds to warblers. Their analysis explicitly considered the full annual life cycle for these migratory species. They classified 10 (20%) of the species as vulnerable to climate change, including six of Minnesota's SGCN: red-necked grebe, common tern, Forster's tern, black tern, eastern whip-poor-will, and black-throated blue warbler. In most cases the breeding season appeared to be when birds were faced with the greatest vulnerability to climate change; however, in some cases including the migratory and wintering habitat exposure increased species' overall vulnerability. Two species, the black tern and Forster's tern, faced high vulnerability on both the winter and breeding grounds. They also distinguished species which had background risk that increased their vulnerability from those that were more specifically more sensitive to climate changes.

In 2019, National Audubon Society conducted an assessment of climate change and anticipated impacts to North American birds entitled [Survival by Degrees: 389 Bird Species on the Brink](#). Using more than 70 data sources, 140 million bird records, and information on vegetative cover, land use, and climate, the researchers modeled distributional changes to birds under three global warming scenarios based on the 2014 Intergovernmental Panel on Climate Change. They include a 1.5°C warming above preindustrial levels (which is considered imminent between 2030 and 2052 if no climate change mitigation measures are taken); a 2° C warming (which could happen as soon as 2050); and a 3°C warming scenario (which could occur by 2080). Distribution maps for each species for each of the three warming scenarios are available and can be used to assess impacts at the national, regional and state level.

Pfannmuller et al. (2024) assessed the predicted results for Minnesota species. They designated a species as extirpated at a given warming scenario when the NAS models portrayed the absence of any current or newly available "thermal breeding range" for the species in Minnesota. At the lowest warming scenario (1.5°C), the models predicted the extirpation of 27 breeding birds from Minnesota. The vast majority of these species are dependent on the northern boreal forests of the Laurentian Mixed Forest Province. They include such SGCN as the spruce grouse (*Canachites canadensis*), great gray owl, olive-sided flycatcher (*Contopus cooperi*), and Cape May warbler. At the 2°C warming scenario Minnesota may lose a total of 41 breeding birds including the SGCN American goshawk, evening grosbeak, and LeConte's sparrow and a total of 66 species at the 3°C warming scenario including the SGCN lesser scaup (*Aythya affinis*), red-necked grebe (*Podiceps grisegena*), and Franklin's Gull (*Leucophaeus pipixcan*).

Turtles

Studies of the Blanding's turtle (*Emydoidea blandingii*) have shown that rising temperatures are correlated with declines in habitat suitability, though these effects may be either mitigated or intensified by human land-use decisions (Millar & Blouin-Demers 2012). Similarly, research on wood turtles (*Glyptemys insculpta*) in Massachusetts found that annual flooding displaced nearly half of a subpopulation, increased mortality rates, and reduced reproductive success. With flood events projected to become more frequent and intense, the situation may be further exacerbated by impervious surfaces and the hardening of upstream riverbanks (Jones & Sievert 2009).

Another important consideration for turtles is temperature-dependent sex determination. As nest temperature plays a key role in determining the sex of hatchlings, there is growing concern that climate change could skew sex ratios toward either females or males, depending on the species' specific thermal

thresholds. However, for some species such as the eastern box turtle (*Terrapene carolina carolina*), the degree of atmospheric warming required to significantly alter nest temperatures and impact sex ratios is not anticipated until later in the century (Savva et al. 2010).

A recent assessment of Adaptive Capacity of Midwest SGCN completed by the University of Michigan in association with other states is described under Principle #3 below.

Species shifting ranges in response to changing climate norms has already been seen and is continuing. Corridors for dispersal and connectivity of habitats is essential for supporting these movements. This is a fundamental function of the Conservation Action Network, to facilitate landscape-scale ecological connectivity. Species movements in response to climate change are already apparent. There is evidence that major changes in species composition and ecological system structure and functions (primary productivity and nutrient cycling) are occurring as a result of plant and animal range shifts (Grimm 2013). Mastrandrea et al. (2010) summarized evidence from 45 studies that indicated significant changes in the timing of life-cycle events for a wide range of plant and animal species. These changes could result in the unavailability of essential food resources during critical life-history stages. The unavailability of a resource during peak migration periods, for example, could reduce the size of the population over time and potentially contribute to its extinction.

When available, we integrate information regarding potential climate impacts to Minnesota SGCN into the SGCN species group sub-chapters and recommend climate-smart conservation actions.

Climate Assessments for Minnesota's Habitats

Each of the habitat sub-chapters in this Plan addresses the context of a changing climate. Where available, we incorporate climate assessments developed through the [State Wildlife Action planning in the Midwest](#) project, conducted by the USGS Midwest Climate Adaptation Science Center (MWCASC). This project was specifically designed to support State Wildlife Action Plans in the Midwest by providing climate-relevant information for major habitats, along with case study profiles for selected SGCN. The assessments offer provide a comprehensive overview of climate stressors, trends, and projected changes across multiple scenarios for each ecoregion and its associated major natural habitat types. Each includes a literature review/synthesis of direct and indirect climate change impacts on habitats and their key or dominant species, a literature-based review of ecological vulnerabilities, and a detailed overview of recommended climate adaptation strategies relevant to each habitat and the threats they face.

The climate assessments are being prepared for seven ecosystem types, listed below, the first four of which are available at the time of this publication and the remaining ones to be incorporated when available:

1. [Temperate Flooded & Swamp Forest](#)
2. [Appalachian Interior Northeastern Forest](#)
3. [North American Bog & Fen](#)
4. [Eastern North American Temperate Freshwater Marsh, Wet Meadow & Shrubland](#)
5. Central and Eastern North American Grassland & Shrubland
6. Central Interior Oak Forest, Woodland, and Savanna
7. Laurentian-Acadian Northern Hardwood – Conifer Forest

We integrated findings from these ecoregional assessments into our corresponding habitat chapters where available. Please note that not all assessments were complete as of July 2025).

Another analysis of projected climate change impacts on Minnesota's grassland and wetland habitats was conducted by MWCASC-affiliated researchers at the University of Wisconsin. A summary of that analysis is presented in the Prairie and Other Grasslands sub-chapter.

There are several great resources regarding our climate change in our forests, including [Climate Change and Minnesota's Forests](#) prepared by the Minnesota Forest Resources Council, which includes sections on recommended management actions management to increase climate adaptability and resilience in our forests. There is work on increasing the adaptive capacity of conifer forests in Minnesota (White et al., 2020)

Climate-smart Strategies and Actions

In this section, we reference the document, *'Voluntary Guidance for States to Incorporate Climate Adaptation into State Wildlife Actions Plans and other Management Plans'* (Association of Fish and Wildlife Agencies 2022). These are the six Principles highlighted in the guidance document:

26. Fully integrate climate change into SWAPs.
27. Adopt forward-looking goals.
28. Explicitly link actions to climate vulnerabilities.
29. Manage for change, not just persistence.
30. Consider broader landscapes and longer timeframes.
31. Address uncertainty by considering future scenarios and use of adaptive management.
32. Engage diverse partners with climate experience and expertise.

1. Fully Integrate Climate Change into SWAPs

As outlined in the sections above, climate change considerations are integrated throughout the SWAP. For Species in Greatest Conservation Need (SGCN), climate change impacts were considered as part of the conservation status assessment (S-rank) process and are also reflected in the list of stressors discussed within each SGCN species group sub-chapter. Within the habitat sub-chapters, projected changes in temperature, precipitation, and hydrology are considered within the standard list of ecological stressors. Climate education and outreach are discussed in Chapter 4. Public Engagement, highlighting the importance of building public awareness and capacity for climate resilience. In Chapter 5. Monitoring, we identify opportunities to better integrate climate considerations into survey, monitoring, and research efforts, supporting a more adaptive and climate-informed approach to conservation.

2. Adopt Forward looking Goals

A key theme in climate-informed ecological conservation is the importance of managing habitats and ecosystems for resilience—the capacity to remain diverse and functionally robust under a wide range of future conditions. Given the projected magnitude of changes in temperature, seasonality, and precipitation, managing for pre-contact or historical conditions may be feasible in some cases, but not universally achievable.

As a result, there is a growing shift in perspective from aiming to restore pre-settlement conditions, toward defining new conservation goals that prioritize biodiversity, ecosystem function, and the delivery of ecosystem

services, even when these outcomes yet may deviate from historic baselines. This adaptive approach acknowledges the dynamic nature of ecosystems in a changing climate and supports more flexible, forward looking management strategies.

3. Explicitly Link Actions to Climate Vulnerabilities

Linking conservation actions to the specific climate vulnerabilities of SGCN, habitats, or ecosystems is a powerful effective way to prioritize efforts where they are likely to have the greatest impact. Climate change vulnerability assessments consider three key components: exposure (the magnitude and rate of climate change a species or habitat is likely to experience), sensitivity (the degree to which it is affected by those changes), and adaptive capacity (ability to cope with, persist through, or adjust to those changes) (Glick et al. 2010; Thurman et al., 2020, see also NatureServe's [Climate Change Vulnerability Index: Species](#)). A more detailed framework for evaluating adaptive capacity was developed by Thurman et al. (2022), who categorized species traits into two adaptive pathways: the capacity to 'persist in place' and the capacity to 'shift in space.'

A major regional effort to assess climate vulnerability and adaptive capacity was conducted by the Michigan Natural Features Inventory in partnership with the Michigan Department of Natural Resources, Nebraska Game and Parks Commission, and the U.S. Geological Survey, with Minnesota DNR serving in an advisory capacity. Funded by a Competitive State Wildlife Grant from the U.S. Fish and Wildlife Service, the project evaluated assessed climate vulnerability and adaptive capacity of 400 Midwest SGCN using a rapid assessment tool focused on 37 life-history traits related to climate vulnerability and resilience (Earl et al., 2024). Of the total, 142 Minnesota of Minnesota's SGCN were included: 6 amphibians, 45 birds, 11 fish, 41 insects, 11 mammals, 22 mollusks, and 6 reptiles. The species groups with the lowest average adaptive capacity scores were invertebrates, with insects scoring lowest, followed by mollusks. Among Minnesota's SGCN, 91 species were ranked as having moderate adaptive capacity, 24 as moderately high, 27 as moderately low, and none were ranked as having low or high adaptive capacity. Notably, a greater number of invertebrates were rated as moderately low in adaptive capacity, while amphibians, birds, and mammals tended to fall into the moderately high category (see Figure 6.14).




Figure 6.14. Graph indicating the overall adaptive capacity assessment scores for Minnesota SGCN evaluated in the Michigan-led adaptive capacity assessment study (Earl et al., 2024).

The 37 life-history traits considered in the adaptive capacity assessment are organized into seven thematic groups: life history, demography, distribution, movement, evolutionary potential, ecological role, and abiotic niche. The traits within each group are evaluated to produce a sub-score for that category, which together inform an overall assessment of a species' adaptive capacity. For example, traits related to distribution and movement reflect a species' ability to shift its range in response to changing climate conditions (i.e., shift in space), whereas attributes within life history and demography often indicate a species' ability to persist in place despite climate stressors (Thurman et al. 2020). The adaptive capacity assessment tool generates visualizations of these trait groupings in a circular graphic format, offering a clear depiction of how different attributes contribute to a species' adaptive capacity (see Figure 6.15).




Figure 6.15. An example of a Summary of Adaptive Capacity for lake sturgeon, which comes out with an average score of 0.5 Moderate (from Earl et al., 2024).

Patterns in adaptive capacity traits across the seven attribute groups can help managers tailor strategies and actions to address the specific factors limiting a species' ability to adapt to climate change. For instance, if a species exhibits greater capacity to shift in space, then actions that enhance habitat connectivity and movement corridors may be most effective. Conversely, for species with limited evolutionary potential, efforts to increase population sizes and promote gene flow may be more effective.

Several high-quality online resources are available to guide conservation actions for wildlife facing stress from climate change. One such tool is the [Wildlife Adaptation Menu](#), developed by the Climate Change Response Framework, which provides adaptation options for wildlife managers. The menu organizes strategies into three categories—resistance, resilience, and transition—and provides specific actions within each. Another valuable resource for climate adaptation strategies is the Great Lakes Indian Fish and Wildlife Commission's (GLIFWC) [Tribal Climate Adaptation Menu](#). This tool offers guidance on incorporating indigenous and traditional knowledge into climate adaptation planning, fostering more holistic and culturally informed conservation approaches.

Another resource is the [Climate Change Bird Atlas](#) hosted by the [Northern Research Station](#) of the United States Department of Agriculture Forest Service (USFS).

In the realm of managing habitats, there are several tools that help one to understand how to tailor management actions to various species traits. One tool is the USFS [Tree Atlas](#) that details models habitat suitability and climate adaptability in terms of reproduction and survival for individual tree species. The

[Northern Forests Climate Hub](#) provides in-depth vulnerability assessments for various forest types and tree species, as well as [adaptation workbooks](#) for managers. A newer addition is the climate adaptation menu for North American grasslands (Bernath-Plaisted et al., 2025), which includes eight overarching strategies that frame 145 specific management actions for implementing climate adaptation in grassland ecosystems.

4. Manage for Change, Not Just Persistence

Traditional restoration and habitat management efforts have focused on recreating ecosystem conditions believed to have existed prior to European settlement. However, the magnitude of climate change interacting with other human-caused stressors is increasingly challenging managers to decide when it is appropriate to manage for the persistence of current or historical conditions, and when it is more effective to manage for ecological change.

The Resist-Accept-Direct (R-A-D) framework was developed in response to the growing mismatch between the scale of ecological change and traditional tools available to address it (Schuurman et al. 2020, Lynch et al. 2021). *R-A-D* offers a structured approach to identifying practical, forward-looking mitigation strategies that prioritize desirable future conditions for both ecosystems and human societies.

The framework provides agencies with: (1) a long-term view of the provisioning of ecosystem services which recognizes agency limitations to manage sources of ecosystem change, (2) a structure to prioritize actions that efficiently achieve management goals, (3) a foundation to re-evaluate current management given new information and resources, and (4) a closer alignment between stakeholder expectations and what an agency is likely to achieve. This last point is key because procedural fairness, a component of good governance (Decker et al. 2016), builds trust in agency policies (Riley et al. 2018, Ford et al. 2020).

The three *R-A-D* strategies are *Resist*, *Accept*, and *Direct* (Schuurman et al. 2020; Figure 6.16). The *Resist* strategy is intended to arrest system change and maintain (or restore) ecosystems to current (or historic) conditions. An *Accept* strategy recognizes that directional change is unavoidable and allows ecosystems to respond without direct intervention on forces that drive that change. A *Direct* strategy recognizes that directional change is unavoidable and aims to steer the ecosystem toward desired novel future conditions while reducing undesired effects on system processes, functions, or composition. An important distinction between *Resist* and *Direct* is that the *Resist* strategy strives to maintain current or historical conditions, whereas the *Direct* strategy strives to promote desired novel ecosystem conditions, often when resisting change is not feasible. In combination, *R-A-D* strategies are intended to cover the full range of policy choices that agencies can evaluate in managing large scale drivers of ecological change. *R-A-D* may best be thought of as a brainstorming tool, often most beneficial when applied at the beginning of a decision-making process, alongside identification of plausible future conditions for the system (Wise et al. 2014; Magness et al. 2022; Bouska et al. 2025). Selecting among *R-A-D* strategies can be done within existing decision-making processes, such as structured-decision making (Runge 2011).

The R-A-D framework can be a valuable tool for guiding conservation planning and action in the face of climate change. Resist strategies are often best suited to areas of climate refugia, where the pace of climate change is relatively slow or in landscapes that have experienced minimal degradation, allowing current conditions to be maintained. Accept strategies involve allowing ecological change to occur while adapting management practices accordingly; examples include diversifying seed mixes, modifying the timing of prescribed fire or grazing, and adjusting to newly emerging conditions. Direct strategies take a more proactive approach by

intentionally guiding ecological transitions. These may include species translocation, assisted migration, or restoring ecosystems to support habitat types that are anticipated in the future, even if they differ from historical or current conditions. An applied example of the R-A-D framework was recently applied to guiding conservation strategies along different reaches of the Mississippi River (Ward et al. 2021).




Figure 6.16. The RAD Framework lays out three approaches for making management decisions for systems undergoing ecosystem transformation: 1) Resist, where managers work to maintain or restore ecosystem composition, structure, processes, or function on the basis of historical or acceptable current conditions, 2) Accept, where managers allow ecosystem composition, structure, process, or function to change autonomously, and 3) Direct, where managers actively shape change in ecosystem composition, structure, processes, or function toward preferred new conditions. (USGS 2025)

5. Consider Broader Landscapes and Longer Timeframes

Providing a place for animals and plants to move, even if very slowly over the course of the next century or more, is an essential part of planning for a resilient landscape. Buffering or expanding protected areas for high-quality habitats may be needed to enable biodiversity to persist by shifting to adapt to changing conditions. Corridors that enable organisms to move among high conservation value areas, such as delineated in the Conservation Action Network, are also likely to play a significant role in enabling plants and animals to adapt to a changing climate.

Climate change over the past 30 years is easily detected and depicted (see Minnesota DNR's [Climate Trends](#)). Projections of future climate track the same warming trajectory into the future. Minnesota's climate is projected to continue changing directionally over the course of the century (see University of Minnesota's [Minnesota CliMAT Climate Mapping and Analysis Tool](#)). Even if all increases in carbon emissions were immediately halted there would be significant residual increases in temperature. Although State Wildlife Plans

are revised every ten years, climate-smart planning for biodiversity conservation must reference a much longer planning horizon to plan for climatic changes across the scale of the century.

6. Address Uncertainty by Considering Future Scenarios and Use of Adaptive Management

Uncertainties associated with climate change stem from four main sources: 1) uncertainty of how much the climate will change, 2) uncertainty in how wildlife, fish, and habitats will react to the change, 3) uncertainty in what management actions to take to address changes, and 4) uncertainty in how people will respond to climate change, both in terms of adaptation as well as mitigation (AFWA 2022).

In the face of this uncertainty, planning and management approaches that enable one to envision multiple potential outcomes such as adaptive management and scenario-based planning are essential. Adaptive management is a strong approach for climate-smart ecological conservation because it allows for the incorporation of uncertainty and the ability to select actions as new conditions unfold over time. The ‘Resist-Accept-Direct’ decision framework, described above, guides one into thinking about pushing a system towards past conditions, supporting it in whatever transitions it might or might not take, or actively managing it toward different future conditions. ‘Scenario management’ allows for planning to include multiple outcomes, which might not be able to be selected immediately due to the dependence on future events that cannot be precisely projected (e.g., drought, wildfire, series of warm winters).

Adaptive management is a structured decision-making approach (see also Chapter 5. Monitoring) that centers on iterative learning and adjusting management decisions on that that learning: “plan, do, monitor, and learn;” it has been described in the climate realm as the ‘climate-smart conservation cycle’ (Stein et al. 2014).

Monitoring is a key part of the adaptive management cycle, that is particularly worth emphasis in the climate-smart conservation realm. It is essential that we continue to collect information about the interaction between climate and SGCN and habitats, as well as invest in robust means to evaluate monitoring and effectiveness of climate-informed conservation (Oakes et al. 2021). As we get a better understanding of how our climate is changing, it becomes evident that we lack information about how species are responding to the changes. We need to build important questions about changes in climate into many of our surveillance studies and consider how climate may interact with our studies of the effectiveness of conservation actions as well. When we design wildlife monitoring studies, we should integrate hypotheses and predictions about how wildlife may be interacting with heat, flooding, drought, or other climatic variables. Adding monitoring components such as microhabitat conditions, such as by using data loggers, may assist in relating behavioral and habitat use profiles to climatic data.

There are several descriptions of adaptive management that explicitly considers a changing climate.

- Investigate climate by integrating into ongoing monitoring programs and projects, such as by including abiotic micro-scale monitoring to provide feedback on models about climate change effects.
- 33. Create ways to test models and make them better.
- 34. Monitor changes in ecosystem composition, structure, and function relative to climate predictions.
- 35. Apply an experimental approach to restoration projects testing hypotheses about which seed sources or practices are most successful.

Scenario-Based Climate Change Adaptation is a strategy for planners to incorporate climate change and its significant impacts to habitats when knowing their exact timing and nature of future impacts is difficult. Scenario-Based Climate Change Adaptation is a strategy that helps planners work with the wide array of future possibilities that climate change may bring (See Figure 6.17). This process uses a small set of possible scenarios that cover a wide range of problems in hope of gaining understanding and practice with these problems, in preparation for when they become a reality. The information in this section is based off of the National Parks Service's [Scenario-Based Climate Change Adaptation Showcase](#). To do this process, a set of scenarios must be identified. Good scenarios are plausible, relevant, divergent (covers a range of future possibilities), and challenging. Scenarios that meet these conditions can help planners understand how current practices will fair in the future, identify important holes in understanding (i.e., places for new research) and define future goals. Characterizing and integrating the uncertainty of possible ecological futures is critical to enhance the robustness of scenario planning. The appropriate process to achieve this could combine quantitative modeling with expert elicitation techniques to describe narrative descriptions of possible alternatives (Lawrence et al., 2021; Rangwala et al., 2021).



Figure 6.17. National Park Service image depicting the difference between forecast-based planning for a single future condition vs. scenario-based climate change adaptation that work with a set of plausible futures to capture a broad range of conditions, providing a framework for making decisions.

Examples of scenario-based climate change adaptation planning in the NPS:

[Apostle Islands National Park](#), 2015. An Apostle Islands scenario-based climate change adaptation planning workshop used regional climate science to craft local scenarios for the park and the surrounding lake (Lake Superior, one of the fastest-warming lakes in the world), to inform both resource management and infrastructure (dock) development.

[Isle Royale National Park](#), 2013. Isle Royale National Park and partners used scenarios to explore the range of ways in which climate, species interactions, ecosystem processes, and lake conditions might change and ultimately inform a decision to supplement a dwindling population of the island's first known wolf population, which had inhabited the island since the 1940s.

Workshops may be an effective way to assist land and natural resource managers in climate-smart planning and resource management. As experts, they know their landscape and ecosystems best and are the ones to design any future planning for management. Local workshops are an effective way for managers and stakeholders to learn about local climate changes, think through likely vulnerabilities, imagine alternative scenarios, and sketch out effective strategies to meet the changing nature of their resource area. Examples include the riparian workshop with the beaver from Wyoming (Voluntary Guidance p. 22) and a series of workshops at U.S. Fish and Wildlife Refuges (Bouska et al. 2025). Similar workshops may be helpful for local managers across the state of Minnesota.

7. Engage diverse partners with climate experience and expertise

Climate science is complex, and it is essential to engage with experts to interpret the data, particularly when working with climate projection models that are globally derived and downscaled to regional and local applications. Depictions of data through graphs, figures, and statistics, are vital toward understanding, applying, and communicating about climate. Sources for climate information are numerous and may change over time, but here are a few prominent ones to note:

36. The [Minnesota State Climatology Office](#) is the state's primary source for current, well-explained climate data.
37. The [Minnesota Climate Adaptation Partnership](#) (MCAP) conducts research and offers resources, tools, and training for organizations, individuals, and communities to respond to and plan for changing climate conditions. They offer climate adaptation training programs, host the [Minnesota CliMAT](#) tool, and keep an up-to-date set of applied well-interpreted resources for those who want to learn more about practicing climate adaptation in their work.
38. The [Midwest Climate Adaptation Science Center](#) (MWCASC) is a regional group of university, Tribal, non-profit, and United States Geological Survey (USGS) scientists within the national USGS Climate Adaptation Science Center network. Scientists team up with natural and cultural resource managers to conduct applied research and develop resources and tools to help people, wildlife, fish, water, and land adapt to a changing climate.
39. The [Northern Forests Climate Hub](#) hosted by the United States Department of Agriculture provides in-depth vulnerability assessments for various forest types and tree species, as well as [adaptation workbooks](#) for managers.
40. The [Northern Research Station](#) of the United States Department of Agriculture Forest Service provides a number of resources that pertain to management of sustainable and healthy forests. There is a [Tree Atlas](#) that details modeled habitat suitability and climate adaptability in terms of reproduction and survival for individual tree species, as well as a [Climate Change Bird Atlas](#).

Case Study: North Shore Stream Connectivity (Climate-informed Fish passage)

Climate is a factor in trout stream health, as warming temperatures and more frequent higher floods put strain on coldwater stream riparian systems. Extremely high floods as have happened in recent years are also putting strain on infrastructure, particularly at stream-road crossings. From flooding events, culverts become less functional due to sedimentation and downcutting and then these become impediments to aquatic habitat connectivity and fish passage. The Division of Fisheries in Minnesota Department of Natural Resources recognized an opportunity to map the coldwater stream network on the north shore of Lake Superior and in the St Louis River watershed to prioritize the replacement of culverts in locations to optimize aquatic connectivity of streams with good groundwater sources that may serve as climate refugia. In these prioritized locations, culverts are replaced with road crossings that are designed to hold up under much larger floods. Also designed with gentle gradients and natural bottoms these provide for superior fish passage at lower water levels that can come with projected late summer droughts.

Case Study: Adapting Bottomland Hardwood Forests to a Changing Climate

[Adaptive Silviculture for Climate Change](#) (ASCC) is a collaborative network of experimental forest management trials to evaluate management options under climate change across a variety of forest types throughout North America. Site-specific treatments are developed for local conditions and tailored to meet site-specific management objectives, while still aligning with a common framework for answering questions about how different forest types will respond to future climate. Trials feature three adaptation options (resistance, resilience, and transition), as well as a ‘no action’ treatment where no management is applied. Monitoring includes overstory, mid-story, and understory forest composition and health and productivity evaluations before and after treatment at 3, 5, and 10 years, providing timely and specific feedback for managers.

One example site in this network is a floodplain ecosystem dominated by ash-elm mixed lowland hardwoods in the [Mississippi National River and Recreation Area](#) (MNRRA) in the Minneapolis - St. Paul urban area. This floodplain forest is experiencing warming temperatures, increased frequency of severe storms, and prolonged floods, all projected to increase with climate change. The warming climate has favored the invasive emerald ash borer and resulted in nearly 100% mortality of ash trees. Increasing temperatures and drought stress are projected to reduce suitability for many of the resident tree species while favoring others. Experimental forestry treatments are *resistance*, which restores native floodplain species resistant to current pests and pathogens; *resilience*, which incorporates a wider diversity of flood-tolerant species native to Minnesota; and *transition*, which incorporates species and genotypes from warmer, southern locations. The project features a strong outreach component, engaging community volunteers in tree planting and monitoring to multiply the impacts of the study. Major partners include Mississippi Park Connection, University of Minnesota, Colorado State University, City of St. Paul Parks and Recreation, MNRRA, Northern Institute of Applied Climate Science, and the U.S. Forest Service. The project has received funding from the Wildlife Conservation Society’s Climate Adaptation Fund, as well as the National Park Foundation, Mississippi Park Connection, Minnesota Pollution Control Agency, the Minneapolis-St. Paul National Science Foundation Long-term Ecological Research Program, and 3M.

Case Study: Pigs Eye Lake Climate Resilient Vegetation Experimentation.

The U.S. Army Corps of Engineers and Ramsey County are partnering on an aquatic and terrestrial habitat project in [Pigs Eye Lake](#) of the Mississippi River. One of the project goals is to restore lost floodplain forest areas. However, the Mississippi River is experiencing more frequent and longer duration flood conditions. Thus, the project is experimenting with different mixes of tree species, including some that are more flood tolerant. It is too early on in the project to know the outcomes, but the project provides an example of how experimentation may be worked in to habitat restoration projects and inform future work.

References for Climate Adaptation

Abatzoglou, J. T. (2013). Development of gridded surface meteorological data for ecological applications and modelling. *International Journal of Climatology*, 33(1), 121–131. <https://doi.org/10.1002/joc.3413>

Abatzoglou, J. T., & Brown, T. J. (2012). A comparison of statistical downscaling methods suited for wildfire applications. *International Journal of Climatology*, 32(5), 772–780. <https://doi.org/10.1002/joc.2312>

Association of Fish and Wildlife Agencies (AFWA). 2022. [Voluntary Guidance for States to Incorporate Climate Adaptation in State Wildlife Action Plans and Other Management Plans. 2nd edition](#). Editors: Climate Adaptation and Wildlife Diversity Conservation and Funding Committees, Voluntary Guidance Revision Work Group. Association of Fish and Wildlife Agencies, Washington, DC.

Bartsch, Will, Cai, Mei, Johnson, Kris, Nixon, Kristi, Sprague, Tiffany, Wright, Chris, Olson, Louis, and Reed, Jane. 2023. Camp Ripley Sentinel Landscape Climate Resilience Analysis and Strategic Plan Amendments. Natural Resources Research Institute, University of Minnesota Duluth, Technical Report NRRI/TR-2023/21. 56 p. + appendices.

Bernath-Plaisted, J. S., Handler, S. D., Ahlering, M., Brandt, L. A., Maresh Nelson, S. B., Niemuth, N. D., Ontl, T., Peterson, C. L., Ribic, C. A., Strohmeier, D., & Zuckerberg, B. (2025). A climate adaptation menu for North American grasslands. *Conservation Science and Practice*, e70017, <https://doi.org/10.1111/csp2.70017>.

Bouska, K.L., Booker, J., Clark, S., Delaney, J., Eash, J., Post van der Burg, M., and Roop, H., 2025, Managing for tomorrow—A climate adaptation decision framework: U.S. Geological Survey Open-File Report 2025–1005, 53 p., <https://doi.org/10.3133/ofr20251005>.

Butler, C. J. (2003). The disproportionate effect of global warming on the arrival dates of short-distance migratory birds in North America. *Ibis*, 145(3), 484–495. <https://doi.org/10.1046/j.1474-919X.2003.00193.x>

Canadell, J. G., Monteiro, P. M. S., Costa, M. H., Cotrim da Cunha, L., Cox, P. M., & other contributing authors. (2021). Chapter 5: Global carbon and other biogeochemical cycles and feedbacks. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press

Coffman, D., Black, K., Boyd, K., Clark, S., Farris, A., Greene, B., Leon, A., Saravana, D., Weske, C. 2024. Climate Change in Minnesota. Prepared for the University of Minnesota Climate Adaptation Partnership. Version 1; September 2024. www.climate.umn.edu/climate-change-in-minnesota

- Contosta, A. R., Casson, N. J., Nelson, S. J., & Garlick, S. (2020). Defining frigid winter illuminates its loss across seasonally snow-covered areas of eastern North America. *Environmental Research Letters*, 15(3), 034020. <https://doi.org/10.1088/1748-9326/ab54f3>
- Culp, L. A., Cohen, E. B., Scarpignato, A. L., Thogmartin, W. E., & Marra, P. P. (2017). Full annual cycle climate change vulnerability assessment for migratory birds. *Ecosphere*, 8(3), e01565. <https://doi.org/10.1002/ecs2.1565>
- Decker, D., Smith, C., Forstchen, A., Hare, D., Pomeranz, E., Doyle-Capitman, C., Schuler, K., & Organ, J. (2016). Governance Principles for Wildlife Conservation in the 21st Century: Governance principles for wildlife conservation. *Conservation Letters*, 9(4), 290–295. <https://doi.org/10.1111/conl.12211>
- Earl, D.J., E.C. Branch, A.A. Cole-Wick, Y. Lee, L.M. Rowe, P.J. Badra, C.M. Wilton, D.L. Cuthrell and N.H. Sexton. 2024. Assessing Climate Vulnerability & Adaptive Capacity of Midwest Species of Greatest Conservation Need. Michigan Natural Features Inventory, Report No. 2024-37, Lansing, MI.
- Environmental Protection Agency. 2022: Climate Change Indicators: Heat-Related Deaths. U.S. Environmental Protection Agency. <https://www.epa.gov/climate-indicators/climate-change-indicators-heat-related-deaths>
- Ford, J. K., Riley, S. J., Lauricella, T. K., & Van Fossen, J. A. (2020). Factors Affecting Trust Among Natural Resources Stakeholders, Partners, and Strategic Alliance Members: A Meta-Analytic Investigation. *Frontiers in Communication*, 5, 9. <https://doi.org/10.3389/fcomm.2020.00009>
- Glick P, Stein BA (eds). 2010. Scanning the Conservation Horizon: A Guide to Climate Change Vulnerability Assessments (Draft). National Wildlife Federation, Washington D.C.
- Grimm NB, Staudinger MD, Staudt A, et al. 2013. Climate-change impacts on ecological systems: introduction to a US assessment. *Frontiers in Ecology and the Environment* 11(9): 456-464.
- Hegewisch, K. C., and J. T. Abatzoglou. 2024. “Historical Climate Tracker Web Tool.” Climate Toolbox. <https://climatetoolbox.org/>.
- Hoffman, J.S., V. Shandas, and N. Pendleton, 2020: The effects of historical housing policies on resident exposure to intra-urban heat: A study of 108 US urban areas. *Climate*, 8 (1), 12. <https://doi.org/10.3390/cli8010012>
- IPCC. 2023: Summary for Policymakers. In: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland, pp. 1-34, doi: 10.59327/IPCC/AR6-9789291691647.001
- Jones, M. T., & Sievert, P. R. (2009). Effects of Stochastic Flood Disturbance on Adult Wood Turtles, *Glyptemys insculpta*, in Massachusetts. *The Canadian Field-Naturalist*, 123(4), 313. <https://doi.org/10.22621/cfn.v123i4.1000>
- Lawrence, D. J., A. N. Runyon, J. E. Gross, G. W. Schuurman, and B. W. Miller. 2021. Divergent, plausible, and relevant climate futures for near- and long-term resource planning. *Climatic Change* 167:38.

LeDee OE, Ribic CA. (2015). Determining climate change management priorities: A case study from Wisconsin. *Journal of Conservation Planning* 11:1-12

Liess, S., Twine, T., Snyder, P., Hutchinson, W., Konar-Steenberg, G., Keeler, B., Brauman, K., 2022. High-Resolution Climate Projections Over Minnesota for the 21st Century. *Earth and Space Science*. 9 (3) <https://doi.org/10.1029/2021EA001893>

Lynch, A. J., Thompson, L. M., Beever, E. A., Cole, D. N., Engman, A. C., Hawkins Hoffman, C., Jackson, S. T., Krabbenhoft, T. J., Lawrence, D. J., Limpinsel, D., Magill, R. T., Melvin, T. A., Morton, J. M., Newman, R. A., Peterson, J. O., Porath, M. T., Rahel, F. J., Schuurman, G. W., Sethi, S. A., & Wilkening, J. L. (2021). Managing for RADical ecosystem change: applying the Resist-Accept-Direct (RAD) framework. *Frontiers in Ecology and the Environment*, 19(8), 461–469. <https://doi.org/10.1002/fee.2377>

Magness, D.R., Hoang, L., Belote, R.T., Brennan, J., Carr, W., Stuart Chapin III, F., Clifford, K., Morrison, W., Morton, J.M. and Sofaer, H.R., 2022. Management foundations for navigating ecological transformation by resisting, accepting, or directing social–ecological change. *BioScience*, 72(1), pp.30-44.

Marvel, K., W. Su, R. Delgado, S. Aarons, A. Chatterjee, M.E. Garcia, Z. Hausfather, K. Hayhoe, D.A. Hence, E.B. Jewett, A. Robel, D. Singh, A. Tripathi, and R.S. Vose, 2023: Ch. 2. Climate trends. In: *Fifth National Climate Assessment*. Crimmins, A.R., C.W. Avery, D.R. Easterling, K.E. Kunkel, B.C. Stewart, and T.K. Maycock, Eds. U.S. Global Change Research Program, Washington, DC, USA. <https://doi.org/10.7930/NCA5.2023.CH2>

Mastrandrea, M. D., Heller, N. E., Root, T. L., & Schneider, S. H. (2010). Bridging the gap: linking climate-impacts research with adaptation planning and management. *Climatic Change*, 100(1), 87–101. <https://doi.org/10.1007/s10584-010-9827-4>

Millar, C. S., & Blouin-Demers, G. (2012). Habitat suitability modelling for species at risk is sensitive to algorithm and scale: A case study of Blanding’s turtle, *Emydoidea blandingii*, in Ontario, Canada. *Journal for Nature Conservation*, 20(1), 18–29. <https://doi.org/10.1016/j.jnc.2011.07.004>

Minnesota Pollution Control Agency (MPCA) and Minnesota Department of Commerce (MDOC). 2025. Greenhouse gas emissions in Minnesota 2005-2022: Biennial inventory report tracking the state’s greenhouse gas emissions contributing to climate change. Report to Legislature, January 2025. Jointly submitted by the Pollution Control Agency and the Department of Commerce. <https://www.pca.state.mn.us/sites/default/files/Iraq-3sy25.pdf>

National Centers for Environmental Information. 2022: U.S. Billion-Dollar Weather and Climate Disasters. National Oceanic and Atmospheric Administration, National Environmental Satellite, Data, and Information Service, National Centers for Environmental Information. <https://www.ncei.noaa.gov/access/billions/>

National Fish, Wildlife, and Plants Climate Adaptation Partnership. 2012. National Fish, Wildlife and Plants Climate Adaptation Strategy. Association of Fish and Wildlife Agencies, Council on Environmental Quality, Great Lakes Indian Fish and Wildlife Commission, National Oceanic and Atmospheric Administration, and U.S. Fish and Wildlife Service. Washington, D.C. <http://www.wildlifeadaptationstrategy.gov/pdf/NFWPCAS-Final.pdf>

- Oakes, L. E., Peterson St-Laurent, G., Cross, M. S., Washington, T., Tully, E., & Hagerman, S. (2022). Strengthening monitoring and evaluation of multiple benefits in conservation initiatives that aim to foster climate change adaptation. *Conservation Science and Practice*, 4(6), e12688. <https://doi.org/10.1111/csp2.12688>
- Pfannmuller, L., Niemi, G. J., & Green, J. C. (2024). *The breeding birds of Minnesota: history, ecology, and conservation* (K. Rewinkel, Ed.). University of Minnesota Press.
- Rangwala, I., Moss, W., Wolken, J., Rondeau, R., Newlon, K., Guinotte, J., & Travis, W. R. (2021). Uncertainty, Complexity and Constraints: How Do We Robustly Assess Biological Responses under a Rapidly Changing Climate? *Climate*, 9(12), 177. <https://doi.org/10.3390/cli9120177>
- Ratcliffe, H., Charton, K., Siddons, T., Lyons, M. and LeDee, O. 2025. Effects of Climate Change on Midwestern Ecosystems: Temperate Flooded and Swamp Forest. Midwest Climate Adaptation Science Center. (110 pages) Available: <https://mwcasc.umn.edu/node/821>
- Riley, S. J., Ford, J. K., Triezenberg, H. A., & Lederle, P. E. (2018). Stakeholder trust in a state wildlife agency. *The Journal of Wildlife Management*, 82(7), 1528–1535. <https://doi.org/10.1002/jwmg.21501>
- Rodenhouse, N. L., Matthews, S. N., McFarland, K. P., Lambert, J. D., Iverson, L. R., Prasad, A., Sillett, T. S., & Holmes, R. T. (2008). Potential effects of climate change on birds of the Northeast. *Mitigation and Adaptation Strategies for Global Change*, 13(5–6), 517–540. <https://doi.org/10.1007/s11027-007-9126-1>
- Runge, M. C. (2011). An Introduction to Adaptive Management for Threatened and Endangered Species. *Journal of Fish and Wildlife Management*, 2(2), 220–233. <https://doi.org/10.3996/082011-JFWM-045>
- Runkle, J., K.E. Kunkel, R. Frankson, D.R. Easterling, S.M. Champion, 2022: Minnesota State Climate Summary 2022. NOAA Technical Report NESDIS 150-MN. NOAA/NESDIS, Silver Spring, MD, 4 pp. <https://statesummaries.ncics.org/chapter/mn/>. Accessed 2/2/2025
- Savva, Y., Swarth, C. W., Gupchup, J., & Szlavecz, K. (2010). Thermal environments of overwintering Eastern Box Turtles (*Terrapene carolina carolina*). *Canadian Journal of Zoology*, 88(11), 1086–1094. <https://doi.org/10.1139/Z10-076>
- Schuurman, G., Cat, H.-H., Cole, D., Lawrence, D., Morton, J., Magness, D., Cravens, A., Covington, S., O'Malley, R., & Fisichelli, N. (2020). *Resist-accept-direct (RAD)—a framework for the 21st-century natural resource manager*. National Park Service. <https://doi.org/10.36967/nrr-2283597>
- Small-Lorenz SL, Walk J, Schneider R. 2013. Climate change vulnerability of fish and wildlife in the US Corn Belt Region. Webinar, December 11, 2013.
- Staudinger, MD, Morelli TL, Bryan AM. (Eds) 2015. In review. Integrating climate change into State Wildlife Action Plans. USGS Cooperative Report. <http://necsc.umass.edu/>
- Stein, B.A., P. Glick, N. Edelson, and A. Staudt (eds.). (2014). *Climate-Smart Conservation: Putting Adaptation Principles into Practice*. National Wildlife Federation, Washington, D.C.

Thurman, L. L., Gross, J. E., Mengelt, C., Beever, E. A., Thompson, L. M., Schuurman, G. W., Hoving, C. L., & Olden, J. D. (2022). Applying assessments of adaptive capacity to inform natural-resource management in a changing climate. *Conservation Biology*, 36(2), e13838. <https://doi.org/10.1111/cobi.13838>

Thurman, L. L., Stein, B. A., Beever, E. A., Foden, W., Geange, S. R., Green, N., Gross, J. E., Lawrence, D. J., LeDee, O., Olden, J. D., Thompson, L. M., & Young, B. E. (2020). Persist in place or shift in space? Evaluating the adaptive capacity of species to climate change. *Frontiers in Ecology and the Environment*, 18(9), 520–528. <https://doi.org/10.1002/fee.2253>

Townsend, A. K., Sillett, T. S., Lany, N. K., Kaiser, S. A., Rodenhouse, N. L., Webster, M. S., & Holmes, R. T. (2013). Warm Springs, Early Lay Dates, and Double Brooding in a North American Migratory Songbird, the Black-Throated Blue Warbler. *PLoS ONE*, 8(4), e59467. <https://doi.org/10.1371/journal.pone.0059467>

United States Geological Survey (USGS) 2025. This is RAD! The Resist-Accept-Direct Framework. <https://www.usgs.gov/programs/cooperative-research-units/rad-resist-accept-direct-framework>. Accessed 4/30/2025.

Vaidyanathan, A., J. Malilay, P. Schramm, and S. Saha, 2020: Heat-related deaths—United States, 2004–2018. *Morbidity and Mortality Weekly Report*, 69 (24), 729–734. <https://doi.org/10.15585/mmwr.mm6924a1>

Ward, N. K., Lynch, A. J., Beever, E. A., Booker, J., Bouska, K. L., Embke, H., Houser, J. N., Kocik, J. F., Kocik, J., Lawrence, D. J., Lemon, M. G., Limpinsel, D., Magee, M. R., Maitland, B. M., McKenna, O., Meier, A., Morton, J. M., Muehlbauer, J. D., Newman, R., ... Wilkening, J. L. (2023). Reimagining large river management using the Resist–Accept–Direct (RAD) framework in the Upper Mississippi River. *Ecological Processes*, 12(1), 48. <https://doi.org/10.1186/s13717-023-00460-x>

White, M. A., Cornett, M. W., Frerker, K., & Etterson, J. R. (2020). Partnerships to Take on Climate Change: Adaptation Forestry and Conifer Strongholds Projects in the Northwoods, Minnesota, USA. *Journal of Forestry*, 118(3), 219–232. <https://doi.org/10.1093/jofore/fvaa005>

Wilson, A.B., J.M. Baker, E.A. Ainsworth, J. Andresen, J.A. Austin, J.S. Dukes, E. Gibbons, B.O. Hoppe, O.E. LeDee, J. Noel, H.A. Roop, S.A. Smith, D.P. Todey, R. Wolf, and J.D. Wood, 2023: Ch. 24. Midwest. In: *Fifth National Climate Assessment*. Crimmins, A.R., C.W. Avery, D.R. Easterling, K.E. Kunkel, B.C. Stewart, and T.K. Maycock, Eds. U.S. Global Change Research Program, Washington, DC, USA. <https://doi.org/10.7930/NCA5.2023.CH24>

Wing, O.E.J., W. Lehman, P.D. Bates, C.C. Sampson, N. Quinn, A.M. Smith, J.C. Neal, J.R. Porter, and C. Kousky, 2022: Inequitable patterns of US flood risk in the Anthropocene. *Nature Climate Change*, 12 (2), 156–162. <https://doi.org/10.1038/s41558-021-01265-6>


Wisconsin Initiative on Climate Change Impacts (WICCI). 2021. *Wisconsin’s Changing Climate: Impacts and Solutions for a Warmer Climate*. Nelson Institute for Environmental Studies, University of Wisconsin–Madison; Wisconsin Department of Natural Resources.


Wise, R.M., Fazey, I., Smith, M.S., Park, S.E., Eakin, H.C., Van Garderen, E.A. and Campbell, B., 2014. Reconceptualising adaptation to climate change as part of pathways of change and response. *Global environmental change*, 28, pp.325-336.

Zumeta, D.C. and Holmes, R.T. (1978) "Habitat Shift and Roadside Mortality of Scarlet Tanagers during a Cold Wet New England Spring," *Wilson Bulletin*: Vol. 90: Iss. 4, Article 9


APPENDIX A – SGCN Designation Flowchart

We developed the following flowchart to determine whether species meet criteria for inclusion on our SGCN or SNI lists. The flowchart has been split into two pieces on the next two pages, indicated by the red line below, for easier reading.





Note: A grey box indicates questions that are more subjective, which require expert knowledge to answer. Dashed arrows provide the choice to take one of multiple paths, based on expert opinion.



APPENDIX B – SGCN and SNI Lists

Tables are presented by taxonomic group. SGCN species are listed first in alphabetical order by scientific name, followed by SNI species in alphabetical order by scientific name.

S-Rank = State conservation status

G-Rank = Global conservation status

1 = Critically Imperiled

2 = Imperiled

3 = Vulnerable

4 = Apparently Secure

5 = Secure

H = Possibly Extinct

X = Presumed Extinct

U = Unrankable (lack of information)

NR = Unranked

S#S# or G#G# = range rank (indicates uncertainty)

Listing Status:

S-SPC = State Special Concern

S-THR = State Threatened

S-END = State Endangered

F-THR = Federally Threatened

F-END = Federally Endangered

F-P-THR = Proposed Federally Threatened

F-P-END = Proposed Federally Endangered

NLW = National Listing Workplan

Amphibians

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Acris blanchardi</i>	Blanchard's cricket frog	S2	G5	S-END	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Ambystoma maculatum</i>	spotted salamander	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, M threats
<i>Anaxyrus cognatus</i>	Great Plains toad	S3	G5	S-SPC	SGCN	2015 SGCN, M threats
<i>Hemidactylium scutatum</i>	four-toed salamander	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN, RFSS, H threats
<i>Lithobates palustris</i>	pickerel frog	S3	G5		SGCN	2015 SGCN, RSGCN Watchlist
<i>Necturus maculosus</i>	mudpuppy	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN, H threats
<i>Notophthalmus viridescens</i>	eastern newt	S3	G5		SGCN	2015 SGCN, RSGCN Watchlist, M threats
<i>Ambystoma mavortium</i>	western tiger salamander	SU	G5		SNI	

Birds

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Accipiter atricapillus</i>	American goshawk	S2	G5	S-SPC	SGCN	2015 SGCN, RFSS, Decline, VH threats
<i>Aechmophorus occidentalis</i>	western grebe	S3	G5		SGCN	2015 SGCN, Decline, H threats
<i>Aegolius funereus</i>	boreal owl	S1	G5	S-SPC	SGCN	2015 SGCN, RFSS, Decline, H threats, Rare
<i>Ammodramus savannarum</i>	grasshopper sparrow	S2	G5		SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Ammospiza leconteii</i>	Leconte's sparrow	S3	G5		SGCN	2015 SGCN, RSGCN, Decline, H threats, Stewardship
<i>Ammospiza nelsoni</i>	Nelson's sparrow	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN, RFSS, H threats
<i>Anas acuta</i>	northern pintail	S3	G5		SGCN	2015 SGCN, Decline, H threats
<i>Anas rubripes</i>	American black duck	S2	G5		SGCN	2015 SGCN, Decline, H threats
<i>Anthus spragueii</i>	Sprague's pipit	SH	G3	S-END	SGCN	2015 SGCN, RSGCN, H threats
<i>Antrastomus vociferus</i>	eastern whip-poor-will	S4	G5		SGCN	2015 SGCN, RSGCN, Decline, M threats
<i>Asio flammeus</i>	short-eared owl	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Athene cunicularia</i>	burrowing owl	SH	G4	S-END	SGCN	2015 SGCN
<i>Aythya affinis</i>	lesser scaup	S3	G5		SGCN	2015 SGCN, H threats
<i>Aythya valisineria</i>	canvasback	S3	G5		SGCN	H threats, limited breeding range, Stewardship
<i>Bartramia longicauda</i>	upland sandpiper	S3	G5		SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Botaurus exilis</i>	least bittern	S4	G4		SGCN	2015 SGCN, Decline
<i>Botaurus lentiginosus</i>	American bittern	S3	G5		SGCN	2015 SGCN, Decline, H threats

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Bucephala albeola</i>	bufflehead	S3	G5		SGCN	H threats, limited breeding range
<i>Buteo lineatus</i>	red-shouldered hawk	S3	G5	S-SPC	SGCN	2015 SGCN, Decline, H threats
<i>Buteo swainsoni</i>	Swainson's hawk	S3	G5		SGCN	2015 SGCN, Decline, H threats
<i>Calcarius ornatus</i>	chestnut-collared longspur	S1	G5	S-END	SGCN	2015 SGCN, RSGCN Watchlist, Decline, H threats, Rare
<i>Calidris canutus</i>	red knot	S3	G4	F-THR	SGCN	2015 SGCN, RSGCN Watchlist, H threats
<i>Calidris subruficollis</i>	buff-breasted sandpiper	S3	G4		SGCN	RSGCN, Decline, H threats, important migratory corridor
<i>Canachites canadensis</i>	spruce grouse	S3	G5		SGCN	2015 SGCN, Decline, H threats
<i>Catharus ustulatus</i>	Swainson's thrush	S3	G5		SGCN	Decline, H threats, Specialist
<i>Centronyx bairdii</i>	Baird's sparrow	SH	G4	S-END	SGCN	2015 SGCN, RSGCN Watchlist, H threats
<i>Centronyx henslowii</i>	Henslow's sparrow	S3	G4	S-END	SGCN	2015 SGCN, RSGCN, H threats
<i>Chaetura pelagica</i>	chimney swift	S3	G4		SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Charadrius melodus</i>	piping plover	S1	G3	S-END, F-END	SGCN	2015 SGCN, RSGCN, Decline, VH threats, Very Rare
<i>Chlidonias niger</i>	black tern	S2	G4		SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Chondestes grammacus</i>	lark sparrow	S3	G5	S-SPC	SGCN	2015 SGCN, Decline, H threats
<i>Chordeiles minor</i>	common nighthawk	S4	G5		SGCN	2015 SGCN, Decline
<i>Circus hudsonius</i>	northern harrier	S3	G5		SGCN	2015 SGCN, Decline, H threats
<i>Cistothorus stellaris</i>	sedge wren	S4	G5		SGCN	2015 SGCN, Decline, H threats, Stewardship
<i>Coccothraustes vespertinus</i>	evening grosbeak	S2	G5		SGCN	2015 SGCN, Decline, M threats
<i>Contopus cooperi</i>	olive-sided flycatcher	S3	G4		SGCN	2015 SGCN, RFSS, Decline, H threats
<i>Coturnicops noveboracensis</i>	yellow rail	S3	G4	S-SPC	SGCN	2015 SGCN, RSGCN, RFSS, Decline, H threats
<i>Dolichonyx oryzivorus</i>	bobolink	S3	G5		SGCN	2015 SGCN, RSGCN, Decline, H threats, Stewardship
<i>Empidonax virescens</i>	Acadian flycatcher	S3	G5	S-SPC	SGCN	2015 SGCN, M threats
<i>Falco peregrinus</i>	peregrine falcon	S3	G4	S-SPC	SGCN	2015 SGCN, M threats
<i>Falco sparverius</i>	American kestrel	S4	G5		SGCN	2015 SGCN, Decline, M threats
<i>Gallinula galeata</i>	common gallinule	S3	G5	S-SPC	SGCN	2015 SGCN, M threats
<i>Gavia immer</i>	common loon	S4	G5		SGCN	2015 SGCN, H threats
<i>Lanius ludovicianus</i>	loggerhead shrike	S2	G4	S-END	SGCN	2015 SGCN, Decline, H threats
<i>Leucophaeus pipixcan</i>	Franklin's gull	S3	G5	S-SPC	SGCN	2015 SGCN, VH threats
<i>Limosa fedoa</i>	marbled godwit	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN, H threats

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Melanerpes erythrocephalus</i>	red-headed woodpecker	S3	G5		SGCN	2015 SGCN, RSGCN, Decline, M threats
<i>Nycticorax nycticorax</i>	black-crowned night-heron	S1	G5		SGCN	2015 SGCN, Decline, M threats
<i>Oporornis agilis</i>	Connecticut warbler	S2	G4		SGCN	2015 SGCN, RSGCN, RFSS, Decline, H threats
<i>Parkesia motacilla</i>	Louisiana waterthrush	S3	G5	S-SPC	SGCN	2015 SGCN, H threats
<i>Pelecanus erythrorhynchos</i>	American white pelican	S3	G4	S-SPC	SGCN	2015 SGCN, Decline, H threats
<i>Phalaropus tricolor</i>	Wilson's phalarope	S3	G5	S-THR	SGCN	2015 SGCN, RFSS, H threats
<i>Picoides arcticus</i>	black-backed woodpecker	S3	G5		SGCN	2015 SGCN, Decline, H threats
<i>Picoides dorsalis</i>	American three-toed woodpecker	S2	G5		SGCN	RFSS, VH threats
<i>Podiceps auritus</i>	horned grebe	SH	G5	S-END	SGCN	2015 SGCN, Decline, H threats
<i>Podiceps grisegena</i>	red-necked grebe	S3	G5		SGCN	2015 SGCN, H threats
<i>Podiceps nigricollis</i>	eared grebe	S3	G5		SGCN	2015 SGCN, H threats
<i>Poecile hudsonicus</i>	boreal chickadee	S3	G5		SGCN	2015 SGCN, H threats
<i>Progne subis</i>	purple martin	S3	G5	S-SPC	SGCN	2015 SGCN, Decline, H threats
<i>Protonotaria citrea</i>	prothonotary warbler	S3	G5		SGCN	2015 SGCN, RSGCN Watchlist, Decline, H threats
<i>Rallus elegans</i>	king rail	SH	G4	S-END	SGCN	2015 SGCN, RSGCN Watchlist, H threats
<i>Setophaga caerulescens</i>	black-throated blue warbler	S3	G5		SGCN	2015 SGCN, H threats
<i>Setophaga castanea</i>	bay-breasted warbler	S3	G5		SGCN	2015 SGCN, H threats
<i>Setophaga cerulea</i>	cerulean warbler	S3	G4	S-SPC	SGCN	2015 SGCN, RSGCN, Decline, M threats
<i>Setophaga tigrina</i>	Cape May warbler	S3	G5		SGCN	2015 SGCN, H threats
<i>Spizella pusilla</i>	field sparrow	S3	G5		SGCN	2015 SGCN, RSGCN Watchlist, Decline, H threats
<i>Sterna forsteri</i>	Forster's tern	S3	G5	S-SPC	SGCN	2015 SGCN, H threats
<i>Sterna hirundo</i>	common tern	S1	G5	S-THR	SGCN	2015 SGCN, Decline, H threats
<i>Strix nebulosa</i>	great gray owl	S3	G5		SGCN	RFSS, H threats, Low breeding densities, Specialist
<i>Sturnella magna</i>	eastern meadowlark	S3	G5		SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Sturnella neglecta</i>	western meadowlark	S3	G5		SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Tringa solitaria</i>	solitary sandpiper	S2	G5		SGCN	H threats, Rare
<i>Tympanuchus cupido</i>	greater prairie-chicken	S3	G4	S-SPC	SGCN	2015 SGCN, RSGCN, H threats
<i>Tympanuchus phasianellus</i>	sharp-tailed grouse	S3	G5		SGCN	2015 SGCN, RSGCN Watchlist, Decline, H threats

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Tyrannus verticalis</i>	western kingbird	S3	G5		SGCN	2015 SGCN, Decline, M threats
<i>Tyto furcata</i>	American barn owl	S2	G5		SGCN	Rare
<i>Vermivora chrysoptera</i>	golden-winged warbler	S4	G4	NLW	SGCN	2015 SGCN, RSGCN, H threats, Stewardship
<i>Vireo bellii</i>	Bell's vireo	S4	G5	S-SPC	SGCN	2015 SGCN, M threats
<i>Vireo philadelphicus</i>	Philadelphia vireo	S2S3	G5		SGCN	2015 SGCN, H threats
<i>Megascops asio</i>	eastern screech-owl	S3	G5		SNI	

Fishes

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Acipenser fulvescens</i>	lake sturgeon	S3	G3	S-SPC	SGCN	2015 SGCN, RSGCN, RFSS, Decline, H threats
<i>Alosa chrysochloris</i>	skipjack herring	SH	G5	S-END	SGCN	2015 SGCN
<i>Ammocrypta clara</i>	western sand darter	S3	G3		SGCN	2015 SGCN, RSGCN, Decline
<i>Aphredoderus sayanus</i>	pirate perch	S3	G5	S-SPC	SGCN	2015 SGCN, Rare
<i>Catostomus catostomus</i>	longnose sucker	S4	G5		SGCN	2015 SGCN
<i>Clinostomus elongatus</i>	redside dace	S3	G3	S-SPC	SGCN	2015 SGCN, RSGCN, Decline, M threats
<i>Cottus cognatus</i>	slimy sculpin	S3	G5		SGCN	H threats
<i>Cottus ricei</i>	spoonhead sculpin	S3	G5		SGCN	2015 SGCN, RSGCN, M threats
<i>Couesius plumbeus</i>	lake chub	S3	G5	S-SPC	SGCN	2015 SGCN, M threats
<i>Crystallaria asprella</i>	crystal darter	S2	G3	S-END	SGCN	2015 SGCN, RSGCN, Decline, Rare
<i>Erimystax x-punctatus</i>	gravel chub	S2	G4	S-THR	SGCN	2015 SGCN, RSGCN Watchlist, Decline, M threats, Rare
<i>Etheostoma chlorosoma</i>	bluntnose darter	S1	G5	S-SPC	SGCN	2015 SGCN, Rare
<i>Etheostoma microperca</i>	least darter	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, Decline, H threats
<i>Fundulus sciadicus</i>	plains topminnow	S3	G4	S-THR	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Hybognathus nuchalis</i>	Mississippi silvery minnow	S3	G5	S-SPC	SGCN	2015 SGCN, Decline
<i>Hybopsis amnis</i>	pallid shiner	S1	G4	S-END	SGCN	2015 SGCN, Decline, M threats, Rare
<i>Ichthyomyzon fossor</i>	northern brook lamprey	S3	G4	S-SPC	SGCN	2015 SGCN, RFSS, Decline
<i>Ichthyomyzon gagei</i>	southern brook lamprey	S3	G5	S-SPC	SGCN	2015 SGCN
<i>Ictiobus niger</i>	black buffalo	S3	G5	S-THR	SGCN	2015 SGCN
<i>Lepomis gulosus</i>	warmouth	S3	G5	S-SPC	SGCN	2015 SGCN
<i>Lepomis peltastes</i>	northern sunfish	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, H threats

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Lythrurus umbratilis</i>	redfin shiner	S2	G5	S-SPC	SGCN	2015 SGCN, Decline
<i>Moxostoma carinatum</i>	river redhorse	S3	G4		SGCN	M threats
<i>Moxostoma duquesnei</i>	black redhorse	S3	G5	S-SPC	SGCN	2015 SGCN
<i>Myoxocephalus thompsonii</i>	deepwater sculpin	S2	G5		SGCN	2015 SGCN
<i>Nocomis biguttatus</i>	hornyhead chub	S4	G5		SGCN	2015 SGCN, Decline, M threats
<i>Notropis anogenus</i>	pugnose shiner	S3	G3	S-THR	SGCN	2015 SGCN, RSGCN, RFSS, Decline, H threats
<i>Notropis nubilus</i>	Ozark minnow	S3	G5	S-SPC	SGCN	2015 SGCN
<i>Notropis texanus</i>	weed shiner	S3	G5		SGCN	2015 SGCN, H threats
<i>Notropis topeka</i>	Topeka shiner	S3	G3	S-SPC, F-END	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Noturus exilis</i>	slender madtom	S1	G5	S-END	SGCN	2015 SGCN, M threats, Very Rare
<i>Percina evides</i>	gilt darter	S3	G4	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, M threats
<i>Phenacobius mirabilis</i>	suckermouth minnow	S3	G5	S-SPC	SGCN	2015 SGCN, Decline
<i>Platygobio gracilis</i>	flathead chub	S1	G5	S-SPC	SGCN	2015 SGCN, RSGCN, Very Rare
<i>Polyodon spathula</i>	paddlefish	S3	G4	S-THR	SGCN	2015 SGCN, Decline, M threats
<i>Salvelinus fontinalis</i>	brook trout (native strains)	S4	G5		SGCN	2015 SGCN, M threats
<i>Coregonus nipigon</i>	Nipigon cisco	SU	G4	S-SPC	SNI	
<i>Coregonus zenithicus</i>	shortjaw cisco	SU	G3	S-SPC	SNI	
<i>Macrhybopsis hyostoma</i>	shoal chub	S4	G5		SNI	
<i>Prosopium coulterii</i>	pygmy whitefish	SU	G5	S-SPC	SNI	
<i>Pungitius pungitius</i>	ninespine stickleback	S3	G5		SNI	

Mammals

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Alces alces</i>	moose	S3	G5	S-SPC	SGCN	2015 SGCN, RFSS, Decline, H threats
<i>Canis lupus</i>	gray wolf	S4	G5	F-THR	SGCN	H threats
<i>Cervus canadensis</i>	elk	S3	G5		SGCN	Decline
<i>Cryptotis parvus</i>	North American least shrew	SH	G5		SGCN	2015 SGCN, RSGCN Watchlist
<i>Eptesicus fuscus</i>	big brown bat	S2	G5		SGCN	2015 SGCN, Decline, VH threats
<i>Glaucomys sabrinus</i>	northern flying squirrel	S3	G5		SGCN	M threats
<i>Lasionycteris noctivagans</i>	silver-haired bat	S2	G3		SGCN	2015 SGCN, RSGCN, Decline, VH threats

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Lasiurus borealis</i>	eastern red bat	S2	G3		SGCN	2015 SGCN, RSGCN, RFSS, Decline, H threats
<i>Lasiurus cinereus</i>	northern hoary bat	S2	G3	NLW	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Lepus americanus</i>	snowshoe hare	S4	G5		SGCN	M threats, cultural significance
<i>Lepus townsendii</i>	white-tailed jackrabbit	S3	G5		SGCN	2015 SGCN, RSGCN, Decline, M threats
<i>Lynx canadensis</i>	Canada lynx	S3	G5	S-SPC, F-THR	SGCN	2015 SGCN, M threats
<i>Microtus chrotorrhinus</i>	rock vole	S3	G5		SGCN	Decline
<i>Microtus ochrogaster</i>	prairie vole	S2	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, H threats
<i>Microtus pinetorum</i>	woodland vole	S2	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, Very Rare
<i>Mustela nivalis</i>	least weasel	S2S3	G5	S-SPC	SGCN	2015 SGCN, Decline
<i>Myotis lucifugus</i>	little brown myotis	S2	G3	S-SPC, NLW	SGCN	2015 SGCN, RSGCN, RFSS, Decline, VH threats
<i>Myotis septentrionalis</i>	northern long-eared bat	S1	G2	S-SPC, F-END	SGCN	2015 SGCN, RSGCN, Decline, VH threats
<i>Napaeozapus insignis</i>	eastern woodland jumping mouse	S3	G5		SGCN	
<i>Onychomys leucogaster</i>	northern grasshopper mouse	S2	G5	S-SPC	SGCN	2015 SGCN, Decline
<i>Perimyotis subflavus</i>	tricolored bat	S1	G3	S-SPC, F-P-END	SGCN	2015 SGCN, RSGCN, Decline, VH threats
<i>Perognathus flavescens</i>	plains pocket mouse	S2	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, H threats
<i>Phenacomys ungava</i>	eastern heather vole	S3	G5	S-SPC	SGCN	2015 SGCN, RFSS
<i>Poliocitellus franklinii</i>	Franklin's ground squirrel	S4	G5		SGCN	2015 SGCN, RSGCN, M threats
<i>Puma concolor</i>	mountain lion	SH	G5	S-SPC	SGCN	2015 SGCN
<i>Reithrodontomys megalotis</i>	western harvest mouse	S3	G5	S-SPC	SGCN	2015 SGCN
<i>Sorex fumeus</i>	smoky shrew	S3	G5	S-SPC	SGCN	2015 SGCN
<i>Spilogale interrupta</i>	plains spotted skunk	S1	G4	S-THR	SGCN	2015 SGCN, RSGCN, Rare
<i>Synaptomys borealis</i>	northern bog lemming	S2	G5	S-SPC, NLW	SGCN	2015 SGCN, RFSS, M threats
<i>Taxidea taxus</i>	American badger	S3	G5		SGCN	2015 SGCN, M threats
<i>Thomomys talpoides</i>	northern pocket gopher	S1	G5	S-THR	SGCN	2015 SGCN, Decline, VH threats, Rare
<i>Urocitellus richardsonii</i>	Richardson's ground squirrel	S2	G5	S-SPC	SGCN	2015 SGCN, Decline, H threats
<i>Neogale frenata</i>	long-tailed weasel	S3	G5		SNI	
<i>Nycticeius humeralis</i>	North American evening bat	SU	G5		SNI	
<i>Sorex haydeni</i>	prairie shrew	SU	G5		SNI	

Reptiles

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Apalone mutica</i>	smooth softshell	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, H threats
<i>Aspidoscelis sexlineatus</i>	six-lined racerunner	S3	G5		SGCN	2015 SGCN, M threats
<i>Coluber constrictor</i>	North American racer	S3	G5	S-SPC	SGCN	2015 SGCN, M threats
<i>Crotalus horridus</i>	timber rattlesnake	S2	G4	S-THR	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Emydoidea blandingii</i>	Blanding's turtle	S2	G4	S-THR, NLW	SGCN	2015 SGCN, RSGCN, RFSS, Decline, VH threats
<i>Glyptemys insculpta</i>	wood turtle	S2	G3	S-THR, NLW	SGCN	2015 SGCN, RSGCN, RFSS, Decline, VH threats
<i>Heterodon nasicus</i>	plains hog-nosed snake	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Heterodon platirhinos</i>	eastern hog-nosed snake	S3	G5		SGCN	2015 SGCN, RSGCN Watchlist, M threats
<i>Opheodrys vernalis</i>	smooth greensnake	S3	G5		SGCN	2015 SGCN, RSGCN, M threats
<i>Pantherophis obsoletus</i>	western ratsnake	S2	G5	S-THR	SGCN	2015 SGCN, H threats, Rare
<i>Pituophis catenifer</i>	gophersnake	S2S3	G5	S-SPC	SGCN	2015 SGCN, Decline, H threats
<i>Plestiodon fasciatus</i>	common five-lined skink	S3	G5	S-SPC	SGCN	2015 SGCN, M threats
<i>Tropidoclonion lineatum</i>	lined snake	S2	G5	S-SPC	SGCN	2015 SGCN, M threats, Very Rare
<i>Graptemys ouachitensis</i>	southern map turtle	S3	G5		SNI	

Bees (Class Insecta; Order Hymenoptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Andrena accepta</i>	two-spotted miner bee	S3	GNR		SGCN	Very Rare, Specialist
<i>Andrena arabis</i>	mustard miner bee	S3	GNR		SGCN	Rare
<i>Andrena bradleyi</i>	Bradley's miner bee	S3	GNR		SGCN	
<i>Andrena brevipalpis</i>	short-tongued miner bee	S3	GNR		SGCN	Decline, Rare
<i>Andrena campanulae</i>	a miner bee	S2	GNR		SGCN	Very Rare
<i>Andrena canadensis</i>	Canada miner bee	S1	GNR		SGCN	Decline, Very Rare
<i>Andrena carolina</i>	Carolina miner bee	S3	GNR		SGCN	
<i>Andrena clarkella</i>	Clark's miner bee	S3	GNR		SGCN	
<i>Andrena distans</i>	distant miner bee	S3	GNR		SGCN	specialist
<i>Andrena erigeniae</i>	spring beauty miner bee	S3	GNR		SGCN	
<i>Andrena erythrogaster</i>	red-bellied miner bee	S3	GNR		SGCN	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Andrena erythronii</i>	trout lily miner bee	S3	GNR		SGCN	
<i>Andrena frigida</i>	cold miner bee	S3	GNR		SGCN	
<i>Andrena haynesi</i>	Haynes's miner bee	S1	GNR		SGCN	H threats, Rare
<i>Andrena helianthiformis</i>	a mining bee	S3	GNR		SGCN	M threats
<i>Andrena integra</i>	intact miner bee	S3	GNR		SGCN	
<i>Andrena lupinorum</i>	lupine miner bee	S3	G5		SGCN	
<i>Andrena mariae</i>	Maria miner bee	S3	GNR		SGCN	
<i>Andrena melanochoa</i>	rose miner bee	S3	GNR		SGCN	Rare
<i>Andrena parnassiae</i>	parnassia miner	S1	GNR		SGCN	M threats, Very Rare
<i>Andrena peckhami</i>	Peckham's miner bee	S1	GNR		SGCN	Proposed RSGCN, Rare
<i>Andrena persimulata</i>	protuberance miner bee	S3	GNR		SGCN	
<i>Andrena placata</i>	peaceful miner bee	S3	GNR		SGCN	
<i>Andrena platyparia</i>	plated miner bee	S3	GNR		SGCN	
<i>Andrena polemonii</i>	an andrenid bee	S1	GNR		SGCN	H threats, Very Rare
<i>Andrena quintilis</i>	eastern scaly miner bee	S3	GNR		SGCN	M threats, specialist
<i>Andrena sigmundi</i>	a mining bee	S3	GNR		SGCN	
<i>Andrena simplex</i>	simple miner bee	S3	GNR		SGCN	
<i>Andrena uvulariae</i>	a mining bee	S2	GNR		SGCN	Very Rare
<i>Andrena vernalis</i>	a mining bee	S3	GNR		SGCN	
<i>Andrena wellesleyana</i>	Wellesley miner bee	S3	GNR		SGCN	Rare
<i>Anthidiellum boreale</i>	a rotund resin bee	S2			SGCN	Rare
<i>Anthidium psoraleae</i>	a leafcutter bee	S2	G3		SGCN	Decline, Very Rare
<i>Bombus affinis</i>	rusty patched bumble bee	S2	G2	F-END	SGCN	2015 SGCN, RSGCN, Decline, VH threats
<i>Bombus ashtoni</i>	a bumble bee	SH	G4		SGCN	2015 SGCN, RSGCN
<i>Bombus fervidus</i>	yellow bumble bee	S3	G3		SGCN	2015 SGCN, Decline, H threats
<i>Bombus fraternus</i>	Southern Plains bumble bee	S2	G3		SGCN	RSGCN, Rare
<i>Bombus frigidus</i>	a bumble bee	S2	G5		SGCN	H threats
<i>Bombus insularis</i>	indiscriminate bumble bee	S2	G3		SGCN	Decline, M threats
<i>Bombus pensylvanicus</i>	American bumble bee	S3	G3	NLW	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Bombus terricola</i>	yellowbanded bumble bee	S2	G3		SGCN	2015 SGCN, RSGCN, RFSS, Decline, H threats
<i>Brachymelecta interrupta</i>	interrupted xeromelecta bee	S2	G4		SGCN	Decline, Very Rare
<i>Calliopsis nebraskensis</i>	a mining bee	S3	GNR		SGCN	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Chelostoma philadelphi</i>	a striped hairy belly bee	S3	GNR		SGCN	Rare
<i>Colletes aberrans</i>	aberrant cellophane bee	S2	GNR		SGCN	Proposed RSGCN Watchlist, H threats
<i>Colletes albescens</i>	white plasterer bee	S1	GNR		SGCN	Decline, Very Rare
<i>Colletes americanus</i>	a cellophane bee	S2	GNR		SGCN	H threats
<i>Colletes andrewsi</i>	Andrews' cellophane bee	S3	GNR		SGCN	
<i>Colletes brevicornis</i>	short-horned plasterer bee	S1	GNR		SGCN	M threats, Rare
<i>Colletes ciliatus</i>	a cellophane bee	S1	GNR		SGCN	M threats, Very Rare
<i>Colletes latitarsis</i>	ground-cherry plasterer bee	S3	GNR		SGCN	Rare
<i>Colletes mandibularis</i>	mandibular plasterer bee	S2	GNR		SGCN	Very Rare
<i>Colletes solidaginis</i>	goldenrod plasterer bee	S2	GNR		SGCN	H threats
<i>Colletes susannae</i>	Susan's plasterer bee	S2	GNR		SGCN	Proposed RSGCN Watchlist, H threats
<i>Colletes willistoni</i>	Williston's cellophane bee	S1	GNR		SGCN	VH threats, Very Rare
<i>Colletes wilmattae</i>	Wilmatte's plasterer bee	S1	GNR		SGCN	H threats, Very Rare
<i>Dianthidium pudicum</i>	modest pebble bee	S3	GNR		SGCN	Rare
<i>Dianthidium simile</i>	a leafcutter bee	S3	GNR		SGCN	
<i>Dieunomia heteropoda</i>	sunflower sweat bee	S2	G5		SGCN	H threats
<i>Dufourea harveyi</i>	a solitary bee	S2	GNR		SGCN	Very Rare
<i>Dufourea marginata</i>	glowing sweat bee	S2	GNR		SGCN	Rare
<i>Dufourea novaeangliae</i>	New England sweat bee	S3	GNR		SGCN	Rare
<i>Epeoloides pilosulus</i>	macropis cuckoo bee	S1	G3		SGCN	M threats, Very Rare
<i>Epeolus ainsliei</i>	a cuckoo bee	S2	GNR		SGCN	RSGCN, M threats, Very Rare
<i>Epeolus lectoides</i>	collector cuckoo nomad bee	S2	GNR		SGCN	Very Rare
<i>Eucera albata</i>	an anthophorid bee	S3	GNR		SGCN	
<i>Heriades leavitti</i>	a resin bee	S3	G5		SGCN	
<i>Hylaeus sparsus</i>	a masked bee	S2	GNR		SGCN	Very Rare
<i>Lasioglossum fedorense</i>	a sweat bee	S1	GNR		SGCN	M threats, Very Rare
<i>Lasioglossum floridanum</i>	a sweat bee	S2	GNR		SGCN	H threats, Rare
<i>Lasioglossum lusorium</i>	a sweat bee	S2	GNR		SGCN	Very Rare
<i>Lasioglossum nelumbonis</i>	water-lily sweat bee	S3	GNR		SGCN	
<i>Lasioglossum swenki</i>	Swenk's sweat bee	S1	GNR		SGCN	H threats, Rare
<i>Lasioglossum texanum</i>	a sweat bee	S2	GNR		SGCN	Very Rare
<i>Macropis nuda</i>	common loosestrife oil bee	S2	GNR		SGCN	Proposed RSGCN, H threats

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Macropis patellata</i>	patellar oil-collecting bee	S1	GNR		SGCN	H threats, Very Rare
<i>Macropis steironematis</i>	an oil-collecting bee	S1	GNR		SGCN	RSGCN, H threats, Rare
<i>Megachile addenda</i>	a leafcutter bee	S3	G5		SGCN	Rare
<i>Megachile circumcincta</i>	a leafcutter bee	S2	G5		SGCN	Very Rare
<i>Megachile fortis</i>	a leafcutter bee	S2	G2		SGCN	Decline, Very Rare
<i>Megachile rugifrons</i>	a leafcutter bee	S2	G2		SGCN	Very Rare
<i>Melissodes apicatus</i>	a long-horned bee	S2	GNR		SGCN	Very Rare
<i>Melissodes bidentis</i>	two-toothed long-horned bee	S2	GNR		SGCN	Very Rare
<i>Melissodes boltoniae</i>	a callirhoe bee	S2	GNR		SGCN	Very Rare
<i>Melissodes confusus</i>	a callirhoe bee	S2	GNR		SGCN	Very Rare
<i>Melissodes denticulatus</i>	a callirhoe bee	S3	GNR		SGCN	
<i>Melissodes dentiventris</i>	a callirhoe bee	S3	GNR		SGCN	Rare
<i>Melissodes illatus</i>	a callirhoe bee	S3	G5		SGCN	
<i>Melissodes menuachus</i>	a callirhoe bee	S3	GNR		SGCN	Rare
<i>Melissodes subagilis</i>	a callirhoe bee	S2	GNR		SGCN	Rare
<i>Melissodes wheeleri</i>	a callirhoe bee	S3	GNR		SGCN	
<i>Neolarra vigilans</i>	a cuckoo bee	S1	GNR		SGCN	H threats, Very Rare
<i>Nomia universitatis</i>	a nomia bee	S3	G3		SGCN	Decline
<i>Osmia cyaneonitens</i>	a mason bee	S3	G3		SGCN	Rare
<i>Osmia distincta</i>	distinct mason bee	S3	G5		SGCN	
<i>Osmia felti</i>	a mason bee	S2	G3		SGCN	H threats
<i>Osmia inspergens</i>	shiny-faced mason bee	S1	G5		SGCN	H threats, Rare
<i>Osmia laticeps</i>	a mason bee	S2	G4		SGCN	M threats, Rare
<i>Osmia nearctica</i>	a mason bee	S2	GU		SGCN	M threats, Rare
<i>Osmia nigriventris</i>	large black-bellied mason bee	S2	G5		SGCN	Very Rare
<i>Osmia virga</i>	a mason bee	S3	G5		SGCN	
<i>Perdita albipennis</i>	shiny green miner bee	S2	GNR		SGCN	M threats, Rare
<i>Perdita bruneri</i>	Bruner's miner bee	S2	GNR		SGCN	Very Rare
<i>Perdita dolichocephala</i>	a fairy bee	S2	GNR		SGCN	Very Rare
<i>Perdita gerhardi</i>	gerhard's perdita	S2	GNR		SGCN	Very Rare
<i>Perdita halictoides</i>	sweatbee-like miner bee	S3	GNR		SGCN	
<i>Perdita ignota</i>	a miner bee	S3	GNR		SGCN	Rare

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Perdita pallidipennis</i>	a tiny bee	S3	GNR		SGCN	
<i>Perdita perpallida</i>	pale miner bee	S3	GNR		SGCN	
<i>Protandrena aestivalis</i>	summer miner bee	S2	G5		SGCN	Very Rare
<i>Protandrena andrenoides</i>	small black miner bee	S3	GNR		SGCN	Rare
<i>Protandrena renimaculata</i>	medium dark bee	S2	GNR		SGCN	Very Rare
<i>Protandrena rudbeckiae</i>	rudbeckia miner bee	S3	GNR		SGCN	Rare
<i>Protandrena simulans</i>	a mining bee	S3	GNR		SGCN	
<i>Sphecodes banksii</i>	a sweat bee	S1	GNR		SGCN	Very Rare
<i>Stelis permaculata</i>	sixteen-spotted cuckoo carder bee	S2	GNR		SGCN	Very Rare
<i>Triepeolus concavus</i>	concave cuckoo nomad bee	S2	GNR		SGCN	Very Rare
<i>Andrena aff. nigrihirta</i>	a miner bee	SU	G5		SNI	
<i>Andrena algida</i>	icy miner bee	SU	G5		SNI	
<i>Andrena aliciae</i>	yellow-faced miner bee	SU	GNR		SNI	
<i>Andrena andrenoides</i>	colorful willow miner bee	SU	GNR		SNI	
<i>Andrena bisalicis</i>	eastern willow miner bee	SU	GNR		SNI	
<i>Andrena colletina</i>	a miner bee	SU	GNR		SNI	
<i>Andrena cragini</i>	a mining bee	SU	GNR		SNI	
<i>Andrena fragilis</i>	fragile miner bee	SU	GNR		SNI	
<i>Andrena illinoiensis</i>	tufted miner bee	SU	GNR		SNI	
<i>Andrena krigiana</i>	an andrenid bee	SU	GNR		SNI	
<i>Andrena macoupinensis</i>	Macoupin County miner bee	SU	GNR		SNI	
<i>Andrena nigrae</i>	black miner bee	SU	GNR		SNI	
<i>Andrena nigrihirta</i>	black-haired miner bee	SU	G5		SNI	
<i>Andrena robertsonii</i>	Robertson's miner bee	SU	G5		SNI	
<i>Andrena robervalensis</i>	an andrenid bee	SU	GNR		SNI	
<i>Andrena runcinatae</i>	planed miner bee	SU	G5		SNI	
<i>Andrena salictaria</i>	small willow miner bee	SU	GNR		SNI	
<i>Anthidium tenuiflorae</i>	scurfy pea carder bee	SU	G5		SNI	
<i>Ashmeadiella buconis</i>	big-cheeked mason bee	SU	GNR		SNI	
<i>Calliopsis coloradensis</i>	Colorado miner bee	SU	GNR		SNI	
<i>Coelioxys alternatus</i>	a leafcutter bee	SU	GNR		SNI	
<i>Coelioxys banksi</i>	a leafcutter bee	SU	GNR		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Coelioxys bisoncornuus</i>	a leafcutter bee	SU	GNR		SNI	
<i>Coelioxys immaculatus</i>	a leafcutter bee	SU	GNR		SNI	
<i>Coelioxys modestus</i>	a leafcutter bee	SU	GNR		SNI	
<i>Coelioxys moestus</i>	sad cuckoo leafcutter bee	SU	GNR		SNI	
<i>Coelioxys octodentatus</i>	a leafcutter bee	SU	GNR		SNI	
<i>Coelioxys porterae</i>	a leafcutter bee	SU	GNR		SNI	
<i>Coelioxys sayi</i>	a leafcutter bee	SU	GNR		SNI	
<i>Coelioxys sodalis</i>	comrade cuckoo leafcutter bee	SU	GNR		SNI	
<i>Colletes compactus</i>	compact plasterer bee	SU	G5		SNI	
<i>Colletes consors</i>	partner plasterer bee	SU	G5		SNI	
<i>Colletes hyalinus</i>	translucent plasterer bee	SU	G5		SNI	
<i>Colletes impunctatus</i>	shiny plasterer bee	SU	G5		SNI	
<i>Colletes nudus</i>	nude plasterer bee	SU	GNR		SNI	
<i>Colletes phaceliae</i>	scorpionweed plasterer bee	SU	G5		SNI	
<i>Colletes rufocinctus</i>	red-belted plasterer bee	SU	GNR		SNI	
<i>Colletes speculiferus</i>	a cellophane bee	SU	GNR		SNI	
<i>Colletes validus</i>	long-faced plasterer bee	SU	GNR		SNI	
<i>Dieunomia triangulifera</i>	a dieunomia bee	SU	G3		SNI	
<i>Dufourea maura</i>	a solitary bee	SU	GNR		SNI	
<i>Epeolus americanus</i>	American cuckoo nomad bee	SU	GNR		SNI	
<i>Epeolus autumnalis</i>	orange cuckoo nomad bee	SU	GNR		SNI	
<i>Epeolus interruptus</i>	a cuckoo bee	SU	GNR		SNI	
<i>Epeolus lectus</i>	Great Plains cuckoo nomad bee	SU	GNR		SNI	
<i>Epeolus pusillus</i>	a cuckoo bee	SU	GNR		SNI	
<i>Eucera aragalli</i>	a long-horned bee	SU	GNR		SNI	
<i>Eucera dubitata</i>	a long-horned bee	SU	GNR		SNI	
<i>Eucera lepida</i>	a long-horned bee	SU	GNR		SNI	
<i>Lasioglossum hartii</i>	a sweat bee	SU	GNR		SNI	
<i>Lasioglossum pectinatum</i>	a sweat bee	SU	GNR		SNI	
<i>Lasioglossum rufulipes</i>	a sweat bee	SU	G5		SNI	
<i>Megachile dakotensis</i>	a leafcutter bee	SU	G2		SNI	
<i>Megachile frugalis</i>	a leafcutter bee	SU	G4		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Megachile inimica</i>	a leafcutter bee	SU	G5		SNI	
<i>Megachile parallela</i>	parallel leaf-cutter bee	SU	G5		SNI	
<i>Megachile petulans</i>	a leafcutter bee	SU	G5		SNI	
<i>Nomada aquilarum</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada banksi</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada bella</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada besseyi</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada depressa</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada erigeronis</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada fervida</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada gracilis</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada graenicheri</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada illinoensis</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada imbricata</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada integerrima</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada lehighensis</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada lepida</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada louisianae</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada obliterateda</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada ovata</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada perplexa</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada rubi</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada rubicunda</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada sayi</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada superba</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada tiftonensis</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada vegana</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada vicina</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomada xanthura</i>	a cuckoo bee	SU	GNR		SNI	
<i>Nomia nortoni</i>	a nomia bee	SU	G5		SNI	
<i>Osmia conjuncta</i>	eastern snail shell mason bee	SU	G5		SNI	
<i>Osmia georgica</i>	a mason bee	SU	G5		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Osmia subarctica</i>	a mason bee	SU	G3		SNI	
<i>Perdita fallax</i>	a miner bee	SU	GNR		SNI	
<i>Perdita maculigera</i>	black-spotted miner bee	SU	GNR		SNI	
<i>Perdita octomaculata</i>	eight-spotted miner bee	SU	GNR		SNI	
<i>Protandrena labrosa</i>	a mining bee	SU	GNR		SNI	
<i>Protandrena parva</i>	small miner bee	SU	GNR		SNI	
<i>Protandrena pauper</i>	poor miner bee	SU	GNR		SNI	
<i>Sphecodes antennariae</i>	a cuckoo bee	SU	GNR		SNI	
<i>Sphecodes atlantis</i>	Atlantic cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes clematidis</i>	clematis cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes coronus</i>	crowned cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes cressonii</i>	Cresson's cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes fattigi</i>	a sweat bee	SU	GNR		SNI	
<i>Sphecodes galerus</i>	a cuckoo bee	SU	GNR		SNI	
<i>Sphecodes hydrangeae</i>	a cuckoo bee	SU	GNR		SNI	
<i>Sphecodes illinoensis</i>	Illinois cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes johnsonii</i>	a cuckoo bee	SU	GNR		SNI	
<i>Sphecodes levis</i>	nimble cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes minor</i>	minor cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes nigricorpus</i>	a cuckoo bee	SU	GNR		SNI	
<i>Sphecodes pecosensis</i>	a cuckoo bee	SU	GNR		SNI	
<i>Sphecodes pimpinellae</i>	eastern cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes prosphorus</i>	black-tipped cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes smilacinae</i>	a sweat bee	SU	GNR		SNI	
<i>Sphecodes solonis</i>	shiny-faced cuckoo sweat bee	SU	GNR		SNI	
<i>Sphecodes townesi</i>	a cuckoo sweat bee	SU	GNR		SNI	
<i>Stelis foederalis</i>	federal cuckoo carder bee	SU	GNR		SNI	
<i>Stelis labiata</i>	kissing cuckoo carder bee	SU	GNR		SNI	
<i>Stelis nitida</i>	healthy cuckoo carder bee	SU	GNR		SNI	
<i>Stelis subemarginata</i>	a cuckoo bee	SU	GNR		SNI	
<i>Svastra atripes</i>	a long-horned bee	SU	GNR		SNI	
<i>Trachusa zebrata</i>	a leafcutter bee	SU	GNR		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Triepeolus distinctus</i>	a cuckoo bee	SU	GNR		SNI	
<i>Triepeolus eliseae</i>	a cuckoo bee	SU	GNR		SNI	
<i>Triepeolus helianthi</i>	sunflower cuckoo nomad bee	SU	GNR		SNI	
<i>Triepeolus occidentalis</i>	a cuckoo bee	SU	GNR		SNI	
<i>Triepeolus rhododontus</i>	red-lipped cuckoo nomad bee	SU	GNR		SNI	
<i>Triepeolus subalpinus</i>	subalpine cuckoo nomad bee	SU	GNR		SNI	
<i>Triepeolus tanneri</i>	a cuckoo bee	SU	GNR		SNI	

Beetles (Class Insecta; Order Coleoptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Adalia bipunctata</i>	two-spotted lady beetle	S2	G5		SGCN	Decline, M threats, Rare
<i>Cicindela denikei</i>	Laurentian tiger beetle	S3	G3		SGCN	2015 SGCN, M threats
<i>Cicindela fulgida</i>	crimson saltflat tiger beetle	S1	G5		SGCN	M threats, Very Rare
<i>Cicindela hirticollis rhodensis</i>	hairy-necked tiger beetle	S1	T4		SGCN	2015 SGCN, M threats, Very Rare
<i>Cicindela lepida</i>	ghost tiger beetle	S1	G3		SGCN	2015 SGCN, Decline, M threats, Very Rare
<i>Cicindela limbata</i>	sandy tiger beetle	SH	G5		SGCN	Very Rare
<i>Cicindela macra</i>	sandy stream tiger beetle	S2	G5		SGCN	M threats, Rare
<i>Cicindela patruela</i>	northern barrens tiger beetle	S3	G3		SGCN	Decline, M threats
<i>Cicindela splendida</i>	splendid tiger beetle	S3	G5		SGCN	Rare
<i>Coccinella novemnotata</i>	nine-spotted lady beetle	S1	G5		SGCN	Decline, H threats, Rare
<i>Coccinella transversoguttata</i>	transverse lady beetle	S1	G5		SGCN	M threats, Very Rare
<i>Coelambus sylvanus</i>	sylvan hygrotus diving beetle	S2	GU		SGCN	Very Rare
<i>Cicindela cursitans</i>	ant-like tiger beetle	SU	G4		SNI	
<i>Cicindela hirticollis</i>	beach-dune tiger beetle	S3	G5		SNI	
<i>Cicindela terricola</i>	variable tiger beetle	SU	G5		SNI	
<i>Lytta nuttali</i>	Nuttall's blister beetle	S3	GNR		SNI	
<i>Nicrophorus guttula</i>	yellow-bellied burying beetle	SU	GNR		SNI	
<i>Nicrophorus hybridus</i>	hybrid burying beetle	SU	G5		SNI	
<i>Nicrophorus investigator</i>	banded sexton beetle	SU	G5		SNI	
<i>Nicrophorus obscurus</i>	obscure burying beetle	SU	G5		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Oberea flavipes</i>	a longhorned beetle	SU	GNR		SNI	
<i>Pyrrhota angulata</i>	Say's firefly	SU	G5		SNI	

Butterflies, Moths (Class Insecta; Order Lepidoptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Agonopterix pergandeella</i>		S2	GNR		SGCN	Proposed RSGCN, Rare
<i>Ambesa laetella</i>		S2	GNR		SGCN	Very Rare
<i>Ancylis semiovana</i>		S2	GNR		SGCN	RSGCN, Very Rare
<i>Anisota manitobensis</i>	Manitoba oakworm moth	SH	G2		SGCN	
<i>Apamea commoda</i>		S3	G5		SGCN	Rare, Specialist
<i>Apamea niveivenosa</i>		S3	G5		SGCN	M threats
<i>Apantesis doris</i>		S2	G4		SGCN	Very Rare
<i>Apantesis margo</i>		S2	GNR		SGCN	M threats, Very Rare
<i>Apodrepanulatrix liberaria</i>		S2	G3		SGCN	RSGCN, Decline, Rare
<i>Argynnis idalia</i>	regal fritillary	S2	G3		SGCN	2015 SGCN, Decline
<i>Aspitates aberrata</i>		SH	G5		SGCN	2015 SGCN
<i>Atrytone arogos</i>	arogos skipper	SH	G2		SGCN	RSGCN
<i>Atrytonopsis hianna</i>	dusted skipper	S2	G4		SGCN	2015 SGCN, RSGCN, Decline
<i>Bagisara rectifascia</i>	straight lined mallow moth	S2	G4		SGCN	Very Rare
<i>Bandera binotella</i>		S3	GNR		SGCN	Rare
<i>Boloria chariclea</i>	Arctic fritillary	S3	G5		SGCN	2015 SGCN
<i>Boloria freija</i>	Freija fritillary	S3	G5		SGCN	H threats
<i>Boloria frigga</i>	Frigga fritillary	S3	G5		SGCN	
<i>Cabera quadrifasciaria</i>		S3	GNR		SGCN	M threats
<i>Capis archaia</i>		S2	GNR		SGCN	Rare
<i>Carmenta anthracipennis</i>		S3	GNR		SGCN	2015 SGCN
<i>Carsia sororiata</i>	Manchester treble-bar moth	S3	G5		SGCN	M threats
<i>Catocala abbreviatella</i>	abbreviated underwing	S3	G3		SGCN	2015 SGCN, RSGCN, Decline
<i>Catocala amestris</i>	three-staff underwing	S3	G4		SGCN	M threats
<i>Catocala nuptialis</i>	married underwing	S3	G3		SGCN	Rare

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Catocala whitneyi</i>	Whitney's underwing	S3	G2		SGCN	2015 SGCN, RSGCN, Decline
<i>Chaetagnlaea cerata</i>		S3	G3		SGCN	
<i>Chlosyne gorgone</i>	gorgone checkerspot	S3	G5		SGCN	Decline
<i>Clepsia illustrana</i>		S2	GNR		SGCN	M threats, Rare
<i>Crambus awemellus</i>		S3	GNR		SGCN	Specialist
<i>Danaus plexippus</i>	monarch	S5	G4	F-P-THR	SGCN	2015 SGCN, RSGCN, RFSS, Decline
<i>Dargida rubripennis</i>	a noctuid moth	S3	G3		SGCN	
<i>Eoreuma crawfordi</i>		S3	GNR		SGCN	M threats
<i>Erastria coloraria</i>		S2	G3		SGCN	Decline, Rare
<i>Erebia discoidalis</i>	red-disked alpine	S3	G5		SGCN	Decline
<i>Erebia mancinus</i>	taiga alpine	S3	G5		SGCN	Endemic, 2015 SGCN, RSGCN Watchlist, RFSS
<i>Erynnis lucilius</i>	columbine duskywing	S3	G3		SGCN	RSGCN Watchlist, Decline
<i>Erynnis martialis</i>	mottled duskywing	SU	G3		SGCN	2015 SGCN, RSGCN, Decline
<i>Erynnis persius</i>	Persius duskywing	S1	G5		SGCN	Decline, Very Rare
<i>Euchloe ausonides</i>	large marble	S3	G5	F-P-THR	SGCN	2015 SGCN, Decline
<i>Euchloe olympia</i>	Olympia marble	S3	G5		SGCN	Decline, M threats
<i>Eucosma giganteana</i>	giant eucosma moth	S3	GNR		SGCN	M threats
<i>Eucosma lapidana</i>		S2	GNR		SGCN	Very Rare
<i>Eucosma millerana</i>	a tortricid moth	S2	GNR		SGCN	M threats, Rare
<i>Eucosma pallidicostana</i>		S3	GNR		SGCN	Rare, Specialist
<i>Euphydryas phaeton</i>	Baltimore checkerspot	S3	G4		SGCN	Decline
<i>Euphyes conspicua</i>	black dash	S3	G4		SGCN	Decline, M threats
<i>Euxoa niveilinea</i>	a dart moth	S3	GNR		SGCN	Decline, Rare, Specialist
<i>Euxoa quadridentata</i>		S2	GNR		SGCN	Very Rare
<i>Euxoa quebecensis</i>		S2	G3		SGCN	Very Rare
<i>Feltia manifesta</i>		S2	G4		SGCN	Very Rare
<i>Habrosyne gloriosa</i>		S3	GNR		SGCN	
<i>Heliothis borealis</i>		S3	G4		SGCN	Specialist
<i>Hemaris gracilis</i>		S3	G3		SGCN	Decline, Rare
<i>Hesperia assiniboia</i>	Assiniboia skipper	SH	G5	S-END	SGCN	2015 SGCN
<i>Hesperia dacotae</i>	Dakota skipper	S1	G2	S-END, F-THR	SGCN	2015 SGCN, RSGCN, Decline, Rare
<i>Hesperia leonardus leonardus</i>	Leonard's skipper	S2	T4	S-SPC	SGCN	Decline

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Hesperia leonardus pawnee</i>	Pawnee skipper	S1	T4	S-SPC	SGCN	Decline, Rare
<i>Hesperia ottoe</i>	Ottoe skipper	SH	G3	S-END	SGCN	2015 SGCN, RSGCN
<i>Hystriophora taleana</i>		S3	GNR		SGCN	Rare, Specialist
<i>Interjectio niviella</i>		S3	GNR		SGCN	Specialist
<i>Lasionycta secedens</i>		S3	G5		SGCN	2015 SGCN
<i>Lasionycta taigata</i>	a noctuid moth	S3	G4		SGCN	2015 SGCN
<i>Lithophane franclemonti</i>		S3	G3		SGCN	Proposed RSGCN, Rare
<i>Lithophane lemmeri</i>		SU	G3		SGCN	Very Rare, Specialist
<i>Lithophane scottae</i>		S3	G3		SGCN	Decline
<i>Lobocleta plemyraria</i>		S3	G3		SGCN	Rare
<i>Melaporphyria immortua</i>		S2	G1		SGCN	Endemic, Proposed RSGCN, Decline, Very Rare
<i>Melitara dentata</i>		S1	GNR		SGCN	M threats, Very Rare
<i>Meroptera mirandella</i>		S2	GNR		SGCN	Very Rare
<i>Microtheoris ophionalis</i>		S2	GNR		SGCN	Very Rare
<i>Mimoschinia rufofascialis</i>		S3	GNR		SGCN	Rare, Specialist
<i>Oarisma garita</i>	Garita skipperling	SH	G5	S-THR	SGCN	2015 SGCN
<i>Oarisma poweshiek</i>	Poweshiek skipperling	SH	G1	S-END	SGCN	2015 SGCN, RSGCN
<i>Oeneis macounii</i>	Macoun's arctic	S3	G5		SGCN	
<i>Oligia bridghamii</i>		S3	G5		SGCN	Rare, Specialist
<i>Oligia egens</i>		S3	GNR		SGCN	Rare, Specialist
<i>Papaipema aweme</i>		S3	G3		SGCN	RSGCN, Rare
<i>Papaipema beeriana</i>	blazing star stem borer	S2	G2		SGCN	RSGCN, Very Rare
<i>Papaipema marginidens</i>		S3	G4		SGCN	Rare
<i>Papaipema nepheleptena</i>		S3	G4		SGCN	RSGCN Watchlist, Rare
<i>Papaipema sciata</i>	Culver's root borer	S3	G3		SGCN	RSGCN, Rare
<i>Pelochrista heathiana</i>		S3	GNR		SGCN	Rare, Specialist
<i>Pelochrista nandana</i>		S3	GNR		SGCN	Rare, Specialist
<i>Pelochrista robinsonana</i>		S3	GNR		SGCN	Rare, Specialist
<i>Pelochrista scintillana</i>		S3	GNR		SGCN	Decline, Specialist
<i>Phoberia ingenua</i>		S2	G3		SGCN	Very Rare
<i>Plagodis kuetzingi</i>		S3	G5		SGCN	
<i>Plebejus idas</i>	northern blue	S2	G5	S-SPC	SGCN	Decline

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Plebejus samuelis</i>	Karner blue	SH	G1	S-END, F-END	SGCN	2015 SGCN, RSGCN, Decline
<i>Poanes massasoit</i>	mulberry wing	S3	G4		SGCN	Decline
<i>Polia propodea</i>		S3	GNR		SGCN	Specialist
<i>Polites origenes</i>	crossline skipper	S2	G5		SGCN	Decline
<i>Ponometia binocula</i>	prairie bird-dropping moth	S3	GNR		SGCN	Rare, Specialist
<i>Proserpinus juanita</i>		S2	G4		SGCN	2015 SGCN, Decline, Very Rare
<i>Protoschinia nuchalis</i>		S3	G5		SGCN	Decline, Rare
<i>Pyrausta pythialis</i>		S3	GNR		SGCN	Endemic, Proposed RSGCN, M threats
<i>Pyrausta scurralis</i>		S3	GNR		SGCN	Rare, Specialist
<i>Pyrausta socialis</i>	sociable pyrausta moth	S3	GNR		SGCN	Rare, Specialist
<i>Schinia bina</i>	bina flower moth	S2	G4		SGCN	RSGCN, Very Rare
<i>Schinia grandimedia</i>	Rockies boneset flower moth	S3	GNR		SGCN	Rare, Specialist
<i>Schinia hulstia</i>		S3	GNR		SGCN	Rare, Specialist
<i>Schinia indiana</i>	phlox moth	S1	G3	S-SPC	SGCN	2015 SGCN, RSGCN, Decline, M threats, Rare
<i>Schinia jaguarina</i>	jaguar flower moth	S3	G4		SGCN	Decline, Rare, Specialist
<i>Schinia lucens</i>	leadplant flower moth	S3	G4	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, M threats
<i>Schinia lynx</i>		S3	G5		SGCN	Rare
<i>Schinia mortua</i>		S3	G4		SGCN	Rare, Specialist
<i>Schinia obscurata</i>		S3	G4		SGCN	Rare, Specialist
<i>Schinia septentrionalis</i>	northern flower moth	S3	G3		SGCN	M threats
<i>Schizura apicalis</i>		S3	G3		SGCN	Rare
<i>Sparganothis violaceana</i>	a tortricid moth	S3	GNR		SGCN	Specialist
<i>Stiria rugifrons</i>	yellow sunflower moth	S3	G4		SGCN	Rare, Specialist
<i>Sympistis piffardi</i>		S3	G4		SGCN	Rare, Specialist
<i>Sympistis stabilis</i>		S3	G5		SGCN	Rare, Specialist
<i>Sympistis viriditincta</i>		S2	GNR		SGCN	Very Rare
<i>Syngrapha montana</i>		S3	G4		SGCN	Rare, Specialist
<i>Tacparia atropunctata</i>		S3	GNR		SGCN	Rare, Specialist
<i>Thaumtopsopsis pectinifer</i>	a crambid snout moth	S2	GNR		SGCN	Rare
<i>Walshia amorphella</i>		S3	GNR		SGCN	Rare, Specialist
<i>Xestia imperita</i>		S2	G5		SGCN	M threats, Very Rare
<i>Xestia mixta</i>		S3	GNR		SGCN	2015 SGCN

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Zenophleps lignicolorata</i>		S2	G5		SGCN	M threats, Rare
<i>Abagrotis cupida</i>	cupid dart moth	S3	G4		SNI	
<i>Abagrotis placida</i>		SU	G5		SNI	
<i>Abrostola ovalis</i>		SU	G4		SNI	
<i>Acasis viridata</i>		SU	G5		SNI	
<i>Acleris arcticana</i>		SU	GNR		SNI	
<i>Acleris clarkei</i>		SU	GNR		SNI	
<i>Acleris effractana</i>	hook-winged tortrix moth	SU	GNR		SNI	
<i>Acleris fragariana</i>		SU	GNR		SNI	
<i>Acleris hudsoniana</i>		S3	GNR		SNI	
<i>Acleris inana</i>		SU	GNR		SNI	
<i>Acleris minuta</i>		SU	GNR		SNI	
<i>Acleris ptychogrammos</i>		S3	GNR		SNI	
<i>Acleris simpliciana</i>		SU	GNR		SNI	
<i>Acleris youngana</i>		SU	GNR		SNI	
<i>Acossus populi</i>	aspen carpenterworm moth	SU	GNR		SNI	
<i>Acrocercops pnosmodiella</i>		SU	GNR		SNI	
<i>Acronicta betulae</i>	birch dagger moth	SU	GNR		SNI	
<i>Acronicta falcata</i>		SU	G3		SNI	
<i>Acronicta lanceolaria</i>		SU	G4		SNI	
<i>Acronicta longa</i>		SU	G5		SNI	
<i>Acronicta radcliffei</i>		S3	G5		SNI	
<i>Acronicta sperata</i>		SU	G5		SNI	
<i>Actebia fennica</i>		S3	G5		SNI	
<i>Aethes baloghi</i>		S3	GNR		SNI	
<i>Aethes louisiana</i>		SU	GNR		SNI	
<i>Aethes monera</i>		SU	GNR		SNI	
<i>Aethes ringsi</i>		SU	G3		SNI	
<i>Aethes spartinana</i>		SU	GNR		SNI	
<i>Agonopterix gelidella</i>		SU	GNR		SNI	
<i>Agonopterix lythrella</i>		SU	GNR		SNI	
<i>Agrotis stigmata</i>		SU	G4		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Amphipyra tragopoginis</i>		SU	G5		SNI	
<i>Anathix aggressa</i>		SU	GNR		SNI	
<i>Ancylis albacostana</i>		SU	GNR		SNI	
<i>Anhimella contrahens</i>		SU	G4		SNI	
<i>Antepione thisoaria</i>		S3	G5		SNI	
<i>Apamea burgessi</i>		SU	G4		SNI	
<i>Apamea cariosa</i>	a noctuid moth	SU	G4		SNI	
<i>Apamea impulsu</i>		S3	G5		SNI	
<i>Apamea indocilis</i>		SU	G5		SNI	
<i>Apamea lutosu</i>		SU	G4		SNI	
<i>Apamea nigrior</i>		SU	G5		SNI	
<i>Apamea relicina</i>		SU	G4		SNI	
<i>Apantesis anna</i>	Anna tiger moth	SU	G4		SNI	
<i>Apantesis arge</i>	arge moth	S3	G5		SNI	
<i>Apantesis phyllira</i>	phyllira tiger moth	S3	G4		SNI	
<i>Apantesis speciosa</i>		S3	G5		SNI	
<i>Apantesis williamsii</i>	William's tiger moth	S3	G5		SNI	
<i>Apotomops wellingtoniana</i>		SU	GNR		SNI	
<i>Archips packardiana</i>		S3	GNR		SNI	
<i>Arctia plantaginis</i>		SU	G5		SNI	
<i>Arequipa turbatella</i>		SU	GNR		SNI	
<i>Argyresthia freyella</i>		SU	GNR		SNI	
<i>Argyria rufesignella</i>		SU	GNR		SNI	
<i>Atascosa glareosella</i>		S3	GNR		SNI	
<i>Autographa flagellum</i>		SU	G5		SNI	
<i>Autographa pseudogamma</i>		SU	G5		SNI	
<i>Autographa rubidus</i>		S3	G5		SNI	
<i>Bactra lancealana</i>		SU	GNR		SNI	
<i>Bactra maiorina</i>		SU	GNR		SNI	
<i>Batrachedra praeangusta</i>		SU	GNR		SNI	
<i>Bellura brehmei</i>		SU	GNR		SNI	
<i>Boloria eunomia</i>	bog fritillary	S4	G5		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Bondia comonana</i>		SU	GNR		SNI	
<i>Brachionycta borealis</i>		SU	G4		SNI	
<i>Brachylomia discinigra</i>		S3	GNR		SNI	
<i>Brenthia pavonacella</i>		SU	GNR		SNI	
<i>Brymbia quadrimaculella</i>		SU	GNR		SNI	
<i>Callophrys eryphon</i>	western pine elfin	S3	G5		SNI	
<i>Caradrina meralis</i>	sand quaker moth	SU	G5		SNI	
<i>Caradrina montana</i>		SU	G5		SNI	
<i>Catocala alabamae</i>		SU	G4		SNI	
<i>Catocala antinympha</i>	sweetfern underwing	S3	G5		SNI	
<i>Catocala clintonii</i>	Clinton's underwing	SU	G4		SNI	
<i>Catocala crataegi</i>	hawthorn underwing	SU	G5		SNI	
<i>Catocala epione</i>	Epione underwing	SU	G5		SNI	
<i>Catocala judith</i>	Judith' s underwing	SU	G5		SNI	
<i>Catocala mira</i>	wonderful underwing	SU	G5		SNI	
<i>Catocala obscura</i>	obscure underwing	SU	G5		SNI	
<i>Catocala residua</i>		SU	G5		SNI	
<i>Catocala resecta</i>		SU	G5		SNI	
<i>Catocala subnata</i>	youthful underwing	SU	G5		SNI	
<i>Catocala umbrosa</i>	an underwing moth	SU	G5		SNI	
<i>Catocala vidua</i>	widow underwing	SU	G5		SNI	
<i>Cerma cora</i>		S3	G3		SNI	
<i>Chersotis juncta</i>		SU	G4		SNI	
<i>Chrysanympa formosa</i>		SU	G5		SNI	
<i>Cissusa spadix</i>		SU	G4		SNI	
<i>Clepsia penetralis</i>		SU	GNR		SNI	
<i>Coenochroa bipunctella</i>		SU	GNR		SNI	
<i>Coleophora maritella</i>		SU	GNR		SNI	
<i>Crambidia pura</i>		SU	G4		SNI	
<i>Crambus hamella</i>		SU	GNR		SNI	
<i>Crambus laqueatellus</i>		SU	GNR		SNI	
<i>Crambus whitmerellus</i>		SU	GNR		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Cucullia omissa</i>		SU	GNR		SNI	
<i>Cydia fletcherana</i>		SU	GNR		SNI	
<i>Cydia ingrata</i>		SU	GNR		SNI	
<i>Decodes macdunnoughi</i>		SU	GNR		SNI	
<i>Depressaria alienella</i>		SU	GNR		SNI	
<i>Diarsia dislocata</i>		SU	G5		SNI	
<i>Dichagyris grotei</i>		SU	G4		SNI	
<i>Digrammia eremiata</i>	a moth	SU	G4		SNI	
<i>Digrammia neptaria</i>		SU	G5		SNI	
<i>Ecliptopera silaceata</i>		SU	G5		SNI	
<i>Enchrysa dissectella</i>		SU	GNR		SNI	
<i>Epiblema minutana</i>		S3	GNR		SNI	
<i>Epinotia cruciana</i>		SU	GNR		SNI	
<i>Epinotia myricana</i>		SU	GNR		SNI	
<i>Epirrhoe sperryi</i>		SU	G5		SNI	
<i>Erynnis brizo</i>	sleepy duskywing	SU	G5		SNI	
<i>Ethmia longimaculella</i>		S3	GNR		SNI	
<i>Ethmia monticola</i>		S3	GNR		SNI	
<i>Ethmia zelleriella</i>		SU	GNR		SNI	
<i>Euchromius ocella</i>		SU	GNR		SNI	
<i>Euclementia bassettella</i>		SU	GNR		SNI	
<i>Eucoptocnemis fimbriaris</i>	fringed dart moth	SU	G4		SNI	
<i>Eucosma autumnana</i>		S3	GNR		SNI	
<i>Eucosma clavana</i>		SU	GNR		SNI	
<i>Eucosma dorsiatomana</i>		SU	GNR		SNI	
<i>Eucosma indagatricana</i>		SU	G5		SNI	
<i>Eucosma infimbriana</i>		SU	GNR		SNI	
<i>Eucosma influana</i>		SU	GNR		SNI	
<i>Eucosma kokana</i>		SU	GNR		SNI	
<i>Eucosma montanana</i>		SU	GNR		SNI	
<i>Eucosma olivaceana</i>		SU	GNR		SNI	
<i>Eucosma verna</i>		SU	GNR		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Eufidonia notataria</i>		SU	G5		SNI	
<i>Euhyponomeutoides gracilariella</i>		S3	GNR		SNI	
<i>Eulithis propulsata</i>		S3	G5		SNI	
<i>Euphyes bimacula</i>	two-spotted skipper	SU	G4		SNI	
<i>Eupithecia mutata</i>		SU	G5		SNI	
<i>Euxoa adumbrata</i>		SU	G5		SNI	
<i>Euxoa albipennis</i>		SU	G4		SNI	
<i>Euxoa castanea</i>		SU	G5		SNI	
<i>Euxoa comosa</i>	hairy euxoa moth	SU	G5		SNI	
<i>Euxoa immixta</i>	mixed dart moth	SU	G4		SNI	
<i>Euxoa intrita</i>		SU	G5		SNI	
<i>Euxoa manitobana</i>		SU	GNR		SNI	
<i>Euxoa mimallonis</i>		S3	G5		SNI	
<i>Euxoa obeliscoides</i>	obelisk dart moth	SU	G5		SNI	
<i>Exaeretia fulva</i>		SU	GNR		SNI	
<i>Exaeretia gracilis</i>		SU	GNR		SNI	
<i>Exyra fax</i>		SU	G4		SNI	
<i>Fagitana littera</i>		S3	G4		SNI	
<i>Feltia mollis</i>	delicate cutworm moth	SU	G5		SNI	
<i>Globia laeta</i>		SU	G4		SNI	
<i>Glyphipterix quadragintapunctata</i>		SU	GNR		SNI	
<i>Gretchena semialba</i>		SU	GNR		SNI	
<i>Hadena capsularis</i>	capsule moth	S3	G5		SNI	
<i>Hahncappsia coloradensis</i>		SU	GNR		SNI	
<i>Hahncappsia pergivalis</i>		SU	GNR		SNI	
<i>Haimbachia squamulella</i>		SU	GNR		SNI	
<i>Hesperia colorado</i>	common branded skipper	S3	G5		SNI	
<i>Hesperia sassacus</i>	Indian skipper	S3	G5		SNI	
<i>Hydraecia perobliqua</i>		S3	GNR		SNI	
<i>Idia majoralis</i>		SU	GNR		SNI	
<i>Incurvaria vetulella</i>		SU	GNR		SNI	
<i>Lacanobia radix</i>		S3	G5		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Landryia impositella</i>		SU	GNR		SNI	
<i>Leucania dia</i>		SU	GNR		SNI	
<i>Leucobrephephos brephoides</i>		SU	G5		SNI	
<i>Lintneria eremitus</i>		S3	G4		SNI	
<i>Lithophane adipel</i>		S3	G4		SNI	
<i>Lithophane georgii</i>		S3	G5		SNI	
<i>Lithophane oriunda</i>		S3	G4		SNI	
<i>Lycia rachelae</i>		SU	G5		SNI	
<i>Macaria abruptata</i>		SU	G5		SNI	
<i>Macaria bitactata</i>		S3	G5		SNI	
<i>Macrochilo bivittata</i>		SU	G3		SNI	
<i>Melanapamea mixta</i>	an apamea moth	SU	GU		SNI	
<i>Meropleon ambifusca</i>		SU	G3		SNI	
<i>Mniotype ducta</i>	a noctuid moth	SU	G5		SNI	
<i>Mycterophora inexplicata</i>		S3	G5		SNI	
<i>Neocoehylis dubitana</i>		SU	GNR		SNI	
<i>Neodactria murellus</i>	a prairie sedge moth	SU	GNR		SNI	
<i>Oligia chlorostigma</i>	a brocade moth	S3	G4		SNI	
<i>Orthofidonia flavivenata</i>		SU	GNR		SNI	
<i>Orthonama evansi</i>		S3	GNR		SNI	
<i>Pachypolia atricornis</i>		SU	G3		SNI	
<i>Paectes oculatrix</i>		SU	G5		SNI	
<i>Paleacrita merriccata</i>		S3	G4		SNI	
<i>Papaipema circumlucens</i>		SU	G3		SNI	
<i>Papaipema maritima</i>		SU	G3		SNI	
<i>Papaipema silphii</i>	silphium borer moth	SU	G3		SNI	
<i>Papestra cristifera</i>		SU	G5		SNI	
<i>Pediasia truncatellus</i>		S3	GNR		SNI	
<i>Pelochrista comatulana</i>		SU	GNR		SNI	
<i>Pelochrista corosana</i>		S3	GNR		SNI	
<i>Pelochrista costastriata</i>		SU	GNR		SNI	
<i>Pelochrista fiskeana</i>		SU	GNR		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Pelochrista pulveratana</i>	a tortricid moth	SU	GNR		SNI	
<i>Pelochrista ridingsana</i>		SU	GNR		SNI	
<i>Peoria bipartitella</i>		SU	GNR		SNI	
<i>Peoria tetradella</i>		SU	GNR		SNI	
<i>Petrophila fulcalis</i>		S3	GNR		SNI	
<i>Photodes defecta</i>	a borer moth	SU	G3		SNI	
<i>Photodes includens</i>		SU	G4		SNI	
<i>Photodes inops</i>		SU	G3		SNI	
<i>Photodes panatela</i>	a moth	SU	GNR		SNI	
<i>Phragmatobia lineata</i>	a tiger moth	SU	G4		SNI	
<i>Phyciodes batesii</i>	tawny crescent	S3	G5		SNI	
<i>Plagiomimicus expallidus</i>	a noctuid moth	SU	GNR		SNI	
<i>Plagiomimicus pityochromus</i>		SU	G5		SNI	
<i>Plebejus melissa</i>	Melissa blue	S4	G5		SNI	
<i>Plebejus saepiolus</i>	greenish blue	S3	G5		SNI	
<i>Plemyria georgii</i>		S3	G5		SNI	
<i>Polia rogenhoferi</i>		SU	GNR		SNI	
<i>Polygonia gracilis</i>	hoary comma	SU	G5		SNI	
<i>Polygonia satyrus</i>	satyr comma	SU	G5		SNI	
<i>Polygrammodes flavidalis</i>	a moth	SU	GNR		SNI	
<i>Ponometia semiflava</i>		SU	G4		SNI	
<i>Ponometia tortricina</i>		SU	GNR		SNI	
<i>Prionapteryx achatina</i>		SU	GNR		SNI	
<i>Prionapteryx nebulifera</i>		SU	GNR		SNI	
<i>Prochoreutis inflatella</i>		SU	GNR		SNI	
<i>Proserpinus flavofasciata</i>		SU	G4		SNI	
<i>Protorthodes incincta</i>		S3	GNR		SNI	
<i>Proxenus mendosa</i>		S3	G5		SNI	
<i>Pyrausta laticlavata</i>		SU	GNR		SNI	
<i>Pyrausta nicalis</i>		S3	GNR		SNI	
<i>Pyrausta volupialis</i>		SU	GNR		SNI	
<i>Pyrgus centaureae</i>	grizzled skipper	S2	G5	S-SPC	SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Scardia anatomella</i>		SU	GNR		SNI	
<i>Schinia cumatilis</i>		SU	G4		SNI	
<i>Schinia gracilentia</i>		SU	G4		SNI	
<i>Schinia nundina</i>		S3	G5		SNI	
<i>Schinia sanguinea</i>	bleeding flower moth	SU	G4		SNI	
<i>Sciota rubescentella</i>		SU	GNR		SNI	
<i>Scopula ancellata</i>		S3	G5		SNI	
<i>Semioscopis merricella</i>		S3	GNR		SNI	
<i>Semioscopis packardella</i>		S3	GNR		SNI	
<i>Sideridis congermana</i>		SU	GNR		SNI	
<i>Sphinx drupiferarum</i>		S3	G4		SNI	
<i>Spragueia leo</i>	common spragueia moth	SU	G5		SNI	
<i>Stamnodes gibbicostata</i>		S3	G4		SNI	
<i>Stereomita andropogonis</i>		SU	GNR		SNI	
<i>Sunira verberata</i>		SU	G5		SNI	
<i>Sympistis chionanthi</i>	a sallow moth	SU	G5		SNI	
<i>Sympistis dinalda</i>		S3	GNR		SNI	
<i>Sympistis riparia</i>		SU	G4		SNI	
<i>Tharsalea epixanthe</i>	bog copper	S4	G5		SNI	
<i>Thera contractata</i>		SU	GNR		SNI	
<i>Trachea delicata</i>		SU	GNR		SNI	
<i>Tricholita notata</i>		SU	G5		SNI	
<i>Trichordestra rugosa</i>		S3	GNR		SNI	
<i>Udea inquinatalis</i>		SU	GNR		SNI	
<i>Ulolonche culea</i>		SU	G5		SNI	
<i>Vernia verna</i>	little glassywing	S3	G5		SNI	
<i>Xanthorhoe decoloraria</i>		SU	G5		SNI	
<i>Xanthorhoe iduata</i>	a geometrid moth	SU	G4		SNI	
<i>Xanthorhoe labradorensis</i>		S3	G5		SNI	
<i>Xanthorhoe packardata</i>		SU	G4		SNI	
<i>Xestia oblata</i>	ruby dart moth	S3	G5		SNI	
<i>Xestia perquiritata</i>		SU	G5		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Xylena cineritia</i>		S3	G5		SNI	
<i>Zanclognatha dentata</i>		SU	G3		SNI	

Flies (Class Insecta; Order Diptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Echthodopa pubera</i>	a robber fly	SU	GNR		SNI	
<i>Laphria huron</i>	a robber fly	SU	GNR		SNI	
<i>Laphystia flavipes</i>	a robber fly	SU	GNR		SNI	
<i>Lasiopogon terricolus</i>	a robber fly	SU	GNR		SNI	
<i>Leptogaster flavipes</i>	a robber fly	SU	GNR		SNI	

Grasshoppers, Crickets (Class Insecta; Order Orthoptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Melanoplus dawsoni</i>	Dawson's spur-throat grasshopper	SU	G5		SNI	
<i>Melanoplus scudderi</i>	Scudder's short-wing grasshopper	SU	G5		SNI	
<i>Nomotettix parvus</i>	low-ridged pygmy grasshopper	SU	G3		SNI	
<i>Orphulella pelidna</i>	spotted-winged grasshopper	SU	G5		SNI	
<i>Phoetaliotes nebrascensis</i>	a grasshopper	S3	G5		SNI	
<i>Pseudopomala brachyptera</i>	bunch grass locust	S3	G5		SNI	

Millipedes (Class Diplopoda)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Achemenides pectinatus</i>	a cave obligate millipede	SU	G3		SNI	

Snails (Class Gastropoda)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Acella haldemani</i>	spindle lymnaea	S4	G3		SGCN	Decline
<i>Gastrocopta rogersensis</i>	Roger's snaggletooth snail	SU	G3	S-SPC	SGCN	2015 SGCN, H threats
<i>Gastrocopta similis</i>	Great Lakes snaggletooth	S2	G5		SGCN	H threats
<i>Planogyra asteriscus</i>	eastern flat-whorl	SU	G4	S-SPC	SGCN	2015 SGCN
<i>Striatura ferrea</i>	black striate	SU	G5	S-THR	SGCN	2015 SGCN
<i>Vallonia perspectiva</i>	thin-lip vallonia	S2	G4		SGCN	H threats
<i>Vertigo gouldi</i>	variable vertigo	S2	G5		SGCN	H threats
<i>Vertigo meramecensis</i>	bluff vertigo	S1	G2	S-THR	SGCN	2015 SGCN, Decline, H threats, Very Rare
<i>Zonitoides limatulus</i>	dull gloss	SU	G4	S-SPC	SGCN	2015 SGCN
<i>Aplexa elongata</i>	lance aplexa	SU	G5		SNI	
<i>Campeloma decisum</i>	pointed campeloma	S4	G5		SNI	
<i>Carychium exiguum</i>	obese thorn	S3	G5		SNI	
<i>Catinella exilis (Mediappendix exilis)</i>	pleistocene catinella	SU	G3		SNI	
<i>Cochlicopa lubricella</i>	thin pillar	SU	G5		SNI	
<i>Euchemotrema fraternum</i>	upland pillsnail	SU	G5		SNI	
<i>Euconulus alderi</i>	a terrestrial snail	SU	G5		SNI	
<i>Gyraulus circumstriatus</i>	disc gyro	SU	G5		SNI	
<i>Helicodiscus parallelus</i>	compound coil	S3	G5		SNI	
<i>Marstonia lustrica</i>	boreal marstonia	SU	G4		SNI	
<i>Novisuccinea ovalis</i>	oval ambersnail	S3	G5		SNI	
<i>Pleurocera acuta</i>	sharp hornsnail	S4	G5		SNI	
<i>Punctum minutissimum</i>	small spot	S3	G5		SNI	
<i>Somatogyrus depressus</i>	sandbar pebblesnail	SU	G2		SNI	
<i>Somatogyrus tryoni</i>	coldwater pebblesnail	SU	G2		SNI	
<i>Strobilops affinis</i>	eightfold pinecone	SU	G4		SNI	
<i>Vallonia costata</i>	costate vallonia	S3	G5		SNI	
<i>Valvata tricarinata</i>	threeridge valvata	S4	G5		SNI	
<i>Vertigo arthuri</i>	Dakota vertigo	S3	G5		SNI	
<i>Vertigo bollesiana</i>	delicate vertigo	S3	G4		SNI	
<i>Vertigo morsei</i>	six-whorl vertigo	SU	G3		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Vertigo nylanderi</i>	deep-throated vertigo	SU	G3		SNI	
<i>Vitrina angelicae</i>	eastern glass-snail	S3	G5		SNI	

Spiders (Class Arachnida; Order Araneae)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Habronattus texanus</i>	a jumping spider	S2	GNR	S-SPC	SGCN	2015 SGCN, Decline, Very Rare
<i>Latrodectus variolus</i>	northern black widow	S3	GNR		SGCN	Very Rare
<i>Neoscona pratensis</i>	an orb weaver	S3	GNR		SGCN	Very Rare
<i>Phidippus apacheanus</i>	a jumping spider	S2	GNR	S-SPC	SGCN	2015 SGCN, Very Rare
<i>Sassacus papenhoei</i>	a jumping spider	S2	GNR	S-SPC	SGCN	2015 SGCN, Very Rare
<i>Sphodros niger</i>	black purse-web spider	S3	G4		SGCN	Rare
<i>Tutelina formicaria</i>	a jumping spider	S1	GNR	S-THR	SGCN	2015 SGCN, M threats, Very Rare

Springtails (Class Collembola)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Arrhopalites dubius</i>	a cave obligate springtail	SU	G2		SNI	
<i>Onychiurus obesius</i>	a cave obligate springtail	SU	G3		SNI	
<i>Tullbergia hades</i>	a cave obligate springtail	SU	G2		SNI	

True Bugs, Cicadas, Leafhoppers (Class Insecta; Order Hemiptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Aflexia rubranura</i>	red-tailed prairie leafhopper	S2	G3	S-SPC	SGCN	2015 SGCN, Decline, H threats
<i>Okanagana balli</i>	prairie cicada	S2	GNR		SGCN	H threats
<i>Attenuipyga joyceae</i>	a leafhopper	SU	GNR		SNI	
<i>Bruchomorpha dorsata</i>	a planthopper	SU	GNR		SNI	
<i>Driotura robusta</i>	a leafhopper	SU	GNR		SNI	
<i>Macrosteles clavatus</i>	caped leafhopper	SU	GNR	S-SPC	SNI	
<i>Paraphlepsius nebulosus</i>	a leafhopper	SU	GNR		SNI	

<i>Polyamia caperata</i>	little bluestem polyamia	SU	GNR		SNI	
<i>Polyamia herbida</i>	prairie panic grass leafhopper	SU	GNR		SNI	
<i>Xerophloea major</i>	a leafhopper	SU	GNR		SNI	

Caddisflies (Class Insecta; Order Trichoptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Agapetus tomus</i>	a caddisfly	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN, Rare
<i>Agrypnia glacialis</i>	a caddisfly	SH	G5		SGCN	
<i>Anabolia ozburni</i>	a caddisfly	S2	G5	S-SPC	SGCN	2015 SGCN, RSGCN, Decline, Rare
<i>Anabolia sordida</i>	a caddisfly	SH	G5		SGCN	
<i>Apatania zonella</i>	a caddisfly	S2	G5		SGCN	Very Rare
<i>Asynarchus rossi</i>	a caddisfly	SU	G4	S-THR	SGCN	RSGCN, Very Rare
<i>Banksiola smithi</i>	a caddisfly	SH	G5		SGCN	
<i>Ceraclea albsticta</i>	a caddisfly	SH	GH		SGCN	Proposed RSGCN
<i>Ceraclea brevis</i>	a caddisfly	SH	G2		SGCN	
<i>Ceraclea misca</i>	a caddisfly	SH	G5		SGCN	
<i>Cheumatopsyche wabasha</i>	a caddisfly	SH	G4		SGCN	
<i>Chilostigma itascaae</i>	Headwaters Chilostigman caddisfly	S2	G2	S-THR	SGCN	Endemic, 2015 SGCN, RSGCN, RFSS
<i>Goera stylata</i>	a caddisfly	S2	G4	S-THR	SGCN	RSGCN, RFSS, H threats
<i>Grammotaulius interrogationis</i>	a caddisfly	SH	G5		SGCN	
<i>Holocentropus glacialis</i>	frozen trumpet-net caddisfly	S2	G3	S-THR	SGCN	RSGCN, Rare
<i>Holocentropus melanae</i>	black river trumpet-net caddisfly	S2	G3		SGCN	Proposed RSGCN
<i>Holocentropus milaca</i>	a caddisfly	S1	G1	S-END	SGCN	RSGCN, Very Rare
<i>Hydropsyche californica</i>	a caddisfly	SH	G5		SGCN	
<i>Hydroptila callia</i>	a caddisfly	SH	G5		SGCN	
<i>Hydroptila delineata</i>	a caddisfly	S2	G5		SGCN	Rare
<i>Hydroptila metoeca</i>	a caddisfly	SH	G4	S-SPC	SGCN	2015 SGCN
<i>Hydroptila quinola</i>	a caddisfly	S3	G5	S-SPC	SGCN	2015 SGCN
<i>Hydroptila rono</i>	a caddisfly	S1	G5	S-THR	SGCN	2015 SGCN, Very Rare
<i>Hydroptila tortosa</i>	a caddisfly	SH	G4	S-SPC	SGCN	2015 SGCN
<i>Ironoquia punctatissima</i>	a caddisfly	S2	G5	S-THR	SGCN	2015 SGCN, M threats, Rare

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Lenarchus keratus</i>	a caddisfly	SH	G4		SGCN	
<i>Lepidostoma libum</i>	a caddisfly	S1	G3	S-THR	SGCN	2015 SGCN, Very Rare
<i>Limnephilus argenteus</i>	a caddisfly	S2	G5		SGCN	
<i>Limnephilus janus</i>	a caddisfly	SH	G5	S-END	SGCN	2015 SGCN
<i>Limnephilus parvulus</i>	a caddisfly	S2	G5		SGCN	Very Rare
<i>Limnephilus sackeni</i>	a caddisfly	S2	G4		SGCN	Proposed RSGCN, Very Rare
<i>Limnephilus secludens</i>	a caddisfly	S1	G5	S-END	SGCN	2015 SGCN, Decline, Very Rare
<i>Limnephilus tarsalis</i>	a caddisfly	SH	G5		SGCN	
<i>Micrasema sprulesi</i>	Sprules's humpleless caddisfly	S3	G3		SGCN	
<i>Oecetis ditissa</i>	a caddisfly	S2	G5	S-THR	SGCN	2015 SGCN, Very Rare
<i>Orthotrichia curta</i>	short orthotrichian microcaddisfly	SH	G4		SGCN	
<i>Oxyethira ecornuta</i>	a caddisfly	S2	G5	S-THR	SGCN	2015 SGCN, RSGCN, Rare
<i>Oxyethira itascaae</i>	a caddisfly	S3	G3	S-SPC	SGCN	2015 SGCN, RSGCN
<i>Parapsyche apicalis</i>	a caddisfly	S2	G5	S-THR	SGCN	2015 SGCN, H threats, Rare
<i>Philarctus bergrothi</i>	Bergroth's northern caddisfly	SH	G5		SGCN	
<i>Protoptila erotica</i>	a caddisfly	S3	G5	S-SPC	SGCN	2015 SGCN
<i>Psilotreta indecisa</i>	uncertain mortarjoint caddisfly	S2	G4		SGCN	
<i>Pycnopsyche scabripennis</i>	a caddisfly	SH	G5		SGCN	
<i>Rhyacophila angelita</i>	a caddisfly	S2	G5		SGCN	M threats, Very Rare
<i>Rhyacophila brunnea</i>	brown free-living caddisfly	S2	G5		SGCN	Very Rare
<i>Rhyacophila minor</i>	smaller free-living caddisfly	S2	G4		SGCN	Rare
<i>Rhyacophila vibox</i>	a rhyacophilan caddisfly	S2	G5		SGCN	H threats, Very Rare
<i>Triaenodes flavescens</i>	a triaenode caddisfly	S3	G5	S-SPC	SGCN	2015 SGCN, Decline, Rare
<i>Triaenodes reuteri</i>	a caddisfly	S2	G5		SGCN	Decline
<i>Wormaldia moesta</i>	a caddisfly	SH	G5		SGCN	
<i>Agrypnia deflata</i>	a caddisfly	SU	G5		SNI	
<i>Arctopora pulchella</i>	a caddisfly	SU	G5		SNI	
<i>Asynarchus mutatus</i>	a caddisfly	SU	G5		SNI	
<i>Banksiola dossuaria</i>	a caddisfly	SH	G5		SNI	
<i>Beothukus complicatus</i>	a caddisfly	SU	G4		SNI	
<i>Brachycentrus fuliginosus</i>	a caddisfly	SU	G5		SNI	
<i>Brachycentrus lateralis</i>	sideways humpleless caddisfly	SU	G4		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Ceraclea neffi</i>	a caddisfly	SH	G5		SNI	
<i>Ceraclea nepha</i>	a ceraclean caddisfly	SH	G5		SNI	
<i>Ceraclea protonepha</i>	a caddisfly	SU	G5		SNI	
<i>Ceraclea slossonae</i>	a caddisfly	SH	G4		SNI	
<i>Cernotina spicata</i>	a caddisfly	SU	G5		SNI	
<i>Fabria inornata</i>	a caddisfly	SU	G4		SNI	
<i>Frenesia missa</i>	a caddisfly	SU	G5		SNI	
<i>Glossosoma lividum</i>	bluish saddle-case caddisfly	SU	G3		SNI	
<i>Holocentropus picicornis</i>	black-horned trumpet-net caddisfly	SU	G5		SNI	
<i>Hydroptila amoena</i>	a caddisfly	SU	G5		SNI	
<i>Hydroptila ampoda</i>	a caddisfly	SU	G5		SNI	
<i>Hydroptila antennopodia</i>	a caddisfly	SU	G3		SNI	
<i>Hydroptila bernerii</i>	Berner's microcaddisfly	SU	G4		SNI	
<i>Hydroptila fiskei</i>	a caddisfly	SU	G4		SNI	
<i>Hydroptila hamata</i>	a caddisfly	SU	G5		SNI	
<i>Hydroptila salmo</i>	a caddisfly	SU	G4		SNI	
<i>Hydroptila waskesia</i>	a caddisfly	SU	G4	S-END	SNI	
<i>Lenarchus crassus</i>	a caddisfly	SU	G5		SNI	
<i>Lepidostoma americanum</i>	a caddisfly	SU	G5		SNI	
<i>Lepidostoma cinereum</i>	a caddisfly	SU	G5		SNI	
<i>Lepidostoma griseum</i>	a caddisfly	SU	G5		SNI	
<i>Lepidostoma prominens</i>	a caddisfly	SU	G5		SNI	
<i>Leptophylax gracilis</i>	a caddisfly	SU	G4		SNI	
<i>Limnephilus partitus</i>	a caddisfly	SU	G5		SNI	
<i>Limnephilus sublunatus</i>	a caddisfly	SU	G5		SNI	
<i>Neophylax aniqua</i>	northeastern little caddisfly	SU	G4		SNI	
<i>Neophylax mitchelli</i>	Mitchell's little caddisfly	SU	G3		SNI	
<i>Neotrichia falca</i>	a caddisfly	SU	G3		SNI	
<i>Neotrichia halia</i>	a caddisfly	SU	G5		SNI	
<i>Nyctiophylax banksi</i>	a caddisfly	SU	G4		SNI	
<i>Ochrotrichia spinosa</i>	a purse casemaker caddisfly	SU	G5	S-END	SNI	
<i>Ochrotrichia stylata</i>	a caddisfly	SU	G5		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Ochrotrichia wojcickyi</i>	a caddisfly	SU	G4		SNI	
<i>Oecetis disjuncta</i>	unyoked long-horned caddisfly	SU	GNR		SNI	
<i>Oecetis sordida</i>	a caddisfly	SU	GNR		SNI	
<i>Oligostomis ocelligera</i>	a caddisfly	SU	G5		SNI	
<i>Oxyethira rossi</i>	a caddisfly	SU	G3		SNI	
<i>Plectrocnemia clinei</i>	Cline's trumpet-net caddisfly	SU	G4		SNI	
<i>Plectrocnemia icula</i>	Laurentides trumpet-net caddisfly	SU	G3		SNI	
<i>Polycentropus centralis</i>	a caddisfly	SU	G5		SNI	
<i>Ptilostomis angustipennis</i>	a caddisfly	SU	G5		SNI	
<i>Pycnopsyche indiana</i>	a caddisfly	SU	G5		SNI	
<i>Stactobiella delira</i>	a caddisfly	SU	G5		SNI	
<i>Triaenodes frontalis</i>	a caddisfly	SU	G5	S-THR	SNI	

Dragonflies and Damselflies (Class Insecta; Order Odonata)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Aeshna sitchensis</i>	zigzag darner	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, M threats
<i>Aeshna subarctica</i>	subarctic darner	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, M threats
<i>Archilestes grandis</i>	great spreadwing	S3	G5		SGCN	Rare
<i>Argia funebris</i>	springwater dancer	S1	G5		SGCN	2015 SGCN, RSGCN Watchlist, M threats, Very Rare
<i>Arigomphus submedianus</i>	jade clubtail	S2	G5		SGCN	Rare
<i>Boyeria grafiana</i>	ocellated darner	S2	G5	S-SPC	SGCN	2015 SGCN, H threats, Rare
<i>Coenagrion angulatum</i>	prairie bluet	S1	G5		SGCN	2015 SGCN, Decline, H threats, Rare
<i>Coenagrion interrogatum</i>	subarctic bluet	S2	G5		SGCN	2015 SGCN, H threats
<i>Enallagma anna</i>	river bluet	S2	G5		SGCN	
<i>Enallagma aspersum</i>	azure bluet	S2	G5		SGCN	Rare
<i>Gomphurus lineatifrons</i>	splendid clubtail	S2	G4		SGCN	2015 SGCN, RSGCN Watchlist, M threats, Rare
<i>Gomphurus ventricosus</i>	skillet clubtail	S3	G3		SGCN	2015 SGCN, RSGCN, M threats
<i>Hylogomphus viridifrons</i>	green-faced clubtail	S3	G3		SGCN	RSGCN, Rare
<i>Nannothemis bella</i>	elfin skimmer	S2	G4		SGCN	2015 SGCN, RSGCN
<i>Ophiogomphus anomalus</i>	extra-striped snaketail	S1	G4	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, H threats, Rare

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Ophiogomphus howei</i>	pygmy snaketail	S2	G3	S-SPC	SGCN	2015 SGCN, RSGCN, M threats, Rare
<i>Ophiogomphus smithi</i>	Sioux snaketail	S1	G2		SGCN	Endemic, 2015 SGCN, RSGCN, Decline, H threats, Rare
<i>Ophiogomphus susbehcha</i>	St. Croix snaketail	S2	G2	S-THR	SGCN	Endemic, 2015 SGCN, RSGCN, Decline, H threats, Rare
<i>Phanogomphus quadricolor</i>	rapids clubtail	S3	G3		SGCN	2015 SGCN
<i>Somatochlora brevicincta</i>	Quebec emerald	S2	G4	S-SPC	SGCN	Endemic, 2015 SGCN, RSGCN, RFSS, H threats, Rare
<i>Somatochlora forcipata</i>	forcipate emerald	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist, H threats
<i>Stylogomphus albistylus</i>	least clubtail	S2	G5		SGCN	2015 SGCN, H threats, Rare
<i>Stylurus notatus</i>	elusive clubtail	S3	G3		SGCN	RSGCN, Decline, H threats
<i>Williamsonia fletcheri</i>	ebony boghaunter	S2	G4		SGCN	2015 SGCN, RSGCN Watchlist, RFSS, H threats, Rare
<i>Zoraena obliqua</i>	arrowhead spiketail	S2	G4		SGCN	2015 SGCN
<i>Amphiagrion abbreviatum</i>	western red damsel	S3	G5		SNI	
<i>Nehallenia gracilis</i>	sphagnum sprite	SU	G5		SNI	
<i>Neurocordulia molesta</i>	smoky shadowdragon	SU	G4		SNI	
<i>Neurocordulia yamaskanensis</i>	stygian shadowdragon	S3	G5		SNI	
<i>Rhionaeschna multicolor</i>	blue-eyed darner	S3	G5		SNI	
<i>Rhionaeschna mutata</i>	spatterdock darner	SH	G4		SNI	
<i>Somatochlora elongata</i>	ski-tailed emerald	S3	G5		SNI	
<i>Stylurus scudderi</i>	zebra clubtail	S3	G5		SNI	

Mayflies (Class Insecta; Order Ephemeroptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Ameletus subnotatus</i>	boreal comb minnow mayfly	S2	G5		SGCN	Rare
<i>Drunella cornuta</i>	big-horned spiny crawler mayfly	SH	G5		SGCN	RSGCN Watchlist
<i>Drunella walkeri</i>	Walker's spiny crawler mayfly	SH	G5		SGCN	
<i>Eurylophella funeralis</i>	brown spiny crawler mayfly	S2	G5		SGCN	RSGCN, H threats
<i>Neophemera bicolor</i>	Canadian large square-gilled mayfly	S1	G1		SGCN	RSGCN, Rare
<i>Siphonurus alternatus</i>	northern primitive minnow mayfly	S1	G5		SGCN	Decline, Very Rare
<i>Siphonurus phyllis</i>	leafy primitive minnow mayfly	S1	G3		SGCN	Proposed RSGCN, Very Rare
<i>Sparbarus maculatus</i>	maculated small square-gilled mayfly	S2	G5		SGCN	M threats
<i>Acanthametropus pecatonica</i>	Pecatonica River mayfly	SU	G3		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Anafroptilum album</i>	white small minnow mayfly	SU	G5		SNI	
<i>Arthroplea bipunctata</i>	American palp-headed mayfly	SU	G5		SNI	
<i>Attenella attenuata</i>	tapered spiny crawler mayfly	SU	G5		SNI	
<i>Attenella margarita</i>	Margaret's spiny crawler mayfly	SU	G5		SNI	
<i>Baetis bundyae</i>	Bundy's small minnow mayfly	SU	G4		SNI	
<i>Baetis phoebeus</i>	a small minnow mayfly	SU	G5		SNI	
<i>Brachycercus harrisella</i>	Curtis's small square-gilled mayfly	SU	G5		SNI	
<i>Brachycercus ojbwe</i>	Ojibwe small square-gilled mayfly	SU	G3		SNI	
<i>Caenis anceps</i>	Traver's small square-gilled mayfly	SU	G5		SNI	
<i>Callibaetis fluctuans</i>	wavy small minnow mayfly	SU	G5		SNI	
<i>Callibaetis pallidus</i>	pale small minnow mayfly	SU	G5		SNI	
<i>Callibaetis pictus</i>	a small minnow mayfly	SU	G5		SNI	
<i>Callibaetis pretiosus</i>	pretty small minnow mayfly	SU	G5		SNI	
<i>Callibaetis skokianus</i>	skokie small minnow mayfly	SU	G4		SNI	
<i>Cercobrachys etowah</i>	etowah small square-gilled mayfly	SU	G4		SNI	
<i>Cercobrachys lilliei</i>	Lillie's small square-gilled mayfly	SU	G2		SNI	
<i>Dannella lita</i>	Burks's spiny crawler mayfly	SU	G5		SNI	
<i>Dannella simplex</i>	simple spiny crawler mayfly	SU	G5		SNI	
<i>Dolania americana</i>	American sand burrowing mayfly	SU	G4		SNI	
<i>Ephemera varia</i>	varied burrowing mayfly	SU	G5		SNI	
<i>Ephemerella dorothea</i>	pale spiny crawler mayfly	SU	G5		SNI	
<i>Ephemerella needhami</i>	Needham's spiny crawler mayfly	SU	G5		SNI	
<i>Eurylophella bicoloroides</i>	Nova Scotia spiny crawler mayfly	SU	G3		SNI	
<i>Eurylophella lutulenta</i>	dirty spiny crawler mayfly	SU	G5		SNI	
<i>Eurylophella minimella</i>	reduced spiny crawler mayfly	SU	G4		SNI	
<i>Eurylophella prudentialis</i>	prudent spiny crawler mayfly	SU	G5		SNI	
<i>Eurylophella verisimilis</i>	plausible spiny crawler mayfly	SU	G5		SNI	
<i>Habrophlebia vibrans</i>	vibrant prong-gilled mayfly	SU	G5		SNI	
<i>Heptagenia pulla</i>	dusky flat-headed mayfly	SU	G5		SNI	
<i>Heptagenia whitingi</i>	Whiting's flat-headed mayfly	SU	G2		SNI	
<i>Heterocloeon curiosum</i>	curious small minnow mayfly	SU	G5		SNI	
<i>Hexagenia rigida</i>	straight hex burrowing mayfly	SU	G5		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Isonychia arida</i>	arid howdy mayfly	SU	G5		SNI	
<i>Isonychia campestris</i>	western howdy mayfly	SU	G4		SNI	
<i>Labiobaetis ephippiatus</i>	saddled small minnow mayfly	SU	G5		SNI	
<i>Leucrocuta aphrodite</i>	Aphrodite's flat-headed mayfly	SU	G5		SNI	
<i>Leucrocuta hebe</i>	Hebe's flat-headed mayfly	SU	G5		SNI	
<i>Leucrocuta juno</i>	Juno's flat-headed mayfly	SU	G4		SNI	
<i>Leucrocuta maculipennis</i>	midwestern flat-headed mayfly	SU	G5		SNI	
<i>Litobrantha recurvata</i>	dark burrowing mayfly	SU	G5		SNI	
<i>Maccaffertium ithaca</i>	Ithaca flat-headed mayfly	SU	G5		SNI	
<i>Maccaffertium pulchellum</i>	pretty flat-headed mayfly	SU	G5		SNI	
<i>Macdunnoa persimlex</i>	a flat-headed mayfly	SU	G4		SNI	
<i>Neocloeon triangulifer</i>	triangle small minnow mayfly	SU	G5		SNI	
<i>Neoleptophlebia adoptiva</i>	Angler's prong-gilled mayfly	SU	G5		SNI	
<i>Nixe inconspicua</i>	inconspicuous flat-headed mayfly	SU	G5		SNI	
<i>Nixe lucidpennis</i>	clear-winged flat-headed mayfly	SU	G4		SNI	
<i>Paracloeodes lotor</i>	Coon Rapids small minnow mayfly	SU	GNR		SNI	
<i>Paracloeodes minutus</i>	minute small minnow mayfly	S4	G5		SNI	
<i>Paraleptophlebia debilis</i>	fragile prong-gilled mayfly	SU	G5		SNI	
<i>Paraleptophlebia guttata</i>	dappled prong-gilled mayfly	SU	G5		SNI	
<i>Paraleptophlebia ontario</i>	Ontario prong-gilled mayfly	SU	G4		SNI	
<i>Paraleptophlebia volitans</i>	hairy prong-gilled mayfly	SU	G5		SNI	
<i>Parameletus croesus</i>	dark-winged primitive minnow mayfly	SU	GH		SNI	
<i>Pentagenia vittigera</i>	common spiny-headed burrowing mayfly	SU	G5		SNI	
<i>Plauditus cingulatus</i>	sepia small minnow mayfly	SU	G5		SNI	
<i>Plauditus elliotti</i>	Elliott's small minnow mayfly	SU	G1		SNI	
<i>Procloeon bellum</i>	aggressive small minnow mayfly	SU	G5		SNI	
<i>Procloeon pennulatum</i>	Eaton's small minnow mayfly	SU	G5		SNI	
<i>Procloeon rubropictum</i>	painted small minnow mayfly	SU	G5		SNI	
<i>Procloeon rufostrigatum</i>	Hobbs's small minnow mayfly	SU	G5		SNI	
<i>Procloeon simplex</i>	simple small minnow mayfly	SU	G5		SNI	
<i>Procloeon viridoculare</i>	cat-eye small minnow mayfly	SU	G5		SNI	
<i>Pseudocentroptiloides morihari</i>	Morihara's small minnow mayfly	SU	G2		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Raptoheptagenia cruentata</i>	predaceous flat-headed mayfly	SU	G4		SNI	
<i>Rhithrogena impersonata</i>	distant flat-headed mayfly	SU	G5		SNI	
<i>Rhithrogena manifesta</i>	central flat-headed mayfly	SU	G5		SNI	
<i>Rhithrogena serpenglena</i>	a flat-headed mayfly	SU	GNR		SNI	
<i>Siphonurus quebecensis</i>	Quebec primitive minnow mayfly	SU	G5		SNI	
<i>Siphloplecton basale</i>	flapped cleft-footed minnow mayfly	SU	G5		SNI	
<i>Siphloplecton interlineatum</i>	flapless cleft-footed minnow mayfly	SU	G5		SNI	
<i>Sparbarus lacustris</i>	a mayfly	SU	G4		SNI	
<i>Sparbarus nasutus</i>	large-nosed small square-gill mayfly	SU	G3		SNI	
<i>Stenacron candidum</i>	candid flat-headed mayfly	SU	G4		SNI	
<i>Stenacron carolina</i>	Carolina flat-headed mayfly	SU	G5		SNI	
<i>Stenacron gildersleevei</i>	Gildersleeve's mountain flat-headed mayfly	SU	G4		SNI	
<i>Stenacron minnetonka</i>	Minnetonka flat-headed mayfly	SU	G4		SNI	
<i>Tortopsis primus</i>	Manitoba white burrowing mayfly	SU	G5		SNI	
<i>Tricorythodes albilineatus</i>	a stout crawler mayfly	SU	G5		SNI	
<i>Tricorythodes allectus</i>	Needham's stout crawler mayfly	SU	G5		SNI	
<i>Tricorythodes cobbi</i>	Cobb's stout crawler mayfly	SU	G3		SNI	
<i>Tricorythodes robacki</i>	Roback's stout crawler mayfly	SU	G4		SNI	

Mussels (Class Bivalvia)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Actinonaias ligamentina</i>	mucket	S2	G5	S-THR	SGCN	2015 SGCN, Decline, H threats
<i>Alasmidonta marginata</i>	elktoe	S2	G4	S-THR	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Arcidens confragosus</i>	rock pocketbook	S2	G4	S-END	SGCN	2015 SGCN, RSGCN Watchlist , Decline, H threats
<i>Cumberlandia monodonta</i>	spectaclecase	S1	G3	S-END, F-END	SGCN	2015 SGCN, RSGCN, Decline, H threats, Rare
<i>Cyclonaias nodulata</i>	wartyback	S3	G4	S-THR	SGCN	2015 SGCN, RSGCN, VH threats
<i>Cyclonaias tuberculata</i>	purple wartyback	S2	G5	S-END	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Ellipsaria lineolata</i>	butterfly	S2	G4	S-THR	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Elliptio complanata</i>	eastern elliptio	S1	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist , M threats, Rare
<i>Elliptio crassidens</i>	elephant-ear	S1	G5	S-END	SGCN	2015 SGCN, RSGCN, Decline, H threats, Very Rare

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Epioblasma triquetra</i>	snuffbox	S1	G3	S-END, F-END	SGCN	2015 SGCN, RSGCN, Decline, H threats, Rare
<i>Eurynia dilatata</i>	spike	S2	G5	S-THR	SGCN	2015 SGCN, RSGCN Watchlist , Decline, H threats
<i>Lampsilis higginsii</i>	Higgins eye	S1	G1	S-END, F-END	SGCN	2015 SGCN, RSGCN, H threats
<i>Lampsilis sietmani</i>	canary kingshell	S1	GNR		SGCN	Proposed RSGCN Watchlist, H threats, Very Rare
<i>Lampsilis teres</i>	yellow sandshell	S1	G5	S-END	SGCN	2015 SGCN, RSGCN, Decline, H threats, Very Rare
<i>Lasmigona compressa</i>	creek heelsplitter	S3	G5	S-SPC	SGCN	2015 SGCN, RSGCN, RFSS, H threats
<i>Lasmigona costata</i>	flutedshell	S3	G5	S-THR	SGCN	2015 SGCN, RFSS, Decline, H threats
<i>Ligumia recta</i>	black sandshell	S3	G4	S-SPC	SGCN	2015 SGCN, RSGCN, RFSS, H threats
<i>Megaloniaias nervosa</i>	washboard	S2	G5	S-END	SGCN	2015 SGCN, H threats
<i>Obovaria olivaria</i>	hickorynut	S2	G4		SGCN	2015 SGCN, Decline, H threats
<i>Plethobasus cyphus</i>	sheepnose	S1	G3	S-END, F-END	SGCN	2015 SGCN, RSGCN, Decline, H threats, Very Rare
<i>Pleurobema sintoxia</i>	round pigtoe	S3	G4	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist , Decline, H threats
<i>Potamilus capax</i>	fat pocketbook	SX	G2		SGCN	2015 SGCN, RSGCN, H threats
<i>Potamilus leptodon</i>	scaleshell	SX	G1		SGCN	RSGCN, H threats
<i>Quadrula fragosa</i>	winged mapleleaf	S1	G1	S-END, F-END	SGCN	2015 SGCN, RSGCN, Decline, H threats, Very Rare
<i>Reginaia eburnus</i>	ebonyshell	S1	G4	S-END	SGCN	2015 SGCN, RSGCN, Decline, H threats, Very Rare
<i>Sagittunio subrostratus</i>	pondmussel	S2	G5	S-THR	SGCN	2015 SGCN, Decline, H threats, Rare
<i>Simpsoniaias ambigua</i>	salamander mussel	S1	G3	S-END, F-P-END	SGCN	2015 SGCN, RSGCN, Decline, H threats, Very Rare
<i>Theliderma metanevra</i>	monkeyface	S2	G4	S-THR	SGCN	2015 SGCN, RSGCN Watchlist , Decline, H threats
<i>Tritogonia nobilis</i>	gulf mapleleaf	SH	G4		SGCN	RSGCN Watchlist
<i>Tritogonia verrucosa</i>	pistolgrip	S2	G4	S-END	SGCN	2015 SGCN, Decline, H threats, Rare
<i>Truncilla donaciformis</i>	fawnsfoot	S2	G5	S-THR	SGCN	2015 SGCN, RSGCN Watchlist , Decline, H threats
<i>Utterbackiana suborbiculata</i>	flat floater	S1	G5	S-SPC	SGCN	2015 SGCN, RSGCN Watchlist , M threats, Very Rare
<i>Venustaconcha ellipsiformis</i>	ellipse	S2	G4	S-THR	SGCN	2015 SGCN, RSGCN, Decline, H threats
<i>Pyganodon lacustris</i>	lake floater	S3	GU		SNI	

Shrimp (Class Branchiopoda)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Eubbranchipus ornatus</i>	ornate fairy shrimp	SU	G3		SNI	

Stoneflies (Class Insecta; Order Plecoptera)

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Alloperla atlantica</i>	a stonefly	S2	G5		SGCN	Proposed RSGCN Watchlist, Decline, Very Rare
<i>Attaneuria ruralis</i>	a stonefly	S2	G4		SGCN	RSGCN
<i>Isogenoides frontalis</i>	a perlid stonefly	S2	G5		SGCN	RSGCN, Rare
<i>Paracapnia opis</i>	a stonefly	S2	G5		SGCN	Proposed RSGCN Watchlist, Rare
<i>Paranemoura perfecta</i>	spotted forestfly	S1	G5		SGCN	Very Rare
<i>Acroneuria carolinensis</i>	Carolina stone	SU	G5		SNI	
<i>Allocaupnia illinoensis</i>	a stonefly	SU	G3		SNI	
<i>Allocaupnia rickeri</i>	a stonefly	SU	G5		SNI	
<i>Allocaupnia vivipara</i>	a stonefly	SU	G5		SNI	
<i>Alloperla leonarda</i>	a stonefly	SU	G4		SNI	
<i>Alloperla usa</i>	Appalachian sallfly	SU	G5		SNI	
<i>Amphinemura delosa</i>	eastern forestfly	SU	G5		SNI	
<i>Amphinemura palmeni</i>	lovely forestfly	SU	G5		SNI	
<i>Amphinemura varshava</i>	Warsaw forestfly	SU	G4		SNI	
<i>Capnia vernalis</i>	a stonefly	SU	G5		SNI	
<i>Capnura manitoba</i>	Manitoba snowfly	SU	G4		SNI	
<i>Haploperla brevis</i>	a stonefly	SU	G5		SNI	
<i>Haploperla orpha</i>	a stonefly	SU	G4		SNI	
<i>Helopicus subvarians</i>	vernal springfly	SU	G5		SNI	
<i>Isogenoides doratus</i>	a stonefly	SU	G3		SNI	
<i>Isogenoides varians</i>	a stonefly	SU	G3		SNI	
<i>Isoperla emarginata</i>	a stonefly	SU	GH		SNI	
<i>Isoperla frisoni</i>	a stonefly	SU	G5		SNI	
<i>Isoperla longiseta</i>	a stonefly	SU	G5		SNI	
<i>Isoperla maxana</i>	a stonefly	SU	GH		SNI	
<i>Leuctra ferruginea</i>	a stonefly	SU	G5		SNI	
<i>Leuctra tenuis</i>	a stonefly	SU	G5		SNI	
<i>Nemoura arctica</i>	arctic forestfly	SU	GNR		SNI	
<i>Neoperla clymene</i>	coastal stone	SU	G5		SNI	
<i>Neoperla robisoni</i>	slender stone	SU	G4		SNI	

Scientific Name	Common Name	S-rank	G-rank	Listing Status	SGCN/SNI	Addl. Factors
<i>Neoperla stewarti</i>	a stonefly	SU	G5		SNI	
<i>Perlesta ephelida</i>	cloudy stonefly	SU	G5		SNI	
<i>Perlesta golconda</i>	two-lined stone	SU	GNR		SNI	
<i>Perlesta lagoi</i>	gulf stone	SU	G3		SNI	
<i>Perlesta ouabache</i>	Wabash stone	SU	GNR		SNI	
<i>Perlinella ephyre</i>	a stonefly	SU	G5		SNI	
<i>Podmosta macdunnoughi</i>	maritime forestfly	SU	G5		SNI	
<i>Prostoia similis</i>	a stonefly	SU	G5		SNI	
<i>Soyedina vallicularia</i>	valley forestfly	SU	G5		SNI	

Vascular Plants

SGCN species are listed first in alphabetical order by scientific name, followed by SNI species in alphabetical order by scientific name.

Physiognomy:

B = broadleaf evergreen

D = broadleaf deciduous

E = needleleaf evergreen

G = graminoid

H = forb

C = climber

K = stem succulent

X = epiphyte

F = floating aquatic

S = submerged aquatic

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Achillea alpina</i>	Siberian yarrow	Asteraceae	H	S1	G5	S-THR	SGCN
<i>Achnatherum hymenoides</i>	Indian ricegrass	Poaceae	G	S1	G5	S-END	SGCN
<i>Adlumia fungosa</i>	Allegheny vine	Fumariaceae	H	S1	G4	S-SPC	SGCN
<i>Agalinis auriculata</i>	eared false foxglove	Orobanchaceae	H	S2	G3	S-END	SGCN
<i>Agalinis gattingeri</i>	round-stemmed false foxglove	Orobanchaceae	H	S1	G4	S-END	SGCN
<i>Agastache nepetoides</i>	yellow giant hyssop	Lamiaceae	H	S2	G5		SGCN
<i>Agrostis hyemalis</i>	winter bentgrass	Poaceae	G	S1	G5	S-END	SGCN
<i>Alisma gramineum</i>	narrow-leaved water plantain	Alismataceae	H,S	S2	G5	S-SPC	SGCN
<i>Allium cernuum</i>	nodding wild onion	Liliaceae	H	S3	G5	S-SPC	SGCN
<i>Allium schoenoprasum</i>	wild chives	Liliaceae	H	S1	G5	S-END	SGCN
<i>Allium textile</i>	white wild onion	Liliaceae	H	S2	G5		SGCN
<i>Amerorchis rotundifolia</i>	small round-leaved orchis	Orchidaceae	H	S3	G5		SGCN
<i>Ammophila breviligulata</i>	beachgrass	Poaceae	G	S2	G5	S-THR	SGCN
<i>Anagallis minima</i>	chaffweed	Myrsinaceae	H	S1	G5		SGCN
<i>Androsace septentrionalis</i>	northern androsace	Primulaceae	H	S3	G5	S-SPC	SGCN
<i>Antennaria microphylla</i>	small-leaf cat's-foot	Asteraceae	H	S2	G5		SGCN
<i>Antennaria parvifolia</i>	small-leaved pussytoes	Asteraceae	H	S2	G5	S-SPC	SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Arisaema dracontium</i>	green dragon	Araceae	H	S3	G5	S-SPC	SGCN
<i>Aristida dichotoma</i>	churchmouse three-awn	Poaceae	G	S3	G5		SGCN
<i>Aristida longespica</i>	slimspike three-awn	Poaceae	G	S1	G5	S-END	SGCN
<i>Aristida purpurea</i>	red three-awn	Poaceae	G	S3	G5	S-SPC	SGCN
<i>Aristida tuberculosa</i>	seaside three-awn	Poaceae	G	S2	G5	S-THR	SGCN
<i>Arnica lonchophylla</i>	long-leaved arnica	Asteraceae	H	S2	G5	S-THR	SGCN
<i>Arnoglossum plantagineum</i>	tuberous Indian plantain	Asteraceae	H	S2	G4	S-THR	SGCN
<i>Arnoglossum reniforme</i>	great Indian plantain	Asteraceae	H	S2	G4	S-THR	SGCN
<i>Asclepias amplexicaulis</i>	clasping milkweed	Apocynaceae	H	S2	G5	S-THR	SGCN
<i>Asclepias hirtella</i>	prairie milkweed	Apocynaceae	H	S1	G5	S-THR	SGCN
<i>Asclepias lanuginosa</i>	woolly milkweed	Apocynaceae	H	S2	G4		SGCN
<i>Asclepias stenophylla</i>	narrow-leaved milkweed	Apocynaceae	H	S1	G4	S-END	SGCN
<i>Asclepias sullivantii</i>	Sullivant's milkweed	Apocynaceae	H	S2	G5	S-THR	SGCN
<i>Asplenium platyneuron</i>	ebony spleenwort	Aspleniaceae	H	S3	G5	S-SPC	SGCN
<i>Asplenium trichomanes</i>	maidenhair spleenwort	Aspleniaceae	H	S2	G5	S-THR	SGCN
<i>Astragalus alpinus</i>	alpine milk-vetch	Fabaceae	H	S1	G5	S-END	SGCN
<i>Astragalus flexuosus</i>	slender milk-vetch	Fabaceae	H	S2	G5	S-SPC	SGCN
<i>Astragalus missouriensis</i>	Missouri milk-vetch	Fabaceae	H	S3	G5	S-SPC	SGCN
<i>Astragalus neglectus</i>	Cooper's milk-vetch	Fabaceae	H	S3	G4		SGCN
<i>Aureolaria pedicularia</i>	fernleaf false foxglove	Chenopodiaceae	H	S1	G5	S-THR	SGCN
<i>Avenula hookeri</i>	spike oat	Poaceae	G	S2	G5	S-SPC	SGCN
<i>Bacopa rotundifolia</i>	waterhyssop	Plantaginaceae	H	S3	G5	S-THR	SGCN
<i>Baptisia bracteata</i>	cream wild indigo	Fabaceae	H	S3	G4	S-SPC	SGCN
<i>Baptisia lactea</i>	white wild indigo	Fabaceae	H	S3	G4	S-SPC	SGCN
<i>Bartonia virginica</i>	yellow bartonia	Gentianaceae	H	S1	G5	S-END	SGCN
<i>Berula erecta</i>	stream parsnip	Apiaceae	H	S2	G4	S-THR	SGCN
<i>Betula nigra</i>	swamp beggarticks	Betulaceae	D	S2	G5		SGCN
<i>Bidens discoidea</i>	discoïd beggarticks	Asteraceae	H	S3	G5	S-SPC	SGCN
<i>Bistorta vivipara</i>	alpine bistort	Polygonaceae	H	S2	G5	S-THR	SGCN
<i>Boechera collinsii</i>	Collins' rock cress	Brassicaceae	H	S1	G5	S-THR	SGCN
<i>Boechera stricta</i>	Drummond's rock cress	Brassicaceae	H	S2	GNR		SGCN
<i>Borodinia laevigata</i>	smooth rock cress	Brassicaceae	H	S3	G5	S-SPC	SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Botrychium angustisegmentum</i>	narrow triangle moonwort	Ophioglossaceae	H	S2	T4	S-THR	SGCN
<i>Botrychium ascendens</i>	upswept moonwort	Ophioglossaceae	H	S2	G4	S-END	SGCN
<i>Botrychium campestre</i> var. <i>campestre</i>	prairie moonwort	Ophioglossaceae	H	S3	G3	S-SPC	SGCN
<i>Botrychium campestre</i> var. <i>lineare</i>	slender moonwort	Ophioglossaceae	H	S1	G3	S-END	SGCN
<i>Botrychium crenulatum</i>	dainty moonwort	Ophioglossaceae	H	S1	G4		SGCN
<i>Botrychium gallicomontanum</i>	Frenchman's bluff moonwort	Ophioglossaceae	H	S1	G2	S-END	SGCN
<i>Botrychium michiganense</i>	Michigan moonwort	Ophioglossaceae	H	S2	G3		SGCN
<i>Botrychium minganense</i>	mingan moonwort	Ophioglossaceae	H	S2	G5	S-SPC	SGCN
<i>Botrychium mormo</i>	goblin fern	Ophioglossaceae	H	S1	G2	S-THR	SGCN
<i>Botrychium neolunaria</i>	new world moonwort	Ophioglossaceae	H	S2	GNR	S-THR	SGCN
<i>Botrychium pallidum</i>	pale moonwort	Ophioglossaceae	H	S3	G3	S-SPC	SGCN
<i>Botrychium simplex</i>	least moonwort	Ophioglossaceae	H	S4	G5	S-SPC	SGCN
<i>Botrychium spathulatum</i>	spatulate moonwort	Ophioglossaceae	H	S1	G3	S-END	SGCN
<i>Botrychium tenebrosum</i>	swamp moonwort	Ophioglossaceae	H	S3	T4		SGCN
<i>Buchloe dactyloides</i>	buffalo grass	Poaceae	G	S3	G4	S-SPC	SGCN
<i>Bulbostylis capillaris</i>	densetuft hairsedge	Cyperaceae	G	S3	G5		SGCN
<i>Calamagrostis montanensis</i>	plains reedgrass	Poaceae	G	S2	G5	S-SPC	SGCN
<i>Calamagrostis purpurascens</i>	purple reedgrass	Poaceae	G	S1	G5	S-END	SGCN
<i>Callirhoe triangulata</i>	clustered poppy-mallow	Malvaceae	H	S1	G3		SGCN
<i>Callitriche hermaphroditica</i>	autumnal water-starwort	Plantaginaceae	F,H	S1	G5		SGCN
<i>Callitriche heterophylla</i>	larger water starwort	Plantaginaceae	F,H	S2	G5	S-THR	SGCN
<i>Caltha natans</i>	floating marsh marigold	Ranunculaceae	H	S2	G5	S-END	SGCN
<i>Calypso bulbosa</i>	fairy slipper	Orchidaceae	H	S2	G5		SGCN
<i>Canadanthus modestus</i>	modest aster	Asteraceae	H	S3	G5		SGCN
<i>Cardamine douglassii</i>	purple cress	Brassicaceae	H	S1	G5		SGCN
<i>Cardamine pratensis</i>	cuckoo flower	Brassicaceae	H	S2	G5	S-THR	SGCN
<i>Carex annectens</i>	yellow-fruit sedge	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Carex aquatilis</i> var. <i>aquatilis</i>	water sedge	Cyperaceae	G	S1	T5		SGCN
<i>Carex capillaris</i>	hair-like sedge	Cyperaceae	G	S3	G5		SGCN
<i>Carex careyana</i>	Carey's sedge	Cyperaceae	G	S2	G4	S-END	SGCN
<i>Carex conjuncta</i>	jointed sedge	Cyperaceae	G	S1	G4	S-THR	SGCN
<i>Carex davisii</i>	Davis' sedge	Cyperaceae	G	S1	G4	S-THR	SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Carex debilis</i>	white-edge sedge	Cyperaceae	G	S3	G5		SGCN
<i>Carex exilis</i>	coastal sedge	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Carex filifolia</i>	thread-leaved sedge	Cyperaceae	G	S3	G5		SGCN
<i>Carex flava</i>	yellow sedge	Cyperaceae	G	S2	G5	S-SPC	SGCN
<i>Carex formosa</i>	handsome sedge	Cyperaceae	G	S1	G4	S-END	SGCN
<i>Carex garberi</i>	Garber's sedge	Cyperaceae	G	S1	G5	S-THR	SGCN
<i>Carex grayi</i>	Gray's sedge	Cyperaceae	G	S3	G4	S-SPC	SGCN
<i>Carex hallii</i>	Hall's sedge	Cyperaceae	G	S3	G4	S-SPC	SGCN
<i>Carex hookerana</i>	Hooker's sedge	Cyperaceae	G	S1	G4	S-SPC	SGCN
<i>Carex jamesii</i>	James' sedge	Cyperaceae	G	S3	G5	S-THR	SGCN
<i>Carex katahdinensis</i>	katahdin sedge	Cyperaceae	G	S2	G2		SGCN
<i>Carex laevivaginata</i>	smooth-sheathed sedge	Cyperaceae	G	S2	G5	S-THR	SGCN
<i>Carex laxiculmis</i>	spreading sedge	Cyperaceae	G	S3	G5	S-THR	SGCN
<i>Carex media</i>	intermediate sedge	Cyperaceae	G	S3	T5	S-SPC	SGCN
<i>Carex michauxiana</i>	Michaux's sedge	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Carex muskingumensis</i>	Muskingum sedge	Cyperaceae	G	S3	G4	S-SPC	SGCN
<i>Carex novae-angliae</i>	New England sedge	Cyperaceae	G	S2	G5	S-THR	SGCN
<i>Carex obtusata</i>	blunt sedge	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Carex ormostachya</i>	necklace sedge	Cyperaceae	G	S2	G4	S-SPC	SGCN
<i>Carex pallescens</i>	pale sedge	Cyperaceae	G	S2	G5	S-END	SGCN
<i>Carex plantaginea</i>	plantain-leaved sedge	Cyperaceae	G	S1	G5	S-END	SGCN
<i>Carex praticola</i>	prairie-dweller sedge	Cyperaceae	G	S2	G5	S-SPC	SGCN
<i>Carex rossii</i>	Ross' sedge	Cyperaceae	G	S1	G5	S-THR	SGCN
<i>Carex scirpoidea</i>	northern single-spike sedge	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Carex sterilis</i>	sterile sedge	Cyperaceae	G	S2	G4	S-THR	SGCN
<i>Carex supina</i>	weak arctic sedge	Cyperaceae	G	S1	G5	S-END	SGCN
<i>Carex torreyi</i>	Torrey's sedge	Cyperaceae	G	S2	G4		SGCN
<i>Carex trichocarpa</i>	hairy-fruited sedge	Cyperaceae	G	S3	G4		SGCN
<i>Carex typhina</i>	cattail sedge	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Carex xerantica</i>	dry sedge	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Castilleja septentrionalis</i>	northern paintbrush	Orobanchaceae	H	S1	G5	S-END	SGCN
<i>Cerastium brachypodium</i>	mouse-ear chickweed	Caryophyllaceae	H	S2	G5		SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Ceratophyllum echinatum</i>	spiny hornwort	Ceratophyllaceae	H,S	S3	G4		SGCN
<i>Chamaerhodos erecta</i>	Nuttall's groundrose	Rosaceae	H	S3	G5		SGCN
<i>Chenopodium standleyanum</i>	Standley's goosefoot	Chenopodiaceae	H	S2	G5		SGCN
<i>Chrysosplenium iowense</i>	Iowa golden saxifrage	Saxifragaceae	H	S1	G4	S-END	SGCN
<i>Cladium mariscoides</i>	twig rush	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Collomia linearis</i>	narrowleaf collomia	Polemoniaceae	H	S2	G5		SGCN
<i>Commelina erecta</i>	slender dayflower	Commelinaceae	H	S1	G5	S-END	SGCN
<i>Conyza ramosissima</i>	dwarf conyza	Asteraceae	H	S1	G5		SGCN
<i>Corispermum americanum</i>	American bugseed	Chenopodiaceae	H	S1	G5		SGCN
<i>Cornus drummondii</i>	rough-leaved dogwood	Cornaceae	D	S1	G5		SGCN
<i>Corydalis micrantha</i>	small-flower corydalis	Fumariaceae	H	S1	G5		SGCN
<i>Coryphantha vivipara</i>	ball cactus	Cactaceae	K	S1	G5	S-END	SGCN
<i>Crassula aquatica</i>	water pygmyweed	Crassulaceae	H	S2	G5	S-THR	SGCN
<i>Crataegus calpodendron</i>	late hawthorn	Rosaceae	D	S2	G5	S-SPC	SGCN
<i>Crataegus douglasii</i>	black hawthorn	Rosaceae	D	S3	G5	S-SPC	SGCN
<i>Crataegus laurentiana</i>	Laurentian hawthorn	Rosaceae	D	S3	GNR		SGCN
<i>Crataegus succulenta</i>	fleshy hawthorn	Rosaceae	D	S2	G5		SGCN
<i>Crocanthemum canadense</i>	Canada frostweed	Cistaceae	H	S3	G5	S-SPC	SGCN
<i>Crotalaria sagittalis</i>	rattlebox	Fabaceae	H	S3	G5	S-SPC	SGCN
<i>Cuscuta cephalanthi</i>	buttonbush dodder	Convolvulaceae	H	S2	G5		SGCN
<i>Cuscuta coryli</i>	hazel dodder	Convolvulaceae	H	S2	G5		SGCN
<i>Cyperus acuminatus</i>	short-pointed umbrella sedge	Cyperaceae	G	S2	G5	S-THR	SGCN
<i>Cyperus houghtonii</i>	Houghton's cyperus	Cyperaceae	G	S3	G4		SGCN
<i>Cypripedium arietinum</i>	ram's head orchid	Orchidaceae	H	S2	G3	S-THR	SGCN
<i>Cypripedium candidum</i>	small white lady's-slipper	Orchidaceae	H	S3S4	G4	S-SPC	SGCN
<i>Cystopteris laurentiana</i>	Laurentian bladder fern	Cystopteridaceae	H	S2	G3		SGCN
<i>Cystopteris tennesseensis</i>	Tennessee bladder fern	Cystopteridaceae	H	S3	G5		SGCN
<i>Dalea candida</i> var. <i>oligophylla</i>	western white prairie-clover	Fabaceae	H	S3	T5	S-SPC	SGCN
<i>Dalea leporina</i>	foxtail prairie-clover	Fabaceae	H	S2	G5		SGCN
<i>Decodon verticillatus</i>	water-willow	Lythraceae	D	S3	G5	S-SPC	SGCN
<i>Dendrolycopodium obscurum</i>	flat-branched tree club-moss	Lycopodiaceae	H	S2	G5		SGCN
<i>Deparia acrostichoides</i>	silvery spleenwort	Athyriaceae	H	S3	G5	S-SPC	SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Deschampsia flexuosa</i>	slender hair grass	Poaceae	G	S1	G5	S-THR	SGCN
<i>Descurainia pinnata</i>	pinnate tansy-mustard	Brassicaceae	H	S2	G5		SGCN
<i>Desmanthus illinoensis</i>	prairie mimosa	Fabaceae	H	S3	G5	S-SPC	SGCN
<i>Desmodium cuspidatum</i>	big tick trefoil	Fabaceae	H	S2	G5	S-THR	SGCN
<i>Desmodium illinoense</i>	Illinois tick-trefoil	Fabaceae	H	S3	G5		SGCN
<i>Desmodium nudiflorum</i>	stemless tick trefoil	Fabaceae	H	S3	G5	S-THR	SGCN
<i>Diarrhena obovata</i>	obovate beakgrain	Poaceae	G	S1	G4	S-END	SGCN
<i>Dicentra canadensis</i>	squirrel corn	Fumariaceae	H	S3	G5	S-SPC	SGCN
<i>Dodecatheon amethystinum</i>	jeweled shooting star	Primulaceae	H	S3	G4		SGCN
<i>Dodecatheon meadia</i>	prairie shooting star	Primulaceae	H	S1	G5	S-END	SGCN
<i>Draba arabisans</i>	Arabian whitlow grass	Brassicaceae	H	S3	G4	S-SPC	SGCN
<i>Draba cana</i>	hoary whitlow grass	Brassicaceae	H	S1	G5	S-END	SGCN
<i>Draba norvegica</i>	Norwegian whitlow grass	Brassicaceae	H	S1	G5	S-END	SGCN
<i>Drosera anglica</i>	English sundew	Droseraceae	H	S3	G5	S-SPC	SGCN
<i>Drosera linearis</i>	linear-leaved sundew	Droseraceae	H	S2	G4	S-SPC	SGCN
<i>Dryopteris filix-mas</i>	male fern	Dryopteridaceae	H	S1	G5		SGCN
<i>Dryopteris goldieana</i>	Goldie's fern	Dryopteridaceae	H	S3	G4	S-SPC	SGCN
<i>Dryopteris marginalis</i>	marginal shield fern	Dryopteridaceae	H	S1	G5	S-END	SGCN
<i>Eclipta prostrata</i>	yerba de tajo	Asteraceae	H	S3	G5		SGCN
<i>Elatine triandra</i>	three-stamened waterwort	Elatinaceae	H,S	S3	G5	S-SPC	SGCN
<i>Eleocharis coloradoensis</i>	dwarf spikerush	Cyperaceae	G	S1	G4	S-SPC	SGCN
<i>Eleocharis engelmannii</i>	Engelmann's spikerush	Cyperaceae	G	S3	G4		SGCN
<i>Eleocharis flavescens</i> var. <i>olivacea</i>	pale spikerush	Cyperaceae	G	S2	G5		SGCN
<i>Eleocharis mamillata</i>	mamillate spike-rush	Cyperaceae	G	S3	G5		SGCN
<i>Eleocharis nitida</i>	neat spikerush	Cyperaceae	G	S3	G4	S-SPC	SGCN
<i>Eleocharis quinqueflora</i>	few-flowered spikerush	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Eleocharis robbinsii</i>	Robbins' spikerush	Cyperaceae	G	S3	G4	S-THR	SGCN
<i>Eleocharis rostellata</i>	beaked spikerush	Cyperaceae	G	S2	G5	S-THR	SGCN
<i>Eleocharis wolfii</i>	Wolf's spikerush	Cyperaceae	G	S2	G4	S-END	SGCN
<i>Elodea bifoliata</i>	two leaf waterweed	Hydrocharitaceae	H,S	SH	G4	S-END	SGCN
<i>Elymus curvatus</i>	beardless wild rye	Poaceae	G	S2	G4		SGCN
<i>Elymus diversiglumis</i>	interrupted wild rye	Poaceae	G	S3S4	G3		SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Elymus riparius</i>	riverbank wildrye	Poaceae	G	S3	G5		SGCN
<i>Empetrum atropurpureum</i>	purple crowberry	Ericaceae	B	S1	T5	S-END	SGCN
<i>Empetrum nigrum</i>	black crowberry	Ericaceae	B	SH	G5	S-END	SGCN
<i>Eragrostis frankii</i>	Frank's lovegrass	Poaceae	G	S2	G5		SGCN
<i>Eragrostis trichodes</i>	sand lovegrass	Poaceae	G	S2	G5		SGCN
<i>Erigeron acris</i>	bitter fleabane	Asteraceae	H	S2	G5	S-END	SGCN
<i>Erigeron lonchophyllus</i>	short ray fleabane	Asteraceae	H	S2	G5	S-THR	SGCN
<i>Eryngium yuccifolium</i>	rattlesnake master	Apiaceae	H	S3	G5	S-SPC	SGCN
<i>Erythronium propullans</i>	dwarf trout lily	Liliaceae	H	S1	G1	S-END, F-END	SGCN
<i>Eupatorium altissimum</i>	tall boneset	Asteraceae	H	S2	G5		SGCN
<i>Eupatorium sessilifolium</i>	upland boneset	Asteraceae	H	S2	G5	S-THR	SGCN
<i>Euphorbia dentata</i>	hairy-fruit spurge	Euphorbiaceae	H	S2	G5		SGCN
<i>Euphorbia missurica</i>	Missouri spurge	Euphorbiaceae	H	SH	G5	S-SPC	SGCN
<i>Euphorbia spathulata</i>	reticulate-seed spurge	Euphorbiaceae	H	S1	G5		SGCN
<i>Euphrasia hudsoniana</i>	Hudson Bay eyebright	Orobanchaceae	H	S2	G5	S-SPC	SGCN
<i>Fimbristylis autumnalis</i>	autumn fimbry	Cyperaceae	G	S3	G5	S-SPC	SGCN
<i>Fimbristylis puberula</i>	hairy fimbry	Cyperaceae	G	S1	G5	S-END	SGCN
<i>Floerkea proserpinacoides</i>	false mermaid	Limnanthaceae	H	S2S3	G5	S-THR	SGCN
<i>Gaillardia aristata</i>	blanketflower	Asteraceae	H	S3	G5	S-SPC	SGCN
<i>Galium circaeans</i>	wild licorice	Rubiaceae	H	S2	G5		SGCN
<i>Galium palustre</i>	marsh bedstraw	Rubiaceae	H	S1	G5		SGCN
<i>Gaylussacia baccata</i>	black huckleberry	Ericaceae	D	S2	G5	S-THR	SGCN
<i>Gentiana affinis</i>	northern gentian	Gentianaceae	H	S3	G5	S-SPC	SGCN
<i>Gentianella amarella</i> ssp. <i>acuta</i>	felwort	Gentianaceae	H	S3	G5	S-SPC	SGCN
<i>Gentianella quinquefolia</i>	stiff gentian	Gentianaceae	H	S2	G5		SGCN
<i>Geranium carolinianum</i>	Carolina crane's-bill	Geraniaceae	H	S2	G5		SGCN
<i>Geum laciniatum</i>	rough avens	Rosaceae	H	S3	G5		SGCN
<i>Gleditsia triacanthos</i>	honeylocust	Fabaceae	D	S1	G5		SGCN
<i>Grindelia hirsutula</i>	hairy gumweed	Asteraceae	H	S1	T5		SGCN
<i>Gymnocarpium robertianum</i>	northern oak fern	Cystopteridaceae	H	S3	G5	S-SPC	SGCN
<i>Gymnocladus dioica</i>	Kentucky coffee tree	Caesalpiniaceae	D	S3	G5	S-SPC	SGCN
<i>Hamamelis virginiana</i>	witch-hazel	Hamamelidaceae	D	S2	G5	S-THR	SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Hasteola suaveolens</i>	sweet-smelling indian plantain	Asteraceae	H	S2	G4	S-END	SGCN
<i>Helianthus nuttallii</i>	Nuttall's sunflower	Asteraceae	H	S3	G5	S-SPC	SGCN
<i>Hesperostipa curisetia</i>	small porcupine grass	Poaceae	G	S1	G5		SGCN
<i>Heteranthera limosa</i>	mud plantain	Pontederiaceae	H,S	S1	G5	S-THR	SGCN
<i>Hieracium longipilum</i>	long-bearded hawkweed	Asteraceae	H	S3	G4		SGCN
<i>Homalosorus pycnocarpus</i>	narrow-leaved spleenwort	Diplaziopsidaceae	H	S2	G5	S-THR	SGCN
<i>Hordeum pusillum</i>	little barley	Asteraceae	G	S2	G5		SGCN
<i>Hudsonia tomentosa</i>	beach heather	Cistaceae	B	S2	G5	S-THR	SGCN
<i>Huperzia appressa</i>	mountain fir moss	Lycopodiaceae	H	S2	G5	S-SPC	SGCN
<i>Huperzia porophila</i>	rock firmoss	Lycopodiaceae	H	S2	G4	S-THR	SGCN
<i>Hybanthus concolor</i>	eastern green-violet	Violaceae	H	S1	G5	S-END	SGCN
<i>Hydrastis canadensis</i>	goldenseal	Ranunculaceae	H	S1	G3	S-END	SGCN
<i>Hydrocotyle americana</i>	American water-pennywort	Araliaceae	H	S3	G5	S-SPC	SGCN
<i>Hypericum ellipticum</i>	pale St. John's-wort	Hypericaceae	H	S2	G5		SGCN
<i>Hypericum punctatum</i>	common St. John's-wort	Hypericaceae	H	S2	G5		SGCN
<i>Iodanthus pinnatifidus</i>	purple rocket	Brassicaceae	H	S1	G5	S-END	SGCN
<i>Isanthus brachiatus</i>	false pennyroyal	Lamiaceae	H	S2	G5		SGCN
<i>Isoetes melanopoda</i>	prairie quillwort	Isoetaceae	H,S	S1	G5	S-END	SGCN
<i>Jeffersonia diphylla</i>	twinleaf	Berberidaceae	H	S3	G5	S-SPC	SGCN
<i>Juglans cinerea</i>	butternut	Juglandaceae	D	S2	G3	S-END	SGCN
<i>Juncus anthelatus</i>	greater poverty rush	Juncaceae	G	S1	GNR		SGCN
<i>Juncus articulatus</i>	jointed rush	Juncaceae	G	S2	G5	S-END	SGCN
<i>Juncus marginatus</i>	marginated rush	Juncaceae	G	S1	G5	S-END	SGCN
<i>Juncus stygius</i>	moor rush	Juncaceae	G	S3	G5	S-SPC	SGCN
<i>Juncus subtilis</i>	slender rush	Juncaceae	G	S1	G5	S-END	SGCN
<i>Juniperus horizontalis</i>	creeping juniper	Cupressaceae	E	S3	G5	S-SPC	SGCN
<i>Lactuca floridana</i>	Florida lettuce	Asteraceae	H	S3	G5		SGCN
<i>Lechea tenuifolia</i>	narrow-leaved pinweed	Cistaceae	H	S1	G5	S-END	SGCN
<i>Leersia lenticularis</i>	catchfly grass	Poaceae	G	S2	G5	S-THR	SGCN
<i>Lepidium virginicum</i>	poor-man's pepper-grass	Brassicaceae	H	S2	G5		SGCN
<i>Lespedeza leptostachya</i>	prairie bush clover	Fabaceae	H	S2	G3	S-THR, F-THR	SGCN
<i>Leucophysalis grandiflora</i>	dwarf ground cherry	Solanaceae	H	S2S3	G4		SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Limosella aquatica</i>	mudwort	Scrophulariaceae	H	S3	G5	S-SPC	SGCN
<i>Linum lewisii</i>	prairie flax	Linaceae	H	S1	G5		SGCN
<i>Listera auriculata</i>	auricled twayblade	Orchidaceae	H	S1	G3	S-END	SGCN
<i>Listera convallarioides</i>	broad-leaved twayblade	Orchidaceae	H	SH	G5	S-SPC	SGCN
<i>Littorella americana</i>	American shore plantain	Plantaginaceae	H,S	S4	G5	S-SPC	SGCN
<i>Lomatium orientale</i>	desert parsley	Apiaceae	H	S2	G5		SGCN
<i>Lotus purshianus</i>		Fabaceae	H	S2	G4		SGCN
<i>Lupinus perennis</i>	wild lupine	Fabaceae	H	S2	G5		SGCN
<i>Luzula parviflora</i>	small-flowered woodrush	Juncaceae	G	S2S3	G5	S-THR	SGCN
<i>Lycopus virginicus</i>	Virginia water horehound	Lamiaceae	H	S4	G5		SGCN
<i>Lysimachia lanceolata</i>	lance-leaved loosestrife	Myrsinaceae	H	S1	G5		SGCN
<i>Lysimachia maritima</i>	sea milkwort	Myrsinaceae	H	S1	G5	S-END	SGCN
<i>Lysimachia quadrifolia</i>	whorled loosestrife	Myrsinaceae	H	S2	G5	S-SPC	SGCN
<i>Malaxis paludosa</i>	bog adder's mouth	Orchidaceae	H	S1	G3	S-END	SGCN
<i>Marsilea vestita</i>	hairy watercress	Marsileaceae	H	S1	G5	S-END	SGCN
<i>Melica nitens</i>	three-flowered melic	Poaceae	G	S1	G5	S-THR	SGCN
<i>Minuartia dawsonensis</i>	rock sandwort	Caryophyllaceae	H	S2	G5	S-THR	SGCN
<i>Moehringia macrophylla</i>	large-leaved sandwort	Caryophyllaceae	H	S2	G5	S-THR	SGCN
<i>Montia chamissoi</i>	montia	Portulacaceae	H	S1	G5	S-END	SGCN
<i>Morus rubra</i>	red mulberry	Moraceae	D	S1	G5		SGCN
<i>Muhlenbergia sylvatica</i>	woodland muhly	Poaceae	G	S1	G5		SGCN
<i>Muhlenbergia uniflora</i>	one-flowered muhly	Poaceae	G	S3	G5	S-SPC	SGCN
<i>Myosotis verna</i>	forget-me-not	Boraginaceae	H	S2	G5		SGCN
<i>Myriophyllum heterophyllum</i>	broadleaf water milfoil	Haloragaceae	H,S	S1	G5	S-SPC	SGCN
<i>Najas gracillima</i>	slender naiad	Najadaceae	H,S	S3	G5	S-SPC	SGCN
<i>Najas guadalupensis ssp. olivacea</i>	olive-colored southern naiad	Najadaceae	H,S	SU	T4	S-SPC	SGCN
<i>Najas marina</i>	sea naiad	Najadaceae	H,S	S2	G5	S-SPC	SGCN
<i>Napaea dioica</i>	glade mallow	Malvaceae	H	S3	G4	S-THR	SGCN
<i>Nothocalais cuspidata</i>	prairie false dandelion	Asteraceae	H	S2	G5		SGCN
<i>Nuttallanthus canadensis</i>	old field toadflax	Plantaginaceae	H	S2S3	G5	S-SPC	SGCN
<i>Nymphaea leibergii</i>	small white waterlily	Nymphaeaceae	F,H	S2	G5	S-THR	SGCN
<i>Oenothera nuttallii</i>	Nuttall's evening primrose	Onagraceae	H	S2	G5		SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Oenothera rhombipetala</i>	rhombic evening primrose	Onagraceae	H	S2	G4	S-SPC	SGCN
<i>Ophioglossum pusillum</i>	adder's tongue	Ophioglossaceae	H	S3	G5		SGCN
<i>Opuntia macrorhiza</i>	devil's tongue	Cactaceae	K	S2S3	G5	S-SPC	SGCN
<i>Orobanche fasciculata</i>	clustered broomrape	Orobanchaceae	H	S2	G4	S-THR	SGCN
<i>Orobanche ludoviciana</i>	Louisiana broomrape	Orobanchaceae	H	S2	G5	S-THR	SGCN
<i>Orobanche uniflora</i>	one-flowered broomrape	Orobanchaceae	H	S2	G5	S-THR	SGCN
<i>Orthocarpus luteus</i>	owl clover	Orobanchaceae	H	S2	G5		SGCN
<i>Osmorhiza berteroi</i>	Chilean sweet cicely	Apiaceae	H	S1	G5	S-END	SGCN
<i>Osmorhiza depauperata</i>	blunt-fruited sweet cicely	Apiaceae	H	S3	G5	S-SPC	SGCN
<i>Oxypolis rigidior</i>	cowbane	Apiaceae	H	S3	G5		SGCN
<i>Oxytropis viscida</i>	sticky locoweed	Fabaceae	H	S1	G5	S-END	SGCN
<i>Packera cana</i>	gray ragwort	Asteraceae	H	S2	G5	S-END	SGCN
<i>Packera indecora</i>	elegant groundsel	Asteraceae	H	S1	G5	S-END	SGCN
<i>Panax quinquefolius</i>	American ginseng	Araliaceae	H	S2	G3	S-SPC	SGCN
<i>Panax trifolius</i>	dwarf ginseng	Araliaceae	H	S2S3	G5		SGCN
<i>Paronychia canadensis</i>	Canada forked chickweed	Caryophyllaceae	H	S1	G5	S-END	SGCN
<i>Paronychia fastigiata</i>	forked chickweed	Caryophyllaceae	H	S1	G5	S-END	SGCN
<i>Parthenium integrifolium</i>	wild quinine	Asteraceae	H	S2	G5	S-END	SGCN
<i>Pellaea atropurpurea</i>	purple cliff brake	Pteridaceae	H	S2	G5	S-SPC	SGCN
<i>Penstemon albidus</i>	white-flower beardtongue	Plantaginaceae	H	S2	G5		SGCN
<i>Persicaria careyi</i>	Carey's smartweed	Polygonaceae	H	S1	G4	S-SPC	SGCN
<i>Phacelia franklinii</i>	Franklin's phacelia	Hydrophyllaceae	H	S2	G5	S-THR	SGCN
<i>Phegopteris hexagonoptera</i>	broad beech fern	Thelypteridaceae	H	S1	G5	S-END	SGCN
<i>Phemeranthus rugospermus</i>	rough-seeded fameflower	Portulacaceae	H	S2	G3	S-THR	SGCN
<i>Phlox maculata</i>	wild sweetwilliam	Polemoniaceae	H	S3	G5	S-SPC	SGCN
<i>Physaria ludoviciana</i>	bladderpod	Brassicaceae	H	S2	G5	S-END	SGCN
<i>Pinguicula vulgaris</i>	butterwort	Lentibulariaceae	H	S3	G5	S-SPC	SGCN
<i>Piptatherum canadense</i>	Canadian ricegrass	Poaceae	G	S2	G4	S-THR	SGCN
<i>Plagiobothrys scouleri</i>	Scouler's popcornflower	Boraginaceae	H	S3	G5	S-SPC	SGCN
<i>Plantago elongata</i>	slender plantain	Plantaginaceae	H	S2	G4	S-SPC	SGCN
<i>Platanthera clavellata</i>	small green wood orchid	Orchidaceae	H	S3S4	G5	S-SPC	SGCN
<i>Platanthera dilatata</i>	tall white bog orchid	Orchidaceae	H	S3	G5		SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Platanthera flava</i> var. <i>herbiola</i>	southern rein-orchid	Orchidaceae	H	S2	G4	S-THR	SGCN
<i>Platanthera hookeri</i>	Hooker's orchid	Orchidaceae	H	S3	G4		SGCN
<i>Platanthera praeclara</i>	western prairie fringed orchid	Orchidaceae	H	S1S2	G3	S-END, F-THR	SGCN
<i>Poa arida</i>	plains bluegrass	Poaceae	G	S3	G5		SGCN
<i>Poa paludigena</i>	bog bluegrass	Poaceae	G	S2	G3	S-THR	SGCN
<i>Poa sylvestris</i>	woodland bluegrass	Poaceae	G	S1	G5		SGCN
<i>Poa wolfii</i>	Wolf's bluegrass	Poaceae	G	SH	G4	S-SPC	SGCN
<i>Polanisia jamesii</i>	James' polanisia	Cleomaceae	H	S1	G5	S-END	SGCN
<i>Polemonium lacustre</i>	Great Lakes Jacob's-ladder	Polemoniaceae	H	S1	G1	S-END	SGCN
<i>Polygala cruciata</i>	cross-leaved milkwort	Polygalaceae	H	S1	G5	S-END	SGCN
<i>Polygonum erectum</i>	erect knotweed	Polygonaceae	H	S1	G5		SGCN
<i>Polystichum acrostichoides</i>	christmas fern	Dryopteridaceae	H	S1	G5	S-END	SGCN
<i>Polystichum braunii</i>	Braun's holly fern	Dryopteridaceae	H	S2	G5	S-THR	SGCN
<i>Polytaenia nuttallii</i>	prairie parsley	Apiaceae	H	SH	G5	S-SPC	SGCN
<i>Potamogeton bicupulatus</i>	snailseed pondweed	Potamogetonaceae	F,H	S2	G4	S-END	SGCN
<i>Potamogeton confervoides</i>	algae-like pondweed	Potamogetonaceae	F,H	S1	G5	S-END	SGCN
<i>Potamogeton diversifolius</i>	diverse-leaved pondweed	Potamogetonaceae	F,H	S1	G5	S-END	SGCN
<i>Potamogeton oakesianus</i>	Oakes' pondweed	Potamogetonaceae	F,H	S2	G5	S-END	SGCN
<i>Potamogeton pulcher</i>	spotted pondweed	Potamogetonaceae	F,H	S1	G5	S-END	SGCN
<i>Potamogeton pusillus</i> ssp. <i>pusillus</i>	slender pondweed	Potamogetonaceae	F,H	S2	T5		SGCN
<i>Potentilla bipinnatifida</i>	tansy cinquefoil	Rosaceae	H	S2	GNR		SGCN
<i>Potentilla hippiana</i>	woolly cinquefoil	Rosaceae	H	S1	G5		SGCN
<i>Potentilla lasiodonta</i>	sandhills cinquefoil	Rosaceae	H	S1	G3		SGCN
<i>Potentilla rivalis</i>	brook cinquefoil	Rosaceae	H	S1	G5		SGCN
<i>Potentilla supina</i>	bushy cinquefoil	Rosaceae	H	S1	G5		SGCN
<i>Prenanthes aspera</i>	hairy rattlesnakeroot	Asteraceae	H	S2	G4		SGCN
<i>Prosartes trachycarpa</i>	rough-fruited fairybells	Liliaceae	H	S1	G5	S-END	SGCN
<i>Pseudognaphalium macounii</i>	clammy cudweed	Asteraceae	H	S1	G5		SGCN
<i>Psoralidium tenuiflorum</i>	slender-leaved scurfpea	Fabaceae	H	S1	G5	S-END	SGCN
<i>Puccinellia nuttalliana</i>	alkali grass	Poaceae	G	S3	G5		SGCN
<i>Pyrola minor</i>	small shinleaf	Ericaceae	H	S3	G5	S-SPC	SGCN
<i>Quercus bicolor</i>	swamp white oak	Fagaceae	D	S2	G5	S-SPC	SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Ranunculus lapponicus</i>	Lapland buttercup	Ranunculaceae	H	S3	G5	S-SPC	SGCN
<i>Ranunculus macounii</i>	Macoun's buttercup	Ranunculaceae	H	S2	G5		SGCN
<i>Rhodiola integrifolia</i>	Leedy's roseroot	Crassulaceae	H	S1	G5	S-END, F-THR	SGCN
<i>Rhynchospora capillacea</i>	hair-like beak rush	Cyperaceae	G	S2	G4	S-THR	SGCN
<i>Rhynchospora capitellata</i>	brownish beak sedge	Cyperaceae	G	S1	G5		SGCN
<i>Rorippa curvipes</i>	bluntleaf yellowcress	Brassicaceae	H	S2	G5		SGCN
<i>Rorippa sessiliflora</i>	sessile-flowered yellow cress	Brassicaceae	H	S2	G5	S-SPC	SGCN
<i>Rorippa sinuata</i>	spreading yellow cress	Brassicaceae	H	S1	G5		SGCN
<i>Rotala ramosior</i>	toothcup	Lythraceae	H	S1	G5	S-THR	SGCN
<i>Rubus acridens</i>	Kennedy's blackberry	Rosaceae	D	S2	G4		SGCN
<i>Rubus chamaemorus</i>	cloudberry	Rosaceae	D	S2	G5	S-THR	SGCN
<i>Rubus dissensus</i>		Rosaceae	D	S2	GNR		SGCN
<i>Rubus fulleri</i>	a bristle-berry	Rosaceae	D	S2	G4	S-THR	SGCN
<i>Rubus groutianus</i>		Rosaceae	D	S1	GNR		SGCN
<i>Rubus missouricus</i>	Missouri bristle-berry	Rosaceae	D	S1	G5	S-END	SGCN
<i>Rubus multiflorus</i>	Kinnickinnick dewberry	Rosaceae	D	SU	G5	S-SPC	SGCN
<i>Rubus quaesitus</i>	Prince Edward Island blackberry	Rosaceae	D	S1	GU	S-SPC	SGCN
<i>Rubus regionalis</i>		Rosaceae	D	S2	G5		SGCN
<i>Rubus satis</i>	dewberry	Rosaceae	D	S2	G4		SGCN
<i>Rubus semisetosus</i>	swamp blackberry	Rosaceae	D	SU	G5	S-THR	SGCN
<i>Rubus stipulatus</i>	a bristle-berry	Rosaceae	D	S1	G4	S-END	SGCN
<i>Rubus vermontanus</i>	Vermont bristle-berry	Rosaceae	D	SU	G5	S-SPC	SGCN
<i>Rubus wheeleri</i>	Wheeler's blackberry	Rosaceae	D	S1	G4		SGCN
<i>Rudbeckia subtomentosa</i>	sweet coneflower	Asteraceae	H	S1	G5		SGCN
<i>Rudbeckia triloba</i>	three-leaved coneflower	Asteraceae	H	S3	G5	S-THR	SGCN
<i>Ruellia humilis</i>	wild petunia	Acanthaceae	H	S1	G5	S-SPC	SGCN
<i>Rumex altissimus</i>	tall dock	Polygonaceae	H	S2	G5		SGCN
<i>Rumex verticillatus</i>	swamp dock	Polygonaceae	H	S1	G5		SGCN
<i>Ruppia cirrhosa</i>	spiral ditchgrass	Ruppiaceae	H,S	S2	G5	S-SPC	SGCN
<i>Sagina nodosa</i>	knotty pearlwort	Caryophyllaceae	H	S1	G5	S-END	SGCN
<i>Sagittaria brevirostra</i>	short-beaked arrowhead	Alismataceae	H	S1	G5	S-END	SGCN
<i>Sagittaria montevidensis</i>	long-lobed arrowhead	Alismataceae	H	S2	T4		SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Salicornia rubra</i>	red saltwort	Chenopodiaceae	H	S2	G5	S-THR	SGCN
<i>Salix maccalliana</i>	Mccalla's willow	Salicaceae	D	S3	G5	S-SPC	SGCN
<i>Salix pellita</i>	satiny willow	Salicaceae	D	S2	G5	S-THR	SGCN
<i>Salix pseudomonticola</i>	false mountain willow	Salicaceae	D	S2	G5	S-SPC	SGCN
<i>Sanicula trifoliata</i>	beaked snakeroot	Apiaceae	H	S3	G4	S-SPC	SGCN
<i>Satureja vulgaris</i>	wild basil	Lamiaceae	H	S3	G5		SGCN
<i>Saxifraga cernua</i>	nodding saxifrage	Saxifragaceae	H	S1	G5	S-END	SGCN
<i>Saxifraga paniculata</i>	encrusted saxifrage	Saxifragaceae	H	S2	G5	S-SPC	SGCN
<i>Sceptridium oneidense</i>	blunt-lobed grapefern	Ophioglossaceae	H	S2	G4	S-THR	SGCN
<i>Sceptridium rugulosum</i>	St. Lawrence grapefern	Ophioglossaceae	H	S3	G3	S-SPC	SGCN
<i>Schedonnardus paniculatus</i>	tumble grass	Poaceae	G	S2	G5	S-SPC	SGCN
<i>Schoenoplectiella purshiana</i> var. <i>purshiana</i>	weak-stalk bulrush	Cyperaceae	G	S3	T4		SGCN
<i>Schoenoplectiella purshiana</i> var. <i>williamsii</i>	weak-stalk bulrush	Cyperaceae	G	S1	TNR		SGCN
<i>Schoenoplectiella smithii</i> var. <i>smithii</i>	blunt-scale bulrush	Cyperaceae	G	S1	TNR		SGCN
<i>Scleria triglomerata</i>	tall nutrush	Cyperaceae	G	S1	G5	S-END	SGCN
<i>Scleria verticillata</i>	whorled nutrush	Cyperaceae	G	S2	G5	S-THR	SGCN
<i>Scutellaria ovata</i>	ovate-leaved skullcap	Lamiaceae	H	S2	G5	S-THR	SGCN
<i>Scutellaria parvula</i>	small skullcap	Lamiaceae	H	S2	G4		SGCN
<i>Selaginella selaginoides</i>	northern spikemoss	Selaginellaceae	H	S1	G5	S-END	SGCN
<i>Senecio integerrimus</i>	entireleaf ragwort	Asteraceae	H	S2	G5		SGCN
<i>Shepherdia argentea</i>	silver buffaloberry	Elaeagnaceae	D	S2	G5		SGCN
<i>Shepherdia canadensis</i>	soapberry	Elaeagnaceae	D	S3	G5	S-SPC	SGCN
<i>Shinnersoseris rostrata</i>	annual skeletonweed	Asteraceae	H	S2	G5	S-THR	SGCN
<i>Silene drummondii</i>	Drummond's campion	Caryophyllaceae	H	S3	G5	S-SPC	SGCN
<i>Silene hitchguirei</i>	mountain campion	Caryophyllaceae	H	S1			SGCN
<i>Silene nivea</i>	snowy campion	Caryophyllaceae	H	S3	G4	S-THR	SGCN
<i>Silphium laciniatum</i>	compass-plant	Asteraceae	H	S3	G5		SGCN
<i>Solidago mollis</i>	soft goldenrod	Asteraceae	H	S3	G5	S-SPC	SGCN
<i>Solidago sciaphila</i>	cliff goldenrod	Asteraceae	H	S3	G3		SGCN
<i>Sparganium androcladum</i>	branching bur-reed	Sparganiaceae	H	S2	G4		SGCN
<i>Sparganium glomeratum</i>	clustered bur-reed	Sparganiaceae	H	S3	G4		SGCN
<i>Spiranthes casei</i>	Case's ladies' tresses	Orchidaceae	H	S1	G4	S-THR	SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Spiranthes magnicamporum</i>	Great Plains ladies'-tresses	Orchidaceae	H	S4	G3		SGCN
<i>Sporobolus vaginiflorus</i>	poverty dropseed	Poaceae	G	S4	G5		SGCN
<i>Stellaria alsine</i>	trailing stitchwort	Caryophyllaceae	H	S1	G5		SGCN
<i>Stellaria crassifolia</i>	fleshy stitchwort	Caryophyllaceae	H	S2	G5		SGCN
<i>Stellaria longipes</i>	long-stalked chickweed	Caryophyllaceae	H	S3	G5	S-SPC	SGCN
<i>Strophostyles helvola</i>	trailing wild bean	Fabaceae	H	S2	G5		SGCN
<i>Strophostyles leiosperma</i>	slick-seed wild bean	Fabaceae	H	S2	G5		SGCN
<i>Stuckenia vaginata</i>	sheathed pondweed	Potamogetonaceae	H,S	S2	G5	S-END	SGCN
<i>Subularia aquatica</i>	awlwort	Brassicaceae	H,S	S2	G5	S-THR	SGCN
<i>Sullivantia sullivantii</i>	reniform sullivantia	Saxifragaceae	H	S2	G4	S-THR	SGCN
<i>Symphyotrichum praealtum</i>	willow aster	Asteraceae	H	S1	G5		SGCN
<i>Symphyotrichum shortii</i>	Short's aster	Asteraceae	H	S3	G5	S-SPC	SGCN
<i>Synthyris bullii</i>	kitten-tails	Plantaginaceae	H	S2	G3	S-THR	SGCN
<i>Taenidia integerrima</i>	yellow pimpernel	Apiaceae	H	S3	G5	S-SPC	SGCN
<i>Tephrosia virginiana</i>	goat's rue	Fabaceae	H	S3	G5	S-SPC	SGCN
<i>Thalictrum revolutum</i>	waxleaf meadowrue	Ranunculaceae	H	S1			SGCN
<i>Thaspium barbinode</i>	hairy-jointed meadow-parsnip	Apiaceae	H	S2	G5	S-SPC	SGCN
<i>Tofieldia pusilla</i>	small false asphodel	Liliaceae	H	S1	G5	S-END	SGCN
<i>Torreyochloa pallida</i> var. <i>pallida</i>	pale manna grass	Poaceae	G	S2	T5	S-SPC	SGCN
<i>Trichophorum clintonii</i>	Clinton's bulrush	Cyperaceae	G	S2	G4	S-THR	SGCN
<i>Trillium nivale</i>	snow trillium	Liliaceae	H	S3	G4	S-SPC	SGCN
<i>Triplasis purpurea</i>	purple sandgrass	Poaceae	G	S2	G4	S-SPC	SGCN
<i>Trisetum spicatum</i>	spike trisetum	Poaceae	G	S2	G5	S-SPC	SGCN
<i>Tsuga canadensis</i>	eastern hemlock	Pinaceae	E	S1	G4	S-END	SGCN
<i>Utricularia geminiscapa</i>	hidden-fruit bladderwort	Lentibulariaceae	H,S	S2	G4	S-THR	SGCN
<i>Utricularia purpurea</i>	purple-flowered bladderwort	Lentibulariaceae	H,S	S2	G5	S-END	SGCN
<i>Utricularia resupinata</i>	lavender bladderwort	Lentibulariaceae	H,S	S2	G4	S-THR	SGCN
<i>Vaccinium uliginosum</i>	alpine bilberry	Ericaceae	D	S1	G5	S-END	SGCN
<i>Valeriana edulis</i>	edible valerian	Valerianaceae	H	S2	G5	S-THR	SGCN
<i>Verbena simplex</i>	narrow-leaved vervain	Verbenaceae	H	S1	G5	S-SPC	SGCN
<i>Viburnum edule</i>	squashberry	Caprifoliaceae	D	S2	G5		SGCN
<i>Viola epipsila</i>	dwarf marsh violet	Violaceae	H	S1	G5		SGCN

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Viola lanceolata</i>	lance-leaf violet	Violaceae	H	S2	G5	S-THR	SGCN
<i>Viola nuttallii</i>	yellow prairie violet	Violaceae	H	S2	G5	S-THR	SGCN
<i>Viola pubescens</i> var. <i>pubescens</i>	downy yellow violet	Violaceae	H	S1	T5		SGCN
<i>Vitis aestivalis</i>	silverleaf grape	Vitaceae	C	S2	G5	S-THR	SGCN
<i>Waldsteinia fragarioides</i>	barren strawberry	Rosaceae	H	S3	G5	S-SPC	SGCN
<i>Wolffia brasiliensis</i>	pointed watermeal	Lemnaceae	F,H	S2	G5		SGCN
<i>Woodsia alpina</i>	alpine woodsia	Woodsiaceae	H	S2	G4	S-THR	SGCN
<i>Woodsia glabella</i>	smooth woodsia	Woodsiaceae	H	S1	G5	S-THR	SGCN
<i>Woodsia obtusa</i>	blunt-lobed woodsia	Woodsiaceae	H	S3	G5		SGCN
<i>Woodsia oregana</i>	western cliff fern	Woodsiaceae	H	S3	G5	S-SPC	SGCN
<i>Woodsia scopulina</i>	Rocky Mountain woodsia	Woodsiaceae	H	S2	G5	S-THR	SGCN
<i>Xanthisma spinulosum</i>	cutleaf ironplant	Asteraceae	H	S2	G5	S-SPC	SGCN
<i>Xyris montana</i>	montane yellow-eyed grass	Xyridaceae	H	S3	G5	S-SPC	SGCN
<i>Xyris torta</i>	twisted yellow-eyed grass	Xyridaceae	H	S1	G5	S-END	SGCN
<i>Zizania palustris</i>	northern wild rice	Poaceae	G	S4	G5		SGCN
<i>Anemone multifida</i>	cut-leaved anemone	Ranunculaceae	H	SH	G5		SNI
<i>Antennaria plantaginifolia</i>	plantain-leaf pussytoes	Asteraceae	H	SU	G5		SNI
<i>Aristida oligantha</i>	prairie three-awn grass	Poaceae	G	SU	G5		SNI
<i>Astragalus racemosus</i>	racemose milk-vetch	Fabaceae	H	SH	G5		SNI
<i>Astragalus tenellus</i>	loose-flowered milk-vetch	Fabaceae	H	SH	G5		SNI
<i>Aureolaria grandiflora</i>	large-flowered false foxglove	Scrophulariaceae	H	SH	G4		SNI
<i>Boechera missouriensis</i>	Missouri rockcress	Brassicaceae	H	SU	G5		SNI
<i>Calopogon oklahomensis</i>	Oklahoma grass pink	Orchidaceae	H	SH	G2		SNI
<i>Carex crus-corvi</i>	raven's foot sedge	Cyperaceae	G	SH	G5		SNI
<i>Carex lucorum</i>	Blue Ridge sedge	Cyperaceae	G	SH	G5		SNI
<i>Carex lurida</i>	shallow sedge	Cyperaceae	G	SU	G5		SNI
<i>Chenopodium berlandieri</i>	pit-seed goosefoot	Amaranthaceae	H	SU	G5		SNI
<i>Corispermum pallasii</i>	Pallas' bugseed	Chenopodiaceae	H	SU	G4		SNI
<i>Corispermum villosum</i>	eastern bugseed	Chenopodiaceae	H	SU	G4		SNI
<i>Crataegus coccinea</i>	scarlet hawthorn	Rosaceae	D	SH	G5		SNI
<i>Crataegus scabrada</i>	rough hawthorn	Rosaceae	D	SH	G5		SNI
<i>Crataegus sheridana</i>	Great Plains hawthorn	Rosaceae	D	SU	GNR		SNI

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Cuscuta indecora</i>	pretty dodder	Cuscutaceae	H	SU	G5		SNI
<i>Cuscuta megalocarpa</i>	large-fruit dodder	Cuscutaceae	H	SU	G5		SNI
<i>Cuscuta obtusiflora</i>	Peruvian dodder	Cuscutaceae	H	SH	G5		SNI
<i>Cuscuta polygonorum</i>	smartweed dodder	Cuscutaceae	H	SU	G5		SNI
<i>Descurainia incana</i>	Richardson's tansy mustard	Brassicaceae	H	SH	G5		SNI
<i>Didiplis diandra</i>	water purslane	Lythraceae	H	SH	G5		SNI
<i>Erigeron pulchellus</i> var. <i>tolsteadii</i>	Ribin's plaintain	Asteraceae	H	SH	TU		SNI
<i>Erysimum asperum</i>	western wallflower	Brassicaceae	H	SH	G5		SNI
<i>Euphorbia hexagona</i>	six-angle spurge	Euphorbiaceae	H	SH	G5		SNI
<i>Euphorbia marginata</i>	snow-on-the-mountain	Euphorbiaceae	H	SU	G5		SNI
<i>Euphorbia vermiculata</i>	worm seeded spurge	Euphorbiaceae	H	SU	G5		SNI
<i>Eutrochium maculatum</i> var. <i>foliosum</i>	spotted joe-pyeweed	Euphorbiaceae	H	SH	T5		SNI
<i>Hedeoma pulegioides</i>	pennyroyal	Lamiaceae	H	SH	G5		SNI
<i>Heterotheca stenophylla</i> var. <i>angustifolia</i>	stiff-leaf false goldenaster	Asteraceae	H	SU	TNR		SNI
<i>Heterotheca villosa</i> var. <i>ballardii</i>	hairy false goldenaster	Asteraceae	H	SU	T5		SNI
<i>Heterotheca villosa</i> var. <i>foliosa</i>	hairy false goldenaster	Asteraceae	H	SU	T5		SNI
<i>Heterotheca villosa</i> var. <i>minor</i>	hairy false goldenaster	Asteraceae	H	SU	T5		SNI
<i>Heterotheca villosa</i> var. <i>villosa</i>	hairy false goldenaster	Asteraceae	H	SU	T5		SNI
<i>Hibiscus laevis</i>	halberd-leaf rosemallow	Malvaceae	H	SU	G5		SNI
<i>Hydrocotyle ranunculoides</i>	floating marsh-pennywort	Apiaceae	H	SU	G5		SNI
<i>Hypericum kalmianum</i>	Kalm's St. John's-wort	Clusiaceae	H	SU	G4		SNI
<i>Juncus gerardii</i>	black grass	Juncaceae	G	SU	G5		SNI
<i>Lemna obscura</i>	little duckweed	Lemnaceae	F, H	SU	G5		SNI
<i>Lemna perpusilla</i>	minute duckweed	Lemnaceae	F, H	SH	G5		SNI
<i>Leucospora multifida</i>	narrowleaf paleseed	Plantaginaceae	H	SU	G5		SNI
<i>Mirabilis linearis</i>	narrowleaf umbrella-wort	Nyctaginaceae	H	SU	G5		SNI
<i>Monolepis nuttalliana</i>	povertyweed	Chenopodiaceae	H	SH	G5		SNI
<i>Muhlenbergia schreberi</i>	dropseed muhly grass	Poaceae	G	SU	G5		SNI
<i>Myriophyllum pinnatum</i>	cutleaf water-milfoil	Haloragaceae	S, H	SU	G5		SNI
<i>Oenothera gaura</i>	biennial gaura	Onagraceae	H	SU	G5		SNI
<i>Oenothera oakesiana</i>	Oake's evening-primrose	Onagraceae	H	SU	G5		SNI
<i>Opuntia humifusa</i>	eastern prickly-pear	Cactaceae	K	SU	G5		SNI

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Peltandra virginica</i>	green arrow-arum	Araceae	H	SU	G5		SNI
<i>Penstemon digitalis</i>	beard-tongue	Scrophulariaceae	H	SU	G5		SNI
<i>Persicaria hydropiperoides</i>	mild water pepper	Polygonaceae	H	SU	G5		SNI
<i>Physalis longifolia</i>	longleaf groundcherry	Solanaceae	H	SU	G5		SNI
<i>Plantago aristata</i>	large-bract plantain	Plantaginaceae	H	SU	G5		SNI
<i>Plantago virginica</i>	hoary plantain	Plantaginaceae	H	SH	G5		SNI
<i>Poa alpina</i>	alpine bluegrass	Solanaceae	H	SU	G5		SNI
<i>Potamogeton x hagstroemii</i>	Hagstrom's pondweed	Potamogetonaceae	H	SU	GNA		SNI
<i>Potamogeton x haynesii</i>	Haynes' pondweed	Potamogetonaceae	H	SU	GNA		SNI
<i>Potentilla pulcherrima</i>	soft cinquefoil	Rosaceae	H	SNR	G5		SNI
<i>Prenanthes crepidinea</i>	nodding rattlesnake-root	Asteraceae	H	SH	G4		SNI
<i>Quercus muhlenbergii</i>	dwarf chestnut oak	Fagaceae	D	SH	G5		SNI
<i>Rhus aromatica</i>	fragrant sumac	Anacardiaceae	D	SU	G5		SNI
<i>Rorippa sphaerocarpa</i>	roundfruit yellowcress	Brassicaceae	H	SU	G5		SNI
<i>Rosa woodsii</i>	Woods' rose	Rosaceae	D	SU	G5		SNI
<i>Rubus canadensis</i>	smooth blackberry	Rosaceae	D	SU	G5		SNI
<i>Rubus junceus</i>		Rosaceae	D	SU	G4		SNI
<i>Rubus plicatifolius</i>		Rosaceae	D	SU	G5		SNI
<i>Rubus spectatus</i>		Rosaceae	D	SU	G5		SNI
<i>Rubus steelei</i>		Rosaceae	D	SU	G4		SNI
<i>Rubus superioris</i>		Rosaceae	D	SU	GNR		SNI
<i>Rubus wisconsinensis</i>	Wisconsin blackberry	Rosaceae	D	SU	G4		SNI
<i>Rumex occidentalis</i>	a species of dock	Polygonaceae	H	SU	T5		SNI
<i>Scirpus georgianus</i>	Georgia bulrush	Cyperaceae	G	SU	G5		SNI
<i>Scirpus pendulus</i>	pendulous bulrush	Cyperaceae	G	SU	G5		SNI
<i>Solidago speciosa</i> var. <i>rigidiuscula</i>	showy goldenrod	Asteraceae	H	SU	T4		SNI
<i>Solidago speciosa</i> var. <i>speciosa</i>	showy goldenrod	Asteraceae	H	SU	T5		SNI
<i>Spermolepis inermis</i>	Red River scaleseed	Apiaceae	H	SU	G5		SNI
<i>Symphyotrichum falcatum</i>	white prairie aster	Asteraceae	H	S2S3	G5		SNI
<i>Symphyotrichum laeve</i> var. <i>geyeri</i>	smooth aster	Asteraceae	H	SH	T5		SNI
<i>Symphyotrichum pilosum</i> var. <i>pilosum</i>	awl aster	Asteraceae	H	SU	T5		SNI
<i>Symphyotrichum pilosum</i> var. <i>pringlei</i>	awl aster	Asteraceae	H	SU	T5		SNI

Scientific Name	Common Name	Family	Physiognomy	S-Rank	G-Rank	Listing Status	SGCN/SNI
<i>Triodanis leptocarpa</i>	Venus' looking-glass	Campanulaceae	H	SH	G5		SNI
<i>Verbena x deamii</i>		Verbena	H	SH	GNA		SNI
<i>Verbena x perriana</i>		Verbena	H	SH	GNA		SNI
<i>Vernonia baldwinii</i>	Baldwin's ironweed	Plantaginaceae	H	SH	G5		SNI
<i>Veronica serpyllifolia</i>	thymeleaf speedwell	Plantaginaceae	H	SU	G5		SNI
<i>Viola affinis</i>	Le Conte's violet	Violaceae	H	SU	G5		SNI

Lichens

SGCN species are listed first in alphabetical order by scientific name, followed by SNI species in alphabetical order by scientific name.

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Abrothallus nephromatis</i>	a lichen	S2	GNR		SGCN
<i>Abrothallus welwitschii</i>	a lichenicolous fungus	SH	GNR		SGCN
<i>Acarospora calcarea</i>	a lichen	S2	GNR		SGCN
<i>Acarospora erythrophora</i>	a cobblestone lichen	S2	GNR		SGCN
<i>Acarospora novomexicana</i>	a lichen	S2	GNR		SGCN
<i>Acarospora obnubila</i>	a lichen	S2	GNR		SGCN
<i>Acarospora rosulata</i>	a cobblestone lichen	S2	GNR		SGCN
<i>Ahtiana aurescens</i>	eastern candlewax lichen	S2	G3G5	S-SPC	SGCN
<i>Allocetraria oakesiana</i>	yellow ribbon lichen	S2	G5	S-THR	SGCN
<i>Amygdalaria panaeola</i>	powdery almond lichen	S2	G3G5	S-SPC	SGCN
<i>Anaptychia crinalis</i>	hanging fringe lichen	S3	G5	S-SPC	SGCN
<i>Anaptychia palmulata</i>	shaggy-fringe lichen	S2	G3G5		SGCN
<i>Arctoparmelia centrifuga</i>	concentric ring lichen	S3	G5	S-SPC	SGCN
<i>Arctoparmelia incurva</i>	finger ring lichen	S2	G5		SGCN
<i>Arctoparmelia subcentrifuga</i>	a species of ring lichen	S3	G4G5	S-SPC	SGCN
<i>Arthrorhaphis citrinella</i>	golden dot-lichen	S3	G4G5	S-THR	SGCN
<i>Arthrorhaphis grisea</i>	a lichen	S1	GNR		SGCN
<i>Biatoropsis angulatae</i>	a lichen	S1	GNR		SGCN
<i>Biatoropsis hirtae</i>	a lichen	S1	GNR		SGCN
<i>Buellia nigra</i>	black disc lichen	S1	G1G2	S-SPC	SGCN
<i>Calicium brevis</i>	a lichen	S1	GNR		SGCN
<i>Calicium denigratum</i>	a stubble lichen	S1	GNR		SGCN
<i>Calicium pinastri</i>	a stubble lichen	S2	GNR		SGCN
<i>Caloplaca parvula</i>	a lichen	S1	G2	S-END	SGCN
<i>Caloplaca stellata</i>	a species of firedot lichen	S1	G3G5	S-SPC	SGCN
<i>Cetraria aculeata</i>	spiny shield lichen	S1	G5		SGCN

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Cetraria arenaria</i>	sand-loving Iceland lichen	S1	G4		SGCN
<i>Cetraria ericetorum</i>	Iceland lichen	S1	G5		SGCN
<i>Cetraria islandica</i>	Iceland lichen	S1	G5		SGCN
<i>Cetraria laevigata</i>	striped Iceland lichen	S1	G5		SGCN
<i>Chaenotheca brachypoda</i>	a lichen	S1	GNR		SGCN
<i>Chaenotheca cinerea</i>	a stubble lichen	S1	G4G5		SGCN
<i>Chaenotheca gracilentia</i>	pin lichen	S1	GNR		SGCN
<i>Chaenotheca nitidula</i>	a lichen	S2	G2G4		SGCN
<i>Chaenothecopsis hendersonii</i>	a lichen	S2	GNR		SGCN
<i>Chaenothecopsis ochroleuca</i>	a lichen	S1	GNR		SGCN
<i>Chaenothecopsis ussuriensis</i>	a lichen	S2	GNR		SGCN
<i>Chaenothecopsis viridialba</i>	a lichen	S1	GNR		SGCN
<i>Cladonia norvegica</i>	baby powderhorn lichen	S1	G4G5		SGCN
<i>Cladonia wainioi</i>	Wainio's reindeer lichen	S3	G5	S-SPC	SGCN
<i>Coccocarpia palmicola</i>	salted shell lichen	S1	G5	S-THR	SGCN
<i>Collema flaccidum</i>	flaking tarpaper lichen	S1	G3G5		SGCN
<i>Collema pustulatum</i>	a lichen	S1	GNR		SGCN
<i>Collema texanum</i>	a lichen	S2	G4		SGCN
<i>Cryptocalicium blascoi</i>	a lichen	S2	GNR		SGCN
<i>Dactylospora amygdalariae</i>	a lichen	S2	GNR		SGCN
<i>Dactylospora pertusariicola</i>	a lichen	S2	GNR		SGCN
<i>Dendroscocaulon intricatulum</i>	olive-thorn lichen	S1	GUQ		SGCN
<i>Dendroscosticta gelida</i>	a lichen	S1	GNR		SGCN
<i>Dermatocarpon moulinsii</i>	shag-belly stippleback lichen	S2	G4	S-END	SGCN
<i>Enchylium bachmanianum</i>	Bachman's jelly lichen	S1	GNR		SGCN
<i>Enchylium coccophorum</i>	tar jelly lichen	S1	G3G5		SGCN
<i>Enchylium expansum</i>	a lichen	S1	GNR		SGCN
<i>Enchylium polycarpon</i>	shaly jelly lichen	S2	GNR		SGCN
<i>Epilichen scabrosus</i>	a lichen	S1	G3G5		SGCN

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Fuscopannaria ahlneri</i>	a lichen	S1	G4		SGCN
<i>Fuscopannaria leucosticta</i>	rimmed shingles lichen	S2	G3G5		SGCN
<i>Heppia conchiloba</i>	a lichen	S2	G4		SGCN
<i>Heterodermia hypoleuca</i>	cupped fringe lichen	S2	G5		SGCN
<i>Heterodermia japonica</i>	Japanese fringe lichen	S2	G5		SGCN
<i>Heterodermia leucomelos</i>	elegant fringe lichen	S1	G5		SGCN
<i>Heterodermia obscurata</i>	orange-tinted fringe lichen	S1	G5	S-SPC	SGCN
<i>Hypotrachyna afrorevoluta</i>	a lichen	S1	G5		SGCN
<i>Hypotrachyna britannica</i>	a loop lichen	S2	GNR		SGCN
<i>Lecanora epanora</i>	a rim lichen	S2	GNR	S-THR	SGCN
<i>Lecanora gisleriana</i>	a lichen	S1	GNR		SGCN
<i>Lecanora handelii</i>	a lichen	S1	GNR		SGCN
<i>Lecidea atrobrunnea</i>	brown tile lichen	S1	G4G5		SGCN
<i>Leptogium apalachense</i>	a lichen	SH	G2	S-END	SGCN
<i>Leptogium arsenei</i>	wrinkled jellyskin lichen	S1	GNR		SGCN
<i>Leptogium corticola</i>	blistered jellyskin	S2	G3G5		SGCN
<i>Leptogium hirsutum</i>	jellyskin lichen	S2	G5		SGCN
<i>Leptogium milligranum</i>	stretched jellyskin lichen	S1	G5		SGCN
<i>Lithothelium phaeosporum</i>	a lichen	S2	GNR		SGCN
<i>Lobaria scrobiculata</i>	textured lungwort	S1	G5	S-END	SGCN
<i>Lobothallia alphoplaca</i>	puffed sunken-disk lichen	S1	G5		SGCN
<i>Melanohalea subolivacea</i>	brown-eyed camouflage lichen	SU	GNR	S-SPC	SGCN
<i>Microcalicium ahlneri</i>	a lichen	S1	GNR		SGCN
<i>Microcalicium arenarium</i>	sandwort microcalicium lichen	S2	G4G5		SGCN
<i>Miriquidica atrofulva</i>	a lichen	S1	GNR		SGCN
<i>Mycocalicium ravenelii</i>	a lichen	S1	GNR		SGCN
<i>Myriolecis perpruinosa</i>	a lichen	S2	GNR		SGCN
<i>Myriolecis torrida</i>	a rim lichen	S2	G3G5		SGCN
<i>Pannaria subfusca</i>	a lichen	S1	G4G5		SGCN

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Pannaria tavaresii</i>	coral-rimmed shingle lichen	S1	G5		SGCN
<i>Parmelia stictica</i>	rock speckled shield lichen	S2	G3G5	S-END	SGCN
<i>Parmelia stuppea</i>	powder-edged ruffle lichen	SH	G4G5	S-THR	SGCN
<i>Parmotrema perlatum</i>	black stone flower	S1	G3G5		SGCN
<i>Peltigera venosa</i>	fan lichen	S3	G5	S-SPC	SGCN
<i>Peltula bolanderi</i>	Bolander's peltula lichen	S1	G4G5	S-THR	SGCN
<i>Pertusaria subobducens</i>	a pore lichen	S1	G3G5		SGCN
<i>Phaeophyscia endococcinodes</i>	red-cored starburst shadow lichen	S2	G4G5		SGCN
<i>Phylliscum demangeonii</i>	black rock-wafer lichen	S1	G5		SGCN
<i>Physcia halei</i>	granite rosette lichen	S4	G3		SGCN
<i>Physconia muscigena</i>	ground frost lichen	S1	G5		SGCN
<i>Physconia subpallida</i>	pale-bellied frost lichen	S1	GNR		SGCN
<i>Placidium arboreum</i>	tree stipplescale lichen	S1	G5		SGCN
<i>Placynthium stenophyllum</i>	pale-bellied ink lichen	S2	G3G4		SGCN
<i>Platismatia glauca</i>	ragbag lichen	S1	G5	S-SPC	SGCN
<i>Plectocarpon peltigerae</i>	a lichen	S2	GNR		SGCN
<i>Plectocarpon triebeliae</i>	a lichen	S1	GNR		SGCN
<i>Polyblastia theleodes</i>	a lichen	S1	G3G5		SGCN
<i>Polychidium muscicola</i>	eyed moss-thorns lichen	S1	G5		SGCN
<i>Porpidia cinereoatra</i>	Hertel's boulder lichen	S2	GNR		SGCN
<i>Protopannaria pezizoides</i>	brown-gray moss-shingle lichen	S2	G5	S-THR	SGCN
<i>Protoparmelia badia</i>	chocolate rim lichen	S1	G4G5		SGCN
<i>Pseudocyphellaria holarctica</i>	yellow specklebelly lichen	S1	GNR	S-END	SGCN
<i>Psora globifera</i>	blackberry scale lichen	S1	G4G5		SGCN
<i>Psora rubiformis</i>	pea-green scale lichen	S2	G3G5		SGCN
<i>Psora russellii</i>	Russell's fishscale lichen	S1	G4G5		SGCN
<i>Punctelia appalachensis</i>	Appalachian speckled shield lichen	S1	G3G4		SGCN
<i>Ramalina obtusata</i>	hooded ribbon lichen	S1	G5		SGCN
<i>Ramalina roesleri</i>	frayed ramalina lichen	S1	G5	S-THR	SGCN

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Ramalina thrausta</i>	angel's hair lichen	S1	G5	S-SPC	SGCN
<i>Rhizocarpon oederi</i>	Oeder's map lichen	S2	GNR		SGCN
<i>Rhizocarpon ridescens</i>	a map lichen	S1	GNR		SGCN
<i>Rhizocarpon umbilicatum</i>	a lichen	S1	G4		SGCN
<i>Rhizoplaca melanophthalma</i>	green rock-posy lichen	S1	G5		SGCN
<i>Rhizoplaca ouimetensis</i>	Ouimet Canyon lichen	S1	G1G2		SGCN
<i>Rinodina adirondackii</i>	Adirondack pepper-spore lichen	SH	GNR		SGCN
<i>Rinodina wetmorei</i>	Wetmore's pepper-spore lichen	S1	GNR		SGCN
<i>Romjularia lurida</i>	imbricate scale lichen	S2	GNR		SGCN
<i>Rostania occultata</i>	crusted tarpaper lichen	S2	GNR		SGCN
<i>Sclerophora amabilis</i>	a lichen	S1	G4G5		SGCN
<i>Sclerophora farinacea</i>	a lichen	S1	GNR		SGCN
<i>Scytinium gelatinosum</i>	rose-petalled jellyskin lichen	S2	G5		SGCN
<i>Scytinium teretiusculum</i>	curly jellyskin lichen	S1	G4		SGCN
<i>Solorina spongiosa</i>	fringed chocolate chip lichen	S1	G5		SGCN
<i>Sphinctrina leucopoda</i>	a lichen	S1	G4		SGCN
<i>Sporodophoron americanum</i>	a lichen	S1	GNR		SGCN
<i>Sticta beauvoisii</i>	fingered moon lichen	S2	G5		SGCN
<i>Sticta fuliginosa</i>	peppered moon lichen	S2	G3G5	S-SPC	SGCN
<i>Synalissa ramulosa</i>	eyed rockgorgon lichen	S1	GNR		SGCN
<i>Thyrea confusa</i>	jelly strap lichen	S2	G3G5		SGCN
<i>Toninia ruginosa</i>	a blister lichen	SH	G3G5		SGCN
<i>Toninia sedifolia</i>	blue blister lichen	S1	G5		SGCN
<i>Toninia squalida</i>	a blister lichen	SH	G3G5		SGCN
<i>Tremella nashii</i>	a jelly fungus	S2	GNR		SGCN
<i>Umbilicaria hirsuta</i>	hairy navel lichen	S1	G5		SGCN
<i>Umbilicaria phaea</i>	emery rocktripe lichen	S1	G5		SGCN
<i>Umbilicaria torrefacta</i>	punctured rock tripe lichen	S1	G5	S-END	SGCN
<i>Usnea angulata</i>	angular beard lichen	S1	G3G5		SGCN

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Usnea ceratina</i>	warty beard lichen	S1	G4G5		SGCN
<i>Usnea dasaea</i>	a beard lichen	S1	GNR		SGCN
<i>Usnea entoviolata</i>	a beard lichen	S1	GNR		SGCN
<i>Usnea glabrescens</i>	a lichen	S2	G5		SGCN
<i>Usnea longissima</i>	Methuselah's beard lichen	S3	G5	S-SPC	SGCN
<i>Usnea mutabilis</i>	bloody beard lichen	S1	G5	S-THR	SGCN
<i>Usnea perhispidella</i>	a beard lichen	S1	GNR		SGCN
<i>Usnea rubicunda</i>	red beard lichen	S3	G4G5	S-SPC	SGCN
<i>Vulpicida juniperina</i>	limestone sunshine lichen	S1	G5		SGCN
<i>Abrothallus halei</i>	a lichenicolous fungus	S3	GNR		SNI
<i>Abrothallus pezizicola</i>	a lichenicolous fungus	S3	GNR		SNI
<i>Acarospora contigua</i>	gold cobblestone lichen	S3	GNR		SNI
<i>Acarospora glaucocarpa</i>	a lichen	S3	G5		SNI
<i>Acarospora sinopica</i>	a lichen	S3	G3G5		SNI
<i>Acarospora strigata</i>	hoary cobblestone lichen	S3	G4G5		SNI
<i>Arthonia fuliginosa</i>	a comma lichen	SH	GNR		SNI
<i>Arthopyrenia cinereopruinosa</i>	shell lichen	SH	GNR		SNI
<i>Arthopyrenia salicis</i>	a lichen	S3S4	GNR		SNI
<i>Bacidina inundata</i>	a lichen	SH	GNR		SNI
<i>Bellemeria cinereorufescens</i>	brown sunken disk lichen	S3	G3G5		SNI
<i>Bryoria kockiana</i>	boreal horsehair lichen	SU	GNR		SNI
<i>Bryoria nadvornikiana</i>	spiny gray horsehair lichen	S4	G5		SNI
<i>Buellia aethalea</i>	darkened button lichen	S3	G3G4		SNI
<i>Buellia spuria</i>	sunken button lichen	S3	G3G5		SNI
<i>Buellia tyrolensis</i>	a disc lichen	S3	GNR		SNI
<i>Calicium notarisii</i>	a lichen	S4	GNR		SNI
<i>Caloplaca borealis</i>	boreal orange lichen	SU	G1G2		SNI
<i>Chaenotheca chlorella</i>	a lichen	S3	GNR		SNI
<i>Chaenothecopsis amurensis</i>	a lichen	S3	GNR		SNI

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Chaenothecopsis asperopoda</i>	a lichen	S4	GNR		SNI
<i>Chaenothecopsis brevipes</i>	a lichen	S3	GNR		SNI
<i>Chaenothecopsis consociata</i>	a lichen	S3	G4		SNI
<i>Chaenothecopsis exilis</i>	a lichen	S3S4	GNR		SNI
<i>Chaenothecopsis viridireagens</i>	a lichen	S3	GNR		SNI
<i>Chrysothrix chlorina</i>	sulphur dust lichen	S3	GNR		SNI
<i>Chrysothrix xanthina</i>	gold dust lichen	SH	GNR		SNI
<i>Cladonia floerkeana</i>	gritty British soldiers lichen	S3	G5		SNI
<i>Cladonia robbinsii</i>	Robbins' cup lichen	S3	G3G5		SNI
<i>Cladonia sulphurina</i>	greater sulphur-cup lichen	S3	G5		SNI
<i>Collema nigrescens</i>	blistered jelly lichen	S4	G5		SNI
<i>Collema pulchellum</i>	blistered jelly lichen	S4	GNR		SNI
<i>Dactylospora lobariella</i>	a lichen	S3	GNR		SNI
<i>Dermatocarpon intestiniforme</i>	quilted stippleback lichen	SU	GNR		SNI
<i>Dermatocarpon meiophyllizum</i>	a stippleback lichen	S3	G3G5		SNI
<i>Didymosphaeria futilis</i>	a lichen	SU	GNR		SNI
<i>Diplotomma venustum</i>	a lichen	S3	GNR		SNI
<i>Enchylium conglomeratum</i>	dotted jelly lichen	S3	G5		SNI
<i>Enchylium limosum</i>	lime-loving tarpaper lichen	SH	GNR		SNI
<i>Enchylium tenax</i>	tar jelly lichen	S3	G5		SNI
<i>Epicladonia sandstedei</i>	a lichen	SH	GNR		SNI
<i>Gyalecta jenensis</i>	rock dimple lichen	S3	G3G4		SNI
<i>Heppia adglutinata</i>	soil ruby lichen	S3	G4G5		SNI
<i>Hypogymnia incurvoides</i>	lattice tube lichen	SU	G5		SNI
<i>Kirschsteiniotelia aethiops</i>	a lichen	SH	GNR		SNI
<i>Lathagrium undulatum</i>	jelly flakes lichen	S3	G5		SNI
<i>Lecanora cateilea</i>	a lichen	SH	GNR		SNI
<i>Lecanora marginata</i>	a lichen	SH	G4G5		SNI
<i>Lecanora nothocaesiella</i>	a rim lichen	SH	GNR		SNI

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Lecanora pseudistera</i>	a rim lichen	S3	GNR		SNI
<i>Lecanora rupicola</i>	white rim lichen	S3	G5		SNI
<i>Lecanora varia</i>	a lichen	SH	G5		SNI
<i>Lecanora wetmorei</i>	Wetmore's rim lichen	S3	GNR		SNI
<i>Lecidea albofuscescens</i>	a lichen	SH	GNR		SNI
<i>Lecidea auriculata</i>	a disk lichen	S3	G5		SNI
<i>Lecidea turgidula</i>	turgid dot lichen	SH	GNR		SNI
<i>Lecidella effugiens</i>	a disk lichen	S3	GNR		SNI
<i>Lempholemma chalazanum</i>	a lichen	S3	GNR		SNI
<i>Lepraria disjuncta</i>	a dust lichen	S3	GNR		SNI
<i>Lepraria membranacea</i>	leproloma lichen	SU	GNR		SNI
<i>Lepraria vouauxii</i>	a dust lichen	S3	GNR		SNI
<i>Leptogium acadense</i>	a lichen	SU	GNR		SNI
<i>Leptogium byssinum</i>	granular jellyskin lichen	SH	GNR		SNI
<i>Leptogium compactum</i>	a lichen	SU	G4		SNI
<i>Leptogium rivulare</i>	flooded jellyskin lichen	S3	G3G5		SNI
<i>Leptogium saturninum</i>	bearded jellyskin lichen	SU	GNR		SNI
<i>Lichenodiplis lecanorae</i>	a lichen	SH	GNR		SNI
<i>Lichenothelia scopularia</i>	a lichen	SH	GNR		SNI
<i>Lichinella iodopulchra</i>	a lichen	S3	GNR		SNI
<i>Lichinella minnesotensis</i>	a lichen	SH	GNR		SNI
<i>Lichinella nigrifella</i>	black rock licorice lichen	S3	G4G5		SNI
<i>Lichinella stipatula</i>	shrubby rock licorice lichen	S3	GNR		SNI
<i>Lobaria quercizans</i>	smooth lungwort	S3	G5		SNI
<i>Megalospora porphyritis</i>	a lichen	SH	GNR		SNI
<i>Melanelia hepaticum</i>	rimmed camouflage lichen	S3	G5		SNI
<i>Merismatium decolorans</i>	a lichen	SH	GNR		SNI
<i>Micarea denigrata</i>	a lichen	SH	G2G4		SNI
<i>Microcalicium conversum</i>	a lichen	S5	GNR		SNI

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Montanelia panniformis</i>	shingled camouflage lichen	S3	G5		SNI
<i>Mycomicrothelia wallrothii</i>	a lichen	SH	GNR		SNI
<i>Myelochroa obsessa</i>	rock axil-bristle lichen	S3	G3G5		SNI
<i>Myriolecis fugiens</i>	a rim lichen	SH	GNR		SNI
<i>Nephroma bellum</i>	naked kidney lichen	S3	G5		SNI
<i>Pannaria conoplea</i>	mealy-rimmed shingle lichen	S4	G4G5		SNI
<i>Parmotrema arnoldii</i>	powdered ruffle lichen	S3	G3G5		SNI
<i>Parmotrema cetratum</i>	ruffle lichen	SH	G5		SNI
<i>Parmotrema hypotropum</i>	powdered ruffle lichen	S3	G5		SNI
<i>Parmotrema margaritatum</i>	southeastern powdered ruffle lichen	SH	GNR		SNI
<i>Parmotrema subtinctorium</i>	mottled ruffle lichen	S3	G5		SNI
<i>Peltigera scabrosa</i>	greater toad pelt lichen	S3	G5		SNI
<i>Peltula patellata</i>	stuffed rock-olive lichen	SH	G3G5		SNI
<i>Pertusaria plittiana</i>	rock wart lichen	S3	G4G5		SNI
<i>Pertusaria rhexostoma</i>	a pore lichen	SH	GNR		SNI
<i>Phaeocalicium minutissimum</i>	red oak stubble lichen	S3	GNR		SNI
<i>Phaeophyscia hispidula</i>	whiskered shadow lichen	S3	G4G5		SNI
<i>Phaeophyscia squarrosa</i>	scaly shadow lichen	S3S4	GNR		SNI
<i>Pilophorus cereolus</i>	a lichen	S3	G3G4		SNI
<i>Placynthium asperellum</i>	a lichen	S3	G4G5		SNI
<i>Plectocarpon lichenum</i>	a lichen	S3	GNR		SNI
<i>Porpidia subsimplex</i>	a lichen	SH	G3G5		SNI
<i>Protoparmelia hypotremella</i>	a rim lichen	SH	GNR		SNI
<i>Psora decipiens</i>	blushing scale lichen	S3	G5		SNI
<i>Psora icterica</i>	yellow scale lichen	SH	G5		SNI
<i>Punctelia borrieri</i>	blinking speckleback lichen	SU	GNR		SNI
<i>Pyrenidium actinellum</i>	a lichen	SH	GNR		SNI
<i>Pyxine soledata</i>	mustard lichen	S4	G5		SNI
<i>Ramalina labiosorediata</i>	powdery twig lichen	S4	G4		SNI

Scientific Name	Common Name	S-Rank	G-Rank	Listing Status	Result
<i>Rhizocarpon lavatum</i>	drenched map lichen	S3	GNR		SNI
<i>Rhizocarpon lecanorinum</i>	a lichen	S3	G4G5		SNI
<i>Rinodina albertana</i>	Albert's pepper-spore lichen	SH	GNR		SNI
<i>Rinodina archaea</i>	brown pepper-spore lichen	SH	G4G5		SNI
<i>Rinodina excrescens</i>	a lichen	S3	GNR		SNI
<i>Rinodina metaboliza</i>	variable pepper-spore lichen	S3	GNR		SNI
<i>Rinodina septentrionalis</i>	a pepper-spore lichen	SH	G3G5		SNI
<i>Sarcosagium campestre</i>	a lichen	SH	GNR		SNI
<i>Schismatomma pericleum</i>	a lichen	SH	GNR		SNI
<i>Sclerophora nivea</i>	a lichen	S3	GNR		SNI
<i>Scytinium dactylinum</i>	brown-buttoned jellyskin lichen	S3	GNR		SNI
<i>Solorina saccata</i>	chocolate chip lichen	S3	G5		SNI
<i>Staurothele elenkinii</i>	Elenkin's rock pimple lichen	SH	G3G5		SNI
<i>Tephromela atra</i>	black-eye lichen	SH	G5		SNI
<i>Thermutis velutina</i>	rockvelvet lichen	S3	G3G5		SNI
<i>Toninia aromatica</i>	aromatic blister lichen	SH	G4G5		SNI
<i>Tremella flavoparmeliae</i>	a lichen	S3	GNR		SNI
<i>Umbilicaria hyperborea</i>	blistered rocktripe lichen	S3	G5		SNI
<i>Usnea amblyoclada</i>	a beard lichen	SH	GNR		SNI
<i>Verrucaria fayettensis</i>	a speck lichen	S3	GNR		SNI
<i>Xanthoparmelia dierythra</i>	a rock-shield lichen	SU	G2G3		SNI
<i>Xanthoparmelia hypomelaena</i>	Ozark rock-shield lichen	SH	G3G4		SNI
<i>Xylographa disseminata</i>	a woodscript lichen	SH	GNR		SNI

APPENDIX C – Animal SGCN Habitat Associations

To associate SGCN with habitat types, species experts classified the SGCN by habitat types they use based on professional knowledge. Vertebrate species and some invertebrate species were classified as using habitats as primary (indicated with a 1) or secondary habitat (indicated with a 2.) Primary habitats are those that species rely on and use most consistently; loss or degradation of these habitats would have the most significant negative impact on their populations. Secondary habitats are used by the species less frequently. Other invertebrate species were associated with habitats without delineating primary vs. secondary habitat (indicated with an X.) Animals with more general habitat requirements were associated with multiple habitat types, while specialists were associated with one or few.

Amphibians

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Non-forested Wetlands	Vernal Pools	Streams	Rivers	Large Rivers	Shallow Lakes	Deep Lakes	Prairie and Other Grasslands	Urban and Other Developed Lands
<i>Acris blanchardi</i>	Blanchard's Cricket Frog					1	1		1	1	1	1			
<i>Ambystoma maculatum</i>	Spotted Salamander	2	1	1	2		1	1							
<i>Anaxyrus cognatus</i>	Great Plains Toad						2							1	2
<i>Hemidactylium scutatum</i>	Four-toed Salamander	2	1	1	2			1							
<i>Lithobates palustris</i>	Pickereel Frog					1	1		1	1					
<i>Necturus maculosus</i>	Mudpuppy								1	1	1	2	1		
<i>Notophthalmus viridescens</i>	Eastern Newt	2	1		2		1	1				2			

Birds

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Vernal Pools	Streams	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Accipiter atricapillus</i>	American Goshawk	1	2												
<i>Aechmophorus occidentalis</i>	Western Grebe										1				
<i>Aegolius funereus</i>	Boreal Owl	2	1				1								
<i>Ammodramus savannarum</i>	Grasshopper Sparrow											1			
<i>Ammospiza leconteii</i>	LeConte's Sparrow							1				2			
<i>Ammospiza nelsoni</i>	Nelson's Sparrow							1							
<i>Anas acuta</i>	Northern Pintail							2			1				
<i>Anas rubripes</i>	American Black Duck							2			1				
<i>Anthus spragueii</i>	Sprague's Pipit											1			
<i>Anurostomus vociferus</i>	Eastern Whip-poor-will	1	1										2		
<i>Asio flammeus</i>	Short-eared Owl							1				1			
<i>Athene cunicularia</i>	Burrowing Owl											1			
<i>Aythya affinis</i>	Lesser Scaup							2			1				
<i>Aythya valisineria</i>	Canvasback							2			1				
<i>Bartramia longicauda</i>	Upland Sandpiper											1			
<i>Botaurus exilis</i>	Least Bittern							1			2				

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Vernal Pools	Streams	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Botaurus lentiginosus</i>	American Bittern							1			2				
<i>Bucephala albeola</i>	Bufflehead							2			1				
<i>Buteo lineatus</i>	Red-shouldered Hawk		2	1	2	1			2						
<i>Buteo swainsoni</i>	Swainson's Hawk		2									1	2		
<i>Calcarius ornatus</i>	Chestnut-collared Longspur											1			
<i>Calidris canutus</i>	Red Knot										1				
<i>Calidris subruficollis</i>	Buff-breasted Sandpiper											1			
<i>Canachites canadensis</i>	Spruce Grouse	2					1								
<i>Catharus ustulatus</i>	Swainson's Thrush	1					2								
<i>Centronyx bairdii</i>	Baird's Sparrow											1			
<i>Centronyx henslowii</i>	Henslow's Sparrow											1			
<i>Chaetura pelagica</i>	Chimney Swift														1
<i>Charadrius melodus</i>	Piping Plover										1				
<i>Chlidonias niger</i>	Black Tern							1			1				
<i>Chondestes grammacus</i>	Lark Sparrow											2	1		
<i>Chordeiles minor</i>	Common Nighthawk											2		1	1

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Vernal Pools	Streams	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Circus hudsonius</i>	Northern Harrier							1				1			
<i>Cistothorus stellaris</i>	Sedge Wren							1				2			
<i>Coccothraustes vespertinus</i>	Evening Grosbeak	1	2												
<i>Contopus cooperi</i>	Olive-sided Flycatcher	1					1								
<i>Coturnicops noveboracensis</i>	Yellow Rail							1				2			
<i>Dolichonyx oryzivorus</i>	Bobolink							2				1			
<i>Empidonax virescens</i>	Acadian Flycatcher		2	1		2									
<i>Falco peregrinus</i>	Peregrine Falcon													1	1
<i>Falco sparverius</i>	American Kestrel											1	2		2
<i>Gallinula galeata</i>	Common Gallinule							1			2				
<i>Gavia immer</i>	Common Loon										1				
<i>Lanius ludovicianus</i>	Loggerhead Shrike											1			2
<i>Leucophaeus pipixcan</i>	Franklin's Gull							1			1				
<i>Limosa fedoa</i>	Marbled Godwit							1				1			
<i>Melanerpes erythrocephalus</i>	Red-headed Woodpecker		2			2							1		2
<i>Nycticorax nycticorax</i>	Black-crowned Night-heron							1			2				

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Vernal Pools	Streams	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Oporornis agilis</i>	Connecticut Warbler	2					1								
<i>Parkesia motacilla</i>	Louisiana Waterthrush			1	2	2				1					
<i>Pelecanus erythrorhynchos</i>	American White Pelican							1			1				
<i>Phalaropus tricolor</i>	Wilson's Phalarope							1			2	2			
<i>Picoides arcticus</i>	Black-backed Woodpecker	2					1								
<i>Picoides dorsalis</i>	American Three-toed Woodpecker	2					1								
<i>Podiceps auritus</i>	Horned Grebe							2			1				
<i>Podiceps grisegena</i>	Red-necked Grebe							2			1				
<i>Podiceps nigricollis</i>	Eared Grebe							2			1				
<i>Poecile hudsonicus</i>	Boreal Chickadee	2					1								
<i>Progne subis</i>	Purple Martin							2			1				2
<i>Protonotaria citrea</i>	Prothonotary Warbler					1									
<i>Rallus elegans</i>	King Rail							1			2				
<i>Setophaga caerulescens</i>	Black-throated Blue Warbler	2	2	1											
<i>Setophaga castanea</i>	Bay-breasted Warbler	1													
<i>Setophaga cerulea</i>	Cerulean Warbler		2	1	2	1									

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Vernal Pools	Streams	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Setophaga tigrina</i>	Cape May Warbler	1					2								
<i>Spizella pusilla</i>	Field Sparrow											1	1		
<i>Sterna forsteri</i>	Forster's Tern							1			1				
<i>Sterna hirundo</i>	Common Tern										1				
<i>Strix nebulosa</i>	Great Gray Owl	2					1								
<i>Sturnella magna</i>	Eastern Meadowlark											1	2		
<i>Sturnella neglecta</i>	Western Meadowlark											1			
<i>Tringa solitaria</i>	Solitary Sandpiper						1	2							
<i>Tympanuchus cupido</i>	Greater Prairie-chicken											1			2
<i>Tympanuchus phasianellus</i>	Sharp-tailed Grouse							2				1			
<i>Tyrannus verticalis</i>	Western Kingbird											1			
<i>Tyto furcata</i>	American Barn Owl											1			2
<i>Vermivora chrysoptera</i>	Golden-winged Warbler	2	2	2	2		2								
<i>Vireo bellii</i>	Bell's Vireo							1				2			
<i>Vireo philadelphicus</i>	Philadelphia Vireo		1	2											

Fish

Scientific Name	Common Name	Streams (SSO 1-4)	Rivers (SSO 5-6)	Large Rivers (SSO 7-8)	Shallow Lakes	Deep Lakes
<i>Acipenser fulvescens</i>	Lake Sturgeon		2	1	2	2
<i>Alosa chrysochloris</i>	Skipjack Herring			1		
<i>Ammocrypta clara</i>	Western Sand Darter		2	1		
<i>Aphredoderus sayanus</i>	Pirate Perch			1**		
<i>Catostomus catostomus</i>	Longnose sucker	1*	2*			1
<i>Clinostomus elongatus</i>	Redside Dace	1	2			
<i>Cottus cognatus</i>	Slimy Sculpin	1*				2*
<i>Cottus ricei</i>	Spoonhead Sculpin					1*
<i>Couesius plumbeus</i>	Lake Chub	1			2	1*
<i>Crystallaria asprella</i>	Crystal Darter		2	1		
<i>Erimystax x-punctatus</i>	Gravel Chub	2	1			
<i>Etheostoma chlorosoma</i>	Bluntnose Darter			1		
<i>Etheostoma microperca</i>	Least Darter				2	1
<i>Fundulus sciadicus</i>	Plains Topminnow	1	2			
<i>Hybognathus nuchalis</i>	Mississippi Silvery Minnow		2	1		
<i>Hybopsis amnis</i>	Pallid Shiner			1		
<i>Ichthyomyzon fossor</i>	Northern Brook Lamprey	1	2			
<i>Ichthyomyzon gagei</i>	Southern Brook Lamprey	1	2			
<i>Ictiobus niger</i>	Black Buffalo			1		
<i>Lepomis gulosus</i>	Warmouth			1**		
<i>Lepomis peltastes</i>	Northern Sunfish				2	1
<i>Lythrurus umbratilis</i>	Redfin Shiner	2	1			
<i>Moxostoma carinatum</i>	River Redhorse		2	1		
<i>Moxostoma duquesnei</i>	Black Redhorse	2	1			
<i>Myoxocephalus thompsonii</i>	Deepwater Sculpin					1*
<i>Nocomis biguttatus</i>	Hornyhead Chub	1	1	2		
<i>Notropis anogenus</i>	Pugnose Shiner				2	1
<i>Notropis nubilus</i>	Ozark Minnow	1	1			
<i>Notropis texanus</i>	Weed Shiner	2	1	1**	2	2

<i>Notropis topeka</i>	Topeka Shiner	1	2			
<i>Noturus exilis</i>	Slender Madtom	1	2			
<i>Percina evides</i>	Gilt Darter		1	2		
<i>Phenacobius mirabilis</i>	Suckermouth Minnow	1	2			
<i>Platygobio gracilis</i>	Flathead Chub		2	1		
<i>Polyodon spathula</i>	Paddlefish			1		
<i>Salvelinus fontinalis</i>	Brook Trout***	1*	2*			1*

*Coldwater specialist

**Backwaters

***Native strains - MN heritage brook trout and Coaster Brook Trout

Mammals

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Caves	Urban and Other Developed Lands
<i>Alces alces</i>	Moose	2	2		2	2		2	2	2					
<i>Canis lupus</i>	Gray Wolf	2	2	2	2	2	2	2				2			2
<i>Cervus canadensis</i>	Elk		2								1	1			
<i>Cryptotis parvus</i>	North American Least Shrew		2	2											2
<i>Eptesicus fuscus</i>	Big Brown Bat	2	2	2	2	2			2	2		2	2	2	2
<i>Glaucomys sabrinus</i>	Northern Flying Squirrel	2	2	2	2		2								
<i>Lasionycteris noctivagans</i>	Silver-haired Bat	2	2	2	2	2			2	2		2			2
<i>Lasiurus borealis</i>	Eastern Red Bat	2	2	2	2	2			2	2		2			2
<i>Lasiurus cinereus</i>	Hoary Bat	2	2	2	2	2			2	2		2			2

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Caves	Urban and Other Developed Lands
<i>Lepus americanus</i>	Snowshoe hare	2	2				2	2							
<i>Lepus townsendii</i>	White-tailed Jackrabbit										1	2			2
<i>Lynx canadensis</i>	Canada Lynx	2	2				2						2		
<i>Microtus chrotorrhinus</i>	Rock Vole	1	1										1		
<i>Microtus ochrogaster</i>	Prairie Vole										1				2
<i>Microtus pinetorum</i>	Woodland Vole			1	1	1									
<i>Mustela nivalis</i>	Least Weasel										1	1			
<i>Myotis lucifugus</i>	Little Brown Myotis	2	2	2	2	2			2	2			2	1	2
<i>Myotis septentrionalis</i>	Northern Long-eared Bat	2	2	2	2	2			2	2			2	1	
<i>Napaeozapus insignis</i>	Eastern Woodland Jumping Mouse	2	2	2									2		
<i>Onychomys leucogaster</i>	Northern Grasshopper Mouse										1	2	2		2
<i>Perimyotis subflavus</i>	Tricolored Bat	2	1	1					2	2		2		1	
<i>Perognathus flavescens</i>	Plains Pocket Mouse										1	1	2		
<i>Phenacomys ungava</i>	Eastern Heather Vole	2	2				1	1					2		
<i>Poliocitellus franklinii</i>	Franklin's Ground Squirrel	2	2	2	2			1			1	2	2		2
<i>Puma concolor</i>	Puma	2	2	2	2	2	2		2	2	2	2	2		2

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Caves	Urban and Other Developed Lands
<i>Reithrodontomys megalotis</i>	Western Harvest Mouse										1	2	2		2
<i>Sorex fumeus</i>	Smoky Shrew	2	2										1		
<i>Spilogale interrupta</i>	Plains Spotted Skunk		2	2							1	1			2
<i>Synaptomys borealis</i>	Northern Bog Lemming						2	1							
<i>Taxidea taxus</i>	American Badger	2	2	2							1	1	2		2
<i>Thomomys talpoides</i>	Northern Pocket Gopher										1	2			1
<i>Urocitellus richardsonii</i>	Richardson's Ground Squirrel										1	2			1

Reptiles

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Non-forested Wetlands	Vernal Pools	Streams	Rivers	Large Rivers	Shallow Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Apalone mutica</i>	Smooth Softshell									1	1					
<i>Aspidoscelis sexlineatus</i>	Six-lined Racerunner		1										1	1	1	2
<i>Coluber constrictor</i>	North American Racer		1	1									1	1	1	2

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Non-forested Wetlands	Vernal Pools	Streams	Rivers	Large Rivers	Shallow Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Crotalus horridus</i>	Timber Rattlesnake		1	1									1	1	1	2
<i>Emydoidea blandingii</i>	Blanding's Turtle					1	1	2	1	1	2*	1	1	1		
<i>Glyptemys insculpta</i>	Wood Turtle	1	1	1	1	1			1	1			1	1		
<i>Heterodon nasicus</i>	Plains Hog-nosed Snake						2						1	1		
<i>Heterodon platirhinos</i>	Eastern Hog-nosed Snake	1	1		1		2						1	1	1	
<i>Opheodrys vernalis</i>	Smooth Greensnake		1				2						1	1		
<i>Pantherophis obsoletus</i>	Western Ratsnake		1	1	1									1	1	
<i>Pituophis catenifer</i>	Gophersnake		1				2						1	1	1	
<i>Plestiodon fasciatus</i>	Common Five-lined Skink		1	1			2						1	1	1	
<i>Tropidoclonion lineatum</i>	Lined Snake												1	1	1	

*Backwaters

Bees (Class Insecta; Order Hymenoptera)

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Non- forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Andrena accepta</i>	two-spotted miner bee							X			
<i>Andrena arabis</i>	mustard miner bee		X	X							
<i>Andrena bradleyi</i>	Bradley's Miner Bee		X								
<i>Andrena brevipalpis</i>	Short-tongued Miner Bee		X					X	X		
<i>Andrena campanulae</i>	a miner bee							X			
<i>Andrena canadensis</i>	Canada Miner Bee							X	X		
<i>Andrena carolina</i>	Carolina Miner Bee		X								
<i>Andrena clarkella</i>	Clark's Miner Bee				X		X			X	
<i>Andrena distans</i>	Distant Miner Bee		X	X					X		
<i>Andrena erigeniae</i>	Spring Beauty Miner Bee			X							
<i>Andrena erythrogaster</i>	Red-bellied Miner Bee				X						
<i>Andrena erythronii</i>	Trout Lily Miner Bee			X							
<i>Andrena frigida</i>	Cold Miner Bee				X						
<i>Andrena haynesi</i>	Haynes's miner bee							X		X	
<i>Andrena helianthiformis</i>	a mining bee							X			
<i>Andrena integra</i>	Intact Miner Bee				X						
<i>Andrena lupinorum</i>	Lupine Miner Bee							X			
<i>Andrena mariaae</i>	Maria Miner Bee				X						
<i>Andrena melanothroa</i>	Rose Miner Bee		X								
<i>Andrena parnassiae</i>	Parnassia Miner				X			X			
<i>Andrena peckhami</i>	Peckham's Miner Bee							X			
<i>Andrena persimulata</i>	Protuberance Miner Bee				X						
<i>Andrena placata</i>	Peaceful Miner Bee							X			
<i>Andrena platyparia</i>	Plated Miner Bee				X						
<i>Andrena polemonii</i>	an andrenid bee			X							
<i>Andrena quintilis</i>	Eastern Scaly Miner Bee							X			
<i>Andrena sigmundi</i>	Andrena sigmundi				X						
<i>Andrena simplex</i>	Simple Miner Bee							X			

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Non- forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Andrena uvulariae</i>	a mining bee			X							
<i>Andrena vernalis</i>	a mining bee							X			
<i>Andrena wellesleyana</i>	Wellesley Miner Bee				X						
<i>Anthidiellum boreale</i>	a rotund resin bee							X			
<i>Anthidium psoraleae</i>	a leafcutter bee							X			
<i>Bombus affinis</i>	Rusty Patched Bumble Bee			X				X			X
<i>Bombus ashtoni</i>	a bumble bee	X	X								
<i>Bombus fervidus</i>	Yellow Bumble Bee							X			
<i>Bombus fraternus</i>	southern plains bumble bee							X			
<i>Bombus frigidus</i>	A Bumble Bee				X						
<i>Bombus insularis</i>	indiscriminate bumble bee										
<i>Bombus pensylvanicus</i>	American Bumble Bee							X			
<i>Bombus terricola</i>	Yellowbanded Bumble Bee	X	X		X						
<i>Brachymelecta interrupta</i>	Interrupted Xeromelecta Bee							X			
<i>Calliopsis nebraskensis</i>	a mining bee							X			
<i>Chelostoma philadelphia</i>	a striped hairy belly bee										X
<i>Colletes aberrans</i>	Aberrant Cellophane Bee							X			
<i>Colletes albescens</i>	White Plasterer Bee							X			
<i>Colletes americanus</i>	a cellophane bee							X			
<i>Colletes andrewsi</i>	Andrews' cellophane bee							X			
<i>Colletes brevicornis</i>	Short-horned Plasterer Bee							X			
<i>Colletes ciliatus</i>	a cellophane bee				X			X			
<i>Colletes latitarsis</i>	Ground-cherry Plasterer Bee							X		X	
<i>Colletes mandibularis</i>	Mandibular Plasterer Bee							X			
<i>Colletes solidaginis</i>	Goldenrod Plasterer Bee							X			
<i>Colletes susannae</i>	Susan's Plasterer Bee							X			
<i>Colletes willistoni</i>	Williston's Cellophane Bee							X			
<i>Colletes wilmattae</i>	Wilmatte's Plasterer Bee							X		X	
<i>Dianthidium pudicum</i>	Modest pebble bee							X			

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Non- forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Dianthidium simile</i>	a leafcutter bee							X			
<i>Dieunomia heteropoda</i>	Sunflower Sweat Bee							X			
<i>Dufourea harveyi</i>	a solitary bee				X			X			
<i>Dufourea marginata</i>	Glowing Sweat Bee										
<i>Dufourea novaeangliae</i>	New England Sweat Bee					X	X				
<i>Epeoloides pilosulus</i>	Macropis Cuckoo Bee							X			
<i>Epeolus ainsliei</i>	a cuckoo bee							X		X	
<i>Epeolus lectoides</i>	Collector Cuckoo Nomad Bee							X		X	
<i>Eucera albata</i>	an anthophorid bee							X			
<i>Heriades leavitti</i>	A resin bee							X			
<i>Hylaeus sparsus</i>	a masked bee			X							
<i>Lasioglossum fedorense</i>	a sweat bee							X		X	
<i>Lasioglossum floridanum</i>	a sweat bee							X		X	
<i>Lasioglossum lusorium</i>	a sweat bee							X			
<i>Lasioglossum nelumbonis</i>	Water-lily Sweat Bee						X				
<i>Lasioglossum swenki</i>	Swenk's Sweat Bee							X		X	
<i>Lasioglossum texanum</i>	a sweat bee							X			
<i>Macropis nuda</i>	Common Loosestrife Oil Bee				X			X			
<i>Macropis patellata</i>	Patellar Oil-collecting Bee				X			X			
<i>Macropis steironematis</i>	an oil-collecting bee				X			X			
<i>Megachile addenda</i>	a leafcutter bee							X		X	
<i>Megachile circumcincta</i>	a leafcutter bee							X			
<i>Megachile fortis</i>	a leafcutter bee							X			
<i>Megachile rugifrons</i>	a leafcutter bee							X			
<i>Melissodes apicatus</i>	a long-horned bee						X				
<i>Melissodes bidentis</i>	Two-toothed Long-horned Bee							X			
<i>Melissodes boltoniae</i>	a callirhoe bee							X		X	
<i>Melissodes confusus</i>	a callirhoe bee							X			
<i>Melissodes denticulatus</i>	A Callirhoe Bee							X			

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Non- forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Melissodes dentiventris</i>	A Callirhoe Bee							X			
<i>Melissodes illatus</i>	A Callirhoe Bee							X			
<i>Melissodes menuachus</i>	A Callirhoe Bee							X			
<i>Melissodes subagilis</i>	A Callirhoe Bee							X			
<i>Melissodes wheeleri</i>	A Callirhoe Bee							X			
<i>Neolarra vigilans</i>	a cuckoo bee							X		X	
<i>Nomia universitatis</i>	a nomia bee							X			
<i>Osmia cyaneonitens</i>	a mason bee							X			
<i>Osmia distincta</i>	Distinct Mason Bee							X	X		
<i>Osmia felti</i>	a mason bee	X	X							X	
<i>Osmia inspergens</i>	Shiny-faced mason bee	X									
<i>Osmia laticeps</i>	a mason bee				X						
<i>Osmia nearctica</i>	a mason bee	X									
<i>Osmia nigriventris</i>	Large Black-bellied Mason Bee				X						
<i>Osmia virga</i>	a mason bee				X						
<i>Perdita albipennis</i>	Shiny Green Miner Bee							X			
<i>Perdita bruneri</i>	Bruner's Miner Bee							X			
<i>Perdita dolichocephala</i>	a fairy bee							X			
<i>Perdita gerhardi</i>	Gerhard's Perdita							X		X	
<i>Perdita halictoides</i>	Sweatbee-like Miner Bee							X			
<i>Perdita ignota</i>	a miner bee							X			
<i>Perdita pallidipennis</i>	a tiny bee							X			
<i>Perdita perpallida</i>	Pale Miner Bee							X		X	
<i>Protandrena aestivalis</i>	Summer Miner Bee							X			
<i>Protandrena andrenoides</i>	Small Black Miner Bee							X			
<i>Protandrena renimaculata</i>	medium dark bee							X			
<i>Protandrena rudbeckiae</i>	Rudbeckia Miner Bee							X			
<i>Protandrena simulans</i>	a mining bee							X			
<i>Sphecodes banksii</i>	a sweat bee							X			

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Non- forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Stelis permaculata</i>	Sixteen-spotted Cuckoo Carder Bee							X			
<i>Triepeolus concavus</i>	Concave Cuckoo Nomad Bee							X			

Beetles (Class Insecta; Order Coleoptera)

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Riparian and Floodplain Forest	Non- forested Wetlands	Streams and Rivers	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Adalia bipunctata</i>	Two-spotted Lady Beetle	X	X	X					X	X		X
<i>Cicindela denikei</i>	Laurentian Tiger Beetle	X									X	
<i>Cicindela fulgida</i>	Crimson Saltflat Tiger Beetle							X				
<i>Cicindela hirticollis rhodensis</i>	Hairy-necked Tiger Beetle							X				
<i>Cicindela lepida</i>	Ghost Tiger Beetle								X	X	X	
<i>Cicindela limbata</i>	Sandy Tiger Beetle									X	X	
<i>Cicindela macra</i>	Sandy Stream Tiger Beetle						X					
<i>Cicindela patruela</i>	Northern Barrens Tiger Beetle									X		
<i>Cicindela splendida</i>	Splendid Tiger Beetle										X	
<i>Coccinella novemnotata</i>	Nine-spotted Lady Beetle				X				X	X		X
<i>Coccinella transversoguttata</i>	Transverse Lady Beetle				X				X	X		X
<i>Coelambus sylvanus</i>	Sylvan Hygrotus Diving Beetle					X						

Butterflies, Moths (Class Insecta; Order Lepidoptera)

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non- forested Wetlands	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Agonopterix pergandeella</i>								X			
<i>Ambesa laetella</i>								X		X	
<i>Ancylis semiovana</i>									X		
<i>Anisota manitobensis</i>	Manitoba Oakworm Moth								X		
<i>Apamea commoda</i>		X							X		
<i>Apamea niveivenosa</i>							X	X			
<i>Apantesis doris</i>									X		
<i>Apantesis margo</i>		X							X		
<i>Apodrepanulatrix liberaria</i>							X		X		
<i>Argynnis idalia</i>	Regal Fritillary							X			
<i>Aspitates aberrata</i>								X			
<i>Atrytone arogos</i>	Arogos Skipper							X			
<i>Atrytonopsis hianna</i>	Dusted Skipper							X	X	X	
<i>Bagisara rectifascia</i>	Straight Lined Mallow Moth								X		
<i>Bandera binotella</i>								X		X	
<i>Boloria chariclea</i>	Arctic Fritillary					X					
<i>Boloria freija</i>	Freija Fritillary	X				X	X				
<i>Boloria frigga</i>	Frigga Fritillary						X				
<i>Cabera quadrifasciaria</i>								X			
<i>Capis archaia</i>							X				
<i>Carmenta anthracipennis</i>								X	X		
<i>Carsia sororiata</i>	Manchester Treble-bar Moth					X	X				
<i>Catocala abbreviatella</i>	Abbreviated Underwing							X	X		
<i>Catocala amestris</i>	Three-staff Underwing							X	X		
<i>Catocala nuptialis</i>	Married Underwing						X	X			
<i>Catocala whitneyi</i>	Whitney's Underwing							X	X		

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non- forested Wetlands	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Chaetagnathaea cerata</i>								X	X		
<i>Chlosyne gorgone</i>	Gorgone Checkerspot							X	X		
<i>Clepsia illustrata</i>		X				X					
<i>Crambus awemellus</i>							X				
<i>Danaus plexippus</i>	Monarch							X	X		X
<i>Dargida rubripennis</i>	a noctuid moth							X	X		
<i>Eoreuma crawfordi</i>								X			
<i>Erastris coloraria</i>									X		
<i>Erebia discoidalis</i>	Red-disked Alpine	X					X				
<i>Erebia mancinus</i>	Taiga Alpine					X					
<i>Erynnis lucilius</i>	Columbine Duskywing							X		X	
<i>Erynnis martialis</i>	Mottled Duskywing								X		
<i>Erynnis persius</i>	Persius Duskywing								X		
<i>Euchloe ausonides</i>	Large Marble	X							X	X	
<i>Euchloe olympia</i>	Olympia Marble							X	X	X	
<i>Eucosma giganteana</i>	Giant Eucosma Moth							X			
<i>Eucosma lapidana</i>								X			
<i>Eucosma millerana</i>	a tortricid moth							X			
<i>Eucosma pallidicostana</i>								X			
<i>Euphydryas phaeton</i>	Baltimore Checkerspot						X				
<i>Euphyes conspicua</i>	Black Dash						X				
<i>Euxoa niveilinea</i>	a dart moth							X		X	
<i>Euxoa quadridentata</i>										X	
<i>Euxoa quebecensis</i>										X	
<i>Feltia manifesta</i>								X		X	
<i>Habrosyne gloriosa</i>			X						X		
<i>Heliothis borealis</i>						X	X		X		
<i>Hemaris gracilis</i>		X				X	X		X		
<i>Hesperia assiniboia</i>	Assiniboia Skipper							X		X	

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non- forested Wetlands	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Hesperia dacotae</i>	Dakota Skipper							X			
<i>Hesperia leonardus leonardus</i>	Leonard's Skipper							X		X	
<i>Hesperia leonardus pawnee</i>	Pawnee Skipper							X			
<i>Hesperia ottoe</i>	Ottoe Skipper							X			
<i>Hystrichophora taleana</i>								X	X		
<i>Interjectio niviella</i>								X			
<i>Lasionycta secedens</i>						X					
<i>Lasionycta taigata</i>	a Noctuid moth					X					
<i>Lithophane franclemonti</i>			X								
<i>Lithophane lemmeri</i>									X	X	
<i>Lithophane scottae</i>			X								
<i>Lobocleta plemyraria</i>								X	X	X	
<i>Melaporphyria immortua</i>								X	X		
<i>Melitara dentata</i>								X			
<i>Meroptera mirandella</i>								X			
<i>Microtheoris ophionalis</i>								X		X	
<i>Mimoschinia rufofascialis</i>								X	X		
<i>Oarisma garita</i>	Garita Skipperling							X		X	
<i>Oarisma poweshiek</i>	Poweshiek Skipperling							X			
<i>Oeneis macounii</i>	Macoun's Arctic	X								X	
<i>Oligia bridghamii</i>							X				
<i>Oligia egens</i>							X	X			
<i>Papaipema aweme</i>							X				
<i>Papaipema beeriana</i>	Blazing Star Stem Borer							X			
<i>Papaipema marginidens</i>			X								
<i>Papaipema nepheleptena</i>							X				
<i>Papaipema sciata</i>	Culvers Root Borer							X	X		
<i>Pelochrista heathiana</i>								X	X		
<i>Pelochrista nandana</i>								X			

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non- forested Wetlands	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Pelochrista robinsonana</i>								X	X	X	
<i>Pelochrista scintillana</i>								X	X		
<i>Phoberia ingenua</i>									X		
<i>Plagodis kuetzingi</i>				X	X						
<i>Plebejus idas</i>	Northern Blue	X									
<i>Plebejus samuelis</i>	Karner Blue								X		
<i>Poanes massasoit</i>	Mulberry Wing						X				
<i>Polia propodea</i>						X					
<i>Polites origenes</i>	Crossline Skipper							X			
<i>Ponometia binocula</i>	Prairie Bird-dropping Moth							X	X		
<i>Proserpinus juanita</i>								X			
<i>Protoschinia nuchalis</i>								X		X	
<i>Pyrausta pythialis</i>								X	X		
<i>Pyrausta scurralis</i>		X							X		
<i>Pyrausta socialis</i>	Sociable Pyrausta Moth							X	X		
<i>Schinia bina</i>	Bina Flower Moth							X			
<i>Schinia grandimedia</i>	Rockies Boneset Flower Moth							X			
<i>Schinia hultia</i>								X			
<i>Schinia indiana</i>	Phlox Moth							X	X		
<i>Schinia jaguarina</i>	Jaguar Flower Moth							X			
<i>Schinia lucens</i>	Leadplant Flower Moth							X	X		
<i>Schinia lynx</i>								X	X	X	
<i>Schinia mortua</i>								X			
<i>Schinia obscurata</i>								X	X	X	
<i>Schinia septentrionalis</i>	Northern Flower Moth							X			
<i>Schizura apicalis</i>		X							X		
<i>Sparganothis violaceana</i>	A Tortricid Moth					X	X				
<i>Stiria rugifrons</i>	Yellow Sunflower Moth							X			

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Deciduous Wet Forest	Riparian and Floodplain Forest	Coniferous Forest Wetland	Non-forested Wetlands	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop	Urban and Other Developed Lands
<i>Sympistis piffardi</i>							X	X	X		
<i>Sympistis stabilis</i>								X	X		
<i>Sympistis viriditincta</i>								X	X		
<i>Syngrapha montana</i>						X					
<i>Tacparia atropunctata</i>						X					
<i>Thaumatopsis pectinifer</i>	a crambid snout moth							X		X	
<i>Walshia amorphella</i>								X	X		
<i>Xestia imperita</i>						X					
<i>Xestia mixta</i>						X					
<i>Zenopheps lignicolorata</i>		X									

Snails (Class Gastropoda)

Scientific Name	Common Name	Upland Conifer Forest and Woodlands	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Coniferous Forest Wetland	Lakes	Prairie and Other Grasslands	Cliff, Talus, Rock Outcrop
<i>Acella haldemani</i>	Spindle Lymnaea						1		
<i>Gastrocopta rogersensis</i>	Rogers Snaggletooth Snail	2	2	2				2	1
<i>Gastrocopta similis</i>	Great Lakes Snaggletooth	2	2	2				2	1
<i>Planogyra asteriscus</i>	Eastern Flat-whorl	1		1		1			1
<i>Striatura ferrea</i>	Black Striate			1		2			1
<i>Vallonia perspectiva</i>	Thin-lip Vallonia	2	1	2					1
<i>Vertigo gouldi</i>	Variable Vertigo	2	1	2					1
<i>Vertigo meramecensis</i>	Bluff Vertigo		1	1	2				1
<i>Zonitoides limatulus</i>	Dull Gloss		2	1					1

Spiders (Class Arachnida; Order Araneae)

Scientific Name	Common Name	Upland Deciduous Forests and Woodlands	Mesic Hardwood Forest	Deciduous Wet Forest	Riparian and Floodplain Forest	Non-forested Wetlands	Lakes	Prairie and Other Grasslands	Savanna	Cliff, Talus, Rock Outcrop
<i>Habronattus texanus</i>	A Jumping Spider							1	1	
<i>Latrodectus variolus</i>	Northern black widow	1	1	2	2			2	1	2
<i>Neoscona pratensis</i>	An orb weaver					1	2	2		
<i>Phidippus apacheanus</i>	A Jumping Spider							1	1	
<i>Sassacus papenhoei</i>	A Jumping Spider							1	1	
<i>Sphodros niger</i>	Black purse-web spider	2	1	2	2					
<i>Tutelina formicaria</i>	A Jumping Spider							1	1	

True Bugs, Cicadas, Leafhoppers (Class Insecta; Order Hemiptera)

Scientific Name	Common Name	Prairie and Other Grasslands
<i>Aflexia rubranura</i>	Red-tailed Prairie Leafhopper	X
<i>Okanagana balli</i>	Prairie Cicada	X

Caddisflies (Class Insecta; Order Trichoptera)

Scientific Name	Common Name	Riverine (general)	Streams (SSO 1-4)	Rivers (SSO 5-6)	Large Rivers (SSO 7-8)	Lakes
<i>Agapetus tomus</i>	A Caddisfly		1*	2*	2*	
<i>Agrypnia glacialis</i>	A Caddisfly				2	2
<i>Anabolia ozburni</i>	A Caddisfly		1	1		1
<i>Anabolia sordida</i>	A Caddisfly		1	1		1
<i>Apatania zonella</i>	A Caddisfly	1				1
<i>Asynarchus rossi</i>	A Caddisfly		1	2		
<i>Banksiola smithi</i>	A Caddisfly	2				1

Scientific Name	Common Name	Riverine (general)	Streams (SSO 1-4)	Rivers (SSO 5-6)	Large Rivers (SSO 7-8)	Lakes
<i>Ceraclea albosticta</i>	A Caddisfly		1			
<i>Ceraclea brevis</i>	A Caddisfly	1				1
<i>Ceraclea misca</i>	A Caddisfly		1	2		
<i>Cheumatopsyche wabasha</i>	A Caddisfly	1				
<i>Chilostigma itascaae</i>	Headwaters Chilostigman Caddisfly	1				1**
<i>Goera stylata</i>	A Caddisfly	1*				
<i>Grammotaulius interrogationis</i>	A Caddisfly	1				1**
<i>Holocentropus glacialis</i>	Frozen Trumpet-net Caddisfly					1
<i>Holocentropus melanae</i>	Black River Trumpet-net Caddisfly					1
<i>Holocentropus milaca</i>	A Caddisfly					1**
<i>Hydropsyche californica</i>	A Caddisfly	1				
<i>Hydroptila callia</i>	A Caddisfly					1**
<i>Hydroptila delineata</i>	A Caddisfly				1	
<i>Hydroptila metoecca</i>	A Caddisfly	1				2
<i>Hydroptila quinola</i>	A Caddisfly		1	1		
<i>Hydroptila rono</i>	A Caddisfly		1			
<i>Hydroptila tortosa</i>	A Caddisfly		1			
<i>Ironoquia punctatissima</i>	A Caddisfly		1			
<i>Lenarchus keratus</i>	A Caddisfly					1
<i>Lepidostoma libum</i>	A Caddisfly	1				
<i>Limnephilus argenteus</i>	A Caddisfly	1				
<i>Limnephilus janus</i>	A Caddisfly		1			
<i>Limnephilus parvulus</i>	A Caddisfly		1	2		2
<i>Limnephilus sackeni</i>	A Caddisfly					1
<i>Limnephilus secludens</i>	A Caddisfly		1			
<i>Limnephilus tarsalis</i>	A Caddisfly	2				1
<i>Micrasema sprulesi</i>	Sprules's Humpless Caddisfly			1	1	
<i>Oecetis ditissa</i>	A Caddisfly	1				1
<i>Orthotrichia curta</i>	Short Orthotrichian Microcaddisfly	2				1
<i>Oxyethira ecornuta</i>	A Caddisfly					1

Scientific Name	Common Name	Riverine (general)	Streams (SSO 1-4)	Rivers (SSO 5-6)	Large Rivers (SSO 7-8)	Lakes
<i>Oxyethira itasca</i>	A Caddisfly		2	1		
<i>Parapsyche apicalis</i>	A Caddisfly		1*			
<i>Philarctus bergrothi</i>	Bergroth's Northern Caddisfly	1				1**
<i>Protophila erotica</i>	A Caddisfly			1	1	
<i>Psilotreta indecisa</i>	Uncertain Mortarjoint Caddisfly		1*	1*		
<i>Pycnopsyche scabripennis</i>	A Caddisfly		1	2		
<i>Rhyacophila angelita</i>	A Caddisfly		1*			
<i>Rhyacophila brunnea</i>	Brown Free-living Caddisfly		1*			
<i>Rhyacophila minor</i>	Smaller Free-living Caddisfly		1*			
<i>Rhyacophila vibox</i>	A Rhyacophilan Caddisfly		1*	1		
<i>Trienodes flavescens</i>	A Trienode Caddisfly		1			
<i>Trienodes reuteri</i>	A Caddisfly	1				
<i>Wormaldia moesta</i>	A Caddisfly		1	1		

*Cold water specialist

**Shallow lakes

Dragonflies and Damselflies (Class Insecta; Order Odonata)

Scientific Name	Common Name	Deciduous Wet Forest	Coniferous Forest Wetland	Non- forested Wetlands	Vernal Pools	Streams (SSO 1-4)	Rivers (SSO 5-6)	Large Rivers (SSO 7-8)	Shallow Lakes	Deep Lakes
<i>Aeshna sitchensis</i>	Zigzag Darner			X						
<i>Aeshna subarctica</i>	Subarctic Darner			X						
<i>Archilestes grandis</i>	Great Spreadwing	X	X							X
<i>Argia funebris</i>	Springwater Dancer				X					X
<i>Arigomphus submedianus</i>	Jade Clubtail					X			X	
<i>Boyeria grafiana</i>	Ocellated Darner				X	X				X
<i>Coenagrion angulatum</i>	Prairie Bluet							X		X
<i>Coenagrion interrogatum</i>	Subarctic Bluet			X						
<i>Enallagma anna</i>	River Bluet									

Scientific Name	Common Name	Deciduous Wet Forest	Coniferous Forest Wetland	Non-forested Wetlands	Vernal Pools	Streams (SSO 1-4)	Rivers (SSO 5-6)	Large Rivers (SSO 7-8)	Shallow Lakes	Deep Lakes
<i>Enallagma aspersum</i>	Azure Bluet				X				X	
<i>Gomphurus lineatifrons</i>	Splendid Clubtail					X	X			
<i>Gomphurus ventricosus</i>	Skillet Clubtail					X				X
<i>Hylogomphus viridifrons</i>	Green-faced Clubtail									X
<i>Nannothemis bella</i>	Elfin Skimmer			X	X					
<i>Ophiogomphus anomalus</i>	Extra-striped Snaketail					X				
<i>Ophiogomphus howei</i>	Pygmy Snaketail						X			
<i>Ophiogomphus smithi</i>	Sioux Snaketail					X				X
<i>Ophiogomphus susbehcha</i>	St. Croix Snaketail					X				
<i>Phanogomphus quadricolor</i>	Rapids Clubtail					X				X
<i>Somatochlora brevicincta</i>	Quebec Emerald			X	X					
<i>Somatochlora forcipata</i>	Forcipate Emerald			X						
<i>Stylogomphus albistylus</i>	Least Clubtail					X				X
<i>Stylurus notatus</i>	Elusive Clubtail					X	X		X	
<i>Williamsonia fletcheri</i>	Ebony Boghaunter			X						
<i>Zoraena obliqua</i>	Arrowhead Spiketail				X					X

Mayflies (Class Insecta; Order Ephemeroptera)

Scientific Name	Common Name	Riverine (general)	Streams (SSO 1-4)	Rivers (SSO 5-6)	Lakes
<i>Ameletus subnotatus</i>	Boreal Comb Minnow Mayfly		1	1	
<i>Drunella cornuta</i>	Big-horned Spiny Crawler Mayfly		1	2	
<i>Drunella walkeri</i>	Walker's Spiny Crawler Mayfly		1	2	
<i>Eurylophella funeralis</i>	Brown Spiny Crawler Mayfly		1		
<i>Neoephemera bicolor</i>	Canadian Large Square-gilled Mayfly	1			
<i>Siphonurus alternatus</i>	Northern Primitive Minnow Mayfly		1		1
<i>Siphonurus phyllis</i>	Leafy Primitive Minnow Mayfly		1		1
<i>Sparbarus maculatus</i>	Maculated Small Square-gilled Mayfly		2	1	

Mussels (Class Bivalvia)

Scientific Name	Common Name	Non-forested Wetlands	Streams (SSO 1-4)	Rivers (SSO 5-6)	Large Rivers (SSO 7-8)	Lakes
<i>Actinonaias ligamentina</i>	Mucket		1	1	2	
<i>Alasmidonta marginata</i>	Elktoe		1	1	1	
<i>Arcidens confragosus</i>	Rock Pocketbook				1	
<i>Cumberlandia monodonta</i>	Spectaclecase			2	1	
<i>Cyclonaias nodulata</i>	Wartyback				1	
<i>Cyclonaias tuberculata</i>	Purple Wartyback			1	1	
<i>Ellipsaria lineolata</i>	Butterfly				1	
<i>Elliptio complanata</i>	Eastern Elliptio		1	2	1	
<i>Elliptio crassidens</i>	Elephant-ear				1	
<i>Epioblasma triquetra</i>	Snuffbox				1	
<i>Eurynia dilatata</i>	Spike		1	1	1	
<i>Lampsilis higginsii</i>	Higgins Eye				1	
<i>Lampsilis sietmani</i>	Canary Kingshell				1	
<i>Lampsilis teres</i>	Yellow Sandshell				1	
<i>Lasmigona compressa</i>	Creek Heelsplitter		1	1	2	
<i>Lasmigona costata</i>	Flutedshell		2	1	1	
<i>Ligumia recta</i>	Black Sandshell		2	1	1	
<i>Megalonaias nervosa</i>	Washboard				1	
<i>Obovaria olivaria</i>	Hickorynut			2	1	
<i>Plethobasus cyphus</i>	Sheepnose			2	1	
<i>Pleurobema sintoxia</i>	Round Pigtoe		2	1	1	
<i>Potamilus capax</i>	Fat Pocketbook				1	
<i>Potamilus leptodon</i>	Scaleshell				1	
<i>Quadrula fragosa</i>	Winged Mapleleaf				1	
<i>Reginaia eburnus</i>	Ebonyshell				1	
<i>Sagittunio subrostratus</i>	Pondmussel		1	2		
<i>Simpsonaias ambigua</i>	Salamander Mussel				1	
<i>Theliderma metanevra</i>	Monkeyface			2	1	
<i>Tritogonia nobilis</i>	Gulf Mapleleaf				1	
<i>Tritogonia verrucosa</i>	Pistolgrip				1	

<i>Truncilla donaciformis</i>	Fawnsfoot				1	
<i>Utterbackiana suborbiculata</i>	Flat Floater	2				1
<i>Venustaconcha ellipsiformis</i>	Ellipse		1	1		

Stoneflies (Class Insecta; Order Plecoptera)

Scientific Name	Common Name	Streams (SSO 1-4)	Rivers (SSO 5-6)
<i>Alloperla atlantica</i>	A stonefly	1*	1*
<i>Attaneuria ruralis</i>	A stonefly	1	1
<i>Isogenoides frontalis</i>	A Perlid Stonefly	1*	1*
<i>Paracapnia opis</i>	A stonefly	1	1
<i>Paranemoura perfecta</i>	Spotted Forestfly	1*	

*Coldwater specialist

APPENDIX D - Plant SGCN Associations to Primary Habitat Types

Plant SGCN are organized here by the SWAP chapter that includes the species' primary Ecological System (ES).

Primary Ecological System:

UP = Upland Prairie

WP = Wet Prairie

FD = Fire-Dependent Forest

MH = Mesic Hardwood

RO = Rock Outcrop

CT = Cliff/Talus

LK = Lakes

OP = Open Rich Peatland

MW = Wet Meadow/Carr

MR = Marsh

APn8 or APn9 type = Acid Peatland

FF = Floodplain Forest

FP = Forested Rich Peatlands

WF = Wet Forest

RV = Riverine

Prairie and Other Grasslands

Scientific Name	Common Name	Family	Primary Ecological System
<i>Achnatherum hymenoides</i>	Indian Ricegrass	Poaceae	UP
<i>Agalinis auriculata</i>	Eared False Foxglove	Orobanchaceae	WP
<i>Agalinis gattereri</i>	Round-stemmed False Foxglove	Orobanchaceae	UP
<i>Agrostis hyemalis</i>	Winter Bentgrass	Poaceae	UP
<i>Allium textile</i>	White Wild Onion	Liliaceae	UP
<i>Antennaria microphylla</i>	Small-leaf Cat's-foot	Asteraceae	UP
<i>Antennaria parvifolia</i>	Small-leaved Pussytoes	Asteraceae	UP
<i>Aristida longespica</i>		Poaceae	UP
<i>Aristida purpurea</i>		Poaceae	UP
<i>Aristida tuberculosa</i>	Seaside Three-awn	Poaceae	UP
<i>Arnoglossum plantagineum</i>	Tuberous Indian Plantain	Asteraceae	UP
<i>Asclepias amplexicaulis</i>	Clasping Milkweed	Apocynaceae	UP
<i>Asclepias hirtella</i>	Prairie Milkweed	Apocynaceae	UP
<i>Asclepias lanuginosa</i>	Woolly Milkweed	Apocynaceae	UP
<i>Asclepias stenophylla</i>	Narrow-leaved Milkweed	Apocynaceae	UP
<i>Asclepias sullivantii</i>	Sullivant's Milkweed	Apocynaceae	WP
<i>Astragalus flexuosus</i>	Slender Milk-vetch	Fabaceae	UP
<i>Astragalus missouriensis</i>	Missouri Milk-vetch	Fabaceae	UP
<i>Astragalus neglectus</i>	Cooper's Milk-vetch	Fabaceae	UP
<i>Avenula hookeri</i>	Spike Oat	Poaceae	UP
<i>Baptisia bracteata</i>	Cream Wild Indigo	Fabaceae	UP
<i>Baptisia lactea</i>	White Wild Indigo	Fabaceae	UP
<i>Botrychium campestre</i> var. <i>campestre</i>	Prairie Moonwort	Ophioglossaceae	UP
<i>Botrychium gallicomontanum</i>	Frenchman's Bluff Moonwort	Ophioglossaceae	UP
<i>Calamagrostis montanensis</i>	Plains Reedgrass	Poaceae	UP
<i>Callirhoe triangulata</i>	Clustered Poppy-mallow	Malvaceae	UP
<i>Carex annectens</i>	Yellow-fruit Sedge	Cyperaceae	UP
<i>Carex filifolia</i>	Thread-leaved Sedge	Cyperaceae	UP
<i>Carex hallii</i>	Hall's Sedge	Cyperaceae	WP
<i>Carex hookerana</i>	Hooker's Sedge	Cyperaceae	UP
<i>Carex obtusata</i>	Blunt Sedge	Cyperaceae	UP
<i>Carex scirpoidea</i>	Northern Single-spike Sedge	Cyperaceae	UP
<i>Carex xerantica</i>	Dry Sedge	Cyperaceae	UP
<i>Chamaerhodos erecta</i>	Nuttall's Groundrose	Rosaceae	UP
<i>Commelina erecta</i>	Slender Dayflower	Commelinaceae	UP

Scientific Name	Common Name	Family	Primary Ecological System
<i>Conyza ramosissima</i>		Asteraceae	UP
<i>Crotalaria sagittalis</i>	Rattlebox	Fabaceae	UP
<i>Cypripedium candidum</i>	Small White Lady's-slipper	Orchidaceae	UP
<i>Dalea candida</i> var. <i>oligophylla</i>	Western White Prairie-clover	Fabaceae	UP
<i>Dalea leporina</i>		Fabaceae	UP
<i>Descurainia pinnata</i>		Brassicaceae	UP
<i>Desmanthus illinoensis</i>	Prairie Mimosa	Fabaceae	UP
<i>Desmodium illinoense</i>	Illinois Tick-trefoil	Fabaceae	UP
<i>Dodecatheon meadia</i>	Prairie Shooting Star	Primulaceae	UP
<i>Eragrostis trichodes</i>		Poaceae	UP
<i>Erigeron lonchophyllus</i>	Short Ray Fleabane	Asteraceae	WP
<i>Eryngium yuccifolium</i>	Rattlesnake Master	Apiaceae	UP
<i>Eupatorium altissimum</i>		Asteraceae	UP
<i>Euphorbia dentata</i>		Euphorbiaceae	UP
<i>Euphorbia missurica</i>	Missouri Spurge	Euphorbiaceae	UP
<i>Euphorbia spathulata</i>		Euphorbiaceae	UP
<i>Gaillardia aristata</i>	Blanketflower	Asteraceae	UP
<i>Gentiana affinis</i>	Northern Gentian	Gentianaceae	WP
<i>Gentianella amarella</i> ssp. <i>acuta</i>	Felwort	Gentianaceae	UP & WP
<i>Gentianella quinquefolia</i>	Stiff Gentian	Gentianaceae	UP
<i>Geranium carolinianum</i>		Geraniaceae	UP
<i>Grindelia hirsutula</i>	Hairy Gumweed	Asteraceae	UP
<i>Helianthus nuttallii</i>	Nuttall's Sunflower	Asteraceae	UP
<i>Hesperostipa curtiseta</i>	Small Porcupine Grass	Poaceae	UP
<i>Hieracium longipilum</i>	Long-bearded Hawkweed	Asteraceae	UP
<i>Hudsonia tomentosa</i>	Beach Heather	Cistaceae	UP
<i>Isanthus brachiatus</i>		Lamiaceae	UP
<i>Juniperus horizontalis</i>	Creeping Juniper	Cupressaceae	UP
<i>Lechea tenuifolia</i>	Narrow-leaved Pinweed	Cistaceae	UP
<i>Lespedeza leptostachya</i>	Prairie Bush Clover	Fabaceae	UP
<i>Linum lewisii</i>		Linaceae	UP
<i>Lomatium orientale</i>	Desert Parsley	Apiaceae	UP
<i>Lotus purshianus</i>		Fabaceae	UP
<i>Lupinus perennis</i>	Wild Lupine	Fabaceae	UP
<i>Minuartia dawsonensis</i>	Rock Sandwort	Caryophyllaceae	UP
<i>Nothocalais cuspidata</i>		Asteraceae	UP
<i>Oenothera nuttallii</i>	Nuttall's Evening Primrose	Onagraceae	UP
<i>Oenothera rhombipetala</i>	Rhombic Evening Primrose	Onagraceae	UP

Scientific Name	Common Name	Family	Primary Ecological System
<i>Ophioglossum pusillum</i>	Adder's Tongue	Ophioglossaceae	WP
<i>Orobanche fasciculata</i>	Clustered Broomrape	Orobanchaceae	UP
<i>Orobanche uniflora</i>	One-flowered Broomrape	Orobanchaceae	UP
<i>Packera cana</i>	Gray Ragwort	Asteraceae	UP
<i>Paronychia canadensis</i>	Canada Forked Chickweed	Caryophyllaceae	UP
<i>Parthenium integrifolium</i>	Wild Quinine	Asteraceae	UP
<i>Penstemon albidus</i>		Plantaginaceae	UP
<i>Phemeranthus rugospermus</i>	Rough-seeded Fameflower	Portulacaceae	UP
<i>Phlox maculata</i>	Wild Sweetwilliam	Polemoniaceae	WP
<i>Platanthera flava</i> var. <i>herbiola</i>	Southern Rein-orchid	Orchidaceae	WP
<i>Platanthera praeclara</i>	Western Prairie Fringed Orchid	Orchidaceae	WP
<i>Poa arida</i>	Plains Bluegrass	Poaceae	WP
<i>Polanisia jamesii</i>	James' Polanisia	Cleomaceae	UP
<i>Polytaenia nuttallii</i>	Prairie Parsley	Apiaceae	UP
<i>Potentilla bipinnatifida</i>		Rosaceae	UP
<i>Potentilla hippiana</i>	Woolly Cinquefoil	Rosaceae	UP
<i>Potentilla lasiodonta</i>	Sandhills Cinquefoil	Rosaceae	UP
<i>Prenanthes aspera</i>	Hairy Rattlesnakeroot	Asteraceae	UP
<i>Pseudognaphalium macounii</i>	Clammy Cudweed	Asteraceae	UP
<i>Psoraleidum tenuiflorum</i>	Slender-leaved Scurfpea	Fabaceae	UP
<i>Puccinellia nuttalliana</i>	Alkali Grass	Poaceae	WP
<i>Rudbeckia subtomentosa</i>	Sweet Coneflower	Asteraceae	UP
<i>Ruellia humilis</i>	Wild Petunia	Acanthaceae	UP
<i>Salix pseudomonticola</i>	False Mountain Willow	Salicaceae	WP
<i>Scleria triglomerata</i>	Tall Nutrush	Cyperaceae	WP
<i>Scutellaria parvula</i>		Lamiaceae	UP
<i>Senecio integerrimus</i>		Asteraceae	UP
<i>Shepherdia argentea</i>		Elaeagnaceae	UP
<i>Shinnersoseris rostrata</i>	Annual Skeletonweed	Asteraceae	UP
<i>Silene drummondii</i>	Drummond's Campion	Caryophyllaceae	UP
<i>Silphium laciniatum</i>	Compass-plant	Asteraceae	UP
<i>Solidago mollis</i>	Soft Goldenrod	Asteraceae	UP
<i>Spiranthes magnicamporum</i>		Orchidaceae	UP
<i>Sporobolus vaginiflorus</i>	Poverty Dropseed	Poaceae	UP
<i>Stellaria longipes</i>	Long-stalked Chickweed	Caryophyllaceae	UP
<i>Strophostyles helvola</i>		Fabaceae	UP
<i>Strophostyles leiosperma</i>		Fabaceae	UP
<i>Symphyotrichum praealtum</i>		Asteraceae	UP

Scientific Name	Common Name	Family	Primary Ecological System
<i>Synthyris bullii</i>	Kitten-tails	Plantaginaceae	UP
<i>Taenidia integerrima</i>	Yellow Pimpernel	Apiaceae	UP
<i>Tephrosia virginiana</i>	Goat's Rue	Fabaceae	UP
<i>Thaspium barbinode</i>	Hairy-jointed Meadow-parsnip	Apiaceae	UP
<i>Trichophorum clintonii</i>	Clinton's Bulrush	Cyperaceae	UP
<i>Valeriana edulis</i>		Valerianaceae	UP
<i>Viola nuttallii</i>	Yellow Prairie Violet	Violaceae	UP
<i>Xanthisma spinulosum</i>	Cutleaf Ironplant	Asteraceae	UP

Savanna

Scientific Name	Common Name	Family	Primary Ecological System
<i>Androsace septentrionalis</i>	Northern Androsace	Primulaceae	UP
<i>Aureolaria pedicularia</i>	Fernleaf False Foxglove	Chenopodiaceae	UP
<i>Crocanthemum canadense</i>	Canada Frostweed	Cistaceae	UP
<i>Eleocharis wolfii</i>	Wolf's Spikerush	Cyperaceae	UP
<i>Nuttallanthus canadensis</i>	Old Field Toadflax	Plantaginaceae	UP
<i>Orthocarpus luteus</i>	Owl Clover	Orobanchaceae	UP
<i>Rubus multiflor</i>	Kinnickinnick Dewberry	Rosaceae	UP
<i>Rubus vermontanus</i>	Vermont Bristle-berry	Rosaceae	UP

Upland Conifer Forest and Woodlands

Scientific Name	Common Name	Family	Primary Ecological System
<i>Achillea alpina</i>	Siberian Yarrow	Asteraceae	FD
<i>Adlumia fungosa</i>	Allegheny Vine	Fumariaceae	FD
<i>Botrychium ascendens</i>	Upswept Moonwort	Ophioglossaceae	FD
<i>Botrychium campestre</i> var. <i>lineare</i>	Slender Moonwort	Ophioglossaceae	FD
<i>Botrychium michiganense</i>	Michigan Moonwort	Ophioglossaceae	FD
<i>Botrychium minganense</i>	Mingan Moonwort	Ophioglossaceae	FD
<i>Botrychium neolunaria</i>	New World Moonwort	Ophioglossaceae	FD
<i>Botrychium pallidum</i>	Pale Moonwort	Ophioglossaceae	FD
<i>Botrychium simplex</i>	Least Moonwort	Ophioglossaceae	FD
<i>Botrychium spathulatum</i>	Spatulate Moonwort	Ophioglossaceae	FD
<i>Canadanthus modestus</i>	Modest Aster	Asteraceae	FD
<i>Carex pallescens</i>	Pale Sedge	Cyperaceae	FD

Scientific Name	Common Name	Family	Primary Ecological System
<i>Crataegus douglasii</i>	Black Hawthorn	Rosaceae	FD
<i>Crataegus laurentiana</i>	Laurentian Hawthorn	Rosaceae	FD
<i>Cyperus houghtonii</i>	Houghton's Cyperus	Cyperaceae	FD
<i>Dendrolycopodium obscurum</i>	Flat-branched Tree Club-moss	Lycopodiaceae	FD
<i>Deschampsia flexuosa</i>	Slender Hair Grass	Poaceae	FD
<i>Gaylussacia baccata</i>	Black Huckleberry	Ericaceae	FD
<i>Leucophysalis grandiflora</i>	Dwarf Ground Cherry	Solanaceae	FD
<i>Listera convallarioides</i>	Broad-leaved Twayblade	Orchidaceae	FD
<i>Moehringia macrophylla</i>	Large-leaved Sandwort	Caryophyllaceae	FD
<i>Persicaria careyi</i>	Carey's Smartweed	Polygonaceae	FD
<i>Piptatherum canadense</i>	Canadian Ricegrass	Poaceae	FD
<i>Platanthera hookeri</i>	Hooker's Orchid	Orchidaceae	FD
<i>Poa wolfii</i>	Wolf's Bluegrass	Poaceae	FD
<i>Prosartes trachycarpa</i>	Rough-fruited Fairybells	Liliaceae	FD
<i>Pyrola minor</i>	Small Shinleaf	Ericaceae	FD
<i>Rubus groutianus</i>		Rosaceae	FD
<i>Rubus quaesitus</i>	Prince Edward Island Blackberry	Rosaceae	FD
<i>Rubus regionalis</i>		Rosaceae	FD
<i>Sceptridium rugulosum</i>	St. Lawrence Grapefern	Ophioglossaceae	FD
<i>Shepherdia canadensis</i>	Soapberry	Elaeagnaceae	FD
<i>Viburnum edule</i>		Caprifoliaceae	FD
<i>Viola epipsila</i>	Dwarf Marsh Violet	Violaceae	FD
<i>Waldsteinia fragarioides</i>	Barren Strawberry	Rosaceae	FD

Upland Deciduous Forest and Woodlands

Scientific Name	Common Name	Family	Primary Ecological System
<i>Achillea alpina</i>	Siberian Yarrow	Asteraceae	FD
<i>Adlumia fungosa</i>	Allegheny Vine	Fumariaceae	FD
<i>Botrychium ascendens</i>	Upswept Moonwort	Ophioglossaceae	FD
<i>Botrychium campestre</i> var. <i>lineare</i>	Slender Moonwort	Ophioglossaceae	FD
<i>Botrychium neolunaria</i>	New World Moonwort	Ophioglossaceae	FD
<i>Botrychium pallidum</i>	Pale Moonwort	Ophioglossaceae	FD
<i>Botrychium simplex</i>	Least Moonwort	Ophioglossaceae	FD
<i>Botrychium spathulatum</i>	Spatulate Moonwort	Ophioglossaceae	FD

Scientific Name	Common Name	Family	Primary Ecological System
<i>Carex pallescens</i>	Pale Sedge	Cyperaceae	FD
<i>Crataegus douglasii</i>	Black Hawthorn	Rosaceae	FD
<i>Crataegus succulenta</i>		Rosaceae	FD
<i>Dendrolycopodium obscurum</i>	Flat-branched Tree Club-moss	Lycopodiaceae	FD
<i>Desmodium nudiflorum</i>	Stemless Tick Trefoil	Fabaceae	FD
<i>Dryopteris filix-mas</i>	Male Fern	Dryopteridaceae	FD
<i>Elymus diversiglumis</i>		Poaceae	FD
<i>Gaylussacia baccata</i>	Black Huckleberry	Ericaceae	FD
<i>Geum laciniatum</i>	Rough avens	Rosaceae	FD
<i>Hamamelis virginiana</i>	Witch-hazel	Hamamelidaceae	FD
<i>Hypericum punctatum</i>		Hypericaceae	FD
<i>Juncus antheratus</i>	Greater Poverty Rush	Juncaceae	FD
<i>Lactuca floridana</i>	Florida Lettuce	Asteraceae	FD
<i>Lepidium virginicum</i>		Brassicaceae	FD
<i>Leucophysalis grandiflora</i>	Dwarf Ground Cherry	Solanaceae	FD
<i>Listera convallarioides</i>	Broad-leaved Twayblade	Orchidaceae	FD
<i>Lysimachia quadrifolia</i>	Whorled Loosestrife	Myrsinaceae	FD
<i>Melica nitens</i>	Three-flowered Melic	Poaceae	FD
<i>Moehringia macrophylla</i>	Large-leaved Sandwort	Caryophyllaceae	FD
<i>Muhlenbergia sylvatica</i>		Poaceae	FD
<i>Paronychia fastigiata</i>	Forked Chickweed	Caryophyllaceae	FD
<i>Rubus acridens</i>		Rosaceae	FD
<i>Rubus fulleri</i>	a bristle-berry	Rosaceae	FD
<i>Rubus quaesitus</i>	Prince Edward Island Blackberry	Rosaceae	FD
<i>Satureja vulgaris</i>	Wild Basil	Lamiaceae	FD

Mesic Hardwood Forest

Scientific Name	Common Name	Family	Primary Ecological System
<i>Allium cernuum</i>	Nodding Wild Onion	Liliaceae	MH
<i>Arnoglossum reniforme</i>	Great Indian Plantain	Asteraceae	MH
<i>Asplenium platyneuron</i>	Ebony Spleenwort	Aspleniaceae	MH
<i>Borodinia laevigata</i>	Smooth Rock Cress	Brassicaceae	MH
<i>Botrychium angustisegmentum</i>	Narrow Triangle Moonwort	Ophioglossaceae	MH
<i>Botrychium crenulatum</i>	Dainty Moonwort	Ophioglossaceae	MH

Scientific Name	Common Name	Family	Primary Ecological System
<i>Botrychium mormo</i>	Goblin Fern	Ophioglossaceae	MH
<i>Cardamine douglassii</i>	Purple Cress	Brassicaceae	MH
<i>Carex careyana</i>	Carey's Sedge	Cyperaceae	MH
<i>Carex debilis</i>		Cyperaceae	MH
<i>Carex formosa</i>	Handsome Sedge	Cyperaceae	MH
<i>Carex jamesii</i>	James' Sedge	Cyperaceae	MH
<i>Carex laxiculmis</i>	Spreading Sedge	Cyperaceae	MH
<i>Carex novae-angliae</i>	New England Sedge	Cyperaceae	MH
<i>Carex ormostachya</i>	Necklace Sedge	Cyperaceae	MH
<i>Carex plantaginea</i>	Plantain-leaved Sedge	Cyperaceae	MH
<i>Carex torreyi</i>	Torrey's Sedge	Cyperaceae	MH
<i>Chenopodium standleyanum</i>		Chenopodiaceae	MH
<i>Cornus drummondii</i>	Rough-leaved Dogwood	Cornaceae	MH
<i>Crataegus calpodendron</i>	Late Hawthorn	Rosaceae	MH
<i>Deparia acrostichoides</i>	Silvery Spleenwort	Athyriaceae	MH
<i>Desmodium cuspidatum</i>		Fabaceae	MH
<i>Diarrhena obovata</i>	Obovate Beakgrain	Poaceae	MH
<i>Dicentra canadensis</i>	Squirrel Corn	Fumariaceae	MH
<i>Dodecatheon amethystinum</i>	Jeweled Shooting Star	Primulaceae	MH
<i>Dryopteris goldieana</i>	Goldie's Fern	Dryopteridaceae	MH
<i>Dryopteris marginalis</i>	Marginal Shield Fern	Dryopteridaceae	MH
<i>Galium circaeans</i>		Rubiaceae	MH
<i>Homalosorus pycnocarpus</i>	Narrow-leaved Spleenwort	Diplaziopsidaceae	MH
<i>Hybanthus concolor</i>	Eastern Green-violet	Violaceae	MH
<i>Hydrastis canadensis</i>	Goldenseal	Ranunculaceae	MH
<i>Jeffersonia diphylla</i>	Twinleaf	Berberidaceae	MH
<i>Juglans cinerea</i>	Butternut	Juglandaceae	MH
<i>Listera auriculata</i>	Auricled Twayblade	Orchidaceae	MH
<i>Lysimachia lanceolata</i>	Lance-leaved Loosestrife	Myrsinaceae	MH
<i>Morus rubra</i>	Red Mulberry	Moraceae	MH
<i>Osmorhiza berteroi</i>	Chilean Sweet Cicely	Apiaceae	MH
<i>Osmorhiza depauperata</i>	Blunt-fruited Sweet Cicely	Apiaceae	MH
<i>Panax quinquefolius</i>	American Ginseng	Araliaceae	MH
<i>Panax trifolius</i>		Araliaceae	MH
<i>Phegopteris hexagonoptera</i>	Broad Beech Fern	Thelypteridaceae	MH
<i>Poa sylvestris</i>	Woodland Bluegrass	Poaceae	MH
<i>Polystichum acrostichoides</i>	Christmas Fern	Dryopteridaceae	MH
<i>Polystichum braunii</i>	Braun's Holly Fern	Dryopteridaceae	MH

Scientific Name	Common Name	Family	Primary Ecological System
<i>Sanicula trifoliata</i>	Beaked Snakeroot	Apiaceae	MH
<i>Sceptridium oneidense</i>	Blunt-lobed Grapefern	Ophioglossaceae	MH
<i>Symphyotrichum shortii</i>	Short's Aster	Asteraceae	MH
<i>Trillium nivale</i>	Snow Trillium	Liliaceae	MH
<i>Tsuga canadensis</i>	Eastern Hemlock	Pinaceae	MH
<i>Verbena simplex</i>	Narrow-leaved Vervain	Verbenaceae	MH
<i>Viola pubescens</i> var. <i>pubescens</i>		Violaceae	MH
<i>Vitis aestivalis</i>	Silverleaf Grape	Vitaceae	MH

Deciduous Wet Forest

Scientific Name	Common Name	Family	Primary Ecological System
<i>Bartonia virginica</i>	Yellow Bartonia	Gentianaceae	WF
<i>Carex laevivaginata</i>	Smooth-sheathed Sedge	Cyperaceae	WF
<i>Floerkea proserpinacoides</i>	False Mermaid	Limnanthaceae	WF
<i>Hydrocotyle americana</i>	American Water-pennywort	Araliaceae	WF
<i>Poa paludigena</i>	Bog Bluegrass	Poaceae	WF
<i>Ranunculus macounii</i>		Ranunculaceae	WF
<i>Rumex verticillatus</i>		Polygonaceae	WF

Coniferous Forest Wetland

Scientific Name	Common Name	Family	Primary Ecological System
<i>Amerorchis rotundifolia</i>	Small Round-leaved Orchis	Orchidaceae	FP
<i>Astragalus alpinus</i>	Alpine Milk-vetch	Fabaceae	FP
<i>Botrychium tenebrosum</i>	Swamp Moonwort	Ophioglossaceae	WF
<i>Calypso bulbosa</i>	Fairy Slipper	Orchidaceae	FP
<i>Cardamine pratensis</i>	Cuckoo Flower	Brassicaceae	FP
<i>Carex aquatilis</i> var. <i>aquatilis</i>		Cyperaceae	FP
<i>Carex capillaris</i>	Hair-like Sedge	Cyperaceae	FP
<i>Cypripedium arietinum</i>	Ram's Head Orchid	Orchidaceae	FP
<i>Gymnocarpium robertianum</i>	Northern Oak Fern	Cystopteridaceae	FP
<i>Luzula parviflora</i>	Small-flowered Woodrush	Juncaceae	FP

Scientific Name	Common Name	Family	Primary Ecological System
<i>Malaxis paludosa</i>	Bog Adder's Mouth	Orchidaceae	FP
<i>Platanthera clavellata</i>	Small Green Wood Orchid	Orchidaceae	APn8
<i>Platanthera dilatata</i>	Tall White Bog Orchid	Orchidaceae	FP
<i>Polemonium lacustre</i>	Great Lakes Jacob's-ladder	Polemoniaceae	FP
<i>Ranunculus lapponicus</i>	Lapland Buttercup	Ranunculaceae	FP
<i>Ranunculus macounii</i>		Ranunculaceae	WF
<i>Rubus chamaemorus</i>	Cloudberry	Rosaceae	APn8
<i>Spiranthes casei</i>	Case's Ladies' Tresses	Orchidaceae	FP
<i>Stellaria crassifolia</i>		Caryophyllaceae	FP
<i>Thalictrum revolutum</i>	Waxleaf Meadowrue	Ranunculaceae	FP
<i>Vaccinium uliginosum</i>	Alpine Bilberry	Ericaceae	FP

Riparian and Floodplain Forest

Scientific Name	Common Name	Family	Primary Ecological System
<i>Agastache nepetoides</i>	Yellow Giant Hyssop	Lamiaceae	FF
<i>Arisaema dracontium</i>	Green Dragon	Araceae	FF
<i>Betula nigra</i>		Betulaceae	FF
<i>Carex conjuncta</i>	Jointed Sedge	Cyperaceae	FF
<i>Carex davisii</i>	Davis' Sedge	Cyperaceae	FF
<i>Carex grayi</i>	Gray's Sedge	Cyperaceae	FF
<i>Carex muskingumensis</i>	Muskingum Sedge	Cyperaceae	FF
<i>Carex typhina</i>	Cattail Sedge	Cyperaceae	FF
<i>Cuscuta cephalanthi</i>		Convolvulaceae	FF
<i>Elymus curvatus</i>		Poaceae	FF
<i>Elymus riparius</i>	Riverbank Wildrye	Poaceae	FF
<i>Erythronium propullans</i>	Dwarf Trout Lily	Liliaceae	FF
<i>Gleditsia triacanthos</i>	Honeylocust	Fabaceae	FF
<i>Gymnocladus dioica</i>	Kentucky Coffee Tree	Caesalpiniaceae	FF
<i>Iodanthus pinnatifidus</i>	Purple Rocket	Brassicaceae	FF
<i>Leersia lenticularis</i>	Catchfly Grass	Poaceae	FF
<i>Lycopus virginicus</i>	Virginia Water Horehound	Lamiaceae	FF

Scientific Name	Common Name	Family	Primary Ecological System
<i>Napaea dioica</i>	Glade Mallow	Malvaceae	FF
<i>Polygonum erectum</i>		Polygonaceae	FF
<i>Quercus bicolor</i>	Swamp White Oak	Fagaceae	FF
<i>Rudbeckia triloba</i>	Three-leaved Coneflower	Asteraceae	FF
<i>Scutellaria ovata</i>	Ovate-leaved Skullcap	Lamiaceae	FF

Non-forested Wetlands

Scientific Name	Common Name	Family	Primary Ecological System
<i>Alisma gramineum</i>	Narrow-leaved Water Plantain	Alismataceae	MR
<i>Berula erecta</i>	Stream Parsnip	Apiaceae	OP
<i>Carex exilis</i>	Coastal Sedge	Cyperaceae	OP
<i>Carex flava</i>	Yellow Sedge	Cyperaceae	OP
<i>Carex garberi</i>	Garber's Sedge	Cyperaceae	WM
<i>Carex michauxiana</i>	Michaux's Sedge	Cyperaceae	OP
<i>Carex sterilis</i>	Sterile Sedge	Cyperaceae	OP
<i>Carex trichocarpa</i>	Hairy-fruited Sedge	Cyperaceae	WM
<i>Cladium mariscoides</i>	Twig Rush	Cyperaceae	OP
<i>Cuscuta coryli</i>		Convolvulaceae	WM
<i>Decodon verticillatus</i>	Water-willow	Lythraceae	MR
<i>Drosera anglica</i>	English Sundew	Droseraceae	OP
<i>Drosera linearis</i>	Linear-leaved Sundew	Droseraceae	OP
<i>Eleocharis coloradoensis</i>	Dwarf Spikerush	Cyperaceae	WM
<i>Eleocharis flavescens</i> var. <i>olivacea</i>		Cyperaceae	WM
<i>Eleocharis mamillata</i>	Mamillate Spike-rush	Cyperaceae	OP
<i>Eleocharis nitida</i>	Neat Spikerush	Cyperaceae	WM
<i>Eleocharis quinqueflora</i>	Few-flowered Spikerush	Cyperaceae	OP
<i>Eleocharis rostellata</i>	Beaked Spikerush	Cyperaceae	OP
<i>Fimbristylis autumnalis</i>	Autumn Fimbry	Cyperaceae	OP
<i>Fimbristylis puberula</i>	Hairy Fimbry	Cyperaceae	OP
<i>Galium palustre</i>	Marsh Bedstraw	Rubiaceae	OP
<i>Hasteola suaveolens</i>	Sweet-smelling Indian Plantain	Asteraceae	MR
<i>Hypericum ellipticum</i>		Hypericaceae	WM

Scientific Name	Common Name	Family	Primary Ecological System
<i>Juncus marginatus</i>	Marginated Rush	Juncaceae	WM
<i>Juncus stygius</i>	Moor Rush	Juncaceae	OP
<i>Lysimachia maritima</i>	Sea Milkwort	Myrsinaceae	WM
<i>Muhlenbergia uniflora</i>	One-flowered Muhly	Poaceae	OP
<i>Oxypolis rigidior</i>	Cowbane	Apiaceae	OP
<i>Polygala cruciata</i>	Cross-leaved Milkwort	Polygalaceae	OP
<i>Potentilla rivalis</i>	Brook Cinquefoil	Rosaceae	WM
<i>Potentilla supina</i>	Bushy Cinquefoil	Rosaceae	OP
<i>Rhynchospora capillacea</i>	Hair-like Beak Rush	Cyperaceae	OP
<i>Rotala ramosior</i>	Toothcup	Lythraceae	OP
<i>Rubus dissensus</i>		Rosaceae	OP
<i>Rubus missouricus</i>	Missouri Bristle-berry	Rosaceae	WM
<i>Rubus semisetosus</i>	Swamp Blackberry	Rosaceae	WM
<i>Rubus stipulatus</i>	A Bristle-berry	Rosaceae	WM
<i>Rubus wheeleri</i>	Wheeler's Blackberry	Rosaceae	WM
<i>Rumex altissimus</i>		Polygonaceae	WM
<i>Sagittaria brevirostra</i>	Short-beaked Arrowhead	Alismataceae	OP
<i>Salicornia rubra</i>	Red Saltwort	Chenopodiaceae	OP
<i>Salix maccalliana</i>	McCalla's Willow	Salicaceae	OP
<i>Scleria verticillata</i>	Whorled Nutrush	Cyperaceae	OP
<i>Sparganium glomeratum</i>	Clustered Bur-reed	Sparganiaceae	OP
<i>Stellaria alsine</i>		Caryophyllaceae	WM
<i>Torreyochloa pallida</i> var. <i>pallida</i>		Poaceae	WM
<i>Utricularia geminiscapa</i>	Hidden-fruit Bladderwort	Lentibulariaceae	WM
<i>Viola lanceolata</i>	Lance-leaf Violet	Violaceae	OP
<i>Xyris montana</i>	Montane Yellow-eyed Grass	Xyridaceae	APn9
<i>Xyris torta</i>	Twisted Yellow-eyed Grass	Xyridaceae	OP

Rivers and Streams

Scientific Name	Common Name	Family	Primary Ecological System
<i>Bidens discoidea</i>	Discoïd Beggarticks	Asteraceae	RV
<i>Eclipta prostrata</i>	Yerba de Tajo	Asteraceae	RV
<i>Eragrostis frankii</i>		Poaceae	RV
<i>Nymphaea leibergii</i>	Small White Waterlily	Nymphaeaceae	RV

<i>Rorippa curvipes</i>	Bluntleaf Yellowcress	Brassicaceae	RV
<i>Rorippa sessiliflora</i>	Sessile-flowered Yellow Cress	Brassicaceae	RV
<i>Sagittaria montevidensis</i>	Long-lobed Arrowhead	Alismataceae	RV
<i>Salix pellita</i>	Satiny Willow	Salicaceae	RV
<i>Silene nivea</i>	Snowy Champion	Caryophyllaceae	RV
<i>Sparganium androcladum</i>		Sparganiaceae	RV

Lakes

Scientific Name	Common Name	Family	Primary Ecological System
<i>Callitriche hermaphrodita</i>		Plantaginaceae	LK
<i>Caltha natans</i>	Floating Marsh Marigold	Ranunculaceae	LK
<i>Carex katahdinensis</i>	Katahdin Sedge	Cyperaceae	LK
<i>Castilleja septentrionalis</i>	Northern Paintbrush	Orobanchaceae	LK
<i>Ceratophyllum echinatum</i>	Spiny Hornwort	Ceratophyllaceae	LK
<i>Draba norvegica</i>	Norwegian Whitlow Grass	Brassicaceae	LK
<i>Eleocharis robbinsii</i>	Robbins' Spikerush	Cyperaceae	LK
<i>Elodea bifoliata</i>	Two Leaf Waterweed	Hydrocharitaceae	LK
<i>Eupatorium sessilifolium</i>	Upland Boneset	Asteraceae	LK
<i>Juncus articulatus</i>	Jointed Rush	Juncaceae	LK
<i>Juncus subtilis</i>	Slender Rush	Juncaceae	LK
<i>Littorella americana</i>	American Shore Plantain	Plantaginaceae	LK
<i>Myriophyllum heterophyllum</i>	Broadleaf Water Milfoil	Haloragaceae	LK
<i>Najas gracillima</i>	Slender Naiad	Najadaceae	LK
<i>Najas guadalupensis ssp. olivacea</i>	Olive-colored Southern Naiad	Najadaceae	LK
<i>Najas marina</i>	Sea Naiad	Najadaceae	LK
<i>Potamogeton bicupulatus</i>	Snailseed Pondweed	Potamogetonaceae	LK
<i>Potamogeton confervoides</i>	Algae-like Pondweed	Potamogetonaceae	LK
<i>Potamogeton diversifolius</i>	Diverse-leaved Pondweed	Potamogetonaceae	LK
<i>Potamogeton oakesianus</i>	Oakes' Pondweed	Potamogetonaceae	LK
<i>Potamogeton pulcher</i>	Spotted Pondweed	Potamogetonaceae	LK
<i>Potamogeton pusillus ssp. pusillus</i>		Potamogetonaceae	LK
<i>Rhynchospora capitellata</i>	Brownish Beak Sedge	Cyperaceae	LK
<i>Ruppia cirrhosa</i>	Spiral Ditchgrass	Ruppiaceae	LK
<i>Schoenoplectiella purshiana</i> var. <i>purshiana</i>	Weak-stalk Bulrush	Cyperaceae	LK
<i>Schoenoplectiella purshiana</i> var. <i>williamsii</i>		Cyperaceae	LK

<i>Schoenoplectiella smithii</i> var. <i>smithii</i>	Blunt-scale Bulrush	Cyperaceae	LK
<i>Selaginella selaginoides</i>	Northern Spikemoss	Selaginellaceae	LK
<i>Stuckenia vaginata</i>	Sheathed Pondweed	Potamogetonaceae	LK
<i>Subularia aquatica</i>	Awlwort	Brassicaceae	LK
<i>Tofieldia pusilla</i>	Small False Asphodel	Liliaceae	LK
<i>Trisetum spicatum</i>	Spike Trisetum	Poaceae	LK
<i>Utricularia purpurea</i>	Purple-flowered Bladderwort	Lentibulariaceae	LK
<i>Utricularia resupinata</i>	Lavender Bladderwort	Lentibulariaceae	LK
<i>Wolffia brasiliensis</i>	Pointed Watermeal	Lemnaceae	LK
<i>Zizania palustris</i>	Northern wild rice	Poaceae	LK

Cliff, Talus, and Rock Outcrop

Scientific Name	Common Name	Family	Primary Ecological System
<i>Allium schoenoprasum</i>	Wild Chives	Liliaceae	RO
<i>Ammophila breviligulata</i>	Beachgrass	Poaceae	CT
<i>Anagallis minima</i>	Chaffweed	Myrsinaceae	CT
<i>Aristida dichotoma</i>	Churchmouse Three-awn	Poaceae	CT
<i>Arnica lonchophylla</i>	Long-leaved Arnica	Asteraceae	CT
<i>Asplenium trichomanes</i>	Maidenhair Spleenwort	Aspleniaceae	CT
<i>Bacopa rotundifolia</i>	Waterhyssop	Plantaginaceae	RO
<i>Bistorta vivipara</i>	Alpine Bistort	Polygonaceae	RO
<i>Boechera collinsii</i>	Collins' Rock Cress	Brassicaceae	CT
<i>Boechera stricta</i>	Drummond's Rock Cress	Brassicaceae	RO
<i>Buchloe dactyloides</i>	Buffalo Grass	Poaceae	RO
<i>Bulbostylis capillaris</i>	Densetuft Hairsedge	Cyperaceae	RO
<i>Calamagrostis purpurascens</i>	Purple Reedgrass	Poaceae	RO
<i>Callitriche heterophylla</i>	Larger Water Starwort	Plantaginaceae	RO
<i>Carex media</i>	Intermediate Sedge	Cyperaceae	CT
<i>Carex praticola</i>	Prairie-dweller Sedge	Cyperaceae	CT
<i>Carex rossii</i>	Ross' Sedge	Cyperaceae	CT
<i>Carex supina</i>		Cyperaceae	RO
<i>Cerastium brachypodum</i>	Mouse-ear Chickweed	Caryophyllaceae	CT
<i>Chrysosplenium iowense</i>	Iowa Golden Saxifrage	Saxifragaceae	CT
<i>Collomia linearis</i>		Polemoniaceae	RO
<i>Corispermum americanum</i>		Chenopodiaceae	RO
<i>Corydalis micrantha</i>		Fumariaceae	RO
<i>Coryphantha vivipara</i>	Ball Cactus	Cactaceae	RO

Scientific Name	Common Name	Family	Primary Ecological System
<i>Crassula aquatica</i>	Water Pygmyweed	Crassulaceae	RO
<i>Cyperus acuminatus</i>	Short-pointed Umbrella Sedge	Cyperaceae	WM
<i>Cystopteris laurentiana</i>	Laurentian Bladder Fern	Cystopteridaceae	CT
<i>Cystopteris tennesseensis</i>	Tennessee Bladder Fern	Cystopteridaceae	RO
<i>Draba arabisans</i>	Arabian Whitlow Grass	Brassicaceae	CT
<i>Draba cana</i>	Hoary Whitlow Grass	Brassicaceae	CT
<i>Elatine triandra</i>	Three-stamened Waterwort	Elatinaceae	RO
<i>Eleocharis engelmannii</i>	Engelmann's Spikerush	Cyperaceae	CT
<i>Empetrum atropurpureum</i>	Purple Crowberry	Ericaceae	CT
<i>Empetrum nigrum</i>	Black Crowberry	Ericaceae	CT
<i>Erigeron acris</i>		Asteraceae	CT
<i>Euphrasia hudsoniana</i>	Hudson Bay Eyebright	Orobanchaceae	CT
<i>Heteranthera limosa</i>	Mud Plantain	Pontederiaceae	RO
<i>Hordeum pusillum</i>	Little Barley	Asteraceae	RO
<i>Huperzia appressa</i>	Mountain Fir Moss	Lycopodiaceae	CT
<i>Huperzia porophila</i>	Rock Firmoss	Lycopodiaceae	CT
<i>Isoetes melanopoda</i>	Prairie Quillwort	Isoetaceae	RO
<i>Limosella aquatica</i>	Mudwort	Scrophulariaceae	RO
<i>Marsilea vestita</i>	Hairy Waterclover	Marsileaceae	RO
<i>Montia chamissoi</i>	Montia	Portulacaceae	LK
<i>Myosotis verna</i>	Forget-me-not	Boraginaceae	RO
<i>Opuntia macrorhiza</i>	Devil's Tongue	Cactaceae	RO
<i>Orobanche ludoviciana</i>	Louisiana Broomrape	Orobanchaceae	RO
<i>Oxytropis viscida</i>	Sticky Locoweed	Fabaceae	RO
<i>Packera indecora</i>	Elegant Groundsel	Asteraceae	RO
<i>Pellaea atropurpurea</i>	Purple Cliff Brake	Pteridaceae	RO
<i>Phacelia franklinii</i>	Franklin's Phacelia	Hydrophyllaceae	RO
<i>Physaria ludoviciana</i>	Bladderpod	Brassicaceae	RO
<i>Pinguicula vulgaris</i>	Butterwort	Lentibulariaceae	RO
<i>Plagiobothrys scouleri</i>	Scouler's Popcornflower	Boraginaceae	RO
<i>Plantago elongata</i>	Slender Plantain	Plantaginaceae	RO
<i>Rhodiola integrifolia</i>	Leedy's roseroot	Crassulaceae	RO
<i>Rorippa sinuata</i>	Spreading Yellow Cress	Brassicaceae	RO
<i>Rubus satis</i>		Rosaceae	RO
<i>Sagina nodosa</i>		Caryophyllaceae	RO
<i>Saxifraga cernua</i>	Nodding Saxifrage	Saxifragaceae	RO
<i>Saxifraga paniculata</i>	Encrusted Saxifrage	Saxifragaceae	RO
<i>Schedonnardus paniculatus</i>	Tumble Grass	Poaceae	RO

Scientific Name	Common Name	Family	Primary Ecological System
<i>Silene hitchguirei</i>	Mountain Campion	Caryophyllaceae	RO
<i>Solidago sciaphila</i>	Cliff Goldenrod	Asteraceae	RO
<i>Sullivantia sullivantii</i>	Reniform Sullivantia	Saxifragaceae	RO
<i>Triplasis purpurea</i>	Purple Sandgrass	Poaceae	RO
<i>Woodsia alpina</i>	Alpine Woodsia	Woodsiaceae	RO
<i>Woodsia glabella</i>	Smooth Woodsia	Woodsiaceae	RO
<i>Woodsia obtusa</i>	Blunt-lobed Woodsia	Woodsiaceae	RO
<i>Woodsia oregana</i>		Woodsiaceae	RO
<i>Woodsia scopulina</i>	Rocky Mountain Woodsia	Woodsiaceae	RO