

Summary of the Superior National Forest's Canada lynx (*Lynx canadensis*) DNA database and population monitoring 2024

December 5, 2024

DAN RYAN – USDA Forest Service, Superior National Forest, Laurentian Ranger District, 318 Forestry Rd., Aurora, MN 55705

DAVE GROSSHUESCH – USDA Forest Service, Superior National Forest, Tofte Ranger District, 7355 W. Hwy 61, P.O. Box 2159 Tofte, MN 55615

CHERON FERLAND - USDA Forest Service, Superior National Forest, Supervisors Office, 8901 Grand Avenue Place, Duluth, MN 55808

KRISTY PILGRIM – USDA Forest Service, National Genomics Center for Wildlife and Fish Conservation, 800 E. Beckwith, Missoula, MT 59801

SHANNON BARBER-MEYER – Pacific Whale Foundation
300 Ma'alaea Road Ste 211, Wailuku, HI 96793

Introduction

Snow tracking and other methods used to obtain genetic samples have confirmed presence of Canada lynx (*Lynx canadensis*) across northeastern Minnesota since December 2000. In 2008, the Superior National Forest (Superior NF) created, and continues to maintain, a database of genetically confirmed Canada lynx (hereafter lynx) to document their occurrence, persistence and reproduction in Minnesota. Genetic samples (typically scat but also hair and tissue) have been collected primarily as part of the Superior NF's survey and monitoring program. Also included in this database are samples collected during an independent genetic research project, a radio telemetry project, mining project surveys, and from specimens that were surrendered to resource agencies (e.g., from animals that had been trapped, shot or killed in vehicle collisions). These samples were submitted to the USDA Forest Service Rocky Mountain Research Station's National Genomics Center for Wildlife and Fish Conservation for testing. Samples that were identified as lynx using mitochondrial DNA analysis were further evaluated using nuclear DNA analysis methods to determine sex (Pilgrim et al. 2005) and individual identification. Further testing was used to determine Canada lynx-bobcat (*Lynx rufus*) hybridization (Schwartz et al. 2004). Field observations combined with DNA analysis have been used to document lynx reproduction.

Summary

The current database contains 3,267 samples that have been submitted for DNA testing. Mitochondrial DNA analysis has identified 3,095 (94.7%) of them to species, 2,785 (90.0%) of those as lynx. Nuclear DNA analysis has determined 659 unique lynx genotypes, 313 female (47.5%), 344 male (52.2%) and 2 of indeterminable sex (0.3%).

Reproduction has been documented on the Superior NF every year since 2001. Since 2010 we have identified a minimum of 95 family groups producing a total of 195 presumed kittens, 104 female (53.3%) and 91 male (46.7%) (Figure 3). Of the 584 individuals that were identified prior to this survey season and were not originally detected as a result of a mortality, 177 (30.3%) are known to have persisted into a second year. Sixteen individuals have persisted for over 6 years, the longest, a female, over 10 years.

During the 2023/2024 survey season 324 samples were collected and submitted for testing. Two hundred and fifty-two (77.8%) were identified as lynx, 224 of those (88.9%) were able to be genotyped identifying 98 individuals: 48 female and 50 male. Fifty-four individuals (55.1%: 30 female and 24 male) were previously recorded in this database (recaptures), and 44 individuals (44.9%: 18 female and 26 male) were new to the database this year including 25 kittens.

Field observations suggest that there were at least 16 family groups with as many as 35 kittens found in the survey area. DNA analyses confirm 11 family groups with 25 individuals (11 female, 14 male) genetically consistent with being offspring. Of the 35 individuals detected during the 2023/2024 survey season whose age can be estimated (24 individuals were new to the database and not thought to be kittens) and that were not kittens, 27 (77.1%) have persisted for 2 years or more (Figure 8). Twenty-four (68.5%) of the 35 have been detected into their third year or more indicating recruitment into the population. Two females have persisted in the survey area for over 10 years but one of those GLNR-S-571 was found dead in January.

The DNA database also contains 44 samples that have been identified as F1 Canada lynx-bobcat hybrids. There are 13 unique lynx-bobcat hybrid genotypes, 5 (38.5%) female and 8 male (61.5%). One Canada lynx-bobcat hybrid, a male, was first detected in February 2011 and was at least 9 years old when he was trapped in December of 2019. There were no Canada lynx-bobcat hybrids detected during the 2023/2024 survey season and the last hybrid detection was the male trapped in 2019.

The USFWS maintains a “take” database of all reported incidents of take in Minnesota (see Take section below for definition). Between 2001 and 2024 there have been 86 reported incidents of lynx take within the state. Sixty-three of those incidents have resulted in mortalities to the animal. There have been 8 mortalities by shooting, 19 incidental trappings where there was lynx mortality, and 22 deaths due to car or truck, snowmobile, or a train collision. There were 17 incidents of mortality where the cause was unknown. Since the February 2023 report, there have been seven incidents of take statewide, six of which were within the Superior National Forest. The six incidents within the National Forest included: two vehicle collisions on Forest Service roads, one collision on a State Road bisecting the forest, two incidents related to accidental trapping, and one mortality due to unknown reasons.

We continue to work with researchers at the University of North Carolina to refine the dynamic occupancy model as described in Hostetter and Gardner (2016). Initial results of this work were published by Hostetter et al. (2020). We also continue to build on the data set used to generate core area population estimates described in Barber-Meyer et al. (2018) to produce ongoing annual estimates and assess potential trends over time. We have two survey seasons of data implementing the Spatial Capture Recapture (SCR) survey method in 2019/2020 and 2022/2023. A grant has been secured to analyze this data and it should get started in 2024.

This genetics database contains all known samples submitted by the Superior NF to the Wildlife Genetics Laboratory since the year 2000. Other contributors to this database are Chippewa National Forest, Franconia Minerals Corporation, Leech Lake Band of Ojibwe, 1854 Treaty Authority, Minnesota Department of Natural Resources, Natural Resources Research Institute (NRRI) of the University of Minnesota-Duluth, NewRange Copper Nickel (formerly PolyMet Mining Inc.), Steve Loch, US Fish and Wildlife Service, US Geological Survey, Voyageur’s National Park and Wolf Ridge ELC.

Canada lynx DNA Database 2000-2024

Survey Effort

For the purposes of this report, the primary survey area is generally considered to be the proclamation boundaries of the Superior National Forest (see attached map). Survey techniques over the years have been predominantly on an ad hoc basis. Survey effort has varied dependent upon funding, personnel availability and suitable snow conditions; biologists usually survey areas on their Districts as time and snow conditions allow. Prior to 2014 records and GPS tracks were not stringently maintained, nor was there consistency between surveyors in how those tracks were collected. However, as part of a recent research effort by North Carolina State University (NCSU) to develop an occupancy model (see DNA Database Research below) and continued development of a monitoring model, we summarized our survey effort into miles of occupancy surveys and trailing miles (miles in which surveyors were following lynx tracks) during survey seasons 2014/2015 – 2023/2024. During surveys season 2012/2013 and 2013/2014 only the trailing miles were recorded and not included in the NCSU work but are included in Table 1 below. These 2 parameters give an index of survey effort by Superior NF personnel/volunteers in collecting genetic samples for this database.

Year	Occupancy miles	Trailing miles	Samples collected	Number lynx samples	Individuals identified	Unknown Individuals**
2013	NA	41.6	149	122	35	16 (46%)
2014	NA	45.6	198	162	68	45 (66%)
2015	1,970	43.1	135	114	49	24 (49%)
2016	2,044	52.8	127	113	38	7 (18%)
2017	2,279	70.2	144	130	42	7 (17%)
2018	2,601	76.8	210	187	68	26 (38%)
2019	2,064	95.6	157	153	59	18 (30%)
2020*	5,032	131.9	221	209	82	22 (27%)
2021	3,532	61.3	171	145	66	9 (14%)
2022	3,794	80.1	236	211	79	27 (34%)
2023*	5,384	165.6	317	298	118	56 (47%)
2024	1,986	88.8	324	252	98	19 (19%)

Table 1. Survey effort for monitoring and DNA collection.

*2020 and 2023 were intensive survey years made possible through a funding agreement with Polymet Mining, Inc.

** New individuals that were not known to be kittens. Percent of all individuals identified that were new.

Species Identification

To date there are 3,267 samples contained in the database, of which 3,095 (94.7%) have been identified to species (Figure 1). Of the samples for which species results were obtained, 2,785 (90.0%) were identified as lynx. Two thousand four hundred and eleven of those samples (86.6%) were able to be genotyped (meaning an individual identification was made), 374 (13.4%) could not be identified to individual (identified as lynx only). One-hundred seventy-two of all the samples contained in the database (5.3%) did not contain enough quality DNA to make a species determination.




Figure 1. Species identification results 2000-2024 ($n = 3,267$)

Of the 2,785 lynx samples:

- 659 individual lynx genotypes were identified, of which there are:
 - 313 females (47.5%), 344 males (52.2%) and 2 (0.3%) undetermined sex (Figure 2)




Figure 2. Total Canada lynx individuals detected 2000-2024 ($n = 785$)

Reproduction

Field observations and DNA analysis have been used to document reproduction of lynx in Minnesota every year since 2001. Areas that contain tracks of family groups (adults and kittens (animals presumed to be <1 year old)) are continually monitored during the survey season to attempt to collect DNA from all individuals. However, genetic samples from each member of the family group may not always be obtained, nor is every family group likely detected each year, so numbers presented here under-represent the total numbers of family groups and kittens. These figures represent only those family groups and kittens for which DNA analysis has shown a parent-offspring relationship.

Field observations of family groups combined with DNA analysis since 2010 have identified a minimum of 95 family groups producing a total of 195 presumed kittens, 104 female (53.3%) and 91 male (46.7%) (Figure 3). Overall, 241 kittens have been identified since 2001 from 60 different mothers and 39 different fathers (52 family groups had genetically unknown fathers).




Figure 3. Family groups and known annual reproduction, birth years 2010-2023

Persistence

Snow tracking and other methods used to obtain genetic samples have confirmed persistence of lynx on the Superior NF and elsewhere across northeastern Minnesota since January 2002 (Figure 4). Of the 584 individuals that were identified prior to this survey season and were not originally detected as a result of a mortality, 177 (30.3%) are known to have persisted into a second year. There is 1 individual, a female, which has persisted for over 10 years (born in 2013) who was last detected in March 2024. GLNR-S-571 was also born in 2013 but she was found dead in May 2024. There are 16 individuals that have been detected over 6 years or more since monitoring began.




Figure 4. Persistence of individuals 2000-2024 (n = 525)

Survivorship/Recruitment

Although not an accurate representation of true kitten survivorship, the persistence of kittens in the database into their second year and longer can be an index of survivorship and recruitment into the overall population. Of the 213 individuals that have been identified as kittens (114 female and 99 male) in the database prior to 2023/2024 and were not initially detected as a mortality, 69 of them (32.2%: 38 female and 31 male) have been recaptured into their second year or beyond and are assumed to have been recruited into the northeastern Minnesota (NE MN) sub-population (Figure 5).




Figure 5. Persistence/recruitment of kittens 2001-2024 (n=213)

It is presumed that not all surviving kittens are detected within the survey area, and other surviving kittens emigrate to other sub-populations so this is a minimum estimate of recruitment. It is not uncommon to find the second year “kittens” in their mother’s home range, and even interacting with the new families of their mothers.

Distribution and Dispersal

Lynx detections are distributed over 12 counties in Minnesota and Canada. The majority occur in St. Louis, Lake and Cook counties in northeastern Minnesota where essentially all field data collection efforts have been focused (Table 2). The attached map at the end of the report represents locations of samples genetically confirmed as lynx within the State of Minnesota since they were listed as a threatened species under the Endangered Species Act (March 24, 2000).

County	No. of individual lynx detections by County	% of total individual lynx (n=609)
Cook	126	19.1%
Lake	391	59.3%
St. Louis	122	18.5%
All other	20	3.0%

Table 2. Distribution of individual lynx detections in Minnesota by county

A new male individual was found in Lake of the Woods County and another new male in Itasca county last winter.

Dispersal and movement of individuals both within and out of the core survey and monitoring area has been documented. Maximum movement distance is 196 miles for males and 46 miles for females.

2023-2024 Monitoring Results

Species Identification

Three hundred and twenty-four samples were collected and submitted for analysis during the period of July 2023 through May 2024. Two hundred and fifty-two samples (77.8%) were identified as lynx, and genotypes were obtained from 224 of these identifying 98 unique individuals, 48 female (49.0%) 50 male (51.0%) (Figure 6). Fifty-four individuals (55.1%: 30 female and 24 male) were previously recorded in this database (recaptures), and 44 individuals (44.9%: 18 female and 26 male) were new to the database this year including 25 kittens.




Figure 6. Canada lynx individuals detected 2024 ($n = 98$)

Reproduction

Field observations during the winter 2023-2024 survey season suggest that there were at least 16 family groups with as many as 35 kittens found in the survey area. DNA analysis did confirm 11 family groups in the survey area with a total of 25 kittens genotyped; 11 female (44.0%) and 14 male (56.0%), genetically consistent with being offspring (Figure 7). Family groups are determined by comparing the genotypes of samples collected from animals using the same area for parent-offspring relationships. For the 11 family groups we collected genetic samples from, litter sizes ranged from 1-2 kittens, with an average of 2.3 kittens/litter.




Figure 7. Family groups and known reproduction birth year 2023

Five of the mothers are from previous Minnesota born litters. Nine of the 11 family groups had a father identified within the database.

Persistence

Persistence age distribution of the current year's individuals may be representative of the overall age structure of the NE MN sub-population. Of the 35 individuals detected during the 2023/2024 survey season whose age can be estimated and that were not kittens, 27 (77.1%) have persisted for 2 years or more (Figure 8). Six of these individuals have been detected over 6 years or more. There were two breeding female that has persisted in the database and remained in the survey area for over 10 years. Both GLNR-S-571 and GLNR-S-551 were born in 2013 and both had kittens in 2023. GLNR-S-571 was found dead in May 2024. There are 24 individuals of an unknown age that are new to the database this year and are therefore not considered in persistence analysis. These 24 were travelling alone, tied to a family group but were unrelated, or were part of a family group for which parent-offspring determinations could not be made.




Figure 8. Persistence of individuals 2024 ($n = 84$)

* Individuals new to the database of an unknown age

Canada Lynx – Bobcat Hybridization

A Canada lynx-bobcat hybrid was first detected in Minnesota from a tissue sample obtained from an animal killed by a train in December 2001. This sample along with 2 other Minnesota samples obtained in November and December 2002 represent the first verified hybridization between Canada lynx and bobcat in the wild (Schwartz et al. 2004). Subsequently, genetic samples from other areas have been tested for hybridization resulting in additional hybrid detections in the State of Maine and New Brunswick, Canada (Homyack et al., 2008). The earliest recorded hybrid in Minnesota comes from a specimen that was reportedly harvested in 1997. All samples submitted that are identified as lynx or bobcat are routinely tested for possible hybridization. As part of this monitoring effort, hybrids have been detected in Cook, Itasca, Lake and Pine Counties in Minnesota, and in Polk County, Wisconsin. One hybrid, a male, had persisted for almost 9 years before he was trapped in December of 2019. All hybrids are F1 and thus far there have been no known offspring from either male or female hybrids. All are a result of female lynx and male bobcat mating.

Although not annually, Canada lynx-bobcat hybrid animals were detected on a regular basis (last detection was December 2019) during surveys for lynx but none have been detected since 2019. To

date the database contains 44 hybrid samples. Forty-three of these have been genotyped representing 13 individuals, 5 female and 8 male (Figure 9).




Figure 9. Total Canada lynx-bobcat hybrid individuals detected ($n = 13$)

Take

Section 9 of the Endangered Species Act (ESA) prohibits the take of endangered and threatened species without special exemption. Take is defined in the Endangered Species Act as “to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct” of a federally listed species (16 U.S.C. Chapter 35 Section 1532). The USDI Fish and Wildlife Service (USFWS) maintains a database of reported incidents of “take” of Canada lynx that have occurred in Minnesota since the year 2001. There have been 85 incidents of reported take of Canada lynx since 2001 in Minnesota (Figure 10). Sixty-two of these incidents have resulted in mortalities to the animal. There have been eight incidents of shooting (all mortalities), 39 trapped (19 mortalities and 20 released alive), and 22 that have been hit by a car or truck, snowmobile or a train (19 mortalities, 3 unknown outcomes (carcasses not recovered)). There were also 16 incidents of take that resulted in the mortality of an animal but the cause is unknown. These include cases of likely predation, recovery of decomposed animals or remains, or the recovery of a radio collar that was no longer attached to a study animal.




Figure 10. Reported Canada lynx take in Minnesota 2001-2024 by type and outcomes ($n = 85$)

Incidental Take - Superior National Forest Plan Implementation

Under the terms of ESA, taking that is incidental to and not intended as part of an agency action is not considered to be prohibited taking under the Act provided that such taking follows the terms and conditions of an Incidental Take Statement issued by USFWS. The risk of incidental take of Canada lynx is not eliminated by provisions in the Superior National Forest Land and Resource Management Plan (Forest Plan). On-going implementation of the Forest Plan is expected to result in the incidental take of Canada lynx. Historically, take that occurred due to vehicle collisions on **all roads on all ownerships within the proclamation boundary of the Forest** was considered incidental to the implementation of the Forest Plan (USDI FWS 2011). According to the original Incidental Take Statement (ITS) of the 2011 Biological Opinion (which “revises and supplants, the 2004 Biological Opinion”), “The Service expects no more than one lynx would be taken annually on the Superior National Forest and no more than 10 would be taken over the [generally] 10-year life of the Forest Plan due to vehicle collision on all roads on all ownerships within the Superior National Forest proclamation boundary” (USDI FWS 2011). In 2021, the USFWS issued a statement which clarified that **“Incidental take due to vehicle collisions will only be counted for the ITS when it occurs on USFS jurisdiction roads.”** (USFWS 2021)

In accordance with the 2011 Biological Opinion Reasonable and Prudent Measures, the Superior National Forest is required to “Document and report to the Service annually any known lynx mortality within the National Forest Proclamation boundaries in Minnesota due to vehicle collision, accidental trapping, or poaching”. All reports of lynx mortality are sent directly and immediately to USFWS when they are received by Forest Service personnel.

Since 2011 there have been two reported incidents of lynx take due to vehicle collisions within the Superior NF’s proclamation boundaries that took place on USFS jurisdiction roads – one in 2023 and one in 2024. Additionally, there was one reported instance of lynx mortality along a USFS jurisdictional

road that took place in 2005 prior to the current ITS. There has also been one mortality due to snowmobile on the National Forest (in 2018).

DNA Database Research

Non-invasive genetic capture-mark-recapture

In 2018, we published a paper in the Canadian Wildlife Biology and Management journal that utilized these DNA data to estimate abundance, trend and density estimates for lynx in northeast Minnesota (Barber-Meyer et al. 2018). Since then, we have continued to use the analytical framework as described in this publication to monitor lynx populations in northeastern Minnesota. Figure 11 shows a population estimate chart generated from these data updated with information from 2023/2024 regarding the larger core areas consistently surveyed since 2015. Figure 12 shows a population estimate chart reflecting the smaller core areas that have been consistently surveyed since 2012. These mark/recap estimates have generally tracked with the estimates from the occupancy survey yearly. This last winter the estimate for the smaller core area showed a decline that wasn't shown by the trend in the occupancy survey or in the larger core area. The reason for this is likely due to the bad snow conditions last winter that precluded snowmobiling and had poor conditions on some of the highways for finding tracks and also logging on a few key roads that limited access to those areas. Estimates from this coming winter could help demonstrate if this is an actual decline in the smaller core area or if it was just a sampling effort issue.




Figure 11. Lynx population point estimates and 95% confidence intervals generated using capture-mark-recapture analysis of genetic samples for larger core areas consistently surveyed from January-March of 2015-2024 in the Superior National Forest, Minnesota, USA. Adapted from Barber-Meyer et al. 2018.

JANUARY-MARCH LYNX POPN ESTIMATE, SMALLER CORE AREA 2012-2024




Figure 12. Lynx population point estimates and 95% confidence intervals generated using capture-mark-recapture analysis of genetic samples for smaller core areas consistently surveyed from January-March of 2012-2024 in the Superior National Forest, Minnesota, USA. Adapted from Barber-Meyer et al. 2018.

Multi-Season Occupancy Surveys

In 2016 we collaborated with researchers from North Carolina State University and USDI Fish and Wildlife Service to develop an occupancy model for lynx in northeastern Minnesota (Hostetter and Gardner 2016). We continued to work with researchers to improve our methods which resulted in the publication of the results in Diversity and Distributions in March 2020 (Hostetter et al. 2020). We continue to work with the authors to refine the model and resulting outputs adding probability analysis for colonization and persistence of cells used in the occupancy analysis. The maps in Figure 13 show the outputs of this model using survey effort and number of detections from survey seasons 2014/2015 to 2023/2024.




Figure 13. Lynx winter-specific occupancy probabilities (medians). Grid cells are 5×5 km and encompass Superior National Forest and designated lynx critical habitat in Minnesota, USA

Figure 14 shows that colonization probability (probability an unoccupied grid cell becomes occupied) was low in all years (<0.20) while persistence probability (probability an occupied grid cell remains occupied) was substantially lower in the first (median <0.30) relative to subsequent years (medians >0.90).




Figure 14. Annual lynx average colonization and persistence probabilities (median, 95% credible intervals) across Superior National Forest and designated lynx critical habitat in Minnesota, USA.

Figure 15 shows the proportion of those cells that were surveyed as being occupied during any given survey season. Winter 2023/2024 was a very low snow season which limited conditions for occupancy surveys. The occupancy estimate for last winter was very high but had a large confidence interval due to the limitations in surveying. Impressions from the field last winter did seem to corroborate those results with lynx showing up in areas they normally don't.




Figure 15. Proportion of sites occupied (median, 95% credible intervals)

Acknowledgements

Thanks to this year’s surveyors; H. Becker, K. Fink, T. Brannock, H. Hill, N. Malinowski, Gary Olson, R. Pennesi, P. Robertsen, S. Swanson, A. Myers, T. Munter, B. Peterson, J. Boyd, M. Swingen,. We thank all those folks that have surveyed and collected DNA over the years. They’ve put on a lot of miles across some difficult country in some tough conditions with what some would call the inexplicable ambition to collect a turd. Wolf Ridge Environmental Learning Center also assisted with trailing and sample collection. Thanks to Dr. M. Schwartz, J. Strand, S. McFarland and I. Ortloff at the RMRS National Genomics Center for all their work and support in conducting the DNA analysis. Thank you to Dr. N. Hostetter for his continued work in conducting the analyses of the occupancy survey data. Thanks to T. Smith and M. Kosterman at the USFWS Field Office in Bloomington, MN for their work in gaining additional funding to support this work over the years. Additional funding to conduct surveys and DNA analysis was provided through an agreement with NewRange Copper and Nickel (formerly Polymet Mining, Inc. during survey years 2020, 2021, 2022, 2023 and 2024. And thanks to Steve Loch for his total dedication to this project from the very beginning.

Literature Cited:

- Barber-Meyer, S., D. Ryan, D. Grosshuesch, T. Catton and S. Malick-Wahls. 2018. Use of non-invasive genetics to generate core-area population estimates of a threatened predator in the Superior National Forest. *Canadian Wildlife Biology & Management*. 7 (1): pp.46–55.
- Homyack, J.A., J.H. Vashon, C. Libby, E.L. Lindquist, S. Loch, D.F. McAlpine, K.L. Pilgrim and M.K. Schwartz. 2008. Canada lynx-bobcat (*Lynx canadensis* X *L. rufus*) hybrids at the southern

periphery of lynx range in Maine, Minnesota and New Brunswick. *American Midland Naturalist*. 159:504-508.

Hostetter, N.J. and B. Gardner. 2016. Detection/non-detection surveys to estimate Canada lynx occupancy in Superior National Forest and critical habitat in Minnesota. Report to USDI Fish and Wildlife Service. 21 pp.

Hostetter, N.J., D. Ryan, D. Grosshuesch, T. Catton, S. Malick-Wahls, T.A. Smith and B. Gardner. 2020. Quantifying spatiotemporal occupancy dynamics and multi-year core-use areas at a species range boundary. *Diversity and Distributions*. 26 (7): pp.795-805.

Pilgrim, K.L., K.S. McKelvey, A.E. Riddle and M.K. Schwartz. 2005. Felid sex identification based on noninvasive genetic samples. *Molecular Ecology Notes*. 5: 60-61.

Schwartz, M.K., K.L. Pilgrim, K.S. McKelvey, E.L. Lindquist, J.J. Claar, S. Loch and L.F. Ruggiero. 2004. Hybridization between Canada lynx and bobcat: Genetic results and management implications. *Conservation Genetics*. 5: 349-355.

USDA Forest Service (USDA FS). 2023. Canada Lynx Incidental Take Database. Excel Workbook. USDA FS Unpublished Data.

USDI Fish and Wildlife Service. 2011. Biological Opinion issued in concurrence of reconsultation of implementation of the 2004 Forest Land and Resource Management Plan for the Superior National Forest. September 16, 2011.

USDI Fish and Wildlife Service. 2021. Email from M. Kosterman to C. Ferland on 3/23/2021, Re:(EXTERNAL) RE: Forest Plan ITS

Minnesota Canada Lynx DNA locations

Legend

- ▲ 2023_2024 Lynx samples
- ▲ 2000_2023 Lynx samples
- Superior National Forest Boundary
- Counties

