

Prepared in cooperation with the Minnesota Department of Natural Resources

Minnesota Lake ID: 69-0810


Area: 742 acres

Watershed Area: 3,695 acres

Ecoregion: Northern Lakes and Forests (NLF)



Figure 2. Elephant Lake Watershed map



Elephant Lake

St. Louis County

Sentinel Lakes

Trophic State: Mesotrophic Maximum Depth: 30 feet Mean Depth: ~15 feet Mixing Status: Polymictic

Figure 2. Elephant Lake bathymetric map




Table 1. Elephant Lake land use as compared to typical range for NLF ecoregion reference lakes

| Land use | Elephant Lake land use percentage | NLF typical land use percentage | |
|-----------------|---|---------------------------------|--|
| Developed | <1 | 0 – 7 | |
| Cultivated (Ag) | 0 | <1 | |
| Pasture & Open | <1 | 0 – 6 | |
| Forest | 74 | 54 – 87 | |
| Water & Wetland | 24 | 14 – 31 | |
| Feedlots (#) | 0 | | |


wq-slice69-0810 February 2009

Table 2. Elephant Lake 2008 as compared to typical range for NLF ecoregion reference lakes

Data from Minnesota Department of Health (MDH) laboratory

| Parameter | Elephant Lake 2008 | NLF | |
|---|----------------------|--------------|--|
| Number of reference lakes | | 32 | |
| Total Phosphorus (µg/L) | 24 | 14 – 27 | |
| Chlorophyll mean (µg/L) | 6.5 | 4 – 10 | |
| Secchi Disk (feet) | 9.6 | 8 -15 | |
| (meters) | 2.9 | 2.4 - 4.6 | |
| Total Kjeldahl Nitrogen (mg/L) | 0.57 | 0.4 - 0.75 | |
| Alkalinity (mg/L) | 38 | 40 – 140 | |
| Color (Pt-Co U) | 27 | 10 – 35 | |
| pH (SU) | 6.84 | 7.2 - 8.3 | |
| Chloride (mg/L) | 1.2 | 0.6 - 1.2 | |
| Total Suspended Solids (mg/L) | 1.7 | <1 – 2 | |
| Total Suspended Inorganic Solids (mg/L) | 1.3 | <1 – 2 | |
| Conductivity (umhos/cm) | 86 | 50 – 250 | |
| TN:TP ratio | 23:1 | 25:1 – 35:1 | |
| μg/L = micrograms per liter | Pt-Co-U = Platinum C | Cobalt Units | |
| mg/L = milligrams per liter | SU = Standard Units | | |
| umhos/cm = micromhos per centimet | er | | |

Figure 3. Elephant Lake dissolved oxygen (DO) and temperature profile



0.0 -0.5 -1.0 -1.5 -2.0 -2.5 -3.0 -3.5

Figure 4. Elephant Lake summer mean secchi transparency data

Water Quality, Fishery and Watershed Management Issues

1993

1994

Elephant Lake, located ten miles northeast of Orr, has a surface area of 742 acres. It forms the headwaters of Elephant Creek, a tributary to the Vermilion River (Figure 1). About 45 percent of the lake area is littoral with an overall average depth of about 15 feet. Water residence time is estimated at 3.5 years. It has a small watershed relative to its surface area (5:1 ratio) that is almost entirely forested and wetland (Table 1); there is a moderate amount of lakeshore development – one resort and about 30 homes or cabins.

1995

2007

2008

Based on DO and temperature profiles from 2008 the lake is well mixed and DO remained above 5 mg/L, a level sufficient to support a warm water fishery (> 5 mg/L), throughout the water column (Figure 3). Total phosphorus (TP), chlorophyll-a, and other water quality variables (Table 2) indicate that Elephant Lake is well within the typical range of Northern Lakes and Forests ecoregion reference ('minimally impacted') lakes. There is limited historical water quality data on Elephant. Citizen Lake Monitoring Program (CLMP) Secchi transparency data indicates summer-mean Secchi has varied from 1.4 - 2.8 meters based on five summers of data. Though the more recent measures are deeper than those reported for the mid-1990's the data record is very short and discontinuous and as such, is insufficient to characterize this as a long-term trend (Figure 4). It will be important that CLMP measurement is continued on the lake as a part of the Sentinel Lake monitoring effort.

Elephant Lake is a nutrient-rich lake with relatively low fish species diversity compared with similar lakes with undisturbed forested watersheds. Elephant maintains a self-sustaining walleye population. The walleye population is currently high (Table 3) with many large individuals, but recruitment has been variable over the years. Like walleye, northern pike populations are currently high quality. Similar to other lakes in the border lakes ecoregion, warm-water centrarchids are becoming increasingly abundant, with abundant populations of black crappie and large smallmouth bass. Climate change should continue to favor warm water fish species perhaps to the detriment of native cool water species. If yellow perch (primary forage of walleye and northern pike) remain abundant, Elephant Lake may be able to support modest populations of cool-water and warm-water species over the long term.

Table 3. Primary species captured during recent surveys and their size and abundance compared with other lakes in the same lake class

| Species | Stocked | Abundance | Size | Notes |
|-----------------|---------|-----------|---------------|--------------------|
| Walleye* | N | High | Average-large | |
| Northern Pike | N | Average | Large | |
| Yellow perch | N | High | Average | |
| Black Crappie | N | High | Average | |
| Smallmouth bass | N | High | Large | |
| Bluegill | N | Average | Average | Discovered in 1977 |
| Pumpkinseed | N | Average | Average | |
| White sucker | N | Average | Average | |

^{*}Management emphasis on this species

Table 4. Aquatic plant summary

| Percent cover of aquatic plants ≤ 15ft deep | | |
|---|--|--|
| Lake depth beyond which most vegetation disappeared | | |
| Number of common species (i.e., ≥ 10% cover) | | |
| Non-native plant infestation | | |