

Prairie Restoration Diversity – Planting and Seed Mixes

Why is Diversity in Prairie Restorations Important?

Diversity gives rise to diversity is a foundational paradigm upon which ecology has been built. More diverse plant communities are resistant to invasions because they can more easily compete for resources. Highly diverse plant communities are also predicted to be more stable in the face of disturbance (Tilman and Downing 1994). In prairie restorations, as we think about diversity, we should expand our vision to include phenological (timing of flowering), structural (plant heights), spatial, and functional heterogeneity in addition to plant species richness. This expanded view of diversity will lead to a diverse system of wildlife that inhabit restored prairies. Diversity is important from an ecological perspective as it is the foundation upon which resilience in prairies is built (Helzer et al. 2010), and it is clearly a priority for many tallgrass prairie land managers (Rowe 2010). This fact sheet is a review of the current literature surrounding ways in which diverse tallgrass prairie restorations (i.e., prairie reconstructions) can be established.



Factors that Increase Diversity in Restorations

Several interrelated factors affect diversity in restorations: planting method, density of seeding (seeds per sq ft), grass: forb ratio, density of dominant species, density of non-dominant species, and inclusion of plant functional guilds (cool-season grasses, warm-season grasses, sedges/rushes, legumes, and non-legume forbs).

Planting Methods

Planting during dormancy by broadcasting seed resulted in the most cover of planted species (Larson et al. 2011) and greater resistance to invasion by exotic cool-season grasses (Norland et al. 2015). Dormant broadcasting appears to be the most preferred method by managers (Rowe 2010). However, planting method (dormant-season broadcasting, growing-season broadcasting, or growing-season drilled) may not affect the establishment and cover of those species that were included in the seed mix (Larson et al. 2017) if dormant broadcasting is not possible. Drill-seeded plantings may favor native, dominant warm-season grasses like big bluestem as well as non-native invasive species (Yurkonis et al. 2010). The spatial patterning of planted species and how they interact with each other at small scales (<1 square meter) can also impact diversity and invasibility. Grouping seeds of the same species during planting may increase the established diversity (Yurkonis and McKenna 2014). Similarly, diversity at small spatial scales better resists invasion (Yurkonis et al. 2011) and it is this scale where the diversity in restorations is significantly lower than in remnant prairies (Martin et al. 2005).

Seed Mix Ratios and Seeding Density

A combination of high forb density, high forb:grass ratio, and high species richness are all important, often interrelated factors in achieving greater forb establishment. One study suggests that increasing the density (seeds per sq ft) of forbs will result in greater forb establishment and abundance (Dickson and Busby 2009). Another study suggests that forb establishment was greater in plantings where the ratio was heavily weighted towards forbs (~80%) (Williams and Smith 2007). However, seeding density may not be as important in resisting invasion as seed mix richness (Nemec et al. 2013). Seed mixes heavily weighted towards forbs (>70%) often have trouble carrying a prescribed fire (Diboll 1997). Knowing exactly which target ratios, seeds per sq ft by species, and number of species to plant to yield a diverse prairie community is an area that still needs more study.

Dominant Native Species

The key is achieving the right balance in the seed mix to ensure that forbs and other less dominant species will persist in the restoration. Research has shown that over time there can be up to a 50% decline in species richness from the original prairie planting date, due to the dominance of big bluestem (Camill et al. 2004). Dominant grasses (Table 1) specifically decrease forb cover and species richness (Dickson and Busby 2009, McCain et al. 2010). Warm-season grasses such as big and little bluestem limit recruitment of other less dominant species such as forbs (Tilman 1997). Dominant warm-season grasses can even facilitate invasions due to increased instability in the low-diversity system (Smith et al. 2004). However, warm-season grasses resist invasions through competition for soil nitrogen and are an important component of restorations, especially in species diverse planting (Fargione and Tilman 2005). Short bunchgrasses such as little bluestem are the best warm-season grasses to enhance forb establishment (Wilsey 2010).

Seed Mix Diversity

Species diversity

High seed mix diversity can successfully resist invasion by smooth brome (DiAllesandro et al. 2013) and Kentucky bluegrass (Norland et al. 2015). Of the main drivers of restoration success (seed mix, land use history, landscape context, management, and site characteristics) high seed mix species richness and high forb density are the dominant factors in determining established diversity (Grman et al. 2013).



Photo by Phil Doll

Functional diversity

Functionally rich (the number of functional groups) plantings are more resistant to invasion (Biondini 2007). Species and functionally rich plantings mimic the diversity in native prairies (Piper et al. 2007). Planted species are better competitors of invaders if they are functionally similar (Fargione and Tilman 2005). Native, cool-season grasses in particular are successful at resisting invasion to cool-season invasives (Symstad 2000). Cool-season forbs are an important guild to include in reconstructions because they begin using resources early in the spring (Losure et al. 2007). Increasing phylogenetic diversity (i.e., including species that are more distantly related evolutionarily) of seed mixes is a way to improve functional diversity of established restorations (Barak et al. 2017). Phenological diversity is another way to increase the functional diversity by providing nectar and pollen resources throughout the growing season for pollinators (Havens and Vitt 2016).

These general rules of thumb can be used when developing seed mixes for prairie restorations to increase the established diversity:

- Minimum total seeding rate of 40 seeds per sq ft
- At least 40% of the total seeding rate should be composed of perennial forbs.
- 7 or more native grass or sedge species with at least 2 species of bunchgrass.
- Limit dominant species (Table 1)
- Fulfill the guilds: cool-season and warm-season grasses; sedges/rushes; legume and non-legume forbs.
- 20 or more native forbs with at least 5 species in each bloom period: early, middle, late

*Site conditions, restoration goals, and budget constraints will ultimately determine seed mix composition.

Table 1. Restoration competitive species. Restoration competitive species are aggressive, easily established, and if planted at too high a rate of seeds per sq. ft. will dominate the planting after a few years, lowering the diversity and function of the restoration. Not all species on this list will be applicable to plant across the prairie region. This list is based on field observation and personal experiences of a variety of restoration professionals within and outside of the Minnesota Department of Natural Resources, and literature.

	Latin Name	Common Name	Notes	Source
Forbs	<i>Galium boreale</i>	Northern bedstraw	Can create dense mats that out-compete other spp.	MNDNR
	<i>Monarda fistulosa</i>	Wild bergamot		Helzer et al. 2010
	<i>Ratibida pinnata</i>	Yellow coneflower		Helzer et al. 2010
	<i>Silphium perfoliatum</i>	Cup plant	Creates colonies from fibrous rhizomes that can crowd out other species.	MNDNR
Grasses	<i>Andropogon gerardii</i>	Big bluestem		Losure et al. 2007 McCain et al. 2010
	<i>Panicum virgatum</i>	Switch grass		McCain et al. 2010
	<i>Sorghastrum nutans</i>	Indian grass		McCain et al. 2010

Published June 2017

For additional information on prairie restoration research, please contact:

Jessica Petersen, MN DNR – Prairie Habitat Biologist

jessica.d.petersen@state.mn.us

507-642-8479

References Cited

- Barak, R. S., E. W. Williams, A. L. Hipp, M. L. Bowles, G. M. Carr, R. Sherman, and D. J. Larkin. 2017. Restored tallgrass prairies have reduced phylogenetic diversity compared with remnants. *Journal of Applied Ecology*.
- Biondini, M. E. 2007. Plant diversity, production, stability, and susceptibility to invasion in restored northern tall grass prairies (United States). *Restoration Ecology* 15:77–87.
- Camill, P., M. J. Mckone, S. T. Sturges, W. J. Severud, E. Ellis, J. Limmer, C. B. Martin, R. T. Navratil, A. J. Purdie, and B. S. Sandel. 2004. Community- and ecosystem-level changes in a species-rich tallgrass prairie restoration. *Ecological Applications* 14:1680–1694.
- DiAllesandro, A., B. P. Kobiela, and M. E. Biondini. 2013. Invasion as a function of species diversity: A case study of two restored North Dakota grasslands. *Ecological Restoration* 13:186–194.
- Diboll, N. 1997. Designing seed mixes. Pages 135–150 in S. Packard and C. Mutel, editors. *The Tallgrass Restoration Handbook: For Prairies, Savannas and Woodlands*.
- Dickson, T. L., and W. H. Busby. 2009. Forb species establishment increases with decreased grass seeding density and with increased forb seeding density in a Northeast Kansas, U.S.A., experimental prairie restoration. *Restoration Ecology* 17:597–605.
- Fargione, J. E., and D. Tilman. 2005. Diversity decreases invasion via both sampling and complementarity effects. *Ecology Letters* 8:604–611.
- Grman, E., T. Bassett, and L. A. Brudvig. 2013. Confronting contingency in restoration: Management and site history determine outcomes of assembling prairies, but site characteristics and landscape context have little effect. *Journal of Applied Ecology* 50:1234–1243.
- Havens, K., and P. Vitt. 2016. The importance of phenological diversity in seed mixes for pollinator restoration. *Natural Areas Journal* 36:531–537.
- Helzer, C., B. Kleiman, C. O’Leary, and B. Glass. 2010. Lessons learned from the grassland restoration network: 2003-2010. Pages 1–1322nd North American Prairie Conference.
- Larson, D. L., J. B. B. Bright, P. Drobney, J. L. Larson, N. Palaia, P. A. Rabie, S. Vacek, and D. Wells. 2011. Effects of planting method and seed mix richness on the early stages of tallgrass prairie restoration. *Biological Conservation* 144:3127–3139.
- Larson, D. L., J. Bright, P. Drobney, J. L. Larson, and S. Vacek. 2017. Persistence of native and exotic plants 10 years after prairie reconstruction. *Restoration Ecology*.
- Losure, D. A., B. J. Wilsey, and K. A. Moloney. 2007. Evenness-invasibility relationships differ between two extinction scenarios in tallgrass prairie. *Oikos* 116:87–98.
- Martin, L. M., K. A. Moloney, and B. J. Wilsey. 2005. An assessment of grassland restoration success using species diversity components. *Journal of Applied Ecology* 42:327–336.
- McCain, K. N. S., S. G. Baer, J. M. Blair, and G. W. T. Wilson. 2010. Dominant grasses suppress local diversity in restored tallgrass prairie. *Restoration Ecology* 18:40–49.
- Nemec, K. T., C. R. Allen, C. F. Heizer, and D. A. Wedin. 2013. Influence of richness and seeding density on invasion resistance in experimental tallgrass prairie restorations. *Ecological Restoration* 31:168–185.
- Norland, J., T. Larson, C. Dixon, and K. Askerooth. 2015. Outcomes of past grassland reconstructions in Eastern North Dakota and Northwestern Minnesota: Analysis of practices. *Ecological Restoration* 33:408–417.
- Piper, J. K., E. S. Schmidt, and A. J. Janzen. 2007. Effects of species richness on resident and target species

- components in a prairie restoration. *Restoration Ecology* 15:189–198.
- Rowe, H. I. 2010. Tricks of the Trade: Techniques and Opinions from 38 Experts in Tallgrass Prairie Restoration. *Restoration Ecology* 18:253–262.
- Smith, M. D., J. C. Wilcox, T. Kelly, A. K. Knapp, A. K. Dominance, and A. K. Kna. 2004. Dominance not richness determines invasibility of tallgrass prairie. *Oikos* 106:253–262.
- Symstad, A. J. 2000. A test of the effects of functional group richness and composition on grassland invasibility. *Ecology* 81:99–109.
- Tilman, D. 1997. Community invasability, recruitment limitations, and grassland biodiversity. *Ecology* 78:81–92.
- Tilman, D., and J. Downing. 1994. Biodiversity and stability in grasslands. *Nature* 367:363–365.
- Williams, D., and D. D. Smith. 2007. The effects varying seeding rates of prairie grasses and forbs on native plant establishment in a prairie reconstruction. Ames, Iowa.
- Wilsey, B. J. 2010. Productivity and subordinate species response to dominant grass species and seed source during restoration. *Restoration Ecology* 18:628–637.
- Yurkonis, K. A., and T. P. McKenna. 2014. Aggregating species at seeding may increase initial diversity during grassland reconstruction. *Ecological Restoration* 32:275–281.
- Yurkonis, K. A., B. J. Wilsey, and K. A. Moloney. 2011. Initial species pattern affects invasion resistance in experimental grassland plots. *Journal of Vegetation Science* 23:4–12.
- Yurkonis, K. A., B. J. Wilsey, K. A. Moloney, and A. G. van der Valk. 2010. The impact of seeding method on diversity and plant distribution in two restored grasslands. *Restoration Ecology* 18:311–321.