

STATUS OF MINNESOTA BLACK BEARS, 2006

Report to Bear Committee

26 February 2007

Dave Garshelis

with contributions from

Karen Noyce

*All data contained herein are subject to revision,
due to updated information, improved analysis
techniques, and/or regrouping of data for analysis.*

TABLE OF CONTENTS

Page no.	Table/Fig.	Data description
SUMMARY		
1-2		Summary of key points for each table & figure
HARVEST DATA		
3	Table 1	Licenses, harvests & success rates: 1985–2006
4	Table 2	Permits available by BMU: 2002–2006
4	Table 3	License applicants and purchases by BMU: 2002–2006
5	Table 4	Harvest tally by BMU: 2006 vs. 2001–2005
6	Table 5	Hunting success by BMU: 2006 vs. 2001–2005
7	Table 6	Harvest by date: 1990–2006
NUISANCE COMPLAINTS		
8	Table 7	Participation in nuisance survey, 1985–2006
9–10	Table 8	Number of complaints, bears killed, and bears moved: 1985–2006
FOOD ABUNDANCE		
11	Table 9	Yearly variation in overall bear food abundance by region: 1984–2006
12	Table 10	Abundance of specific foods by region: 2006 vs. previous years
13	Table 11	Yearly variation in fall food abundance by region: 1984–2006
14	Fig. 1	Fall food production 2006 – geographic distribution
POPULATION INDICATORS		
15	Fig. 2	Actual harvests vs. harvests predicted by regression: 1984–2006
16	Fig. 3	Harvest sex ratios by BMU: 2000–2006
17	Fig. 4	Median ages of harvested females by BMU: 2000–2006
18	Fig. 5	Statewide harvest age structure: median ages by sex, 1982–2006
19	Fig. 6	Statewide harvest age structure: proportion in age categories, 1982–2006

Key points: 2006 bear harvest, nuisance activity, foods & population status

Table 1	The number of permit applications was the lowest since 1984. Applications have been declining since 1998. The estimated number of hunters in the field (12,400) was the same as last year.
Tables 2-3	Permits were reduced in 2006 in 4 BMUs that have consistently been undersubscribed, mainly to reduce hunter crowding. Six of 11 BMUs were still undersubscribed, but nearly all surplus licenses were purchased.
Table 4	Estimated harvest (accounting for lost registration data) was 3290, which is close to the 5-year (3436) and 10-year (3389) means. The harvest has been much more stable in the past 4 years than in other 4-year periods. However, harvest by BMU has fluctuated greatly from year-to-year. In 2005, the northwestern no-quota zone (BMU 11) had a record harvest; this year (2006) the harvest was low in that area. However, this year the southern no-quota area (BMU 52) had a record high harvest of 400 bears.
Table 5	Statewide hunting success (25-26% depending on how it is measured; see also Table 1) has been the same for the past 4 years. Within the quota zone, hunting success was significantly higher than normal in BMUs 22, 31 & 51, and lower than normal in 12, 13, 41 & 44 (western areas).
Table 6	As typical for a year with overall "average" fall food abundance, ~70% of the harvest occurred during the first week of the season (this does not vary with the day of the week for opening day).
Tables 7-8	The number of wildlife and enforcement personnel submitting bear nuisance tally forms each month was about normal. However, the number of bear complaints investigated on-site was the lowest ever recorded (57; down from >1500 in 1995), as was the number of bears killed as nuisances (21, including early hunting kills).
Tables 9-11	Overall food conditions (summer–fall) were not particularly high or low in any parts of the bear range. However, several summer foods tended to have low fruit abundance (due to drought conditions in June-July), whereas a few fall foods had above-average production. The various fruits differ in their impacts on harvest and nuisance activity.
Fig. 1	Three primary fall foods tended, as a group, to be lowest in the northeast and highest in the central part of the state. Especially high acorn production in the northwest accounted for poor hunting success in that area, whereas poor oak production in the southeastern bear range accounted for the record harvest there.

Fig. 2	A combination of two key factors, fall food abundance and number of hunters, accounts for 88% of the yearly variation in the harvest. In each of the past 5 years, however, the regression based on these 2 variables predicted a slightly higher harvest than actually occurred.
Fig. 3	Sex ratios of harvested bears reflect both the sex ratio of the living population as well as the relative vulnerability of the sexes to hunters. Harvest sex ratios tend to be more male-dominated and also more variable in the northwestern part of the range (BMUs 11,12,13). BMU 41 also is particularly variable because of its small size and because many bears there are killed near cropfields. In years with poor natural foods, more bears are attracted to cropfields and hunters' baits, and the harvest is less male-biased. In 2006, natural foods were exceptionally good in the area around BMU 41 (Fig. 1), so the harvest there was very male-biased.
Fig. 4	Ages of harvested bears also reflect both the age structure of the living population as well as the relative vulnerability of bears to hunters (including hunter selection for larger, older bears). Harvest ages of females (shown in this figure) are more variable than for males, reflecting differing vulnerability to hunting by food conditions (older females increasing in vulnerability in poorer food years). The more heavily-hunted, southerly BMUs have a younger age structure. The northern BMUs show high year-to-year variation due to fluctuating food resources.
Fig. 5-6	Ages of harvested bears of both sexes steadily declined for about 2 decades (decline in median age and increase in proportion of 1-2 year olds in the harvest), reflecting increasingly higher harvest levels over this period. More consistent harvests of about 3400 bears during the past 4 years (Table 1) seem to have stabilized the age structure (with the hint of a recent slight increase in ages of harvested bears).

Table 1. Bear permits, licenses, hunters, harvests, and success rates, 1985–2006.

	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Permit applications	22954	20694	19687	25879	24096	24861	25890	26428	27365	30127	29922	30405	27353	30245	29384	29275	26824	21886	16431	16466	16153	15725
Permits available	4290	4730	4810	5310	5520	6370	7140	7920	8630	9400	11950	12030	11370	18210	20840	20710	20710	20610	20110	16450	15950	14850
Licenses purchased (total) ^a	3948	4188	6054	5643	5901	7094	7757	8485	9224	9826	12448	12414	11440	16737	18355	19304	16510	14639	14409	13669	13199	13164
Quota area ^a	3948	4188	4213	4297	4628	5568	6257	6845	7528	8125	10304	10592	9655	14941	16563	17021	13632	12350	9833	10063	9340	9169
Quota surplus/military ^a																	235	209	2554	1356	1591	1561
No-quota area ^a			1841	1346	1273	1526	1500	1640	1696	1701	2144	1822	1785	1796	1792	2283	2643	2080	2022	2238	2268	2434
% Licenses bought ^b																						
Of permits available ^b	92.0	88.5	87.6	80.9	83.8	87.4	87.6	86.4	87.2	86.4	86.2	88.0	84.9	82.0	79.5	82.2	67.0	60.9	61.6	69.4	68.5	72.3
Of permits issued ^b														84.4	87.2	83.9	69.8	66.3	65.7	68.3	67.1	68.9
Estimated no. hunters ^c	3700	3900	5600	5100	5500	6600	7200	7900	8600	9100	11600	11500	10300	14500	15900	16800	15500	13700	13500	12800	12400	12400
Harvest	1340	1438	1577	1509	1930	2381	2143	3175	3003	2329	4956	1874	3212	4110	3620	3898	4936	1915	3598	3391	3340 ^d	3290 ^d
Harvest sex ratio (%M) ^e	53	59	60	58	57	52	59	50	56	62	47	62	55	55	53	58	56	61	58	57	59	58
Success rate (%) ^f																						
Total harvest/hunters	36	37	28	30	35	36	30	40	35	26	43	16	31	28	23	23	29	14	26	26	26	26
Quota harvest/licenses			33	28	36	35	30	41	34	26	42	15	29	25	20	20	28	14	25	26	25	25

^a Quota area established in 1982. No-quota area established in 1987. Surplus licenses from undersubscribed quota areas sold beginning in 2000; originally open only to unsuccessful permit applicants, but beginning in 2003, open to all. Total licenses = quota + quota surplus + no-quota + military (no permit needed).

^b Quota licenses bought (including surplus)/permits available, or licenses bought (prior to surplus)/permits issued (permits issued more relevant for years when some areas were undersubscribed; see Table 3).

^c Number of licensed hunters x percent of license-holders hunting. Percent hunting is based on data from bear hunter surveys conducted during 1981–91, 1998 (86.8%), and 2001(93.9%).

^d Harvest estimated from tallied registration + lost registration data (ascertained from tooth envelopes received without matching registration data)..

^e Sex ratio as reported by hunters; hunters classify about 10% of female bears as males, so the actual harvest has a lower %M than shown here. In good food years, the harvest is more male-biased.

^f Success rates in 2001–2004 were calculated as number of successful hunters/total hunters, rather than bears killed/total hunters, because hunters could take 2 bears. This was complicated even more in 2005 and 2006 because the total harvest was estimated (footnote d), and hunters could take 1 bear in the quota area plus 2 bears in the no-quota area. From the registration tally and tooth envelopes received in 2006, 50 hunters took more than 1 bear (45 took 2 bears on NQ license, 2 hunters took 1 quota and 1 NQ bear, and 3 hunters took 2 bears on a quota license [illegally]); thus, there were 3290-50 = 3240 successful hunters.

Table 2. Number of bear hunting permits available per year, 2002–2006 (aligned with permit applications in Table 3 below; highlighted numbers show drop from previous year).

BMU	2006	2005	2004	2003	2002
12	550	550	700	700	700
13	800	900	900	1100	1100
22	150	150	150	250	250
24	1000	1200	1200	1500	1500
25	1900	1900	1900	2400	2400
26	1500	1500	1500	1500	1500
31	2100	2100	2100	2660	2660
41	450	450	500	500	500
44	1700	1700	2000	2500	3000
45	1200	1500	1500	2000	2000
51	3500	4000	4000	5000	5000
Total	14850	15950	16450	20110	20610

Table 3. Number of bear hunting license applicants, and number and percent of available surplus licenses bought, 2002–2006^a.

BMU	2006		2005		2004		2003		2002	
	Apps	Surplus bought	Apps	Surplus bought	Apps	Surplus bought	Apps	Surplus bought	Apps	Surplus bought
12	1005		864		808		837		1061	
13	680	120 100%	714	186 100%	670	129 56%	668	167 39%	831	41 18%
22	92	58 100%	65	46 54%	73	47 61%	88	26 16%	124	5 4%
24	624	367 98%	749	270 60%	766	259 60%	756	193 26%	979	40 8%
25	1789	112 100%	1923		1793	111 100%	1716	317 46%	1985	41 11%
26	1915		1997		2110		2280		2873	
31	2290		2097	4 100%	2006	92 100%	1996	412 62%	2503	26 23%
41	683		653		601		688		810	
44	2838		2884		2934		2855		4043	
45	840	360 100%	927	346 60%	1092	332 81%	1069	461 50%	1535	56 14%
51	2969	531 100%	3276	726 100%	3613	386 100%	3467	978 64%	5141	
None	0		0		0		2		1	
Total	15725	1548 ~100%	16149	1578 78%	16466	1356 78%	16431	2554 50%	21886	209 12%

^a Surplus licenses available beginning in 2001, but restricted to permit applicants in 2001 & 2002.

Undersubscribed

Nearly undersubscribed

Table 4. Minnesota bear harvest tally^a for 2006 by Bear Management Unit (BMU) and sex compared to harvests during 2001-2005 and record high harvests.

BMU	2006					2005	2004	2003	2002	2001	5 year mean	Record high harvest (yr)
	M	(%M)	F	U	Total							
Quota												
12	48	(69)	22	0	70	165	165	174	104	263	174	263 (01)
13	98	(65)	53	0	151	205	197	185	116	241	189	258 (95)
22	6	(40)	9	0	15	8	10	3	7	6	7	41 (89)
24	102	(53)	92	0	194	144	212	163	101	273	179	288 (95)
25	196	(47)	225	0	421	404	546	510	328	584	474	584 (01)
26	189	(60)	124	1	314	285	320	303	171	397	295	513 (95)
31	320	(66)	162	0	482	445	484	436	301	697	473	697 (01)
41	27	(68)	13	0	40	104	83	100	51	201	108	201 (01)
44	120	(62)	72	0	192	273	283	444	183	553	347	643 (95)
45	60	(51)	57	1	118	107	118	143	36	178	116	178 (01)
51	411	(57)	308	2	721	505	544	667	300	895	582	895 (01)
Total	1577	(58)	1137	4	2718	2759 ^b	2962	3128	1698	4288	2967	4288 (01)
No Quota^c												
11	87	(72)	33	0	120	335	177	200	112	321	229	351 (05)
52	216	(54)	183	1	400 ^d	223	252	270	105	327	235	382 (93)
Total	303	(58)	216	1	520	581 ^b	429	470	217	648	469	678 (95)
State	1880	(58)	1353	5	3290^b	3340^b	3391	3598	1915	4936	3436	4956 (95)

^a Harvest data were obtained from registration slips electronic registration, and tooth envelopes. The following table shows the number of tooth envelopes that had no corresponding registration slip or e-registration.

Year	Quota area	No-quota area
2001	56	7
2002	46	7
2003	84	13
2004	96	39
2005	179	31
2006	63	15

^b The estimated registered harvest, including those in which registration data were lost and no tooth envelope was received. Value for 2006 does not match column or row total because other data on table are uncorrected for estimated lost registration data.

^c Some hunters with no-quota licenses hunted in the quota area. Some were drawn for the quota area but received NQ licenses. Others hunted in the wrong area purposefully or out of ignorance ($n = 48$ in 2006).

^d Record high harvest in area 52 in 2006. Last column on this line shows previous record.

Table 5. Bear hunting success (%) by BMU, measured as the registered harvest (excluding second bear) divided by the number of licenses sold^a, 2001–2006.

BMU	Mean success 2001-2005	2006		2005 ^b		2004		2003		2002		2001	
		% Success	% Taking 2 bears ^c	% Success	% Taking 2 bears ^c	% Success	% Taking 2 bears ^c	% Success	% Taking 2 bears ^c	% Success	% Taking 2 bears ^c	% Success	% Taking 2 bears ^c
Quota	24	25	—	25	—	26	—	25	—	14	—	28	(11)
12	35	19	—	41	—	33	—	35	—	22	—	44	(17)
13	29	24	—	32	—	33	—	31	—	19	—	31	(9)
22	8	14	—	10	—	11	—	4	—	8	—	7	(0)
24	23	25	—	20	—	27	—	25	—	15	—	28	(8)
25	32	30	—	30	—	38	—	34	—	23	—	34	(11)
26	29	30	—	34	—	31	—	29	—	17	—	32	(10)
31	28	33	—	31	—	33	—	25	—	17	—	34	(15)
41	27	13	—	31	—	23	—	29	—	14	—	40	(16)
44	21	16	—	24	—	20	—	26	—	9	—	23	(10)
45	11	14	—	13	—	12	—	13	—	4	—	13	(7)
51	18	28	—	18	—	19	—	21	—	9	—	24	(10)
No Quota	19	22 ^d	(9)	23	(9)	18	(7)	21	(10)	10	(7)	23	(9)
Statewide	23	25	—	25	—	25	—	25	—	13	—	27	(11)

^a Harvest/licenses instead of harvest/hunters because BMU-year-specific estimates for the rate of hunting by licensed hunters are unreliable. Statewide estimates of harvest/hunters are presented in Table 1.

^b For 2005, estimated registered harvest was used instead of known registered harvest due to a large loss of registration data.

^c Percent of successful hunters that shot 2 bears; 2nd bear is not included in the calculation of hunting success. The taking of 2 bears was legal statewide in 2001, but only in the no-quota area in 2002–2006.

^d Although BMU 52 had a record harvest (see Table 1), there is no way to split BMUs 11 and 52 to examine hunting success because the number of hunters in each area is unknown (a single NQ license covers both BMUs).

Table 6. Cumulative bear harvest (% of total harvest) by date, 1990–2006.

Year	Day of week for opener	Aug 22/23 – Aug 31 (9–10 days)	Sep 1 – Sep 7 (7 days)	Sep 8 – Sep 14 (7 days)	Sep 15 – Sep 30 (16 days)
1990	Sat		69	82	96
1991	Sun		64	76	93
1992	Tue		72	86	96
1993	Wed		67	80	94
1994	Thu		67	78	92
1995	Fri		72	87	97
1996	Sun		56	70	87 ^a
1997	Mon		76	88	97
1998	Tue		76	87	96
1999	Wed		69	81	95
2000	Wed	57	72	82	96
2001	Wed	67	82	88	98
2002	Sun		57	69	90 ^a
2003	Mon		72	84	96
2004	Wed		68	82	95
2005	Thu		72	81	94
2006	Fri		69	83	96

^a The large proportion of the harvest taken late in the season in 1996 and 2002 (e.g., >10% in October) was related to the high abundance of food in those years.

Table 7. Number of people participating in nuisance bear survey, 1985^a – 2006.

	Apr	May	Jun	Jul	Aug	Sep	Oct
1985	17	29	37	30	26	23	20
1986	37	52	52	51	47	46	32
1987	45	71	75	65	62	52	37
1988	68	74	77	75	73	68	69
1989	67	84	80	85	81	79	66
1990	75	79	80	81	78	74	70
1991	82	83	87	85	82	85	67
1992	74	79	81	85	83	74	62
1993	83	84	82	88	82	81	68
1994	77	88	82	86	83	68	61
1995	74	77	79	83	80	72	61
1996	71	83	84	77	75	67	54
1997	61	69	69	64	62	60	43
1998	34	67	71	63	55	41	33
1999	52	52	40	47	44	39	16
2000	60	58	50	54	42	37	33
2001 ^b	52	54	50	49	42	32	21
2002	50	44	43	46	35	29	19
2003	36	39	34	29	27	25	14
2004	28	33	34	32	32	24	13
2005	35	36	42	36	35	26	20
2006	28	39	46	43	30	29	24

^a Monthly tallies of complaints were required of Conservation Officers and Wildlife Managers beginning in 1984.

^b Electronic submission of monthly complaint tally beginning in 2001.

Table 8. Number of nuisance bear complaints registered by Conservation Officers and Wildlife Managers during 1985–2006, including number of nuisance bears killed and translocated, and bears killed in vehicular collisions.

	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Number of personnel participating in survey ^a	37	52	75	77	85	81	87	85	88	86	83	84	69	71	52	60	54	50	39	34	42	46
Complaints examined on site ^b	1115	972	789	771	1117	1890	935	1562	1010	696	1568	337	661	226	189	105	122	75	81	75	61	57
Complaints handled by phone ^c	1744											959	2196	743	987	618	660	550	424	507	451	426
Total complaints received	2859											1296	2857	969	1176	723	782	625	505	582	512	483
• % Handled by phone	61%											74%	77%	77%	84%	85%	84%	88%	84%	87%	88%	88%
Bears killed by:																						
• Private party or DNR	364	221	150	134	157	321	97	187	111	67	232	27	93	31	25	25	22	12	13	25	28	11
• Hunter before season ^d																						
– from nuisance survey	15	21	9	44	27	69	14	38	21	28	81	6	32	23	5	7	4	0	3	3	6	2
– from registration file	15	11	9	35	15	50	15	52	30	25	138	18	35	31	24	43	20	11	8	4	13	6
• Hunter during/after season ^e	4	3	6	11	15	21	16	19	8	3	13	0	4	3	0	1	1	0	0	0	1	0
• Permittee ^f							20	28	6	3	57	4	7	11	7	2	6	4	6	1	5	4
Bears translocated	116	123	152	109	257	358	214	342	180	171	295	64	115	24	29	1	6	3	1	3	3	3
• % bears translocated ^g	10	13	19	14	23	19	23	22	18	25	19	19	17	11	15	1	5	4	1	4	5	5
Bears killed by cars	119	95	75	46	69	74	50	90	54	40	68	42	52	61	60	39	43	26	25	16	22 ^h	17 ^h

Table 8 footnotes:

- ^a Maximum number of people turning in a nuisance bear report each month (from Table 7). Monthly reports were required beginning in 1984.
- ^b Adjusted for low and variable survey participation during 1981–86.
- ^c Tallies of complaints handled by phone were made only during the indicated years.
- ^d The discrepancy between the number recorded on the nuisance survey and the number registered before the opening of the season indicates incomplete data.
- ^e Data only from nuisance survey because registration slips do not indicate whether bear was a nuisance.
- ^f A permit for non-landowners to take a nuisance bear before the bear season was officially implemented in 1992, but some COs individually implemented this program in 1991. Data are based on records from the nuisance survey, not directly from permit receipts.
- ^g Percent of on-site investigations resulting in a bear being captured and translocated.
- ^h Car kill data were reported on the monthly nuisance form for the first time in 2005 (value shown). In all previous years, car kill data were from confiscation records. Confiscation records in 2005 indicated 18 car kills.

Table 9. Bear food index values for five survey areas (see map below) in northern Minnesota's bear range, 1984 – 2006. Pink-shaded blocks indicate particularly low index values (<45); green blocks indicate particularly high index values (>70).

Year	Survey Area					Entire Range ^a
	NW	NC	NE	WC	EC	
1984	32.3	66.8	48.9	51.4	45.4	51.8
1985	43.0	37.5	35.3	43.5	55.5	42.7
1986	83.9	66.0	54.7	74.7	61.1	67.7
1987	62.7	57.3	46.8	67.4	69.0	61.8
1988	51.2	61.1	62.7	54.4	47.3	56.0
1989	55.4	58.8	48.1	47.8	52.9	51.6
1990	29.1	39.4	55.4	44.0	47.9	44.1
1991	59.7	71.2	64.8	72.1	78.9	68.4
1992	52.3	59.9	48.6	48.1	63.3	58.2
1993	59.8	87.8	75.0	73.9	76.8	74.3
1994	68.6	82.3	61.3	81.5	68.2	72.3
1995	33.8	46.5	43.9	42.0	50.9	44.4
1996	89.5	93.2	88.4	92.2	82.1	87.6
1997	58.2	55.5	58.8	62.0	70.1	63.9
1998	56.9	72.8	66.4	72.3	84.5	71.1
1999	63.7	59.9	61.1	63.2	60.6	62.0
2000	57.7	68.0	54.7	69.2	67.4	62.3
2001	40.6	48.7	55.6	62.2	66.0	55.8
2002	53.1	63.4	60.4	68.6	68.3	66.8
2003	59.1	57.5	55.2	58.6	49.7	58.8
2004	57.0	60.5	61.1	70.3	67.9	64.4
2005	53.4	65.9	61.4	59.9	72.6	62.3
2006	51.0	64.9	53.4	51.0	52.1	56.9

^a Values represent the sums of mean statewide index values for 14 species surveyed. Means were calculated using all surveys completed in the state, not by averaging values from the 5 food survey areas.

Fig 1. Boundaries of Minnesota's 5 bear food survey areas.




Table 10. Index values of bear food abundance^a in 2006 compared to the previous 22-year mean (1984-2005) in 5 survey areas across Minnesota’s bear range. Pink-shaded blocks indicate poor fruit abundance (abundance index ≥ 1 point lower than average); green blocks indicate high fruit abundance (≥ 1 point higher than average).

FRUIT	NW		NC		NE		WC		EC		Entire Range	
	22yr \bar{x}	2006 $n = 10^b$	22yr \bar{x}	2006 $n = 15$	22yr \bar{x}	2006 $n = 16$	22yr \bar{x}	2006 $n = 13$	22yr \bar{x}	2006 $n = 12$	22yr \bar{x}	2006 $n = 56^c$
SUMMER												
Sarsaparilla	4.2	3.3	5.9	5.6	5.4	4.9	4.7	3.3	5.7	4.6	5.1	4.7
Pincherry	3.2	4.0	4.4	4.9	4.1	3.1	4.0	4.0	3.7	3.1	3.8	3.8
Chokecherry	5.5	3.8	5.1	4.3	4.2	3.3	5.5	3.8	4.6	3.2	5.0	3.6
Juneberry	4.7	3.6	4.6	4.6	4.7	4.3	3.6	3.6	4.0	2.5	4.2	3.9
Elderberry	1.5	1.6	3.0	5.5	3.5	3.6	3.3	1.6	3.3	3.3	3.0	3.7
Blueberry	4.9	2.6	5.2	2.3	4.7	2.2	3.6	2.6	3.7	1.1	4.3	2.2
Raspberry	6.5	4.8	8.0	5.9	8.0	5.4	7.1	4.8	7.0	4.5	7.3	5.5
Blackberry	1.4	0.8	2.3	3.1	0.8	1.6	3.5	0.8	4.6	3.1	2.8	2.8
FALL												
Wild Plum	1.9	1.9	1.7	2.0	0.8	1.8	2.6	1.9	2.3	2.0	2.0	2.0
HB Cranberry	5.2	3.6	4.2	4.7	3.4	3.7	3.7	3.6	3.5	3.3	3.9	3.5
Dogwood	5.8	5.0	5.8	5.5	5.1	4.1	5.7	5.0	6.1	5.3	5.6	4.9
Oak	3.2	5.3	2.7	3.3	1.4	2.3	5.8	5.3	6.0	2.9	4.3	4.4
Mountain Ash	1.4	3.2	2.3	5.7	4.3	5.4	1.7	3.2	1.9	5.5	2.4	4.2
Hazel	6.0	7.5	7.5	7.5	7.2	7.7	8.2	7.5	7.8	7.7	7.4	7.7
TOTAL	55.4	51.0	62.7	64.9	57.6	53.4	63.0	51.0	64.2	52.1	61.1	56.9

^a Food abundance indices were calculated by multiplying species abundance ratings x fruit production ratings.

^b n = Number of surveys used to calculate 2006 mean index values.

^c Sample size for the entire bear range does not equal the sum of the sample sizes of the 5 areas because some surveys were conducted on the border of 2 or more areas and were included in tabulations for each area.

Table 11. Regional productivity indices (summed) for oak, hazel, and dogwood, 1984 – 2006. Shaded blocks indicate particularly low (≤ 5.0 , yellow) or high (≥ 7.5 , tan) fall food productivity.

Year	Survey Area					Entire Range ^a
	NW	NC	NE	WC	EC	
1984	4.2	7.6	7.0	6.2	7.0	6.5
1985	4.9	2.8	4.2	4.7	5.3	4.4
1986	7.2	5.0	4.0	7.0	6.2	6.2
1987	8.0	7.8	7.3	7.6	8.0	7.7
1988	5.5	7.2	7.3	6.8	6.1	6.7
1989	6.0	5.3	4.1	5.7	6.4	5.8
1990	3.3	4.2	6.4	5.7	6.4	5.2
1991	6.2	6.2	5.4	7.2	7.7	6.7
1992	4.7	5.0	4.4	4.4	6.8	5.1
1993	5.3	7.1	6.7	6.2	7.7	6.5
1994	7.1	7.8	5.8	7.8	7.1	7.2
1995	4.8	4.8	5.1	4.6	5.3	4.9
1996	8.7	8.6	8.1	9.2	8.5	8.6
1997	5.8	5.4	5.1	6.8	6.5	6.2
1998	5.8	6.0	6.3	7.1	7.8	6.7
1999	6.4	5.1	5.9	6.6	6.0	6.2
2000	5.8	7.7	7.2	7.5	8.5	7.0
2001	3.4	4.1	5.7	6.0	6.5	5.2
2002	8.7	7.1	6.6	8.8	8.2	8.1
2003	6.3	6.0	5.5	6.2	6.0	6.1
2004	6.1	5.4	5.4	6.4	6.1	5.9
2005	5.8	5.8	6.1	6.4	7.0	6.2
2006	6.7	6.1	6.0	6.7	5.8	6.3

^a This value represents the sum of mean statewide productivity index values for hazel, oak, and dogwood. Means were calculated using all surveys completed in the state, not by averaging values from the 5 food survey areas.

Fig. 1. Fall production of primary bear foods, 2006.




Fig 2. Number of bears killed vs. number predicted, based on fall food abundance and hunter numbers. Prediction for 2006 based on regression from 1984–2005 ($R^2 = 0.88$).




Fig 3. Sex ratios of harvested bears by BMU, 2000–2006





Fig 4. Median ages of harvested female bears by BMU, 2000–2006



* Sample size too small in 2006

