



2019 MINNESOTA BOWHUNTER OBSERVATION SURVEY REPORT

Tyler R. Obermoller, Wildlife Research Biologist
Eric S. Michel, Ungulate Research Scientist
Lou Cornicelli, Wildlife Research Manager

INTRODUCTION

White-tailed deer (*Odocoileus virginianus*) hunting season recommendations should incorporate objective and reliable information to move populations towards a desired density goal. Because the Minnesota Department of Natural Resources (MNDNR) adjusts regulatory decisions (seasons and bag limits) annually, agencies require current information. In Minnesota, deer are managed by individual deer permit areas (DPAs; $N = 130$) with traditional firearm season lengths of 9 (200-series areas), 16 (100-series areas), or 18 (300-series areas; 2 seasons) days. Bag limits also vary by permit area and range from bucks only (1 antlered deer) to intensive (1 buck and up to 3 antlerless deer) management designations. Additionally, early antlerless seasons are used in limited situations. To inform these annual decisions, the MNDNR incorporates mandatory hunter-reported harvest, hunter effort, winter severity, and vital rate parameters (survival, fecundity, etc) into a population model to make population trend inferences (i.e., λ ; Norton and Giudice 2017). Population model indices are sensitive to varying hunting season regulations and changes in the relationship between winter severity and deer survival. Confidence in the population model is improved by collecting annually recurrent information to independently estimate the population trend. The Office of Legislative Auditors conducted an independent evaluation of the MNDNR deer population management program (OLA 2016) and recommended additional data collection to improve deer population estimates. Winter aerial surveys can provide an index, but logistical and environmental (adequate snow cover) constraints limit their use to every 5- to 10-years. Furthermore, aerial surveys are not considered reliable across much of northern Minnesota where predominant coniferous cover results in insufficient detection probability (Haroldson 2014) or across southwestern Minnesota where deer movements vary throughout the year (winter migrations).

Several Midwestern states have explored the use of annual hunter observation surveys for monitoring white-tailed deer population trends (Rolley et al. 2016). Early archery season (Saturday closest to September 15 to the Saturday closest to November 6) observation surveys are desirable because they are longer than firearm seasons (9 or 16-day season starting the Saturday closest to November 6). Bowhunters also typically employ stationary hunting methods (tree stand, ground blind) which allow more time to observe undisturbed wildlife (Norton and Clark 2016). Thus, our objective was to evaluate the utility of bowhunter observation surveys in Minnesota for monitoring trends in white-tailed deer and other wildlife populations. Our secondary objective was to compare trends in fawn:adult female ratios from bowhunter observations to other recruitment metrics. In Minnesota, landscape types vary more than other Midwestern states. Because of the variability of habitat, we chose to evaluate results among three ecozones: 1) farmland, 2) transition, and 3) forest (Figure 1). To evaluate the most efficient data collection strategy, we developed both a mail and online survey instrument.

METHODS

We modeled our survey after the Iowa Department of Natural Resources (IADNR) bowhunter observation survey (Norton and Clark 2016). The primary differences between our survey and the IADNR bowhunter survey were the species monitored, age-sex classification of deer, and

the addition of a separate online survey. Specifically, we asked hunters to document badger (*Taxidea taxus*), bear (*Ursus americanus*), bobcat (*Lynx rufus*), coyote (*Canis latrans*), fisher (*Martes pennanti*), gray fox (*Urocyon cinereoargenteus*), gray wolf (*Canis lupus*), and wild turkey (*Meleagris gallopavo*) observations (Appendix I) and differentiate between antlered, adult female, fawn, and unknown white-tailed deer age-sex classes. We also asked hunters to record DPA for hunting trip observations and provide a distance and direction from the nearest town. On the online survey, we also collected more precise locations (latitude and longitude), weather information, antler points of harvested deer, and inside antler spread of harvested deer.

Our sampling frame included individuals that purchased an archery deer hunting license for 2 consecutive seasons (2017 and 2018; $n = 65,628$). In Minnesota, collection of email addresses is not mandatory and only 31% ($n = 20,233$) of bowhunters had an email on file. Therefore, to draw our samples we first randomly selected 18,000 individuals to receive a mail survey, regardless of whether they provided an email. We mailed a one-page, front and back, hunting diary log with a cover letter and postage paid return envelope to paper survey participants (Appendix I). The remaining individuals with an email on file received the online survey ($n = 16,404$). We also provided online survey participants the option to print and then mail the observation diary log (Appendix I). Sampling rates for the mailed survey were higher than the online survey in all ecozones except the transition zone because of sampling design (Table 1). We also mailed and emailed follow-up letters to survey individuals prompting them to complete and then return their survey.

We evaluated differences in mean respondent age and response rates between the mail and online surveys. We also evaluated differences between date of hunting trips, hunting trips per hunter, hours hunted per trip, and observation rates between the mailed and online respondents. We clustered the survey design for response data by individual hunter and provided separate estimates for each ecozone. We estimated variances using Taylor series linearization and constructed 95% confidence intervals using the Normal approximation. We used t-tests to compare all responses between mail and online respondents. We used an alpha value of 0.05 to determine significant differences between mail and online response groups. For observation rates, we applied a Bonferroni correction to account for 13 species or cohort categories resulting in a critical alpha value of 0.004. We estimated hours hunted per hunting trip and observation rates per hour using Program R and the survey library (Lumley 2004, R Development Core Team 2016).

We did not compare hunter observation rates among ecozones because hunter distribution, similar to deer populations, is not randomly distributed. Thus, hunter observation rates among ecozones vary by hunter distribution. For example, deer densities are highest in the transition ecozone (Norton and Giudice 2017), but hunter observation rates per 1,000 hours were greatest in the farmland ecozone. Therefore, we only compared the relative proportion of species hunters observed across ecozones.

RESULTS

After removing undeliverable samples, we administered 17,725 mail and 16,404 online surveys. Of those, we received 1,723 mail and 457 online responses, which resulted in adjusted response rates of 0.097 and 0.028, respectively (Table 1). Response rates were comparable among regions; however, they differed between survey modes (ranges: mail response rate, 0.095–0.101; online response rates, 0.022–0.029; Table 1). Mean age of respondents was older than the sample for both the mail ($\bar{x} = 54$ vs 44 years) and online ($\bar{x} = 48$ vs 39) surveys. Online respondents also averaged 38% fewer trips per hunter ($\bar{x} = 5.71$, SE = 0.28) compared to mail respondents ($\bar{x} = 9.21$, SE = 0.17; Figure 2). Mean hunting observation dates occurred later for mailed ($\bar{x} = 16$ October) vs online ($\bar{x} = 11$ October) responses (Figure 3). Despite lower response rates and fewer hunter observations later in the season, hours hunted per trip

(online $\bar{x} = 3.22$, SE = 0.07, mail $\bar{x} = 3.23$, SE = 0.03) and hunter observation rates per hour among species did not differ between survey modes (Figures 4-10; Table 2, Appendix II).

Overall, the percent of antlered deer among total deer observations was similar to previous years and comparable among regions with the greatest observations occurring in the transition ecozone ($\bar{x} = 0.20$), followed by the farmland ecozone ($\bar{x} = 0.19$), then the forest ecozone ($\bar{x} = 0.17$). The greatest observed fawn:doe ratio was in the transition ecozone ($\bar{x} = 0.77$), followed by the farmland ecozone ($\bar{x} = 0.68$) and forest ecozone ($\bar{x} = 0.63$, Figures 5–7). Among other species surveyed, diversity was greater in the forest ecozone with relatively more bear, bobcat, wolf, fisher, and gray fox observations compared to the transition and farmland ecozones. Turkeys had the highest proportion reported (compared to all other species) in the transition ecozone (Appendix II).

For the hunter-harvested data recorded on the online responses, 63 hunters harvested 64 adult bucks. The adult bucks averaged 7.5 points (SE = 0.31, range = 2–12, $n = 64$) with an inside spread of 13.9 inches (SE = 0.57, range = 5–25, $n = 25$).

DISCUSSION

Although mean hunter age, response rates, and trips per hunter were significantly different between online and mail respondents, similar observation rates suggested that inferences about population trends could be obtained from either survey mode. However, the low response rates and low number of trips per hunter from the online survey results in a reduced amount of information. Online respondents recorded fewer observations later in the season than mailed respondents, leading to fewer overall trips per hunter. Because online respondents could input data throughout the hunting season, we postulate they inputted their early season observations, but failed to input their late season observations, whereas, mailed respondents recorded observations on their datasheet until the survey period was done, and then sent in their datasheet.

In the future, we will continue to explore methods that increase response rates, especially for the online survey, such as sending additional reminder emails throughout the observation season or adding an incentive for participation. Also, we will continue to share the results from this survey with future participants to generate interest and increase participation. More importantly, we do not currently know whether trends in observation rates among years will be similar between survey modes.

We intend to conduct a formal analysis on all three years to evaluate the trend inference between survey mode and region. We can also determine whether this survey will contribute to our knowledge of population trends and, if so, determine the minimum spatial scale required to provide reliable inferences. We will compare correlations for each species and cohorts between the two survey modes. We will also compare the correlation between the results from the bowhunter survey indices and recruitment rates with estimates from harvest modeling techniques and antlered harvest catch-per-unit-effort. Finally, we will simulate a stochastic stage-structured population projection model through 50 years, parameterized based on relevant literature or data collected in Minnesota, and use the 5th and 95th percentiles of λ to determine lower and upper bounds for population growth rates. We will use the bounds to evaluate biological believability of the index provided by the bowhunter observation surveys. For example, if the index were to suggest the population doubled in a single year, we would know this growth rate is unrealistic for a wild deer population based on white-tailed deer ecology.

The cost to conduct the mailed surveys was higher than online surveys because the mailed surveys required letters and postage. In addition, inputting the returned mailed surveys was labor intensive while the online surveys required minimal time and cost and did not require data entry. Because of the low cost and feasibility of the online survey, we have decided to only

conduct the online survey and incorporate several reminder emails and include an incentive for completing the survey.

ACKNOWLEDGMENTS


We thank all bowhunters that responded to the survey. We also thank the MNDNR Farmland Wildlife Populations and Research Group staff that provided feedback on earlier versions of the survey, and Katie Steffl and Tonya Klinkner for entering and proofing all data from the survey responses. Finally, we thank Veronique St-Louis for assistance with the sampling design and statistical analysis.

Table 1. Sampling statistics from the bowhunter observation survey in Minnesota, USA, 14 September – 8 November 2019.

Mode	Ecozone	Sampling Frame	Sample	Sampling Rate	Number of Responses	Sampling Frame Mean Age (SE)	Sample Mean Age (SE)	Responses Mean Age (SE)
Online	State	65628	16404	0.25	457	43 (0.07)	39 (0.13)	48 (0.31)
	Forest	17485	4298	0.25	124	46 (0.13)	41 (0.26)	51 (0.67)
	Transition	38424	9684	0.25	279	42 (0.08)	38 (0.16)	47 (0.37)
	Farmland	9719	2422	0.25	54	42 (0.17)	37 (0.33)	49 (0.90)
Mail	State	65628	17725	0.27	1723	43 (0.07)	44 (0.13)	54 (0.12)
	Forest	17485	5337	0.31	512	46 (0.13)	46 (0.24)	57 (0.22)
	Transition	38424	6344	0.17	638	42 (0.08)	42 (0.21)	52 (0.19)
	Farmland	9719	6044	0.62	573	42 (0.17)	43 (0.22)	53 (0.22)

Table 2. Statewide mean (\pm standard error) and 95% confidence intervals of online and mailed responses for hours hunted per hunting trip and observation rates per 1,000 hours from the bowhunter observation survey in Minnesota, USA, 14 September – 8 November 2019.

Parameter	Survey Mode			
	Online		Mail	
	Mean (SE)	95% CI	Mean (SE)	95% CI
Hours/Trip	3.22 (0.07)	3.07 - 3.36	3.23 (0.03)	3.16 - 3.29
Antlered Deer/1,000 Hours	193.07 (13.49)	166.63 - 219.51	193.78 (6.36)	181.32 - 206.24
Adult Female Deer/1,000 Hours	404.24 (22.77)	359.60 - 448.88	433.20 (13.04)	407.65 - 458.76
Fawn Deer/1,000 Hours	291.92 (21.11)	250.55 - 333.3	308.27 (11.59)	285.55 - 331
Unknown Deer/1,000 Hours	87.90 (8.81)	70.63 - 105.17	77.09 (4.14)	68.98 - 85.19
Total Deer/1,000 Hours	977.13 (52.59)	874.06 - 1080.21	1012.35 (29.63)	954.27 - 1070.43
Turkeys/1,000 Hours	315.75 (31.68)	253.65 - 377.84	305.13 (15.81)	274.15 - 336.12
Bears/1,000 Hours	2.98 (0.85)	1.32 - 4.64	2.50 (0.40)	1.71 - 3.28
Coyotes/1,000 Hours	15.96 (2.36)	11.34 - 20.58	15.53 (1.11)	13.35 - 17.72
Bobcats/1,000 Hours	0.12 (0.12)	0 - 0.35	0.86 (0.14)	0.58 - 1.13
Wolves/1,000 Hours	3.45 (1.50)	0.51 - 6.40	4.06 (0.69)	2.70 - 5.41
Fisher/1,000 Hours	1.07 (0.35)	0.38 - 1.77	1.54 (0.23)	1.08 - 2.00
Gray Foxes/1,000 Hours	2.50 (0.67)	1.18 - 3.82	2.75 (0.63)	1.51 - 3.99
Badgers/1,000 Hours	0.24 (0.17)	0 - 0.57	0.53 (0.14)	0.26 - 0.80



Political Boundaries Source: Minnesota DNR Quick Layers
 Prepared by: Minnesota DNR Farmland Wildlife Populations and Research Group



Figure 1. Deer management zones used to describe results of bowhunter observation surveys in Minnesota, USA during 2019. Red circles depict hunter locations ($n = 2,512$) during the early archery season (14 September – 8 November 2019). Generally, forested deer permit areas (DPAs) were composed of $\geq 60\%$ woody cover, transition DPAs were composed of 6%-50% woody cover, and farmland DPAs were composed of $\leq 5\%$ woody cover




Figure 2. Mean hunting observation trips per bowhunter by ecozone and survey type with 95% Confidence Intervals during the early archery season (14 September – 8 November 2019) in Minnesota, USA. Trips per hunter were different between mail and online survey respondents ($P < 0.001$).




Figure 3. Date of hunting observation trips for mail and online respondents during the early archery season (14 September – 8 November 2019) in Minnesota, USA.




Figure 4. Mean hours hunted per trip with 95% Confidence Intervals for mail and online respondents during the early archery season (14 September – 8 November 2019) in Minnesota, USA. Online and mail respondents did not differ ($P = 0.73$) in number of hours hunted per trip.




Figure 5. Mean deer observation rates per 1,000 hours with 95% Confidence Intervals in the forest ecozone during the early archery season (14 September – 8 November 2019) in Minnesota, USA.




Figure 6. Mean deer observation rates per 1,000 hours with 95% Confidence Intervals in the transition ecozone during the early archery season (14 September – 8 November 2019) in Minnesota, USA.




Figure 7. Mean deer observation rates per 1,000 hours with 95% Confidence Intervals in the farmland ecozone during the early archery season (14 September – 8 November 2019) in Minnesota, USA.




Figure 8. Mean observation rates of other species per 1,000 hours with 95% Confidence Intervals in the forest ecozone during the early archery season (14 September – 8 November 2019) in Minnesota, USA.




Figure 9. Mean observation rates of other species per 1,000 hours with 95% Confidence Interval in the transition ecozone during the early archery season (14 September – 8 November 2019) in Minnesota, USA.




Figure 10. Mean observation rates of other species per 1,000 hours with 95% Confidence Intervals in the farmland ecozone during the early archery season (14 September – 8 November 2019) in Minnesota, USA.

LITERATURE CITED

- Haroldson, B. S. 2014. 2014 white-tailed deer surveys. Pages 29-34 *in* M. H. Dexter, editor. Status of wildlife populations, fall 2014. Unpub. Rep., Division of Fish and Wildlife, Minn. Dept. Nat. Res., St. Paul, Minnesota. 328 pp.
- Lumley, T. 2004. Analysis of complex survey samples. *Journal of Statistical Software* 9:1–19.
- Norton, A. S., and W. R. Clark. 2016. 2016 Bow Hunter Observation Survey. Unpub. Rep, Iowa Dept. of Nat. Res., Boone, Iowa.
- Norton, A. S. and J. H. Giudice. 2017. Monitoring population trends of white-tailed deer in Minnesota – 2017. Unpub. Rep., Division of Fish and Wildlife, Minn. Dept. Nat. Res., St. Paul, Minnesota.
- Office of Legislative Auditor (OLA). 2016. Evaluation report, Department of Natural Resources: Deer population management. Program Evaluation Division, St. Paul, Minnesota.
- R Development Core Team. 2016. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/> (accessed 1 March 2018).
- Rolley, R. E., D. J. Storm, K. B. Wallenfang, and M. J. Tonkovich. 2016. Midwest deer metrics: What, how, and why we measure. Unpub. Rep, Wisc. Dept. of Nat. Res., Madison, Wisconsin 22 pp.

APPENDIX I. Mailed bowhunter observation survey for the early archery season (14 September – 8 November 2019) in Minnesota, USA.



August 6, 2019

2019 Bowhunter Observation Survey
HUNTER NAME
ADDRESS LINE 1
ADDRESS LINE 2

Dear Hunter,

You have been selected from a list of dedicated bowhunters to participate in the “2019 Bowhunter Observation Survey.” This survey is designed to enlist bowhunters to help monitor deer population trends. You were randomly selected from a list of people who purchased an archery license over the last 2 years. We chose dedicated bowhunters because of the amount of time you spend hunting deer. The valuable information you provide for this survey promotes better management of Minnesota’s deer herd, in addition to a better understanding of trends in other wildlife populations.

This survey is being conducted only during the early bow season, **Sept. 14 – Nov. 8, 2019**. Your help with this survey is very important, as it is new and we would like to conduct annually if we can collect good data.

All you have to do is record when and where you hunt, how many hours you hunt, and the number of animals you see while bowhunting. It is important to return your completed form in the postage paid envelope enclosed, and place it in the mail by November 15, 2019. If you finish all of your bowhunting prior to this date, please return the form earlier.

We have provided 4 columns for hunt locations. If you hunt more than 4 locations, pick your most frequent 4. If you hunted fewer than 4, just leave the other columns blank. For each column, please provide the following,

- **Deer Permit Area (DPA).** This is the 3-digit area you are hunting.
- **Nearest Town.** Please record the closest Incorporated town to your hunting location
- **Direction from Town.** Please use one of the 8 possible directions (N, NE, E, SE, S, SW, W, NW)
- **Distance from Town.** Please estimate the straight line distance between the town you listed and your hunting location

For the rest of the survey, the first row shows you an example of how we’d like you to complete the log. Some key points,

- **Area #.** The number that corresponds to the location information you listed above
- **Hours Hunting.** Please round to the nearest half hour. We ask that you use decimal points (ex – 2.5)
- **Deer Observed.** Please record what you see that day. Please record a ‘0’ if you didn’t see anything
- **Other Species.** Only write something if you see one of these animals. If left blank, we’ll assume it’s ‘0’

Please use 1 row for each day you hunted. In other words, if you hunted 6 days, you’d have 6 rows of data. This is the third year of the survey. If you participated in the survey in the past 2 years, we thank you. Results from the first 2 years of the survey can be found at <https://www.dnr.state.mn.us/mammals/deer/management/statistics.html> under the bowhunter survey heading. Thank you for your dedication to the sport of bowhunting, and we wish you a safe, enjoyable, and successful hunting season.

Sincerely,

A handwritten signature in cursive script that reads "Eric Michel".

Eric Michel, Ungulate Project Leader
eric.michel@state.mn.us; 507-578-8918

HUNTER NAME
 ADDRESS LINE 1
 ADDRESS LINE 2

BOWHUNTER OBSERVATION SURVEY 2019
 Minnesota Department of Natural Resources

MDNR Number:
 999999999

Thank you for participating in the 2019 Bowhunter Observation Survey. Please return this original form when you have finished bowhunting or by November 15, 2019, whichever comes first. When finished, place the form in the postage-paid, self-addressed return envelope. For questions, please call the Madelia wildlife research office at (507) 578-8912.

Hunt Location Information: Please record up to 4 locations where you will bowhunt. Please fill out the table below for those areas, along with the other information. When you record observations, you will use the location number (1, 2, 3, 4) to fill out the appropriate line of data. If you hunt more than 4 areas, please use your MOST FREQUENT 4. We realize some data may be lost if you hunt a lot of different areas.

1) DPA (3 digit number): _____ Nearest town: _____ Direction from town: _____ Distance from town: _____	2) DPA (3 digit number): _____ Nearest town: _____ Direction from town: _____ Distance from town: _____	3) DPA (3 digit number): _____ Nearest town: _____ Direction from town: _____ Distance from town: _____	4) DPA (3 digit number): _____ Nearest town: _____ Direction from town: _____ Distance from town: _____
---------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------

Month	Day	Area # (1,2,3,4)	# Hours Hunting (rounded to nearest 1/2 hour)	DEER OBSERVED (Record 0 if not seen)				OTHER SPECIES OBSERVED (Only write if you see one of these)							
				Antlered	Adult Female	Fawn	Not Sure	Wild Turkey	Coyote	Black Bear	Bobcat	Gray Wolf	Fisher	Gray Fox	Badger
Ex) 9	16	2	2.5	1	1	0	2				1		1		

APPENDIX II. Mean observation rates of other species per 1,000 hours and hours per trip with 95% Confidence Intervals by ecozone during the early archery season (14 September – 8 November 2019) in Minnesota, USA.

Parameter	Ecozone	Survey Mode	Mean	95% CI
Hours/Trip	Forest	Online	3.54 (SE = 0.17)	3.21 - 3.87
Antlered Deer/1,000 Hours	Forest	Online	113.59 (SE = 19.86)	74.65 - 152.52
Adult Female Deer/1,000 Hours	Forest	Online	261.08 (SE = 30.45)	201.41 - 320.75
Fawn Deer/1,000 Hours	Forest	Online	201.02 (SE = 25.62)	150.81 - 251.24
Not Sure Deer/1,000 Hours	Forest	Online	62.10 (SE = 9.58)	43.33 - 80.88
Total Deer/1,000 Hours	Forest	Online	637.79 (SE = 71.32)	498.01 - 777.57
Turkeys/1,000 Hours	Forest	Online	194.89 (SE = 35.71)	124.91 - 264.88
Bears/1,000 Hours	Forest	Online	8.17 (SE = 2.73)	2.81 - 13.53
Coyotes/1,000 Hours	Forest	Online	6.54 (SE = 2.15)	2.32 - 10.76
Bobcats/1,000 Hours	Forest	Online	0 (SE = 0)	0 - 0
Wolves/1,000 Hours	Forest	Online	11.85 (SE = 5.13)	1.80 - 21.90
Fisher/1,000 Hours	Forest	Online	0.82 (SE = 0.58)	0 - 1.95
Gray Foxes/1,000 Hours	Forest	Online	1.63 (SE = 1.01)	0 - 3.61
Badgers/1,000 Hours	Forest	Online	0 (SE = 0)	0 - 0
Hours/Trip	Forest	Mail	3.60 (SE = 0.08)	3.45 - 3.75
Antlered Deer/1,000 Hours	Forest	Mail	94.58 (SE = 7.36)	80.16 - 109.01
Adult Female Deer/1,000 Hours	Forest	Mail	260.66 (SE = 16.38)	228.55 - 292.77
Fawn Deer/1,000 Hours	Forest	Mail	159.21 (SE = 10.65)	138.34 - 180.08
Not Sure Deer/1,000 Hours	Forest	Mail	47.29 (SE = 5.25)	37.00 - 57.58
Total Deer/1,000 Hours	Forest	Mail	561.75 (SE = 31.44)	500.12 - 623.38
Turkeys/1,000 Hours	Forest	Mail	191.59 (SE = 19.8)	152.79 - 230.39
Bears/1,000 Hours	Forest	Mail	6.67 (SE = 1.22)	4.27 - 9.07
Coyotes/1,000 Hours	Forest	Mail	10.73 (SE = 1.37)	8.04 - 13.42
Bobcats/1,000 Hours	Forest	Mail	2.49 (SE = 0.43)	1.65 - 3.32
Wolves/1,000 Hours	Forest	Mail	13.21 (SE = 2.29)	8.73 - 17.70
Fisher/1,000 Hours	Forest	Mail	2.55 (SE = 0.58)	1.42 - 3.69
Gray Foxes/1,000 Hours	Forest	Mail	3.92 (SE = 1.20)	1.57 - 6.28
Badgers/1,000 Hours	Forest	Mail	0.20 (SE = 0.11)	0 - 0.42
Hours/Trip	Transition	Online	3.08 (SE = 0.08)	2.92 - 3.25
Antlered Deer/1,000 Hours	Transition	Online	211.86 (SE = 17.5)	177.56 - 246.16
Adult Female Deer/1,000 Hours	Transition	Online	445.92 (SE = 30.88)	385.38 - 506.45
Fawn Deer/1,000 Hours	Transition	Online	321.88 (SE = 30.47)	262.16 - 381.60
Not Sure Deer/1,000 Hours	Transition	Online	101.45 (SE = 12.93)	76.11 - 126.79
Total Deer/1,000 Hours	Transition	Online	1081.10 (SE = 72.04)	939.91 - 1222.30
Turkeys/1,000 Hours	Transition	Online	325.38 (SE = 35.10)	256.59 - 394.18
Bears/1,000 Hours	Transition	Online	0.97 (SE = 0.44)	0.11 - 1.83
Coyotes/1,000 Hours	Transition	Online	17.72 (SE = 3.33)	11.19 - 24.25
Bobcats/1,000 Hours	Transition	Online	0.19 (SE = 0.19)	0 - 0.58
Wolves/1,000 Hours	Transition	Online	0 (SE = 0)	0 - 0
Fisher/1,000 Hours	Transition	Online	1.17 (SE = 0.47)	0.25 - 2.09
Gray Foxes/1,000 Hours	Transition	Online	2.92 (SE = 0.91)	1.15 - 4.69
Badgers/1,000 Hours	Transition	Online	0.39 (SE = 0.27)	0 - 0.92

Appendix II. Page 2.

Parameter	Ecozone	Survey Mode	Mean	95% CI
Hours/Trip	Transition	Mail	3.15 (SE = 0.05)	3.06 - 3.24
Antlered Deer/1,000 Hours	Transition	Mail	226.96 (SE = 9.20)	208.92 - 244.99
Adult Female Deer/1,000 Hours	Transition	Mail	446.47 (SE = 19.59)	408.07 - 484.86
Fawn Deer/1,000 Hours	Transition	Mail	346.76 (SE = 19.43)	308.68 - 384.83
Not Sure Deer/1,000 Hours	Transition	Mail	86.06 (SE = 6.54)	73.23 - 98.88
Total Deer/1,000 Hours	Transition	Mail	1106.24 (SE = 45.14)	1017.77 - 1194.71
Turkeys/1,000 Hours	Transition	Mail	409.97 (SE = 29.88)	351.40 - 468.53
Bears/1,000 Hours	Transition	Mail	1.20 (SE = 0.38)	0.45 - 1.94
Coyotes/1,000 Hours	Transition	Mail	13.65 (SE = 1.51)	10.7 - 16.60
Bobcats/1,000 Hours	Transition	Mail	0.28 (SE = 0.13)	0.02 - 0.53
Wolves/1,000 Hours	Transition	Mail	0.14 (SE = 0.08)	0 - 0.29
Fisher/1,000 Hours	Transition	Mail	1.38 (SE = 0.31)	0.77 - 1.98
Gray Foxes/1,000 Hours	Transition	Mail	2.57 (SE = 0.81)	0.98 - 4.16
Badgers/1,000 Hours	Transition	Mail	0.28 (SE = 0.13)	0.02 - 0.53
Hours/Trip	Farmland	Online	3.20 (SE = 0.18)	2.86 - 3.54
Antlered Deer/1,000 Hours	Farmland	Online	313.65 (SE = 51.14)	213.43 - 413.88
Adult Female Deer/1,000 Hours	Farmland	Online	571.96 (SE = 78.01)	419.06 - 724.85
Fawn Deer/1,000 Hours	Farmland	Online	376.38 (SE = 59.88)	259.02 - 493.74
Not Sure Deer/1,000 Hours	Farmland	Online	79.95 (SE = 27.28)	26.49 - 133.41
Total Deer/1,000 Hours	Farmland	Online	1341.94 (SE = 164.38)	1019.77 - 1664.12
Turkeys/1,000 Hours	Farmland	Online	618.70 (SE = 202.68)	221.44 - 1015.95
Bears/1,000 Hours	Farmland	Online	0 (SE = 0)	0 - 0
Coyotes/1,000 Hours	Farmland	Online	33.21 (SE = 9.22)	15.15 - 51.27
Bobcats/1,000 Hours	Farmland	Online	0 (SE = 0)	0 - 0
Wolves/1,000 Hours	Farmland	Online	0 (SE = 0)	0 - 0
Fisher/1,000 Hours	Farmland	Online	1.23 (SE = 1.23)	0 - 3.64
Gray Foxes/1,000 Hours	Farmland	Online	2.46 (SE = 2.49)	0 - 7.34
Badgers/1,000 Hours	Farmland	Online	0 (SE = 0)	0 - 0
Hours/Trip	Farmland	Mail	3.00 (SE = 0.06)	2.89 - 3.11
Antlered Deer/1,000 Hours	Farmland	Mail	249.72 (SE = 15.13)	220.06 - 279.38
Adult Female Deer/1,000 Hours	Farmland	Mail	598.51 (SE = 29.57)	540.55 - 656.46
Fawn Deer/1,000 Hours	Farmland	Mail	409.72 (SE = 25.45)	359.84 - 459.61
Not Sure Deer/1,000 Hours	Farmland	Mail	95.41 (SE = 9.28)	77.23 - 113.60
Total Deer/1,000 Hours	Farmland	Mail	1353.36 (SE = 68.68)	1218.75 - 1487.97
Turkeys/1,000 Hours	Farmland	Mail	266.82 (SE = 25.02)	217.77 - 315.86
Bears/1,000 Hours	Farmland	Mail	0 (SE = 0)	0 - 0
Coyotes/1,000 Hours	Farmland	Mail	23.57 (SE = 2.91)	17.86 - 29.28
Bobcats/1,000 Hours	Farmland	Mail	0 (SE = 0)	0 - 0
Wolves/1,000 Hours	Farmland	Mail	0.21 (SE = 0.16)	0 - 0.52
Fisher/1,000 Hours	Farmland	Mail	0.70 (SE = 0.31)	0.09 - 1.32
Gray Foxes/1,000 Hours	Farmland	Mail	1.76 (SE = 1.42)	0 - 4.54
Badgers/1,000 Hours	Farmland	Mail	1.27 (SE = 0.44)	0.41 - 2.12